

Research Article

Cite this article: Wang N, Xu H, Xu G (2025). Seasonal responses of periphytic protozoan fauna to the antibiotic nitrofurazone at sensitive concentration in marine environments. *Journal of the Marine Biological Association of the United Kingdom* **105**, e29, 1–8. <https://doi.org/10.1017/S0025315425000074>

Received: 16 September 2023

Revised: 14 December 2024

Accepted: 20 December 2024

Keywords:

community structure; ecotoxicity; nitrofurazone; periphytic protozoa; seasonal variability

Corresponding author:

Henglong Xu;

Email: henglongxu@126.com

Seasonal responses of periphytic protozoan fauna to the antibiotic nitrofurazone at sensitive concentration in marine environments

Ning Wang¹, Henglong Xu¹ and Guangjian Xu²

¹Laboratory of Microbial Ecology, Ocean University of China, Qingdao 266003, China and ²College of Environment and Safety Engineering, Qingdao University of Science & Technology, Qingdao 266042, China

Abstract

In order to evaluate the seasonal responses of periphytic protozoan fauna to the antibiotic nitrofurazone at sensitive concentration, a 1-year baseline survey was carried out in Chinese coastal waters of the Yellow Sea. To assess the nitrofurazone (NFZ)-induced toxicokinetics in different season, the test protozoan samples were collected using microscope slides and exposed to the sensitive NFZ concentration of 8 mg ml⁻¹. Differences in species composition and typical species were observed in the test organism fauna in the control and treatment among four seasons. However, the community patterns were significantly shifted under the sensitive concentration, with a part of stressed test samples significantly departed from a respected taxonomic pattern. Therefore, it is suggested that periphytic protozoan fauna may be significantly changed at the same sensitive concentration in both the species composition and community pattern, although there were significant differences in tolerant species among four seasons in marine environments.

Introduction

As widely used pharmaceuticals, antibiotics can enter the environment through a variety of pathways, including hospital wastewater treatment plant, uncontrolled disposal of un-used drugs, runoff from agricultural fields and wastewater discharges from livestock facilities (Isidori *et al.*, 2005; Bhagat *et al.*, 2020; Anh *et al.*, 2021; Wang *et al.*, 2024). The extensive use of antibiotics has led to the detection of antibiotic residues in the marine environment worldwide, and thus causing serious damage to the meat-derived food, soil and water, the ecological environment, and public health (Puckowski *et al.*, 2016; Zhou *et al.*, 2021; Si *et al.*, 2022). Broad-spectrum veterinary antibiotics commonly used in aquaculture and animal husbandry are primarily used to treat protozoan and bacterial infections. Broad-spectrum veterinary antibiotics mainly include furazolidone (FZD), nitrofurazone (NFZ), nitrofurantoin (NFT) and furaltadone (FTD), all of which are nitrofurans (NFs). Due to their carcinogenic and mutagenic properties, these compounds are potentially hazardous to human health (Du *et al.*, 2014; Ghosh *et al.*, 2021; Kazmi *et al.*, 2022b). As a result, NF compounds have been classified as prohibited additives for food and animal production additives by the European Union (in 1995) and the USA (in 2002). As mariculture is flourishing globally, the use of antibiotics in the culturing process is lack of restrictions, and has caused serious ecological problems. Currently, research focusing on the presence of antibiotic residues in the mariculture environment is limited (Han *et al.*, 2020). NFZ is the most common and widely used one among these NF compounds (Chang *et al.*, 2016; Wang *et al.*, 2020). Hence, there is an increasing need to assess the ecotoxicological impact of antibiotics, especially NFZ, on environmental quality (Vutukuru *et al.*, 2007; Puckowski *et al.*, 2016).

Ecotoxicology is an integrated approach used to assess the toxic effects of toxicants and chemical pollutants on ecosystems and their inhabiting biota. Bioassays are considered to be the most reliable, feasible and cost-effective method of toxicity assessment in ecotoxicology. The most critical aspect of such toxicological study is the selection of suitable model organisms. The model organisms for bioassays should be abundant, ubiquitous, easy to manipulate and ecologically relevant (Dahms *et al.*, 2011). Periphytic protozoan communities generally meet these criteria and have therefore been chosen as model organisms in several ecotoxicological studies (Girling *et al.*, 2000; Niemeyer *et al.*, 2010; Kazmi *et al.*, 2022a).

Protozoans are the primary components of microbial fauna, and play an important role in driving the functional process of microbial food webs linking both planktonic and benthic ecosystems (Trielli *et al.*, 2007; Tan *et al.*, 2010; Xu *et al.*, 2014; Kazmi *et al.*, 2022b). In addition, they employ the indispensable contributor in maintaining/improving water quality of aquatic ecosystem by removing organic pollutants and various other water contaminants (Xu *et al.*, 2014; Kazmi *et al.*, 2020). Due to their simple life cycle, they are more sensitive to environmental changes than post-zoobenthos, so changes in their community pattern of protozoan fauna may significantly drive the functional process of marine ecosystems (Kathol *et al.*, 2009; Xu *et al.*, 2011a, 2011b; Xu *et al.*, 2014; Sikder *et al.*, 2020b).



It has been recognized that there is a significant seasonal variation in the community structure of periphytic protozoan fauna in marine ecosystems (Wey *et al.*, 2009; Jiang *et al.*, 2013; Guo *et al.*, 2020). Recent studies have demonstrated that the relative species number, taxonomic distinctness indices and body-size distinctness indices of periphytic protozoan fauna are sensitive to NFZ at the concentration of 8 mg ml^{-1} in autumn season (Kazmi *et al.*, 2022a, 2022b). However, with the seasonal responses of the periphytic protozoan fauna to the toxin at this concentration, little information was reported.

In this study, a 1-year baseline survey on seasonal responses of periphytic protozoan fauna to NFZ at the sensitive concentration was conducted. The objectives are (1) to reveal the variation in community pattern of periphytic protozoan fauna under the sensitive NFZ concentration; (2) to clarify whether there was seasonal variability in ecotoxicology of NFZ; and (3) to confirm the departure of the test protozoan communities from the expected community pattern in marine ecosystems.

Materials and methods

Sampling site and collection of test samples

Protozoan samples were collected from the coastal waters of the Yellow Sea near the mouth of Jiaozhou Bay, Qingdao, northern China in spring, summer, autumn and winter (Figure 1). The sampling site is a clean/slightly polluted area with an average water depth of $\sim 9 \text{ m}$, a tidal interval of 3 m and transparency of 2–3 m (Hassan Kazmi *et al.*, 2021).

The protozoan assemblages as test organisms were collected through glass slides measuring $2.5 \times 7.5 \text{ cm}$ according to the

method of Xu *et al.* (2011a, 2011b, 2012). Briefly, a (polyvinyl chloride) frame can hold 10 glass slides. Four frames were immersed at a depth of 2 m from the water surface and were left for 14 days to allow the protozoans (mainly ciliates) to colonize the slides. The collected samples were then transported to the laboratory via *in situ* water and stored in a cooler (Xu *et al.*, 2012). After the samples were domesticated for 3 days by setting the laboratory conditions in an illumination cabinet (temperature 25°C , illumination 3960 lx), 30 slides with protozoan colony colonization were randomly selected for the next experiment.

Experimental design

Nitrofurazone (5-nitro-2-furfural semicarbazone) in the form of yellow crystalline powder from Sigma-Aldrich Co., Ltd. (Shanghai, China, CAS No. 59870) was model antibiotic. A stock solution of 300 mg l^{-1} nitrofurazone was prepared according to Hong *et al.* (2015). Briefly, 300 mg of nitrofurazone powder was dissolved in artificial seawater (AMW; in 1000 ml distilled water, pH 8.2, salinity 28‰, 28 g of NaCl, 0.8 g of KCl, 5 g of $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ and 1.2 g of CaCl_2) and then further diluted in artificial seawater to prepare experimental concentrations (Kazmi *et al.*, 2022a).

All bioassay experiments were carried out in Petri dishes for 10 days according to the method of Li *et al.* (2014). For each season prepare a control group (C): 0 and a treatment group (T): 8 mg ml^{-1} , respectively. Each glass slide with protozoan communities was placed in a separate Petri dish. The Petri dishes contained 1 vs 1 solution of habitat water and NFZ in a final volume of 20 ml. Three independent replicates of each treatment were used as parallel tests. The species composition and individual abundances of the protozoans were observed throughout the experiment.

Identification and enumeration

The test protozoan communities were observed through $10\text{--}400\times$ magnification with a bright-field microscope. The enumeration and identification of protozoa were based on Xu *et al.* (2014) and Song *et al.* (2009), respectively.

Data analysis

The taxonomic breadth was derived from the average taxonomic distinctness (Δ^+) and variation in taxonomic distinctness (Λ^+), calculated as follows (Kazmi *et al.*, 2022a):

$$\Delta^+ = [\sum \sum_{i < j} \omega_{ij}^2 x_i x_j] / [S(S-1)/2]$$

$$\Lambda^+ = [\sum \sum_{i < j} (\omega_{ij} - \Delta^+)] / [S(S-1)/2]$$

where, ω_{ij} = distinctness weight given to the path length linking species (i and j); x_i ($i = 1, 2, \dots, S$) = abundance of the i^{th} species; N = total number of individuals in the sample and S is the number of species (Warwick and Clarke, 1995).

PRIMER v7 with PERMANOVA+ calculated the toxicodynamics in protozoan communities (Clarke and Gorley, 2015). The variations in species composition and toxic dynamics of periphytic protozoa in the control and treatment groups were presented by shade plotting with cluster analysis (Anderson *et al.*, 2008). In addition, distance-based redundancy analyses (dbRDA) revealed the community patterns of periphytic protozoa across seasons for treatment and control groups. Moreover, TAXTDTEST ellipse plotting was used to present the significance of deviation from an expectation at different groups (Clarke and Gorley, 2015).




Figure 1. Sampling station, for the collection of test protozoan communities, located in the coastal waters of the Yellow Sea, northern China.




Figure 2. Shade plotting with clustering analysis on the index of association showed seasonal variability in species distribution of periphytic protozoa and relative abundance in the controls (C) and treatments (T, NFZ concentration of 8 mg ml⁻¹) (1, spring; 2, summer; 3, autumn; 4, winter; I-IV, Groups I-IV).

Table 1. Typical species to the test organism communities in control and treatment during four seasons

Species	Spring			Summer			Autumn			Winter		
	C	T	Ctb%	C	T	Ctb%	C	T	Ctb%	C	T	Ctb%
<i>Litonotus paracygnus</i>	++++	+	20.8	+	–	3.5	–	–	–	+	+	1.9
<i>Acineta foetida</i>	+++	–	18.2	–	–	–	–	–	–	–	–	–
<i>Euplotes raikovi</i>	+++	+	14.4	–	–	–	++++	+	13.8	–	–	–
<i>Diophrys appendiculata</i>	++	–	13.1	++	–	9.7	–	–	–	+	+	1.3
<i>Uronychia transfuga</i>	+	–	4.7	–	–	–	–	–	–	–	–	–
<i>Diophrys hystrix</i>	+	+	4.5	–	–	–	–	–	–	–	–	–
<i>Pseudovorticella anomala</i>	+	–	3.8	–	–	–	–	–	–	–	–	–
<i>Pseudovorticella parakenti</i>	+	–	3.3	–	–	–	–	–	–	–	–	–
<i>Aspidisca aculeata</i>	+	+	3.0	++++	+	14.7	++	–	5.3	+	+	2.6
<i>Vorticella utriculus</i>	+	+	2.7	–	–	–	–	–	–	–	–	–
<i>Zoothamnium vermicola</i>	+	+	2.4	–	–	–	–	–	–	–	–	–
<i>Thuricola parafolliculata</i>	+	–	1.8	–	–	–	–	–	–	–	–	–
<i>Loxophyllum simplex</i>	+	–	1.8	–	–	–	–	+	2.9	–	–	–
<i>Holosticha bradburyae</i>	+	–	1.3	–	–	–	–	–	–	–	–	–
<i>Chlamydonella derouxi</i>	–	–	–	++++	–	16.0	–	–	–	–	–	–
<i>Tachysoma dragescoi</i>	–	–	–	+++	+	13.2	–	–	–	–	–	–
<i>Acineta tuberosa</i>	–	–	–	+++	+	10.2	+++	–	12.1	–	–	–
<i>Acineta compressa</i>	–	–	–	+	–	4.7	–	–	–	–	+	1.2
<i>Coeloperix sleighi</i>	–	–	–	+	–	3.8	–	–	–	+	–	1.9
<i>Holosticha heterofoissneri</i>	–	–	–	+	–	3.4	–	–	–	–	–	–
<i>Pseudovorticella paraformicata</i>	–	–	–	–	+	3.0	–	–	–	–	–	–
<i>Stephanopogon paramesnili</i>	–	–	–	–	+	2.8	–	–	–	–	–	–
<i>Dysteria pusilla</i>	–	–	–	+	–	2.7	–	–	–	–	–	–
<i>Aspidisca steini</i>	–	–	–	+	+	1.9	+	–	4.4	++++	+	23.0
<i>Orthodonella apohamatus</i>	–	–	–	+	–	1.7	–	–	–	–	–	–
<i>Stephanopogon minuta</i>	–	–	–	+	–	1.6	–	–	–	–	–	–
<i>Tachysoma ovata</i>	–	–	–	+	+	1.6	+	+	2.1	–	–	–
<i>Protogastrostyla pulchra</i>	–	–	–	+	+	1.5	+++	+	10.7	–	–	–
<i>Euplotes minuta</i>	–	–	–	–	–	–	+++	+	10.9	–	–	–
<i>Ephelota crustaceorum</i>	–	–	–	–	–	–	++	–	7.4	–	–	–
<i>Aspidisca magna</i>	–	–	–	–	–	–	+	–	4.5	+	–	1.6
<i>Discocephalus ehrenbergi</i>	–	–	–	–	–	–	+	–	4.2	–	–	–
<i>Metaurostyloopsis salina</i>	–	–	–	–	–	–	+	–	3.5	–	–	–
<i>Pseudovorticella marina</i>	–	–	–	–	–	–	+	–	3.5	–	–	–
<i>Zoothamnium plumula</i>	–	–	–	–	–	–	–	+	3.1	–	–	–
<i>Euplotes vannus</i>	–	–	–	–	–	–	+	+	1.9	+	+	5.2
<i>Strombidium paracalkinsi</i>	–	–	–	–	–	–	+	–	1.8	–	–	–
<i>Philasterides armatalis</i>	–	–	–	–	–	–	+	–	1.7	–	–	–
<i>Thigmokeronopsis rubra</i>	–	–	–	–	–	–	+	+	1.6	–	–	–
<i>Litonotus blattereri</i>	–	–	–	–	–	–	–	–	–	+++++	+	44.0
<i>Hartmannula derouxi</i>	–	–	–	–	–	–	–	–	–	+	–	5.5
<i>Litonotus yinae</i>	–	–	–	–	–	–	–	–	–	+	+	3.2
<i>Pseudokeronopsis rubra</i>	–	–	–	–	–	–	–	–	–	+	+	1.5
<i>Uronychia binucleata</i>	–	–	–	–	–	–	–	–	–	+	+	1.3
<i>Hartmannula angustipilosa</i>	–	–	–	–	–	–	–	–	–	+	+	1.1

Average abundance: '–' = 0; '+' = 0–1, '++' = 1–5, '+++ = 5–10, '++++' = 15–20, '+++++' > 20; C, control; T, treatment; Ctb, Contribution.

The *t*-test was used to signify the differences in abundance between the treatment and the control using the program SPSS (v22) (Xu *et al.*, 2014).

Result

Species composition and changes

Figure 2 shows the species composition and changes in terms of average abundances, and ecological types of the test periphytic protozoan communities. A total of 60 protozoan species were identified. A total of 14, 15, 16 and 13 species were identified in the controls (C), while 6, 8, 8 and 11 species were observed in the treatments (T) from spring to winter, respectively (Figure 2 and Table 1).

These species were roughly divided into four groups using clustering analysis with the SIMPROF test (Figure 2). The shade plotting with clustering analysis showed a clear seasonal variation in species distribution, and four groups (I–IV) dominated spring, summer, autumn and winter, respectively (Figure 2a, b). From spring to winter, the relative abundance and relative number of species of dominant contributors changed in the following order: Group I→II→III→IV, and sharply dropped from the controls and treatments (Figures 2 and 3). Species number and individual abundance followed the same variation (Figure 4). For example, Group I, *Acineta foetida*, was very sensitive, which was mainly present in the control group. *Diophrys hystrix*, which was ubiquitous in the treatment group, was tolerant to 8 mg ml⁻¹ nitrofurazone (Figure 2 and Table 1).

It should be noted that there were significant differences in both species number and individual abundance between the treatment and the control ($P < 0.05$).

Table 1 summarizes the abundance, frequency of occurrence and contribution of typical protozoan species of the test organisms in different seasons. The dominant contributors in each season showed different contribution rates in the treatment and control groups. For example, in spring, *Litonotus paracygnus* was the largest contributor with a contribution of 20.8%. *Chlamydonella derouxi* dominated the summer with its contribution of 16%. In addition, *Euplotes raikovi* with 13.8% and *Litonotus blattereri* with 44% contributed in autumn and winter, respectively.

Variation in protozoan community pattern

Distance-based redundancy analysis (dbRDA) ordinations showed that there were different colonization patterns of the protozoan communities among the four seasons (Figure 5). Taxonomic patterns at Group C were separated from Group T by dbRDA1.

It can be clearly observed that the vectors of six species are pointed toward the data points of Group C, whereas only *D. hystrix* and *Aspidisca aculeata* pointed toward the data clouds of Group T (Figure 5a); the vectors of nine species point to the data of Group C and only three species (*Stephanopogon paramesnilii*, *Aspidisca steini*, *Pseudovorticella paraformicata*) point to the Group T data (Figure 5b). In Figure 5c, there are eight and three species vectors in the Groups T and C, respectively; in Figure 5d, there are seven vectors of species pointing to the Group C data and four species pointing to the Group T.

Variations in taxonomic distinctness

Variations in taxonomic distinctness and average taxonomic distinctness (Λ^+ and Δ^+) are summarized in Figure 6. Ellipse tests on the 95% probability regions have a range with sublist sizes (10, 20




Figure 3. Seasonal variability in relative species numbers (a) and relative abundance (b) of Yellow Sea coastal periphytic protozoa in the controls and treatments (I–IV, Groups I–IV).




Figure 4. Seasonal variability in species numbers (a) and individual abundance (b) of Yellow Sea coastal periphytic protozoa in Groups C and T (I–IV, Groups I–IV).




Figure 5. Distance-based redundancy analysis (dbRDA), showing the seasonal variation of protozoan community patterns in Groups C and T in the coastal waters of the Yellow Sea (a, spring; b, summer; c, autumn; d, winter).

and 30 species) of the protozoan samples for all seasonal controls and treatments (Figure 6). It was clear that there were differences in taxonomic pattern of the protozoan communities between the controls and treatments, for example, all samples were fallen in 10, 20 and 30 species contour in the controls (Figure 6a), whereas a part of these showed a significant departure from the expected community pattern (Figure 6b).

Discussion

Antibiotics are deposited in surface water through multiple sources, thus posing a serious ecological threat (Puckowski *et al.*, 2016; Bawa-Allah and Ehimiyen, 2022). Aquatic ecosystems primarily serve as the main repository for various kinds of antibiotics, posing ecological risks to aquatic organisms (freshwater algae, microphytes, macrophytes, zooplankton and fishes) (Kovalakova *et al.*, 2020; Anh *et al.*, 2021; Zhou *et al.*, 2024). Protozoa, as important hubs in marine ecosystems, necessitate the study of their response to antibiotic toxicity. Recent investigations have demonstrated that the periphytic protozoan

communities are sensitive particularly to NFZ at 8 mg ml^{-1} in the concentration (Kazmi *et al.*, 2022a, 2022b).

Due to environmental heterogeneity, differences in food availability between seasons significantly influenced the protozoan colonization dynamics, with significant seasonal changes in community structure and functioning (Sikder *et al.*, 2020a, 2020b). In our study, it was found that 8 mg ml^{-1} caused a decrease in the relative species number and relative abundance in each season and therefore NFZ was toxic in each season. However, each season has different dominant species, and SIMPROF analysis allowed to divide the 60 protozoa observed into four groups. The contribution of these four groups was different in each season, Group I, II, III and IV were occupying the four seasons of spring, summer, autumn and winter, respectively, which indicated that there were seasonal differences in the toxic effects of NFZ. It is probably because NFZ affects the food supply of ciliates, and the difference in food supply can have a significant effect on ciliate colonization dynamics, which needs further confirmation.

The dbRDA analysis showed that nitrofurazone toxicity led to changes in the community structure of periphytic protozoa in




Figure 6. Ellipse plots of 95% probability regions with a range of three sublist sizes (10, 20 and 30 species) for the taxonomic distinctness indices, i.e. average taxonomic distinctness (Δ^+) and variation taxonomic distinctness (Λ^+), of the protozoan communities in the treatment and control group, showing the deviation of the protozoan communities from an expected range of 10, 20 and 30 species contours (a, 0; b, 8 mg ml⁻¹).

different seasons, which may be due to differences in tolerance between species as well as seasonal variation. This reflects that 8 mg ml⁻¹ concentration of NFZ respond to the protozoa tested with some variation depending on the season.

Taxonomic distinctness indices for analysing variability in taxonomic breadth of a community have the advantage of being low sensitivity to sampling effort and sample size, and being able to test the significance of departure from expectation within a statistical framework (Clarke and Warwick, 1998; Leonard *et al.*, 2006; Somerfield *et al.*, 2008; Prato *et al.*, 2009; Sikder *et al.*, 2020a). Thus, in the present study these taxonomic indices (Δ^+ and Λ^+) may have played an auxiliary role exploring the adaptation of protozoa to the same concentration of NFZ across seasons. Ellipse plots of these metrics indicate that protozoan communities deviated significantly from the expected taxonomic width when NFZ concentrations were 8 mg ml⁻¹, whereas in controls no samples deviated from the expected taxonomic pattern.

Thus, our study confirms that periphytic protozoa can be used as biomarkers for evaluating NFZ ecotoxicity, which is consistent with previous reports. In total, 8 mg ml⁻¹ of NFZ does not lose its effect on protozoa toxicity due to seasonal variability. Because the major contributors differed in each season, when the season changed, new populations were substituted to cope with the NFZ toxicity effects. This finding could explain the ability of protozoa to transport material and energy from plankton to benthos when exposed to environmental pollutants, playing an important role in the functioning of microbial food webs and maintaining ecosystem stability.

In summary, differences in species composition and typical species were observed in the test organism fauna in the control and treatment among four seasons. However, the community patterns were significantly shifted under the sensitive concentration, with a part of test samples showed a significant departure from the respected taxonomic pattern. Therefore, it is suggested that periphytic protozoan fauna may be significantly changed at the sensitive concentration of 8 mg ml⁻¹ in four seasons, although there were significant differences in both species composition and community pattern in marine environments.

Author contributions. All authors wrote the paper and approved the final manuscript.

Financial support. This work was supported by 'The Natural Science Foundation of China' (project numbers: 31672308 and 41076089).

Competing interest. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethical standards. This work meets ethical standards.

Data availability. The data that support the findings of the study are available from the corresponding author upon reasonable request.

References

- Anderson MJ, Gorley RN, Clarke KS, Anderson MS, Gorley RN, Clarke KR and Anderson M (2008) *PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods*. Plymouth: PRIMER-E Ltd.
- Anh HQ, Le TPQ, Da Le N, Lu XX, Duong TT, Garnier J, Rochelle-Newall E, Zhang S, Oh N, Oeurng C, Ekkawatpanit C, Nguyen TD, Nguyen QT, Nguyen TD, Nguyen T, Tran TL, Kunisue T, Tanoue R, Takahashi S, Minh TB, Le HT, Pham TMH and Nguyen TAH (2021) Antibiotics in surface water of east and southeast Asian countries: a focused review on contamination status, pollution sources, potential risks, and future perspectives. *Science of the Total Environment* **764**, 142865.
- Bawa-Allah KA and Ehimiyein AO (2022) Ecotoxicological effects of human and veterinary antibiotics on water flea (*daphnia magna*). *Environmental Toxicology and Pharmacology* **94**, 103932.
- Bhagat C, Kumar M, Tyagi VK and Mohapatra PK (2020) Proclivities for prevalence and treatment of antibiotics in the ambient water: a review. *NPI Clean Water* **3**, 42.
- Chang G, Chen H and Lin F (2016) Analysis of banned veterinary drugs and herbicide residues in shellfish by liquid chromatography-tandem mass spectrometry (lc/ms/ms) and gas chromatography-tandem mass spectrometry (gc/ms/ms). *Marine Pollution Bulletin* **113**, 579–584.
- Clarke K and Gorley R (2015) *Primer version 7: user manual/tutorial*. PRIMER-E 192.
- Clarke KR and Warwick RM (1998) A taxonomic distinctness index and its statistical properties. *Journal of Applied Ecology* **35**, 523–531.

- Dahms HU, Hagiwara A and Lee JS (2011) Ecotoxicology, ecophysiology, and mechanistic studies with rotifers. *Aquatic Toxicology* **101**, 1–12.
- Du N, Chen M, Sheng L, Chen S, Xu H, Liu Z, Song C and Qiao R (2014) Determination of nitrofurans metabolites in shrimp by high performance liquid chromatography with fluorescence detection and liquid chromatography–tandem mass spectrometry using a new derivatization reagent. *Journal of Chromatography A* **1327**, 90–96.
- Ghosh S, Majumder S and Roychowdhury T (2021) Impact of microbial multi-metal and broad spectrum antibiotic tolerance in urban sw (adi ganga, kolkata) on adjacent groundwater: a future threat. *Groundwater for Sustainable Development* **14**, 100608.
- Girling AE, Pascoe D, Janssen CR, Peither A, Wenzel A, Schäfer H, Neumeier B, Mitchell GC, Taylor EJ, Maund SJ, Lay JP, Jüttner I, Crossland NO, Stephenson RR and Persoone G (2000) Development of methods for evaluating toxicity to freshwater ecosystems. *Ecotoxicology and Environmental Safety* **45**, 148–176.
- Guo C, Gui Y, Bai X, Sikder MNA and Xu H (2020) Seasonal variation in biological trait distribution of periphytic protozoa in coastal ecosystem: a baseline study for marine bioassessment. *Marine Pollution Bulletin* **160**, 111593.
- Han QF, Zhao S, Zhang XR, Wang XL, Song C and Wang SG (2020) Distribution, combined pollution and risk assessment of antibiotics in typical marine aquaculture farms surrounding the yellow sea, north China. *Environment International* **138**, 105551.
- Hassan Kazmi SSU, Xuexi T, Xu G and Xu H (2021) An approach to optimizing sampling effort for bioassessment surveys based on periphytic ciliates according to water depths in marine ecosystems. *Ecological Indicators* **122**, 107222.
- Hong Y, Lin X, Cui X, Zhou L, Al-Rasheid KAS and Li J (2015) Comparative evaluation of genotoxicity induced by nitrofurazone in two ciliated protozoa by detecting DNA strand breaks and DNA–protein crosslinks. *Ecological Indicators* **54**, 153–160.
- Isidori M, Lavorgna M, Nardelli A, Pascarella L and Parrella A (2005) Toxic and genotoxic evaluation of six antibiotics on non-target organisms. *Science of the Total Environment* **346**, 87–98.
- Jiang Y, Xu H, Zhu M and Al-Rasheid KAS (2013) Temporal distributions of microplankton populations and relationships to environmental conditions in Jiaozhou bay, northern China. *Journal of the Marine Biological Association of the United Kingdom* **93**, 13–26.
- Kathol M, Norf H, Arndt H and Weitere M (2009) Effects of temperature increase on the grazing of planktonic bacteria by biofilm-dwelling consumers. *Aquatic Microbial Ecology* **55**, 65–79.
- Kazmi SSUH, Uroosa, Warren A, Zhong X and Xu H (2022a) Insights into the ecotoxicity of nitrofurazone in marine ecosystems based on body-size spectra of periphytic ciliates. *Marine Pollution Bulletin* **174**, 113217.
- Kazmi SSUH, Uroosa, Xu H and Xuexi T (2022b) An approach to determining the nitrofurazone-induced toxic dynamics for ecotoxicity assessment using protozoan periphytons in marine ecosystems. *Marine Pollution Bulletin* **175**, 113329.
- Kazmi SSUH, Xuexi T, Xu G, Sikder MNA and Xu H (2020) Vertical variability in taxonomic breadth of biofilm-dwelling ciliates in marine bioassessment surveys. *Regional Studies in Marine Science* **38**, 101366.
- Kovalakova P, Cizmas L, McDonald TJ, Marsalek B, Feng M and Sharma VK (2020) Occurrence and toxicity of antibiotics in the aquatic environment: a review. *Chemosphere* **251**, 126351.
- Leonard DRP, Robert Clarke K, Somerfield PJ and Warwick RM (2006) The application of an indicator based on taxonomic distinctness for UK marine biodiversity assessments. *Journal of Environmental Management* **78**, 52–62.
- Li J, Zhou L, Lin X, Yi Z and Al-Rasheid KAS (2014) Characterizing dose–responses of catalase to nitrofurazone exposure in model ciliated protozoan *Euplotes vannus* for ecotoxicity assessment: enzyme activity and mRNA expression. *Ecotoxicology and Environmental Safety* **100**, 294–302.
- Niemeyer JC, Moreira-Santos M, Nogueira MA, Carvalho GM, Ribeiro R, Da Silva EM and Sousa JP (2010) Environmental risk assessment of a metal-contaminated area in the tropics. Tier i: screening phase. *Journal of Soils and Sediments* **10**, 1557–1571.
- Prato S, Morgana JG, La Valle P, Fioia MG, Lattanzi L, Nicoletti L, Ardizzone GD and Izzo G (2009) Application of biotic and taxonomic distinctness indices in assessing the ecological quality status of two coastal lakes: Caprolace and Fogliano lakes (central Italy). *Ecological Indicators* **9**, 568–583.
- Puckowski A, Mioduszevska K, Bukaszewicz P, Borecka M, Caban M, Maszkowska J and Stepnowski P (2016) Bioaccumulation and analytics of pharmaceutical residues in the environment: a review. *Journal of Pharmaceutical and Biomedical Analysis* **127**, 232–255.
- Si R, Yao Y, Liu X, Lu Q and Liu M (2022) Role of risk perception and government regulation in reducing over-utilization of veterinary antibiotics: evidence from hog farmers of China. *One Health* **15**, 100448.
- Sikder MNA, Xu G and Xu H (2020a) Seasonal variability in taxonomic breadth of biofilm-dwelling ciliates in colonization surveys for marine bioassessment. *Marine Pollution Bulletin* **151**, 110828.
- Sikder MNA, Xu H, Xu G and Warren A (2020b) Seasonal variability in trophic-functional patterns of marine biofilm-dwelling ciliates during the process of colonization. *Regional Studies in Marine Science* **35**, 101236.
- Somerfield PJ, Clarke KR, Warwick RM and Dulvy NK (2008) Average functional distinctness as a measure of the composition of assemblages. *ICES Journal of Marine Science* **65**, 1462–1468.
- Song W, Warren A and Hu X (2009) *Free-living Ciliates in the Bohai and Yellow Seas*. Beijing: Science Press. In both Chinese and English.
- Tan X, Shi X, Liu G, Xu H and Nie P (2010) An approach to analyzing taxonomic patterns of protozoan communities for monitoring water quality in Songhua river, northeast China. *Hydrobiologia* **638**, 193–201.
- Trielli F, Amaroli A, Sifredi F, Marchi B, Falugi C and Corrado MUD (2007) Effects of xenobiotic compounds on the cell activities of *Euplotes crassus*, a single-cell eukaryotic test organism for the study of the pollution of marine sediments. *Aquatic Toxicology* **83**, 272–283.
- Vutukuru SS, Prabhath NA, Raghavender M and Yerramilli A (2007) Effect of arsenic and chromium on the serum amino-transferases activity in Indian major carp, *Labeo rohita*. *International Journal of Environmental Research and Public Health* **4**, 224–227.
- Wang K, Guo C, Li J, Wang K, Liang S, Wang W and Wang J (2024) A critical review of the adsorption-desorption characteristics of antibiotics on microplastics and their combined toxic effects. *Environmental Technology & Innovation* **35**, 103729.
- Wang Y, Guo Y, Pan K, Lin X and Ni Y (2020) Electrochemical reaction mechanism of nitrofurazone at poly-ACBK/GCE and its analytic application. *Chemistry Africa* **3**, 727–734.
- Warwick R and Clarke K (1995) New 'biodiversity' measures reveal a decrease in taxonomic distinctness with increasing stress. *Marine Ecology Progress Series* **129**, 301–305.
- Wey JK, Norf H, Arndt H and Weitere M (2009) Role of dispersal in shaping communities of ciliates and heterotrophic flagellates within riverine biofilms. *Limnology and Oceanography* **54**, 1615–1626.
- Xu H, Jiang Y, Al-Rasheid KAS, Al-Farraj SA and Song W (2011a) Application of an indicator based on taxonomic relatedness of ciliated protozoan assemblages for marine environmental assessment. *Environmental Science and Pollution Research* **18**, 1213–1221.
- Xu H, Zhang W, Jiang Y and Yang EJ (2014) Use of biofilm-dwelling ciliate communities to determine environmental quality status of coastal waters. *Science of the Total Environment* **470–471**, 511–518.
- Xu H, Zhang W, Jiang Y, Zhu M and Al-Rasheid KAS (2012) Sampling sufficiency for analyzing taxonomic relatedness of periphytic ciliate communities using an artificial substratum in coastal waters. *Journal of Sea Research* **72**, 22–27.
- Xu H, Zhang W, Jiang Y, Zhu M, Al-Rasheid KA, Warren A and Song W (2011b) An approach to determining the sampling effort for analyzing biofilm-dwelling ciliate colonization using an artificial substratum in coastal waters. *Biofouling* **27**, 357–366.
- Zhou X, Shi Y, Lu Y, Song S, Wang C, Wu Y, Liang R, Qian L, Xu Q, Shao X and Li X (2024) Ecological risk assessment of commonly used antibiotics in aquatic ecosystems along the coast of China. *Science of the Total Environment* **935**, 173263.
- Zhou W, Tang Y, Du X, Han Y, Shi W, Sun S, Zhang W, Zheng H and Liu G (2021) Fine polystyrene microplastics render immune responses more vulnerable to two veterinary antibiotics in a bivalve species. *Marine Pollution Bulletin* **164**, 111995.