

# Application of network analysis to identify interactive systems of eating disorder psychopathology

K. T. Forbush\*, C. S. Q. Siew and M. S. Vitevitch

Department of Psychology, University of Kansas, Lawrence, KS 66045, USA

**Background.** Traditional approaches for the classification of eating disorders (EDs) attribute symptoms to an underlying, latent disease entity. The network approach is an alternative model in which mental disorders are represented as networks of interacting, self-reinforcing symptoms. This project was the first to use network analysis to identify interconnected systems of ED symptoms.

**Method.** Adult participants ( $n = 143$ ; 77.6% women) with a Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DSM-5) ED were recruited from the community to take part in a larger ongoing longitudinal study. The Structured Clinical Interview for DSM Disorders (SCID-I) was used to establish diagnoses. An undirected network of ED symptoms was created using items from the Eating Pathology Symptoms Inventory (EPSI) and the R package qgraph.

**Results.** Body checking emerged as the strongest and most important single symptom in the entire network by having the shortest average distance to other symptoms in the network, and by being the most frequent symptom on the path between any two other symptoms. Feeling the need to exercise every day and two symptoms assessing dietary restraint/restricting emerged as ‘key players’, such that their removal from the network resulted in maximal fracturing of the network into smaller components.

**Conclusions.** Although cognitive-behavioral therapy for EDs focuses on reducing body checking to promote recovery, our data indicate that amplified efforts to address body checking may produce stronger (and more enduring) effects. Finally, results of the ‘key players analysis’ suggested that targeting interventions at these key nodes might prevent or slow the cascade of symptoms through the ‘network’ of ED psychopathology.

Received 2 August 2015; Revised 6 January 2016; Accepted 7 January 2016; First published online 8 July 2016

**Key words:** Body checking, disordered eating symptoms, DSM, eating disorders, network analysis.

## Introduction

Issues related to the nosology of mental disorders remains at the forefront of clinical science and practice (Regier *et al.* 2013; Cuthbert, 2014; Insel, 2014). Although the identification of new ways of conceptualizing mental disorders is a topic that cuts across diagnostic categories, there has been particular interest in developing innovative classification schemes for eating disorders (EDs) (for reviews, see Keel *et al.* 2012; Wildes & Marcus, 2013). Indeed, researchers have contended that there are ‘few mental disorders for which novel conceptualizations are needed more’ than for EDs (Wildes *et al.* 2013, p. 1031). Here we present the first application of network analysis to identify ‘causal’

systems<sup>†</sup> of ED symptoms in a community-recruited sample of adults with EDs; we conclude by discussing diagnostic and treatment implications.

## Limitations with categorical conceptualizations of EDs

The publication of the fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5; American Psychiatric Association, 2013) led to several improvements in ED diagnoses compared with previous editions. Specifically, binge eating disorder (BED) was recognized as an official diagnostic category and the frequency criteria for bulimia nervosa (BN) and BED were lowered, which reduced the prevalence of subthreshold diagnoses (and provided full-threshold diagnoses to more individuals who were suffering from serious, clinically significant EDs). However, as

\* Address for correspondence: K. T. Forbush, Department of Psychology, University of Kansas, Fraser Hall, 1400 Jayhawk Boulevard, Lawrence, KS 66045, USA.  
(Email: kforbush@ku.edu)

† The notes appear after the main text.

we describe below, remaining issues with current ED categorizations pose certain limitations to their validity and clinical utility.

First, one of the most common ED diagnoses is ‘other specified feeding or eating disorder’ (OSFED), which is comprised of various clinically significant ED symptom constellations. Recent studies indicate that approximately 30–60% of persons with DSM-5 EDs are diagnosed with an OSFED (Allen *et al.* 2013; Stice *et al.* 2013); other research indicates that it is not possible to distinguish full-threshold EDs from OSFEDs on the basis of psychosocial impairment or genetic risk factors (Fairweather-Schmidt & Wade, 2014). As a result of relegating OSFEDs to a ‘residual’ diagnostic category, there are few studies describing its long-term course and outcome, and even fewer treatment studies (for a review of this issue, see Fairburn & Bohn, 2005).

Second, there is substantial symptom overlap across ED diagnoses. Few symptoms distinguish among ED diagnoses; thus, a change in a single symptom (e.g. weight gain or loss) can lead to a change in diagnosis [e.g. anorexia nervosa (AN) to BN or vice versa]. Finally – due to symptom overlap among EDs – diagnostic ‘migration’ is common with approximately 30–50% of individuals with an ED being diagnosed with a different ED at follow-up (Eddy *et al.* 2008; Castellini *et al.* 2011; Allen *et al.* 2013). Taken together, these issues indicate that it may be more clinically and scientifically informative to elucidate EDs at the symptom-level (rather than at the level of predefined diagnostic groupings of disorders).

### *Are all ED symptoms created equally?*

The DSM-5 considers ED symptoms to be interchangeable. Within each ED diagnosis, certain criteria can be met in more than one way. For example, criterion B for AN can be met if a person endorses: (1) intense fear of gaining weight or becoming fat, even though underweight; or (2) persistent behaviors that interfere with weight gain. Criterion B for BN can be met if the person engages in one (or more) of the following to compensate for binge-eating episodes at least once per week for 3 months: (1) fasting; (2) excessive exercise; (3) self-induced vomiting; (4) laxative, diuretic, enema, suppository, or syrup of ipecac use; or (5) insulin omission (for persons with diabetes mellitus) or misuse of thyroid medication. Similarly, BED requires a person to meet three out of five behavioral or cognitive symptoms associated with binge-eating episodes (criterion B).

Grouping symptoms as ‘equivalent’, for the purpose of determining whether someone meets a specific criterion for a diagnosis makes sense if the behaviors

and cognitions that count toward the criterion are functionally equivalent. Yet, there are few empirical studies to indicate whether symptoms actually are similar in terms of their clinical utility, reliability and validity. Published research studies indicate that certain criterion B symptoms of BED, including: eating much more rapidly than normal (criterion B1) and eating alone because of being embarrassed by how much one is eating (criterion B4), do not load on a latent binge eating factor (Forbush *et al.* 2013). Other research indicates that these same criteria have lower reliabilities ( $\kappa$ ) compared with other symptoms of BED (Brody *et al.* 1994), which suggests that these symptoms are weaker than other criterion B symptoms in predicting diagnoses.

With regard to BN, Thelen *et al.* (1991) found that self-reported laxative and diuretic use did not predict future diagnosis. These authors attributed the lack of predictive validity to the low base rates of these behaviors in their sample, and suggested that although these behaviors are rare, they are still clinically important. Recently, however, Hovrud & De Young (2015) tested a community sample of individuals with EDs, and found that nearly half of the variance in psychosocial impairment was uniquely accounted for by binge-eating frequency, weight/shape concerns and dietary restriction. Purging frequency (which included self-induced vomiting and misuse of laxatives or diuretics) and fear of ‘fatness’ were not unique predictors of impairment. Rather than suggesting that purging is an unimportant symptom, these studies highlight the possibility that certain ED symptoms may be more ‘central’ than others in terms of predicting other clinically relevant variables, such as impairment and relapse.

### *Trans-diagnostic models of ED psychopathology*

In contrast to the DSM system, Fairburn’s enhanced cognitive-behavioral model of EDs (CBT-E; Fairburn *et al.* 2008) theorizes that overvaluation of weight, shape or their control represents the ‘core’ psychopathology shared across all EDs. In this context, the term ‘overvaluation’ refers to basing one’s self-worth on one’s body weight or shape, and/or ability to exert strict control over one’s eating. An important feature of CBT-E is that overvaluation is theorized to lead to – and reinforce – further expressions of ED psychopathology.

Tabri *et al.* (2015) tested the theory underlying CBT-E in a large sample of individuals with EDs who were tested weekly over a 2-year period. Their findings indicated that overvaluation on a given week led to increases in non-compensatory weight-control behaviors the following week, and vice versa. The results of Tabri *et al.* (2015) were the same regardless of

ED diagnosis, which provided support for the trans-diagnostic nature of the CBT-E model. These data suggested that: (1) ED symptoms function similarly across ED diagnoses, and (2) certain ED symptoms are mutually self-reinforcing.

### *Application of network analysis to ED psychopathology*

The study of Tabri *et al.* (2015) provides insights into the temporal sequencing of two key DSM symptom clusters ('overvaluation' and non-compensatory weight control behaviors). However, additional studies are needed to identify patterns of interaction among symptoms across the full range of ED behaviors. Network analysis is a relatively new statistical technique that is ideally suited to understanding the dynamic interactions among psychopathological symptoms (Borsboom & Cramer, 2013). From the network perspective, individual symptoms are represented as nodes and connections are placed between symptoms that tend to co-occur, forming a web-like structure (or network) of a particular psychopathology. Symptom nodes in the network are viewed as interconnected clusters that may or may not share an underlying, latent cause. For example, if one symptom within the network arises (e.g. prolonged fasting – which leads to increased physiological hunger drives), that symptom may activate other symptoms within the network (e.g. large eating episodes).

Network analysis is also able to test important chains of symptoms, such as those postulated in the CBT-E model. For example, 'overvaluation' of weight, shape or their control might directly lead to efforts to restrain one's eating (i.e. fasting), which over time leads to a negative energy balance that directly precipitates binge-eating episodes. Binge eating may, in turn, feedback to overvaluation due to actual or feared weight gain (i.e. overvaluation → fasting → binge-eating → overvaluation).

Finally, as we previously described, it appears that not all ED symptoms were created equally. Network analysis offers powerful insights into this issue by identifying the most central symptom(s), which has (have) many connections to other symptoms, and peripheral symptoms, which have few or no connections to other symptoms in the network. Because central symptoms activate many other symptom 'nodes' within the network, they represent symptoms that are particularly clinically important and may represent good treatment targets.

### *Current study*

Based on our literature review and previous theoretical models of ED psychopathology, we hypothesized that

symptoms that reflect body dissatisfaction or overvaluation of weight, shape or their control would represent the most impactful (central) symptoms within the ED 'network'.

## **Method**

### *Participants*

Participants were recruited from the community to take part in an ongoing longitudinal study of ED psychopathology and co-morbidity course and maintenance. Participants were recruited from fliers, newspaper advertisements, bus advertisements and mass emails sent to faculty, staff and students at two large universities in the Midwest of the USA. The current study reports data from participants' baseline (initial) study visit.

Because a goal of the larger study for which these data were collected was to test the longitudinal course of both full- and subthreshold EDs, our inclusion/exclusion criteria were intentionally broad to identify a representative community sample of individuals with EDs. Inclusion criteria were: (1) aged  $\geq 14$  years; (2) endorsement of at least two symptoms of an ED on the Eating Disorder Diagnostic Scale (Stice *et al.* 2000); (3) a score  $\geq 16$  on the Clinical Impairment Assessment Questionnaire (CIA; Bohn *et al.* 2008) or diagnosis of a current full-threshold DSM-5 ED; and (4) body mass index (BMI)  $\geq 14$  kg/m<sup>2</sup>. A cut score of 16 was chosen on the CIA because this score has optimal sensitivity and specificity for distinguishing ED cases from non-cases (Bohn *et al.* 2008). We chose 14 years as the lower age limit because the peak age of ED onset is mid-adolescence to early adulthood and EDs are rare among younger adolescents and children (Swanson *et al.* 2011). Exclusion criteria were assessed using a screening questionnaire, and included: (1) BMI  $< 14$  kg/m<sup>2</sup>; (2) intellectual developmental disability, neurological disorder, or current psychosis; (3) inability to read/write fluently in English.

Participants ( $n = 143$ ; 77.6% women) ranged in age from 18 to 55 years, with a mean age of 25.0 (s.d. = 7.7) years. Participants' BMI ranged from 15.9 to 56.98 kg/m<sup>2</sup> (mean = 27.53, s.d. = 8.2 kg/m<sup>2</sup>). Participants self-reported the following ethnic and racial identities (participants were allowed to select multiple categories): Caucasian (67.1%), Asian (15.4%), African-American (7.7%), multi-racial (5.6%), Native American or Alaskan Native (2.1%), Hawaiian or Pacific Islander (0.7%), or another race or ethnicity (3.5%); 9.1% of participants identified as Hispanic (of any race). Most participants were employed ( $n = 97$ ; 67.8%) and had  $\geq 12$  years of education ( $n = 142$ ; 99.3%; note: these data were missing for one participant).

ED diagnoses were applied using DSM-5 criteria. Participants were diagnosed with the following current EDs: AN ( $n=4$ ; 2.8%), BN ( $n=58$ ; 40.6%), BED ( $n=7$ ; 4.9%), or an OSFED ( $n=74$ ; 51.7%). The mean self-reported duration of an ED was 7.80 (s.d. = 6.97) years. Few participants were receiving psychotherapy/counseling ( $n=31$ ; 27.1%) or were prescribed psychotropic medications ( $n=26$ ; 18.2%).

### Procedure

The institutional review board approved all study procedures. Baseline procedures took place in person, and participants provided their written, informed consent prior to engaging in study-related procedures. Participants' weight and height were measured using a digital scale and wall-mounted stadiometer. Interviewers were trained research clinicians with a minimum bachelor's-level education, and were supervised by the first author (K.T.F.) during weekly diagnostic consensus meetings. Interviews were audiotaped (with participants' written permission); we randomly selected 10% of interviews that were rated by an independent research clinician to establish diagnostic reliability.

### Measures

#### Psychiatric diagnoses

The Structured Clinical Interview for DSM-IV-TR Axis I Disorders (SCID-I; First *et al.* 2007 revision) was adjusted for DSM-5 criteria to derive ED diagnoses. Inter-rater reliabilities for ED diagnoses were excellent (Conger's  $\kappa=0.89$  for AN and OSFED to  $\kappa=0.90$  for BN and BED).

#### Eating Pathology Symptoms Inventory (EPSI)

The EPSI is a 45-item self-report measure that was developed to comprehensively assess clinically relevant dimensions of ED psychopathology (Forbush *et al.* 2013). EPSI items reflect the frequency of various ED cognitions (e.g. 'I wished the shape of my body was different') and behaviors (e.g. 'I made myself vomit in order to lose weight'). Each item is rated on a five-point Likert scale that ranges from 0 (never) to 4 (very often). The EPSI has eight factor-analytically derived scales, including: Body Dissatisfaction, Binge Eating, Cognitive Restraint, Purging, Excessive Exercise, Restricting, Muscle Building, and Negative Attitudes toward Obesity. The EPSI has demonstrated evidence for strong convergent and discriminant validity, and good-to-excellent internal consistency and test-retest reliability across a range of different samples (Forbush *et al.* 2013, 2014). Moreover, the eight-factor structure of the EPSI has replicated between Western and non-Western cultures (Tang *et al.* 2015). We

chose to use the EPSI, rather than DSM-5 ED symptoms, because within the SCID-I there are numerous symptoms embedded within a single criterion. Because it is possible that certain symptoms within a DSM-5 criterion may have stronger or weaker importance in the overall network, we chose to use a better-differentiated symptom measure.

### Statistical analyses

SPSS (IBM Corp., 2011) and AgreeStat (Advanced Analytics LLC, 2010) were used to compute descriptive analyses and Conger's  $\kappa$ , respectively. The R package qgraph (Epskamp *et al.* 2012) was used to compute measures of centrality (see description in the next paragraph). Each 'node' represented one of the 45 EPSI items. We used a weighted association network<sup>2</sup> that reflected the Pearson's zero-order correlation strength between pairs of items in the EPSI. A potential concern with using Pearson's correlations is that there is a possibility that they may result in biased estimates for rank-ordered (ordinal) scales. To address this concern, we reanalysed the network using the 'cor\_auto()' function in qgraph to determine whether the assumption of multivariate normality (required for Pearson's  $r$ ) was violated. Results showed that the same set of nodes was highly central regardless of correlation choice, and the interpretation of findings did not change as a result of whether Pearson's or polychoric correlations were used. We chose to use Pearson's  $r$  because this approach is consistent with recent network studies of psychopathology (Bringmann *et al.* 2015; McNally *et al.* 2015).

Following McNally *et al.* (2015), each correlation, or edge, represented the strength of association between two symptoms. Consistent with prior literature (McNally *et al.* 2015), we computed the following centrality measures: strength, closeness, and 'betweenness'. For all centrality measures, higher values reflect a node's greater importance to the network. 'Strength' represents the summed weights of connections to a particular node from other nodes.

'Closeness centrality' is defined as the inverse of the sum of distances from a particular node to all other nodes in the network. Note that 'distance' is the number of links (edges) that are traversed from one node to the next. Closeness centrality is an index of how 'quickly' information (or, in this case, ED symptomatology) flows through a network to reach a given node (Borgatti, 2005). Therefore, high closeness indicates a short average distance between a given node and all other nodes in the network.

Betweenness centrality measures the number of times a node appears in the shortest path between two other nodes. A node is said to be high in

betweenness centrality if it lies along many shortest paths, and a node is said to be low in betweenness centrality if it lies along few shortest paths. Manipulating (removing) nodes with high betweenness centrality would slow or halt the flow of information through the network (Borgatti, 2005).

Finally, we used key player analysis (Borgatti, 2008) as another way to define 'importance' by identifying which node(s) that, when removed from the network, resulted in maximal fracturing of the network into several smaller components. By fracturing a connected network into smaller, disconnected components one can disrupt the spread of information through the network. Although measures of centrality and key player analysis are both ways to identify 'important' nodes within the network, key player analysis identifies nodes that maintain the cohesiveness of the network. In other words, 'key players' are the nodes that, when removed, would result in a residual network with the least amount of cohesion possible. Removing nodes with high centrality, however, does not necessarily reduce the cohesion of the network into smaller sets of components. Thus, measures of centrality and key player analysis provide different and unique information about potential clinical applications.

The optimal set of 'key players' is determined by a measure known as 'fragmentation', which is defined as the ratio between the number of pairs of nodes that are not connected once the set of key players have been removed, and the total number of pairs in the original fully connected network (i.e. prior to the removal of the set of key players). The minimum fragmentation value of zero indicates that the network consists of a single component (i.e. it has not fractured), and the maximum fragmentation value of one indicates that the network has been completely fractured, solely consisting of isolates, or nodes with no connections (Borgatti, 2006).

### *Ethical standards*

All procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

### **Results**

Fig. 1 shows the suppressed network that contains links with correlations  $\geq 0.30$ . Abbreviations are used to designate each of the 45 EPSI items (see Table 1); these abbreviations are used in the figures depicting the centrality values of nodes.


As shown in the centrality plots (Fig. 2), EPSI items that have the highest strength (in descending order) include: trying on different outfits because one did not like how one looked (outfit), excessive exercise (ex\_hard), eating when not physically hungry (not\_hungry), being disgusted by the sight of obese people (disgust), and not liking how one's body looked (body). Nodes that had the highest closeness centrality (in descending order) included: trying on different outfits because one did not like how one looked (outfit), trying to avoid foods with high calorie content (cal\_avoid), not liking how clothes fit the shape of one's body (clothes), not liking how one's body looked (body), and not liking the size of one's thighs (thigh).

Nodes that had the highest betweenness centrality (in descending order) included: trying on different outfits because one did not like how one looked (outfit), considered taking diuretics to lose weight (diur\_consider), excessive exercise (ex\_hard), trying to avoid foods with high calorie content (cal\_avoid), and planned one's days around exercising (ex\_plan). It is noteworthy that trying on different outfits (outfit) emerged as the symptom with the highest value on all three centrality measures, suggesting that this symptom is of particular importance in the ED network.

In the key player analysis we attempted to find a set of nodes whose removal would partition the network into smaller, disconnected components. The optimal set of three nodes included: people encouraged me to eat more (eatmore), the need to exercise nearly every day (ex\_everyday), and trying to avoid foods with high calorie content (cal\_avoid). Note that this is not an ordered set, meaning one node in the set is not more 'important' than another. Rather, together the nodes represent the set of three nodes that maximally disconnects the network, as evident by a fragmentation value of 0.79. This set of three nodes provided maximal fragmentation with the minimum number of nodes in the set; smaller sets did not produce more fragmentation, and larger sets produced only incremental increases in fragmentation.

### **Discussion**

This project was the first to use network analysis to identify interactive systems of ED symptoms. We hypothesized that symptoms that reflected body dissatisfaction or overvaluation of weight, shape or their control would represent the most central symptoms within the network. Our results supported our hypothesis; body checking (as indicated by trying on different outfits because one did not like how one looked) and various other body dissatisfaction items emerged as some of the most important symptoms within the



**Fig. 1.** Eating disorders network. Each node represents an item on the Eating Pathology Symptoms Inventory (EPSI), and each link represents the zero-order correlation between each pair of items. Only links with  $r \geq 0.30$  are shown. The thickness of a link represents the magnitude of the correlation. Abbreviations for each node are provided in Table 1. Louvain community detection (Blondel *et al.* 2008) was used to identify the presence of densely connected subgroups of nodes within the network (Newman, 2006) or ‘communities’. Louvain community detection revealed the presence of seven ‘communities’, which approximated the covariance structure of the EPSI (Forbush *et al.* 2013). These communities reflected: Restricting (Community 1); Muscle Building scale items that assess protein supplement and steroid misuse (Community 2); Body Dissatisfaction (Community 3); Excessive Exercise, Cognitive Restraint, and an item that reflects desire for increased muscularity (Community 4); Negative Attitudes toward Obesity (Community 5); purging behaviors, excluding self-induced vomiting (Community 6); and binge eating items, plus the self-induced vomiting item (Community 7).

network, which was reflected by their high values on measures of strength, closeness and betweenness. In support of these findings, Lavender *et al.* (2013) found that daily body-checking frequency was associated with increased dietary restriction on the same (and next) day. Other research has shown that the frequency of body checking significantly predicted quality-of-life impairment in both mental and physical health, above-and-beyond depression symptoms and relevant demographic variables (Latner *et al.* 2012).

Given that body checking emerged as one of the most important nodes within the network, it may be useful for future diagnostic conceptualizations of EDs to consider including body checking as a symptom.

Currently the DSM-5 includes ‘undue influence of body weight or shape on self-evaluation’ as a symptom of both AN and BN, but ‘body checking’ itself is not included as a symptom of EDs. Because it is probably easier to report overt behaviors rather than complex cognitive concepts, the inclusion of body checking within future diagnostic manuals may lead to improved diagnostic validity for persons with limited insight or poor abstract thinking abilities, such as children and adolescents.

Excessive-exercise symptoms had connections to desire for muscle building, and other studies have found positive associations between muscularity concerns and excessive exercise (Furnham *et al.* 2002;

**Table 1.** Abbreviations for EPSI items<sup>a</sup>

Abbreviations	EPSI item
ate_large	I ate a very large amount of food in a short period of time (e.g., within 2 hours)
stuffed	I stuffed myself with food to the point of feeling sick
uncomf_full	I ate until I was uncomfortably full
offer_food	If someone offered me food, I felt that I could not resist eating it
not_hungry	I ate when I was not hungry
auto_pilot	I ate as if I was on auto-pilot
snack	I snacked throughout the evening without realizing it
notice	I did not notice how much I ate until after I had finished eating
body	I did not like how my body looked
clothes	I did not like how clothes fit the shape of my body
shape	I wished the shape of my body was different
outfit	I tried on different outfits, because I did not like how I looked
muscle	I thought my muscles were too small
hips	I was not satisfied with the size of my hips
thigh	I did not like the size of my thighs
butt	I thought my butt was too big
unhealthy	I tried to exclude “unhealthy” foods from my diet
cal_avoid	I tried to avoid foods with high calorie content
cal_count	I counted the calories of foods I ate
surprise	People would be surprised if they knew how little I ate
eatmore	People encouraged me to eat more
notmuch	People told me that I do not eat very much
disgust	I was disgusted by the sight of obese people
lazy	I felt that overweight people are lazy
selfcontrol	I thought that obese people lack self-control
unattractive	I felt that overweight people are unattractive
tight	I was disgusted by the sight of an overweight person wearing tight clothes
vomit	I made myself vomit in order to lose weight
laxative	I thought laxatives are a good way to lose weight
steroids	I thought about taking steroids as a way to get more muscular
suppl_used	I used muscle building supplements
suppl_consider	I considered taking a muscle building supplement
diet_pills	I used diet pills
suppl_used_p	I used protein supplements
diur_use	I used diuretics in order to lose weight
diur_consider	I considered taking diuretics to lose weight
diet_tea	I used diet teas or cleansing teas to lose weight
ex_hard	I pushed myself extremely hard when I exercised
ex_strenuous	I engaged in strenuous exercise at least five days per week

**Table 1** (cont.)

Abbreviations	EPSI item
ex_exhaust	I exercised to the point of exhaustion
ex_plan	I planned my days around exercising
ex_everyday	I felt that I needed to exercise nearly every day
skip	I skipped two meals in a row
full_easy	I got full more easily than most people
full_small	I got full after eating what most people would consider a small amount of food

EPSI, Eating Pathology Symptoms Inventory.

<sup>a</sup>To download a copy of the EPSI, please visit <https://psych.ku.edu/kelsie-t-forbush>. Please note that the EPSI is free for non-commercial research or clinical use. Any modification of items must be approved in writing from the copyright holder (Kelsie Forbush). Copyright 2011 Kelsie Forbush.

Tiggemann *et al.* 2007), with men reporting higher levels of excessive exercise (albeit for different reasons) than women (Forbush *et al.* 2013, 2014). Excessive exercise also had connections to symptoms of cognitive restraint, indicating that excessive exercise may trigger (or be triggered by) goals to achieve a bigger or thinner body image ideal. It was noteworthy that certain attitudes toward others' bodies (e.g. believing that overweight people are unattractive) were connected to excessive-exercise behaviors. Although additional research is needed, findings indicate that reducing dietary restraint and cognitions related to desire for muscularity and negative attitudes toward overweight persons could be important treatment targets for patients with compulsive-exercise patterns. Interestingly, one exercise behavior – feeling the need to exercise nearly every day (ex\_everyday) – was identified as a key player (along with eatmore and cal\_avoid). We suggest that these key symptoms may be useful for generating hypotheses for future treatment development: interventions targeted at these nodes could create ‘firewalls’ in the symptom network, preventing or at least slowing the cascading of symptoms that might be brought on by certain ‘trigger’ symptoms (i.e. symptoms high in centrality measures).

Similar to Hovrud & De Young (2015), we found that self-induced vomiting had lower ‘importance’ within the network, as indicated by low values of betweenness. Other forms of purging (i.e. diuretic and laxative misuse) had low centrality values compared with other ED symptoms. These data do not indicate that purging behaviors are unworthy of clinical attention; it is clear that purging behaviors have deleterious health consequences. However, our data




Fig. 2. Centrality measures for the eating disorders network representing the betweenness, closeness and strength of each node. Abbreviations for each node are provided in Table 1.

suggest that purging behaviors have fewer connections to other ED symptoms. For example, although self-induced vomiting was most strongly associated with binge-eating behaviors, it had relatively weak connections to other symptoms within the network. Misuse of diuretics or laxatives was connected to restricting behaviors, with weaker connections to other symptoms. Our results indicate that targeting other, non-purging symptoms in the network could have indirect effects on purging behaviors; directly focusing on reducing binge-eating and restricting behaviors is probably all that is needed to ameliorate purging behaviors. These findings support the DSM-5's classification of purging as a method of compensating for perceived or actual binge-eating episodes.

Certain limitations of this study are worth noting. First, the prevalence of certain ED diagnoses in the current study was low. However, lifetime diagnoses of AN and BED were higher, and other research supports the trans-diagnostic relevance of the EPSI in individuals with low-weight EDs (Forbush *et al.* 2013, 2014). On the other hand, because Mountford *et al.* (2007) found that body-checking behavior is less prevalent among individuals with AN (*v.* BN), it will be important for future studies to replicate our findings in samples with greater numbers of individuals with AN. Second, because our study tested associations among ED symptoms at one point in time, directionality cannot be inferred. Third, although our sample size is low for certain latent variable models, Monte Carlo



research indicates that samples between 100 and 200 provide reliable estimates of the data in network analysis (Scott, 2012). Fourth, consistent with recent research (McNally *et al.* 2015), our study used an association network, but research testing partial correlation networks could be informative because a correlation between two nodes might be high, but only because a third variable causes them both. Finally, although studies have shown high correlations between body checking and body image dissatisfaction (Reas *et al.* 2002; Shafran *et al.* 2004; Shafran *et al.* 2007; Berg *et al.* 2012), future research is needed to test whether the specific EPSI Body Dissatisfaction scale item 'I tried on different outfits because I did not like how my body looked' is associated with greater body checking or avoidance behaviors. Despite limitations, the current study has several strengths. This study was the first to apply network analysis to understanding ED psychopathology. Our study sample had more ethnic-racial diversity than most other published community samples of individuals with EDs. We also used a measure with good discriminant validity, which provided an opportunity to identify the unique contribution of symptoms within the network (in contrast to diagnostic measures that include numerous symptoms within each criterion).

In conclusion, the results of our study support the key mechanisms delineated in CBT-E, given that the most central symptoms within the network represented overt behavior that reflect overvaluation with weight, shape or their control (i.e. body checking) and body dissatisfaction. We believe our findings may be useful for generating hypotheses to be tested in future treatment studies. For example, although CBT-E already includes techniques to address overvaluation of weight, shape or their control, focused work on body checking is introduced after the first eight sessions of therapy (Fairburn, 2008). Our findings suggest that it may be informative to test whether introducing body-checking 'experiments' earlier in CBT-E improves treatment response, given that rapid, early change among patients with an ED has been shown to be an important predictor of outcome (Agras *et al.* 2000; Raykos *et al.* 2013). More targeted interventions for reducing body checking may also be useful for producing improved outcomes such as focused mirror exposure, which has shown promise for reducing long-term behavioral and cognitive expressions of overvaluation in non-clinical women with high levels of pre-treatment body dissatisfaction (Jansen *et al.* 2015). Additional future directions for research include understanding neuropsychological or personality-based mechanisms that underlie associations between body checking and other ED symptoms. Targeted treatments that disrupt mechanisms

underlying body checking may lead to reductions in ED symptomatology and improvements in quality of life for individuals with EDs.

### Supplementary material

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S003329171600012X>

### Acknowledgements

This work was supported by the Clifford B. Kinley Trust, and the University of Kansas New Faculty Research Fund. The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding agencies.

### Declaration of Interest

None.

### Notes

<sup>1</sup> The scientific community has used three general criteria by which to support the use of the term causality, including association, temporal precedence, and isolation/internal validity. The term 'causal' within the network analysis literature has recently been used to illustrate a major point of philosophical divergence from traditional latent modeling approaches (Borsboom & Cramer, 2013). For example, traditional latent variable modeling approaches assume that symptoms of a single disorder (e.g. major depression) share a single, underlying (latent) cause, which is consistent with a disease or medical model. As a fundamental point of departure, however, network analysis makes the conceptual/theoretical assumption that symptoms are inter-connected, rather than indicators of the effects of a latent disorder. Thus, the network approach uses the term 'causal' to denote a symptom that activates other symptoms in circular and self-reinforcing ways, and cannot be thought of as directional (particularly when using undirected networks). As described by McNally *et al.* (2015), the term 'causal' refers to the fact that clinical interventions that affect one symptom will probably affect other symptoms that are directly connected to the target symptoms, and only produce indirect effects elsewhere in the network (Kindermann & Snell, 1980; Costantini *et al.* 2015). Nevertheless, given that the term 'causal' has traditionally been used to refer to directionality, we avoided this term in the current paper.

<sup>2</sup> Given that a correlation between two nodes might be high, but only because a third variable causes them both, we computed a 'concentration network' using partial correlations with edges >0.30 depicted. The resulting network was highly fractured with fewer connections among symptoms. Similar issues were identified by McNally *et al.* (2015), in which the partial correlation network was less

densely connected than the association (zero-order correlations) network. Although we chose to report the association network in the present paper, the partial correlation network can be obtained in the online Supplementary material.

## References

- Advanced Analytics LLC** (2010). *AgreeStat for Excel (Mac)*. Advanced Analytics LLC: Gaithersburg, MD.
- Agras WS, Crow SJ, Halmi KA, Mitchell JE, Wilson GT, Kraemer HC** (2000). Outcome predictors for the cognitive behavior treatment of bulimia nervosa: data from a multisite study. *American Journal of Psychiatry* **157**, 1302–1308.
- Allen KL, Byrne SM, Oddy WH, Crosby RD** (2013). DSM-IV-TR and DSM-5 eating disorders in adolescents: prevalence, stability, and psychosocial correlates in a population-based sample of male and female adolescents. *Journal of Abnormal Psychology* **122**, 720–732.
- American Psychiatric Association** (2013). *Diagnostic and Statistical Manual of Mental Disorders*, 5th edn. American Psychiatric Association: Washington, DC.
- Berg KC, Peterson CB, Frazier P, Crow SJ** (2012). Psychometric evaluation of the Eating Disorder Examination and Eating Disorder Examination-Questionnaire: a systematic review of the literature. *International Journal of Eating Disorders* **45**, 428–438.
- Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E** (2008). Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment* **10**, P10008.
- Bohn K, Doll HA, Cooper Z, O'Connor M, Palmer RL, Fairburn CG** (2008). The measurement of impairment due to eating disorder psychopathology. *Behaviour Research and Therapy* **46**, 1105–1110.
- Borgatti SP** (2005). Centrality and network flow. *Social Networks* **27**, 55–71.
- Borgatti SP** (2006). Identifying sets of key players in a social network. *Computational and Mathematical Organization Theory* **12**, 21–34.
- Borgatti SP** (2008). *Keyplayer 1.44*. Analytic Technologies: Lexington, KY.
- Borsboom D, Cramer AO** (2013). Network analysis: an integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology* **9**, 91–121.
- Bringmann LF, Lemmens LH, Huibers MJ, Borsboom D, Tuerlinckx F** (2015). Revealing the dynamic network structure of the Beck Depression Inventory-II. *Psychological Medicine* **45**, 747–757.
- Brody ML, Walsh BT, Devlin MJ** (1994). Binge eating disorder: reliability and validity of a new diagnostic category. *Journal of Consulting and Clinical Psychology* **62**, 381–386.
- Castellini G, Lo Sauro C, Mannucci E, Ravaldi C, Rotella CM, Faravelli C, Ricca V** (2011). Diagnostic crossover and outcome predictors in eating disorders according to DSM-IV and DSM-V proposed criteria: a 6-year follow-up study. *Psychosomatic Medicine* **73**, 270–279.
- Costantini G, Epskamp S, Borsboom D, Perugini M, Mõttus R, Waldorp LJ, Cramer AO** (2015). State of the aRt personality research: a tutorial on network analysis of personality data in R. *Journal of Research in Personality* **54**, 13–29.
- Cuthbert BN** (2014). The RDoC framework: facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry* **13**, 28–35.
- Eddy KT, Dorer DJ, Franko DL, Tahilani K, Thompson-Brenner H, Herzog DB** (2008). Diagnostic crossover in anorexia nervosa and bulimia nervosa: implications for DSM-V. *American Journal of Psychiatry* **165**, 245–250.
- Epskamp S, Cramer AO, Waldorp LJ, Schmittmann VD, Borsboom D** (2012). qgraph: Network visualizations of relationships in psychometric data. *Journal of Statistical Software* **48**, 1–18.
- Fairburn CG** (2008). Shape concern, shape checking, feeling fat, and mindsets. In *Cognitive Behavior Therapy and Eating Disorders*, pp. 96–123. The Guilford Press: New York.
- Fairburn CG, Bohn K** (2005). Eating disorder NOS (EDNOS): an example of the troublesome “not otherwise specified” (NOS) category in DSM-IV. *Behaviour Research and Therapy* **43**, 691–701.
- Fairburn CG, Cooper Z, Shafran R** (2008). Enhanced cognitive behavior therapy for eating disorders (“CBT-E”): an overview. In *Cognitive Behavior Therapy and Eating Disorders* (ed. C. G. Fairburn), pp. 23–34. Guilford Press: New York.
- Fairweather-Schmidt AK, Wade TD** (2014). DSM-5 eating disorders and other specified eating and feeding disorders: is there a meaningful differentiation? *International Journal of Eating Disorders* **47**, 524–533.
- First MB, Spitzer RL, Gibbon M, Williams JBW** (2007 Revision). *Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition With Psychotic Screen (SCID-I/P W/PSY SCREEN)*. Biometrics Research, New York State Psychiatric Institute: New York.
- Forbush KT, Wildes JE, Hunt TK** (2014). Gender norms, psychometric properties, and validity for the Eating Pathology Symptoms Inventory. *International Journal of Eating Disorders* **47**, 85–91.
- Forbush KT, Wildes JE, Pollack LO, Dunbar D, Luo J, Patterson K, Pollpeter M, Stone A, Bright A, Watson D** (2013). Development and validation of the Eating Pathology Symptoms Inventory (EPSI). *Psychological Assessment* **25**, 859–878.
- Furnham A, Badmin N, Sneade I** (2002). Body image dissatisfaction: gender differences in eating attitudes, self-esteem, and reasons for exercise. *Journal of Psychology* **136**, 581–596.
- Hovrud L, De Young KP** (2015). Unique contributions of individual eating disorder symptoms to eating disorder-related impairment. *Eating Behaviors* **18**, 103–106.
- IBM Corp.** (2011). *IBM SPSS Statistics for Windows, Version 20.0*. IBM Corp: Armonk, NY.
- Insel TR** (2014). The NIMH research domain criteria (RDoC) project: precision medicine for psychiatry. *American Journal of Psychiatry* **171**, 395–397.

- Jansen A, Voorwinde V, Hoebink Y, Rekkers M, Martijn C, Mulkens S** (2015). Mirror exposure to increase body satisfaction: should we guide the focus of attention towards positively or negatively evaluated body parts? *Journal of Behavior Therapy and Experimental Psychiatry* **50**, 90–96.
- Keel PK, Brown TA, Holland LA, Bodell LP** (2012). Empirical classification of eating disorders. *Annual Review of Clinical Psychology* **8**, 381–404.
- Kindermann R, Snell JL** (1980). *Markov Random Fields and their Applications*. American Mathematical Society Providence, RI.
- Latner JD, Mond JM, Vallance JK, Gleaves DH, Buckett G** (2012). Body checking and avoidance in women: associations with mental and physical health-related quality of life. *Eating Behaviors* **13**, 386–389.
- Lavender JM, De Young KP, Wonderlich SA, Crosby RD, Engel SG, Mitchell JE, Crow SJ, Peterson CB, Le Grange D** (2013). Daily patterns of anxiety in anorexia nervosa: associations with eating disorder behaviors in the natural environment. *Journal of Abnormal Psychology* **122**, 672–683.
- McNally RJ, Robinaugh DJ, Wu GW, Wang L, Deserno MK, Borsboom D** (2015). Mental disorders as causal systems a network approach to posttraumatic stress disorder. *Clinical Psychological Science* **3**, 836–849.
- Mountford V, Haase AM, Waller G** (2007). Is body checking in the eating disorders more closely related to diagnosis or to symptom presentation? *Behaviour Research and Therapy* **45**, 2704–2711.
- Newman ME** (2006). Finding community structure in networks using the eigenvectors of matrices. *Physical Review E* **74**, 036104.
- Raykos BC, Watson HJ, Fursland A, Byrne SM, Nathan P** (2013). Prognostic value of rapid response to enhanced cognitive behavioral therapy in a routine clinic sample of eating disorder outpatients. *International Journal of Eating Disorders* **46**, 764–770.
- Reas DL, Whisenhunt BL, Netemeyer R, Williamson DA** (2002). Development of the body checking questionnaire: a self-report measure of body checking behaviors. *International Journal of Eating Disorders* **31**, 324–333.
- Regier DA, Kuhl EA, Kupfer DJ** (2013). The DSM-5: classification and criteria changes. *World Psychiatry* **12**, 92–98.
- Scott J** (2012). Lines, neighbourhoods, and densities. In *Social Network Analysis*, pp. 63–82. Sage: Thousand Oaks, CA.
- Shafran R, Fairburn CG, Robinson P, Lask B** (2004). Body checking and its avoidance in eating disorders. *International Journal of Eating Disorders* **35**, 93–101.
- Shafran R, Lee M, Payne E, Fairburn CG** (2007). An experimental analysis of body checking. *Behaviour Research and Therapy* **45**, 113–121.
- Stice E, Marti CN, Rohde P** (2013). Prevalence, incidence, impairment, and course of the proposed DSM-5 eating disorder diagnoses in an 8-year prospective community study of young women. *Journal of Abnormal Psychology* **122**, 445–457.
- Stice E, Telch CF, Rizvi SL** (2000). Development and validation of the Eating Disorder Diagnostic Scale: a brief self-report measure of anorexia, bulimia, and binge-eating disorder. *Psychological Assessment* **12**, 123–131.
- Swanson SA, Crow SJ, Le Grange D, Swendsen J, Merikangas KR** (2011). Prevalence and correlates of eating disorders in adolescents: results from the National Comorbidity Survey Replication Adolescent Supplement. *Archives of General Psychiatry* **68**, 714–723.
- Tabri N, Murray H, Thomas J, Franko D, Herzog D, Eddy K** (2015). Overvaluation of body shape/weight and engagement in non-compensatory weight-control behaviors in eating disorders: is there a reciprocal relationship? *Psychological Medicine* **45**, 2951–2958.
- Tang X, Forbush KT, Lui PP** (2015). Development and validation of the Chinese-Language Version of the Eating Pathology Symptoms Inventory (CEPSI). *International Journal of Eating Disorders* **48**, 1016–1023.
- Thelen MH, Farmer J, Wonderlich S, Smith M** (1991). A revision of the Bulimia Test: The BULIT-R. *Psychological Assessment* **3**, 119–124.
- Tiggemann M, Martins Y, Kirkbride A** (2007). Oh to be lean and muscular: body image ideals in gay and heterosexual men. *Psychology of Men and Masculinity* **8**, 15–24.
- Wildes JE, Forbush KT, Markon KE** (2013). Characteristics and stability of empirically derived anorexia nervosa subtypes: towards the identification of homogeneous low-weight eating disorder phenotypes. *Journal of Abnormal Psychology* **122**, 1031–1041.
- Wildes JE, Marcus MD** (2013). Alternative methods of classifying eating disorders: models incorporating comorbid psychopathology and associated features. *Clinical Psychology Review* **33**, 383–394.