

Exogenous nucleosides accelerate differentiation of rat intestinal epithelial cells

Jose María Vieites¹, Raúl de la Torre¹, María del Carmen Ramírez¹, María Isabel Torres², Antonio Sánchez-Pozo¹, Angel Gil¹ and Antonio Suárez^{1*}

¹Department of Biochemistry and Molecular Biology II, School of Pharmacy, University of Granada, Granada, 18071, Spain

²Department of Experimental Biology, University of Jaen, Jaen, Spain

(Received 27 March 2007 – Revised 25 June 2007 – Accepted 20 August 2007)

Several studies have reported differing data on the effect of exogenous nucleosides and nucleotides on the proliferation and differentiation in various intestinal cell lines and explants. To study whether exogenous nucleosides modulate intestinal cell differentiation, IEC-6 cells were differentiated in the presence or absence of a nucleoside mixture (cytidine, uridine, guanosine and inosine, 30 µM each), and the concentrations of nucleoside derivatives were determined by HPLC. Cell differentiation was assessed by electron microscopy, alkaline phosphatase activity and *Rnd3* gene expression. The present results showed that uridine, guanosine and inosine were cleared from culture media (up to 32, 63 and 100 % in proliferating cells, and 31, 80 and 94 % in differentiated cells, respectively) whereas cytidine concentrations increased. Differentiation of IEC-6 cells was associated with a significant increase in intracellular nucleotide concentrations. Clearance of nucleosides correlated with a significant increase in the intracellular nucleotide pool in proliferating and differentiated IEC-6 cells. Intracellular guanosine nucleotides increased 2.5- and 5-fold in nucleoside-supplemented proliferating and differentiated cells, respectively. At 24 h, nucleoside-supplemented differentiated IEC-6 cells had significantly higher energy charge and GTP levels than non-supplemented ones. These modifications paralleled changes in cell differentiation as indicated by increased alkaline phosphatase activity, prolonged microvilli formation and accelerated down-regulation of *Rnd3* gene expression. The present findings suggest that exogenous nucleosides were selectively taken up by IEC-6 cells, increased the intracellular nucleotide pool, GTP and energy charge, and favoured cell morphological and functional changes during differentiation.

Differentiation: Intestinal epithelial cells: Exogenous nucleosides: Intracellular nucleotides

Nucleotides (NT) are the building blocks of DNA and RNA and are normal components of the human diet and the non-protein-nitrogen fraction of milk¹. Nucleosides (NS) and NT are not considered essential since cells possess biochemical pathways for their *de novo* synthesis and salvage. However, in certain circumstances, NT may become conditionally essential nutrients that modulate different biological processes, such as lipid metabolism and immune function, and have an important role in cell proliferation, maturation, and apoptosis^{2–4}. Besides, dietary NT have been shown to enhance gut recovery after different kinds of intestinal injury^{5–9}. Because of the biochemical and trophochemical properties of dietary NT and NS, the European Commission has permitted some of them to be used as a supplement in the manufacture of infant and follow-on formulas.

Although NT deficiencies have not been related to any particular disease, various studies have assessed the effect of exogenous NT and NS on the intracellular pool of NT and on cell proliferation and differentiation. In hepatic cells, our group showed that exogenous NS were actively incorporated by hepatocytes and stimulated hepatocyte proliferation and

expression of *c-myc*, *h-ras* genes and α -fetoprotein^{10,11}. Gil *et al.*¹² recently showed that intestinal explants hydrolyse RNA to NT, and NT to NS that were efficiently incorporated into intestinal cells, except for cytidine. Reports on the effect of NT in intestinal cell proliferation and differentiation are contradictory. Sato *et al.*¹³ reported that a 10:1:1:1:1 CMP–AMP–UMP–IMP–GMP mixture enhanced proliferation, sucrase and maltase activities in IEC-6 cells but not in Caco-2 cells. In the same line, addition of NT (AMP, CMP, UMP, IMP, GMP) to a glutamine-free medium stimulated intestinal cell proliferation and increased alkaline phosphatase and sucrose-isomaltase activities in differentiated cells¹⁴. However, Boza *et al.*¹⁵ showed that the proliferative effect of NT on Caco-2 cells was associated with the presence of glutamine in cell culture medium. Moreover, Tanaka *et al.*¹⁶ and Chen *et al.*¹⁷ observed that AMP supplementation decreased cellular proliferation and increased apoptosis in organ cultures and IEC-6 cells, respectively.

The goal of the present study was to reconcile the conflicting data published on the effect of NS on intestinal cells. The aim was to test the hypothesis that exogenous NS are taken up

Abbreviations: cDNA, first-strand complementary DNA; D – N, differentiated IEC-6 cells; D + N, differentiated IEC-6 cells grown with exogenous nucleosides; NS, nucleosides; NT, nucleotides; P – N, proliferating IEC-6 cells; P + N, proliferating IEC-6 cells grown with exogenous nucleosides.

* **Corresponding author:** Dr Antonio Suárez, fax +34 958248960, email asuarez@ugr.es

and modify the concentrations of intracellular NT and the energy charge with effects on the morphological, functional and gene expression changes associated to cell differentiation. Overall, these data would determine whether selective NS uptake by intestinal cells alter the intracellular NT pool, thereby enhancing the differentiation on enterocytes. For this purpose, we used IEC-6 cells that undergo differentiation when grown on an extracellular protein matrix. IEC-6 cells were grown in the presence or absence of an inosine–guanosine–cytidine–uridine supplement at physiologic levels (30 μM each, a concentration selected to match NS concentrations in human milk, which range from 0.5 to 37 μM)¹⁸.

Materials and methods

Materials

Plastic culture ware was purchased from Nunc (Roskilde, Denmark). Cell culture medium (Dulbecco's modified Eagle's medium), trypsin, insulin, NS and NT were from Sigma Aldrich (St. Louis, MO, USA). SuperScript II, RNase Out, Trizol, L-glutamine, penicillin/streptomycin, fetal bovine serum and PBS were from Invitrogen (Carlsbad, CA, USA). [α -³²P]dCTP, Matrigel and dispase were purchased from Beckton Dickinson (Bedford, MA, USA). DNase I FPLCpure and TaqGold polymerase were from Amersham Pharmacia Biotech (Piscataway, NJ, USA). NIH Scion Image Beta 4.0.2 software (Scion Corporation, MD, USA) was used to quantify band intensity.

Cell culture

IEC-6 cell line was obtained from the American Type Culture Collection (Rockville, MD, USA) and was used between passages 20 and 25 in most experiments. A stock culture of IEC-6 was maintained using Dulbecco's modified Eagle's medium with 4.5 g/l glucose, 10% fetal bovine serum, bovine insulin (0.1 U/ml), penicillin/streptomycin and L-glutamine (0.6 g/l). The stock was passaged twice a week and fed three times a week.

In order to determine the effects of NS on cell proliferation and differentiation, IEC-6 cells (5×10^5) were plated on 35 mm dishes coated with a thin layer of Matrigel. Media were supplemented with or without NS (cytidine, uridine, guanosine, inosine; 30 μM each). The culture media were replaced with fresh NS every 12 h. Cells were recovered from Matrigel by using dispase according to manufacturer's instructions.

Transmission electron microscopy studies

IEC-6 cells were grown on 3.0 μm pore size Transwell-clear polyester membranes precoated with Matrigel, or on 3.0 μm Transwell-COL type I and III collagen-coated membranes (Corning, Cultek, Spain). At 12 and 24 h, monolayers were washed and fixed in 2.5% glutaraldehyde in cacodylate buffer (0.1 M, pH 7.3) and post-fixed in 1% osmium tetroxide. Monolayers were then dehydrated in ethanol and embedded in Epon 812 resin. Ultrathin sections (50 nm) were contrasted with uranyl acetate and lead citrate and examined under a Zeiss 902 transmission electron microscope.

HPLC analysis of nucleosides and nucleotides

NS present in culture media were separated and quantified by means of an ion-exchange HPLC method with some changes¹⁹. Briefly, media were recovered using a syringe and filtered through a 0.45 μm filter to eliminate cell debris. Cleared media (40 μl) were injected into the chromatograph. Two mobile phases were used: (1) 0.1 M-potassium acetate, pH 4.5, 2 mM-hexasulphonic acid and 1% acetonitrile; and (2) 0.1 M-potassium acid, pH 4.5, 2 mM-hexasulphonic acid and 6% acetonitrile. NS were identified and quantified at 254 nm and compared with HPLC standards.

NT intracellular pools were analysed by using a HPLC method²⁰. Briefly, cells were rinsed with PBS and frozen using liquid nitrogen to avoid NT degradation. Cells were then treated with perchloric acid and incubated for 45 min to precipitate proteins. Supernatants were neutralized with potassium hydroxide, cooled, centrifuged and filtered. Cleared supernatants (60 μl) were injected into the chromatograph. Two mobile phases were used: (1) 0.2 M-ammonium phosphate, 0.1 M-triethanolamine, pH 6; and (2) 0.2 M-ammonium phosphate, 0.1 M-triethanolamine, pH 6, and 15% acetonitrile. NT were identified and quantified at 254 nm and compared with HPLC standards.

Enzyme activities

Alkaline phosphatase (EC 3.1.3.1) activity was measured on crude cell homogenates using a spectrophotometric method based on the use of *p*-nitrophenylphosphate, which is transformed to phosphate and *p*-nitrophenol. The latter is a coloured compound that can be measured at 405 nm²¹. Units were expressed per milligram protein. The protein concentration was determined by the bicinchoninic acid method, using bovine albumin serum as standard²².

RNA extraction and reverse transcription-PCR

Total RNA was extracted from cells grown on 35 mm dishes using Trizol according to the manufacturer's instructions. In order to avoid contamination by DNA, total RNA was treated with DNase I. The reaction took place at 37°C for 30 min. Finally, RNA was extracted using phenol–chloroform and precipitated with sodium acetate and ethanol. The pellet was washed with 70% ethanol and stored dry at –80°C. First-strand complementary DNA (cDNA) was synthesized with 0.1 μg DNase-treated RNA and 0.5 μg oligo-dT₁₆ in 20 μl of a solution containing 1 \times first-strand buffer, 10 mmol/l dithiothreitol, 500 $\mu\text{mol/l}$ each dNTP and 200 U Moloney murine leukaemia virus RT. PCR reactions were carried out with primers for sucrase-isomaltase cDNA amplification forward primer, 5'-CAGAACCACCTGGTCTGAG-3'; reverse primer, 5'-TGGTTGTCATCTGCAGCAAC-3'. Glyceraldehyde-3-phosphate dehydrogenase was used as an internal control of a 'housekeeping' gene which was amplified with a forward primer (5'-TAAAGGGCATCCTGGGCT-3') and a reverse primer (5'-TTACTC CTTGGAGGCCATG-3'). PCR reactions were performed in the same set with an Applied Biosystems DNA thermal cycler model 9700 as follows: an initial denaturing for 5 min at 94°C; forty cycles: 94°C for 30 s, 55°C for 30 s and 72°C for 30 s; final extension at 72°C for 1 min.

DNA synthesis was resolved by standard agarose gel electrophoresis.

Northern blot analysis

A total of 10 mg of denatured RNA was loaded per lane in a formaldehyde–1% agarose gel, electrophoresed, transferred to nylon filters, and fixed with a UV crosslinker. The integrity of each sample was confirmed by staining the RNA with ethidium bromide. Filters were sequentially hybridized with a *Rnd3* [α - 32 P]dCTP-labelled cDNA probe, stripped and hybridized with a [α - 32 P]dCTP-labelled rat glyceraldehyde-3-phosphate dehydrogenase cDNA probe to validate Northern blot results.

Statistical analysis

Results are means and their standard errors (n 3). Since data were not normally distributed, comparisons between different groups were assessed by non-parametric tests. Mean comparisons between independent groups were calculated using the Mann–Whitney *U*-test. Mean comparisons within a group for different times were done using a Wilcoxon test for paired samples. Differences with $P < 0.05$ were considered significant. Data were analysed using the Statistical Package for Social Sciences software (SPSS Inc., Chicago, IL, USA).

Results

Uptake of nucleosides by IEC-6 cells

The time course of exogenous NS clearance in medium was determined. The culture media were replaced with fresh NS (30 μ M each) every 12 h. The present results show that uridine and purine NS were taken up by IEC-6 cells (Table 1). Purine NS were cleared at higher rates than pyrimidine NS. The clearance rate of uridine concentrations was similar in proliferating and differentiated cells, the values decreasing to 69% of the initial concentration at 12 h and remaining at that level at 24 and 48 h. Guanosine was initially taken up by differentiated cells at a higher rate than proliferating cells although the contrary was observed at 48 h. Inosine was the NS with the highest clearance rate in IEC-6 cells. In proliferating cells, inosine was completely cleared from culture media at 24 and 48 h. At all time-points, cytidine concentrations were

higher than the dose in culture media (30 μ M) in proliferating and differentiated cells. No uric acid was detectable in culture media.

Effects of nucleosides on the morphology of differentiated IEC-6 cells

Exogenous NS did not change the external morphology of proliferating or differentiated IEC-6 cells. Intestinal cells grown on Matrigel showed a polarized morphology, typical of differentiated intestinal cells, with presence of microvilli at the apical pole. Interestingly, the addition of NS had a positive effect on the maturation of intestinal cells. Differentiated IEC-6 cells supplemented with NS (D + N) showed more microvilli on their surface than did non-supplemented differentiated IEC-6 cells (D – N), and microvilli were still observed at 24 h in D + N but not D – N cells (Fig. 1). At 48 h, no difference between D + N and D – N cells was observed (data not shown).

Intracellular pool of nucleotides in IEC-6 cells

Intracellular TMP, GDP and UDP were not detectable by HPLC analysis in proliferating IEC-6 cells. Thymidine NT, GDP and uridine diphosphoglucose were not detectable in differentiated cells. Except for TTP, concentrations of intracellular NT decreased from 12 to 24 h in proliferating IEC-6 cells (P – N) and in proliferating cells supplemented with NS (P + N; Table 2). The differentiation state resulted in an increase in the pool of intracellular NT in comparison with the proliferating state.

Addition of NS significantly changed the concentrations of intracellular NT in proliferating and differentiated cells. At 12 and 24 h, concentrations of intracellular NT and total intracellular NT were significantly higher in D + N *v.* D – N cells ($P < 0.05$) with the exception of AMP and CMP. Intracellular IMP was detectable only in D + N cells. Interestingly, the pool of guanosine NT was 5.0-fold higher in D + N cells than in D – N cells and 2.4-fold higher in P + N *v.* P – N cells, and these differences were significant. Although intracellular ATP did not change, the GTP levels and energy charge were significantly higher in D + N *v.* D – N cells (Fig. 2). It is noteworthy that intracellular NT levels in D + N cells at 12 h were similar to or higher than levels in D – N cells at 24 h.

Table 1. Nucleoside concentrations in media of proliferating and differentiated IEC-6 cells†
(Mean values with their standard errors for three determinations)

	Proliferating cells (mm)						Differentiated cells (mm)					
	12 h		24 h		48 h		12 h		24 h		48 h	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Cytidine	50.9	1.4	35.7	3.1	36.9	1.6	37.6*	1.6	40.6	0.4	39.9	1.1
Uridine	20.4	0.1	21.1	1.0	21.8	0.2	20.8	1.0	21.4	0.5	24.3*	0.1
Guanosine	11.2	0.6	5.3	0.3	6.8	0.1	6.1*	2.5	6.3	0.8	12.0*	0.4
Inosine	1.3	0.2	–	–	–	–	2.1	0.4	1.7*	0.1	6.2*	1.5

Mean values were significantly different from those of proliferating cells at the same time-point: * $P < 0.05$.
† IEC-6 cells were cultured in media supplemented with uridine, cytidine, guanosine and inosine (30 μ M each).




Fig. 1. Transmission electron micrographs of differentiated IEC-6 cells grown with (A) or without (B) nucleosides. Cell media was supplemented with uridine, cytidine, guanosine and inosine ($30\ \mu\text{M}$ each). \rightarrow , Microvilli.

Alkaline phosphatase activity in IEC-6 cells

On the functional level, differentiated cells showed greater alkaline phosphatase activity, a marker of enterocyte differentiation, compared with proliferating cells. Interestingly, the activity of alkaline phosphatase was significantly higher in D + N v. D - N cells (Fig. 3).

Analysis of *Rnd3* gene expression in differentiated IEC-6 cells

The differential display technique had been used to identify genes whose expression was regulated during enterocyte differentiation. Results obtained showed that *Rnd3* mRNA levels were lower in differentiated than in proliferating IEC-6 cells³⁶. In the present study, addition of NS accelerated the reduction in *Rnd3* mRNA levels in D + N v. D - N cells (Fig. 4).

Discussion

Contradictory reports have been published on the effect of NT or NS on the proliferation and differentiation of intestinal cells, probably due to differences in experimental approach, including the use of different mixtures of nucleot(s)ides at varying concentrations, and to the use of distinct cancer cell lines (Caco-2, HT-29)^{13–17}. Cancer cell lines are known to have higher rates of *de novo* NS synthesis than normal cells. To avoid the interference of cell transformation, the effect of NS on cell differentiation was studied in IEC-6 cells

because it is a non-transformed intestinal cell line that undergoes differentiation when grown on an extracellular protein matrix. The only published report on the effect of NT on IEC-6 cell differentiation showed that addition of NT enhanced proliferation but not differentiation, as assessed by alkaline phosphatase and sucrose-isomaltase activities¹⁴. However, some aspects of the latter study can be questioned. First, NT rather than NS were used, and the high level of alkaline phosphatase activity in mature intestine is known to degrade NT to phosphate and NS that enter cells via specific transporters²³. Second, the possibility of a hormonal effect due to adenosine generation in the culture media cannot be ruled out. For these reasons, IEC-6 cells were grown in the presence or absence of an inosine-guanosine-cytidine-uridine supplement at physiologic levels ($30\ \mu\text{M}$ each, a concentration selected to match NS concentrations in human milk, which range from 0.5 to $37\ \mu\text{M}$)¹⁸. Finally, Sterling & Cutroneo²⁴ detected no sucrase-isomaltase activity in differentiated IEC-6 cells, consistent with the present study, in which no expression of the sucrase-isomaltase gene was detected by reverse transcription-PCR in these cells.

According to the present findings, proliferating and differentiated IEC-6 cells control the selective uptake of NS (Table 1). Our group recently reported that hepatocytes^{10,11} and intestinal explants¹² are capable of selective NS uptake. The present results showed that the rate of uridine and purine NS clearance in IEC-6 cells was similar to that of adult hepatocytes¹⁰ and of intestinal explants¹². IEC-6 cells took up uridine, guanosine and inosine, up to 32, 63 and 100% in proliferating cells, and 31, 80 and 94% in differentiated cells, respectively. To date, three Na^+ -dependent concentrative and two Na^+ -independent equilibrative transporters have been identified²⁵. In IEC-6 cells, Aymerich *et al.*²⁶ showed that these cells express high levels of equilibrative NS transporters (with low substrate selectivity) and of the concentrative NS transporter CNT2 (highly selective for purine NS and uridine). Differentiated IEC-6 cells mainly express CNT2 and, slightly, CNT1 (selective for pyrimidine NS and adenosine) with a decrease in the activity of equilibrative transporters. The apparent K_m values for rat CNT2-mediated transport of uridine, guanosine and inosine are 24, 34 and $12\ \mu\text{M}$, respectively²⁷. We suggest that the predominance of CNT2 transporter activity would explain the preferential clearance of the former NS by proliferating and differentiated IEC-6 cells.

Concerning cytidine, Ohyanagi *et al.*²⁸ and Arnaud *et al.*¹⁰ observed that hepatocytes hardly took up cytidine. In the present study and in intestinal explants¹², there was an efflux of cytidine to the culture medium. The apparent K_m for rat CNT2-mediated transport of cytidine is $505\ \mu\text{M}$, 16.8-fold higher than its initial concentration in culture media. Therefore, cytidine would be poorly transported by CNT2 into IEC-6 cells. The results suggest that *de novo* synthesis and salvage of uridine by uridine kinase is preferentially used to meet intracellular pyrimidine NT needs. Cells produce cytidine, in the form of CTP, from UTP by cytidine triphosphate synthetase, using ATP and glutamine. The increase of cytidine in culture media may be the result of cytidine export due to excessive biosynthesis from UTP.

The present study is the first report on the effect of exogenous NS on the intracellular NT pool in differentiated intestinal

Table 2. Intracellular nucleotide concentrations in proliferating and differentiated IEC-6 cells cultured in the presence or absence of a nucleoside supplement†

(Mean values with their standard errors for three determinations)

	P - N (nmol/10 ⁶ cells)				P + N (nmol/10 ⁶ cells)				D - N (nmol/10 ⁶ cells)				D + N (nmol/10 ⁶ cells)			
	12 h		24 h		12 h		24 h		12 h		24 h		12 h		24 h	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
CDP-choline	3.2 ^c	0.2	—	—	4.7 ^{ab}	0.3	1.4 ^e	0.2	2.1 ^d	0.1	4.0 ^b	0.3	5.3 ^a	0.2	5.0 ^a	0.3
CMP	5.8 ^b	0.5	1.7 ^d	0.4	10.5 ^a	0.6	4.6 ^c	0.5	2.4 ^d	0.4	4.6 ^c	0.4	4.9 ^{bc}	0.6	2.3 ^d	0.3
CDP	2.7 ^e	0.5	—	—	—	—	—	—	45.7 ^c	3.6	75.5 ^b	6.1	97.9 ^a	6.4	101.0 ^a	3.9
CTP	8.6 ^c	0.4	0.6 ^f	0.1	10.4 ^b	0.8	3.1 ^e	0.4	5.9 ^d	0.5	13.2 ^a	2.6	14.5 ^a	2.6	13.6 ^a	1.6
Total cytosine nucleotides	20.3 ^d	2.5	2.3 ^f	0.5	25.6 ^d	2.9	9.1 ^e	1.0	56.1 ^c	4.6	97.3 ^b	8.8	122.6 ^a	9.5	121.9 ^a	10.3
UDP-glucose	4.2 ^b	0.3	1.0 ^d	0.3	6.9 ^a	1.0	3.0 ^c	0.2	—	—	—	—	—	—	—	—
UMP	6.3 ^d	0.5	1.9 ^e	0.4	14.8 ^c	1.4	6.0 ^d	0.5	6.8 ^d	0.9	5.8 ^d	0.6	40.9 ^a	3.9	33.1 ^b	4.2
UTP	14.8 ^b	2.0	4.1 ^d	0.2	25.9 ^a	3.7	10.3 ^c	1.3	11.0 ^c	1.8	16.6 ^b	2.1	30.9 ^a	3.1	30.5 ^a	3.4
Total uridine nucleotides	25.2 ^c	2.7	7.0 ^e	0.7	47.6 ^b	5.8	19.4 ^c	1.6	17.7 ^d	2.6	22.4 ^{cd}	2.1	71.8 ^a	6.0	63.7 ^a	6.4
NAD ⁺	—	—	0.4 ^b	0.1	1.2 ^a	0.2	0.5 ^b	0.1	—	—	1.7 ^a	0.2	1.3 ^a	0.2	1.5 ^a	0.4
AMP	—	—	—	—	—	—	—	—	5.0 ^b	0.4	8.0 ^a	1.0	8.4 ^a	0.8	7.5 ^a	0.5
ADP	6.4 ^b	0.4	3.7 ^c	0.3	8.4 ^a	0.8	4.8 ^{bc}	0.9	5.8 ^b	0.9	9.9 ^a	1.2	9.2 ^a	0.3	10.5 ^a	1.6
ATP	10.7 ^c	0.7	4.2 ^d	0.9	9.9 ^c	1.3	4.2 ^d	0.6	20.0 ^b	3.5	32.9 ^a	1.7	27.8 ^a	3.3	34.3 ^a	3.4
Total adenine nucleotides	17.1 ^c	2.9	8.3 ^d	1.2	19.5 ^c	2.5	9.5 ^d	1.4	30.8 ^b	5.4	52.5 ^a	6.3	46.7 ^a	6.4	53.9 ^a	7.7
GMP	32.6 ^e	5.8	15.2 ^f	2.5	120.3 ^b	11.1	38.9 ^{de}	4.2	47.0 ^d	7.2	65.8 ^c	9.7	420.3 ^a	42.8	428.5 ^a	51.2
GTP	2.2 ^f	0.5	2.2 ^f	0.4	10.1 ^e	1.4	2.8 ^f	0.6	30.2 ^d	6.7	49.4 ^c	7.0	99.8 ^b	12.3	137.5 ^a	18.5
Total guanine nucleotides	34.8 ^c	5.1	17.4 ^d	1.8	130.3 ^b	15.7	41.7 ^c	6.6	77.2 ^c	3.3	115.2 ^b	10.6	520.0 ^a	51.1	565.9 ^a	45.4
TTP	1.29 ^c	0.1	3.16 ^b	0.2	1.68 ^c	0.2	4.79 ^a	0.7	—	—	—	—	—	—	—	—
IMP	—	—	—	—	—	—	—	—	—	—	—	—	9.6 ^b	1.9	14.0 ^a	1.8

CDP-choline, cytidine diphosphocholine; D - N, differentiated cells; D + N, differentiated cells grown with nucleosides; P - N, proliferating cells; P + N, proliferating cells grown with nucleosides; UDP-glucose, uridine diphosphoglucose.

† Proliferating and differentiated IEC-6 cells were cultured in the presence or absence of a nucleoside supplement: uridine, cytidine, guanosine and inosine (30 μM each).

^{a-f} Mean values within a row with unlike superscript letters were significantly different ($P < 0.05$).

cells (Table 2). It is worth noting that the total intracellular NT pool was significantly higher in differentiated cells than in proliferating ones (D - N *v.* P - N). A similar result was observed in differentiated Caco-2 cells¹⁴, and it was concluded that differentiation may activate NT biosynthesis in intestinal cells. Analysis of the intracellular NT pool in IEC-6 cells showed that differentiated cells had no detectable intracellular thymidine NT compared to proliferating ones, probably related to the arrest of cell division.

Exogenous NS significantly altered the concentration and distribution of soluble intracellular NT. Incubation of IEC-6 cells with NS produced an increase in intracellular NT concentrations of proliferating (P + N *v.* P - N) and differentiated

(D + N *v.* D - N) cells. This result agrees with the work of many authors in primary cells or cell lines^{13-17,29-32}. We believe that salvage of NS contributed to *de novo* synthesis, thereby increasing the intracellular NT pool. Uptake and salvage of guanosine and inosine significantly increased the intracellular pool of guanosine NT rather than adenosine derivatives. Total guanosine NT, which play an important role in GTPase-mediated cell structure organization and signalling as well as in protein synthesis^{33,34}, increased 2.5-fold in P + N cells and 5-fold in D + N cells with respect to unsupplemented counterparts. The detection of intracellular IMP may indicate that the synthesis of purine NT was saturated in differentiated cells supplemented with exogenous NS (D + N). Indeed, the highest concentrations of intracellular purine NT were reached in these cells. The present data




Fig. 2. Energy charge values in proliferating (P) and differentiated (D) IEC-6 cells cultured with (+N) or without nucleosides (-N) at 12 h (■) and 24 h (▨). Cell media was supplemented with uridine, cytidine, guanosine and inosine (30 μM each). Values are means with their standard errors depicted by vertical bars for three calculations per group. * Significant effect of time, $P < 0.05$.




Fig. 3. Alkaline phosphatase activities in proliferating (P) and differentiated (D) IEC-6 cells cultured with (+N) or without nucleosides (-N). Cell media was supplemented with uridine, cytidine, guanosine and inosine (30 μM each). Values are means with their standard errors depicted by vertical bars for three determinations per group. * Significant effect of nucleoside supplementation, $P < 0.05$. † Significant effect of cell state, $P < 0.05$.




Fig. 4. Effect of nucleoside supplementation on the levels of *Rnd3* mRNA in differentiated IEC-6 cells. Cells were cultured with (○) or without (●) uridine, cytidine, guanosine and inosine (30 mM each). mRNA levels were analysed by Northern blot and normalized to the expression of glyceraldehyde-3-phosphate dehydrogenase. Values are means with their standard errors depicted by vertical bars for three determinations per group. The mean of undifferentiated cells (time 0 h) was set to 1.

showed that intracellular ATP and NAD^+ concentrations are tightly controlled in IEC-6 cells. Although exogenous NS modified NAD^+ and adenosine NT values, the amount of total adenine NT and ATP levels were significantly different only at 12 h in D + N cells compared with D - N cells. These data contrasted with the energy charge values that were significantly higher in D + N than in D - N cells at 24 h (Fig. 2), and suggested that energy was more efficiently stored in NS-supplemented differentiated IEC-6 cells. It is noteworthy that uric acid was not detected in culture media although intestinal cells express xanthine oxidase. The result indicated that purine NS were incorporated into the intracellular NT pool rather than being catabolized to uric acid.

In parallel, we observed that exogenous NS enhanced the differentiation of IEC-6 cells. Morphologically, microvilli were more abundant in D + N cells and persisted longer than in D - N cells (Fig. 1). Moreover, at the enzymatic level, D + N cells had higher alkaline phosphatase activity than D - N cells (Fig. 3). The present results agree with those of Sato *et al.*³⁵, who observed that NS supplementation promoted the formation of tight junctions, the expression of microvilli, and the activities of maltase and sucrose-isomaltase. Concerning gene expression, we used the differential display technique to identify genes regulated during the initial stages of enterocyte differentiation. One of these differentiation-regulated genes was *Rnd3*³⁶, a member of the Rho family of small GTPases involved in cytoskeleton arrangements³⁵. Northern blot analyses showed that *Rnd3* mRNA levels significantly diminished during IEC-6 differentiation. We are aware that, although statistically modest, the present results showed that NS supplementation accelerated the changes in *Rnd3* mRNA levels during enterocyte differentiation (Fig. 4).

In conclusion, exogenous NS were selectively taken up by IEC-6 cells and increased the intracellular NT pool in proliferating and differentiated cells. Because *de novo* synthesis of NT is an energy-intensive process, cells fulfil their NT requirements by NS salvage that spares energy and intermediary metabolites. The increase in energy charge and intracellular GTP concentrations in differentiated cells may favour the turnover of proteins and nucleic acids required for the transition from the proliferative to differentiation state in IEC-6 cells. These changes associated with an enhanced maturation of intestinal cells, improving their differentiation at morphological, functional and gene expression levels. The present findings agree with the work of many authors in cells and

animal models of injury^{2-9,37-39}, and provide new evidence of the beneficial effect of exogenous NS on the functionality of intestinal epithelial cells.

Acknowledgements

This work was financed by Spanish Science Ministry Research Project PM97-0171 and by ABBOTT research project 71 400. Raul de la Torre was the recipient of a Junta de Andalucía pre-doctoral fellowship.

References

- Schlimme E, Martin D & Meisel H (2000) Nucleosides and nucleotides: natural bioactive substances in milk and colostrum. *Br J Nutr* **84**, Suppl. 1, S59–S68.
- Sánchez-Pozo A & Gil A (2002) Nucleotides as semiessential nutritional components. *Br J Nutr* **87**, Suppl. 1, S135–S137.
- Gil A (2002) Modulation of the immune response mediated by dietary nucleotides. *Eur J Clin Nutr* **56**, Suppl. 3, S1–S4.
- Evans ME, Jones DP & Ziegler TR (2005) Glutamine inhibits cytokine-induced apoptosis in human colonic epithelial cells via the pyrimidine pathway. *Am J Physiol Gastrointest Liver Physiol* **289**, G388–G396.
- Nunez MC, Bueno JD, Ayudarte MV, Almendros A, Rios A, Suarez MD & Gil A (1996) Dietary restriction induces biochemical and morphometric changes in the small intestine of nursing piglets. *J Nutr* **126**, 933–944.
- Arnaud A, Lopez-Pedrosa JM, Torres MI & Gil A (2003) Dietary nucleotides modulate mitochondrial function of intestinal mucosa in weanling rats with chronic diarrhea. *J Pediatr Gastroenterol Nutr* **37**, 124–131.
- Mabley JG, Pacher P, Liaudet L, Soriano FG, Hasko G, Marton A, Szabo C & Salzman AL (2003) Inosine reduces inflammation and improves survival in a murine model of colitis. *Am J Physiol Gastrointest Liver Physiol* **284**, G138–G144.
- Dowdall JF, Winter DC & Bouchier-Hayes DJ (2002) Inosine modulates gut barrier dysfunction and end organ damage in a model of ischemia-reperfusion injury. *J Surg Res* **108**, 61–68.
- Belo A, Marchbank T, Fitzgerald A, Ghosh S & Playford RJ (2006) Gastroprotective effects of oral nucleotide administration. *Gut* **55**, 165–171.
- Arnaud A, Fontana L, Angulo AJ, Gil A & Lopez-Pedrosa JM (2003) Exogenous nucleosides alter the intracellular nucleotide pool in hepatic cell cultures. Implications in cell proliferation and function. *Clin Nutr* **22**, 391–399.
- Saez-Lara MJ, Manzano M, Angulo AJ, Suarez A, Torres MI, Gomez-Llorente C, Gil A & Fontana L (2004) Exogenous nucleosides stimulate proliferation of fetal rat hepatocytes. *J Nutr* **134**, 1309–1313.
- Gil A, Gómez-León C & Rueda R (2007) Exogenous nucleic acids and nucleotides are efficiently hydrolysed and taken up as nucleosides by intestinal explants from suckling piglets. *Br J Nutr* **98**, 285–291.
- Sato N, Nakano T, Kawakami H & Idota T (1999) In vitro and in vivo effects of exogenous nucleotides on the proliferation and maturation of intestinal epithelial cells. *J Nutr Sci Vitaminol (Tokyo)* **45**, 107–118.
- He Y, Chu SH & Walker WA (1993) Nucleotide supplements alter proliferation and differentiation of cultured human (Caco-2) and rat (IEC-6) intestinal epithelial cells. *J Nutr* **123**, 1017–1027.
- Boza JJ, Moennoz D, Bournot CE, Blum S, Zbinden I, Finot PA & Ballevre O (2000) Role of glutamine on the *de novo* purine nucleotide synthesis in Caco-2 cells. *Eur J Nutr* **39**, 38–46.

16. Tanaka M, Lee K, Martinez-Augustin O, He Y, Sanderson IR & Walker WA (1996) Exogenous nucleotides alter the proliferation, differentiation and apoptosis of human small intestinal epithelium. *J Nutr* **126**, 424–433.
17. Chen X, Wang R, Deng M & Ran X (2001) Effects of exogenous adenosine 5'-monophosphate on proliferation and apoptosis of small intestinal epithelial cells. *Wei Sheng Yan Jiu* **30**, 165–167.
18. Duchon K & Thorell L (1999) Nucleotide and polyamine levels in colostrum and mature milk in relation to maternal atopy and atopic development in the children. *Acta Paediatr* **88**, 1338–1343.
19. Gehrke C & Kuo K (editors) (1990) *Chromatography and Modification of Nucleosides, Part A, B and C Analytical Methods for Major and Modified Nucleosides HPLC, GC, MS, NMR and FT-IR*. Amsterdam: Elsevier.
20. Perret D (1993) Nucleotides, nucleosides and bases. In *High Performance Liquid Chromatography in Neuroscience Research*, pp. 163–165 [RB Holman, AJ Cross and MH Joseph, editors]. London: Wiley.
21. Goldstein R, Klein T, Freier S & Mencil J (1971) Alkaline phosphatase and disaccharidase activities in the rat intestine from birth to weaning. I. Effect of diet on enzyme development. *Am J Clin Nutr* **24**, 1224–1231.
22. Brown RE, Jarvis KL & Hyland KJ (1989) Protein measurement using bicinchoninic acid: elimination of interfering substances. *Anal Biochem* **180**, 136–139.
23. Aymerich I, Pastor-Anglada M & Casado FJ (2004) Long term endocrine regulation of nucleoside transporters in rat intestinal epithelial cells. *J Gen Physiol* **124**, 505–512.
24. Sterling KM Jr & Cutroneo KR (2002) Differentiation-dependent induction of CYP1A1 in cultured rat small intestinal epithelial cells, colonocytes, and human colon carcinoma cells: basement membrane-mediated apoptosis. *J Cell Biochem* **86**, 440–450.
25. Aymerich I, Duflot S, Fernandez-Veledo S, Guillen-Gomez E, Huber-Ruano I, Casado FJ & Pastor-Anglada M (2005) The concentrative nucleoside transporter family (SLC28): new roles beyond salvage? *Biochem Soc Trans* **33**, 216–219.
26. Aymerich I, Fougelle F, Ferre P, Casado FJ & Pastor-Anglada M (2006) Extracellular adenosine activates AMP-dependent protein kinase (AMPK). *J Cell Sci* **119**, 1612–1621.
27. Larrayoz IM, Fernandez-Nistal A, Garcés A, Gorraitz E & Lostao MP (2006) Characterization of the rat Na⁺/nucleoside cotransporter 2 and transport of nucleoside-derived drugs using electrophysiological methods. *Am J Physiol Cell Physiol* **291**, C1395–C1404.
28. Ohyanagi H, Nishimatsu S, Kanbara Y, Usami M & Saitoh Y (1989) Effects of nucleosides and a nucleotide on DNA and RNA syntheses by the salvage and de novo pathway in primary monolayer cultures of hepatocytes and hepatoma cells. *JPEN J Parenter Enteral Nutr* **13**, 51–58.
29. Mayer D, Natsumeda Y, Ikegami T, Faderan M, Lui M, Emrani J, Reardon M, Olah E & Weber G (1990) Expression of key enzymes of purine and pyrimidine metabolism in a hepatocyte-derived cell line at different phases of the growth cycle. *J Cancer Res Clin Oncol* **116**, 251–258.
30. Moyer JD, Oliver JT & Handschumacher RE (1981) Salvage of circulating pyrimidine nucleosides in the rat. *Cancer Res* **41**, 3010–3017.
31. Pels Rijcken WR, Overdijk B, Van den Eijnden DH & Ferwerda W (1995) The effect of increasing nucleotide-sugar concentrations on the incorporation of sugars into glycoconjugates in rat hepatocytes. *Biochem J* **305**, 865–870.
32. Karle JM, Anderson LW & Cysyk RL (1984) Effect of plasma concentrations of uridine on pyrimidine biosynthesis in cultured L1210 cells. *J Biol Chem* **259**, 67–72.
33. Hallett MA, Dagher PC & Atkinson SJ (2003) Rho GTPases show differential sensitivity to nucleotide triphosphate depletion in a model of ischemic cell injury. *Am J Physiol Cell Physiol* **285**, C129–C138.
34. Hucul JA, Henshaw EC & Young DA (1985) Nucleoside diphosphate regulation of overall rates of protein biosynthesis acting at the level of initiation. *J Biol Chem* **260**, 15585–15591.
35. Sato N, Kawakami H & Idota A (2000) Nucleotide and nucleoside supplementation may morphologically promote the differentiation of human caco-2 cells. *J Nutr Sci Vitaminol (Tokyo)* **46**, 175–179.
36. Vieites JM, Sanchez-Pozo A, Gil A & Suarez A (2005) Development of a protocol for the analysis of genetic expression by differential display. *Ars Pharm* **46**, 193–204.
37. Perez MJ, Suarez A, Gomez-Capilla JA, Sanchez-Medina F & Gil A (2002) Dietary nucleotide supplementation reduces thioacetamide-induced liver fibrosis in rats. *J Nutr* **132**, 652–657.
38. Perez MJ, Sanchez-Medina F, Torres M, Gil A & Suarez A (2004) Dietary nucleotides enhance the liver redox state and protein synthesis in cirrhotic rats. *J Nutr* **134**, 2504–2508.
39. Grimble GK (1994) Dietary nucleotides and gut mucosal defence. *Gut* **35**, S46–S51.