

Factors influencing hydroids (Cnidaria: Hydrozoa) biodiversity and distribution in Arctic kelp forest

MARTA RONOWICZ, MARIA WŁODARSKA-KOWALCZUK AND PIOTR KUKLINSKI

Institute of Oceanology, Polish Academy of Sciences ul. Powstancow Warszawy 55, Sopot 81-712, Poland

*The biodiversity and distribution patterns of epiphytic hydroids were studied in kelp forests (composed of *Laminaria digitata*, *Saccharina latissima* and *Alaria esculenta*) located in an Arctic glaciated fiord (Hornsund, west Spitsbergen). In total, twenty-eight species were found colonizing algae, stones connected to holdfast, and overgrowing the surface of other animals associated with kelps. The characteristics of the algal host (e.g. algae species, age, rhizoid volume or biomass) did not show any effect upon hydroid species richness or species composition. High hydroid biodiversity was strongly dependent on microsubstrate heterogeneity. The highest biodiversity as well as frequency of hydroid occurrence were noted at a site located furthest from the glacier and characterized by the lowest sediment concentration and sedimentation rate. Sexual reproduction also seemed to be inhibited by glacier-derived disturbance. Of ten fertile species found at the 'clearest' site only two were fertile at sites under the strong influence of such perturbations. Potential physical drivers of species occurrence were linked to the activity of tidal glaciers, particularly to high loads of mineral sedimentation and iceberg scouring.*

Keywords: benthic Hydrozoa, Arctic, biodiversity, kelp forest, sedimentation, distribution

Submitted 6 November 2007; accepted 5 February 2008; first published online 22 July 2008

INTRODUCTION

Dense kelp forests can be found in the shallow hard-bottom sublittoral (5–30 m) in Spitsbergen fjords (Lippert *et al.*, 2001). These macroalgae communities consist mostly of large (up to 2–3 m high) kelps: *Laminaria digitata* (Hudson) Lamouroux, 1813, *Saccharina latissima* (Linnaeus) C.E. Lane, C. Mayes, Druehl & G.W. Saunders, 2006, and *Alaria esculenta* (Linnaeus) Greville, 1830. Macroalgae are an important component of ecosystems, not only as primary producers, but also as ecosystem engineers offering a habitat for a number of macrofaunal species. They provide substrate for a large number of sedentary animals (Kuklinski *et al.*, 2006; Carlsen *et al.*, 2007) as well as a shelter for mobile invertebrates (Martin-Smith, 1993).

In polar regions, the development of seaweeds, and therefore of epifaunal assemblages is limited by the very short period of favourable light conditions (the polar night lasts about four months), by icebergs and sea ice scouring the bottom, and particularly by the high loads of fine sediment discharged with glacial or glaciofluvial melt water that restrict light penetration (Wiencke *et al.*, 2007). The patterns of increasing disturbance by high levels of sedimentation in glacial fjords is related to continual glacier retreat (at a rate of up to 0.5 km y⁻¹), which is a consequence of global warming (Svendsen, 1996; Włodarska-Kowalczyk & Weslawski, 2001). High inorganic suspension concentration increases water turbidity, thus influencing light conditions.

The seasonal fresh glacial water inflow disturbs the physical regime of fjord waters (Weslawski *et al.*, 1995) and impacts both macroalgal and macrofaunal communities. The presence of sediments in the water column is regarded as a severe stress agent for hard-bottom macro-organisms, especially the suspension feeders (Airoldi, 2003). It is most likely that hydroids, the main focus of this study, as filter feeders are influenced by these factors.

Studies from other regions indicate that, among sessile organisms, a large part of epifauna is formed by hydrozoans (Nishihira, 1965; Hughes *et al.*, 1991; Llobet *et al.*, 1991; Watson, 1992; Faucci & Boero, 2000; Frascchetti *et al.*, 2002, 2006). However, knowledge of Arctic epifaunal hydrozoans is very scanty, and there are only a few studies describing macroalgae and its epifaunal communities in the polar regions (Różycki & Gruszczynski, 1986; Hansen & Haugen, 1989; Lippert *et al.*, 2001). The hydroids' inconspicuous colonies and the problematic identification of species are probably the most likely reason for the low number of records in the published literature cited above. This study intends to fill in some of the gaps in the knowledge of the ecology of Arctic hydrozoans associated with macroalgae.

Our aim was to describe the hydroid community associated with kelp forest in the specific environment of an Arctic glacial fjord. The objectives of the study were:

- (1) to present an updated species list of hydroids associated with the kelp forest in Hornsund;
- (2) to analyse the relations between hydroid biodiversity and substrate characteristics (species, age, biomass of algae and microsubstrate heterogeneity); and
- (3) to describe patterns of hydroid community composition and diversity in relation to mostly glacial-driven

Corresponding author:

M. Ronowicz

Email: martigor1@o2.pl




Fig. 1. Sampling sites in Hornsund fjord; dark grey, land; light grey, glaciers.

environmental factors (mineral suspension concentration and sedimentation rate).

STUDY AREA

The 30 km-long Hornsund Fjord is the most southerly situated fjord off west Spitsbergen, the largest of about 150 islands of the Svalbard Archipelago located on the edge of the Barents Sea shelf, 74°–81°N 10°–34°E (Figure 1). The fjord is under the influence of two water masses: the relatively warm, saline Atlantic water ($T > 2^{\circ}\text{C}$, $S > 34.7$ psu) of the West Spitsbergen Current and the colder Arctic waters of the Sorkapp Current ($T < 0^{\circ}\text{C}$, $S = 34.3$ – 34.8 psu) (Loeng, 1991).

Fourteen tidal glaciers that enter the fjord waters, together with rivers running from the melting glaciers, strongly modify the hydrology of the Spitsbergen fjords especially in the summer season (Weslawski *et al.*, 1995). Due to the fresh water inflow (about 1.8 km^3 annually in Hornsund; Jania & Pulina, unpublished data, after Weslawski *et al.*, 1995), the temperature and salinity oscillations are high and a gradient of inorganic sedimentation is created (Swerpel, 1985; Görlich *et al.*, 1987). In July in Isbjørnhamna (Hornsund Fjord), over a distance of 3 km the surface salinity may vary from 15 to 31 psu and suspension concentration from 48 to 367 mg dm^{-3} (Görlich *et al.*, 1987).

Three sites at different distances to active glacier outflows were selected in order to compare contrasting sedimentary and hydrology regimes (Table 1; Figure 1):

Isbjørnhamna: situated close to the fjord entrance, close to the Hansbreen glacier outflow, with very high inorganic particle sedimentation;

Hyrneodden: situated in the inner part of the fjord, close to several active tidal glacier outflows, with intermediate level of inorganic particle sedimentation; and

Gåshamna: situated close to the fjord entrance, far from the glacier outflows, with very low inorganic particle sedimentation.

MATERIALS AND METHODS

Sampling and environmental parameters collection

Three hundred and forty samples of algal thalli were collected in Hornsund fjord (Spitsbergen) in July 2003 at the three selected sites (Figure 1; Table 1), with 133 samples from Isbjørnhamna, 113 from Gåshamna and 94 from Hyrneodden.

Specimens of the most common macroalgal species (*Laminaria digitata*, *Saccharina latissima* and *Alaria esculenta*) were carefully collected at each site from 5 and 10 m depths by SCUBA divers. Each algae sample was placed in a separate bag with 1 mm mesh size. In the laboratory, the age of each laminarian algae was determined (after Kain, 1963), and the algae were divided into cauloid, phylloid and rhizoid. The length, wet weight, and volume (measured as the volume of displaced water after immersion in the cylinder) of each part were measured. The associated hydroid fauna was fixed in 4% formalin and was identified under stereomicroscope and microscope. Other species of algae (*Desmarestia* sp., *Phycodryas rubens* (Linnaeus) Batters, 1902, and *Odonthalia* sp.) were collected less frequently in the study area. Due to the small number of samples collected, they were excluded from most of the analysis and comparisons.

The environmental parameters measured at each site included STD profiles, current meter measurements with a Sensordata Bergen SD6000 (deployed in Isbjørnhamna for 48 minutes, Gåshamna for 242 minutes and Hyrneodden for 68 minutes), and the collection of four replicates of water samples for inorganic and organic suspended matter concentrations, as well as the deployment of sediment traps made of PVC cylinders (opening diameter, 8 cm; height, 50 cm). In order to measure the sedimentation rate at each site, three traps were installed on the seabed. After 24 hours they were recovered and taken to the laboratory and left for another day and night, until the sediments had stabilized. Then, the supernatant was decanted, and the remaining mixture of

Table 1. Physical conditions at three sites (mean \pm SD).

Site	Isbjørnhamna		Gåshamna		Hyrneodden	
Depth (m)	5	10	5	10	5	10
Temperature ($^{\circ}\text{C}$)	1.29	1.77	2.45	2.45	1.34	1.72
Salinity (PSU)	32.66	32.21	30.3	30.3	31.59	31.55
Icebergs	numerous		none		numerous	
Sedimentation rate—mineral ($\text{mg dm}^{-3} \text{ d}^{-1}$)	146.53 ± 103.68		12.73 ± 4.75		45.96 ± 11.96	
Sedimentation rate—organic ($\text{mg dm}^{-3} \text{ d}^{-1}$)	20.36 ± 12.24		5.27 ± 2.59		7.18 ± 2.28	
Suspension concentration—mineral (mg dm^{-3})	10.10 ± 1.51		8.02 ± 5.34		9.43 ± 0	
Suspension concentration—organic (mg dm^{-3})	2.83 ± 0.52		2.26 ± 0.41		2.71	
Current velocity (cm s^{-1})	2.4 ± 0.6		1.75 ± 0.5		2.3 ± 1.0	
Secchi disc transparency (m)	1.6 ± 0.8		10 ± 0.3		1.7 ± 0.4	

water and sediment was filtered through the filters (Whatman GF/C, 47 mm). These were then dried at 60°C for 24 hours and weighed. The weighed filters loaded with sediment were placed in an oven (450°C) for 24 hours and then weighed again. The same procedure was used to analyse the organic/inorganic suspension concentrations.

Data analysis

Multivariate analysis (done with the PRIMER package) was performed to identify the patterns of hydroids occurrence. The similarity between samples was calculated with the Bray–Curtis index. Only presence/absence data were used, as it was often not possible to distinguish, and so to enumerate, separate colonies. The differences between species composition among the three sites, three kelp species and two depths were tested with the pair-wise ANOSIM test. The differences

between the sample species richness among the same combinations of sample groups were tested using the non-parametric Kruskal–Wallis test. Pair-wise Mann–Whitney *U*-tests were used for post hoc multiple comparisons.

RESULTS

Species list

A total of 28 taxa of hydroids belonging to two orders and eight families were found associated with kelp forest (Table 2). The order Thecata was represented by 24 species, whereas Athecata was represented by only four species. Among taxa noted during the present study, merely three species (*Cuspidella* sp., *Obelia longissima* and *Sarsia* cf. *loveni*) generate the medusa stage for

Table 2. List of all taxa recorded during the present study. Occurrence at the studied sites, depths and substrates, as well as reproductive mode for each taxon, are also presented.

Species	Site	Depth		Substrate				Sexual reproduction	
		5 m	10 m						
THECATA									
Campanulinidae									
<i>Calycella syringa</i> (Linnaeus, 1758)	I G H	+	+	A	B	H	R	S	G
<i>Campanulina pumila</i> (G.O. Sars, 1874)	I G H	+	+	A	B				G
<i>Cuspidella</i> sp.	I G H	+	+	A	B	H		S	M
Phialellidae									
<i>Opercularella lacerata</i> (Johnston, 1847)	I	+		A					G
Lafoeidae									
<i>Filellum serpens</i> (Hassall, 1848)	I G		+	A	B				G
<i>Grammaria abietina</i> (M. Sars, 1850)	I H	+	+	-					G
<i>Lafoea dumosa</i> (Fleming, 1820)	I G H	+	+	A	B		R	S	G
Haleciidae									
<i>Halecium curvicaule</i> Lorenz, 1886	I G	+	+	A	B	H		S	G
<i>Halecium arcticum</i> Ronowicz & Schuchert, 2007	I G H	+		A	B	H	R	S	G
<i>Halecium minutum</i> Broch, 1903	I	+		-					G
<i>Halecium mirabile</i> Schydlovsky, 1902	I G H	+	+	A	B	H		S	G
<i>Halecium muricatum</i> (Ellis & Solander, 1786)	I	+			B				G
<i>Halecium</i> cf. <i>speciosum</i> Nutting, 1901	I	+		-					G
<i>Halecium textum</i> Kramp, 1911	G	+	+	-					G
Sertulariidae									
<i>Sertularella rugosa</i> (Linnaeus, 1758)	I G	+	+	A	B				G
<i>Sertularella tenella</i> (Alder, 1856)	I		+		B	H			G
<i>Sertularia argentea</i> Linnaeus, 1758	I G H	+	+	A	B			S	G
<i>Sertularia cupressoides</i> Clark, 1876	I G	+	+	-					G
<i>Sertularia schmidti</i> Kudelin, 1914	I	+		-					G
<i>Symplectoscyphus tricuspoidatus</i> (Alder, 1856)	I G	+	+	A	B			S	G
Campanulariidae									
<i>Campanularia integra</i> MacGillivray, 1842	I G		+	A	B			S	G
<i>Campanularia volubilis</i> (Linnaeus, 1758)	I G H	+	+	A	B	H		S	G
<i>Gonothyrea loveni</i> (Allman, 1859)	I G H	+	+	A	B	H		S	Em
<i>Obelia longissima</i> (Pallas, 1766)	I G	+	+	A	B				M
ATHECATA									
Eudendriidae									
<i>Eudendrium annulatum</i> Norman, 1864	I G	+	+	A					G
<i>Eudendrium</i> cf. <i>insigne</i> Hincks, 1861	G	+		-					G
<i>Eudendrium rameum</i> (Pallas, 1766)	H		+				R		G
Corynidae									
<i>Sarsia</i> cf. <i>loveni</i> (M. Sars, 1846)	I	+		-					M

Site: I, Isbjørnhamna; G, Gåshamna; H, Hyrneodden; Substrate: A, algae; B, Bryozoa; H, Hydrozoa; R, rock; S, shell; Sexual reproduction: g, fixed gonophores; m, medusae; em, eumedusoids.




Fig. 2. Numbers of hydroid species (mean \pm 0.95 CI) per sample recorded on different algae (L.dig., *Laminaria digitata*; S.lat., *Saccharina latissima*; A.esc., *Alaria esculenta*; P.rub., *Phycodrys rubens*).

sexual reproduction and one species (*Gonothyrea loveni*) creates eumedusoids (Table 2).

Primary substrate characteristics

Seventeen hydroids taxa grew directly on the algae. The remaining 11 species colonized other types of substrata connected with the algae (Table 2), or were found unattached to any substrate.

The frequency of hydroid occurrence was very similar regardless of laminarian species, ranging from 21% on *S. latissima* to 27% on *L. digitata*.

Each laminarian species hosted an average of 2.5 (*A. esculenta*) to 3.5 (*L. digitata* and *P. rubens*) hydroid species (Figure 2). The highest hydroid species richness per sample (= per algae specimen) was found on *L. digitata* (8 species). Altogether ten species of hydroids were found on this algae species. Maximum species richness per sample was slightly lower on *S. latissima* (7 species) and *A. esculenta* (6 species), but the total number of hydroids found on these two kelp species was higher than on *L. digitata* (14 and 13 species, respectively).

No differences were noted in species richness per sample among the sampled algal species (Kruskal–Wallis test, $P > 0.05$).

Table 3. One-way ANOSIM pairwise analysis testing differences among species composition of kelp species, sites and depths.

ANOSIM	R	P (%)
Algae		
<i>Laminaria digitata</i> – <i>S. latissima</i>	0.043	10.5
<i>L. digitata</i> – <i>Alaria esculenta</i>	–0.046	93.9
<i>Saccharina latissima</i> – <i>A. esculenta</i>	0.004	31.1
Sites		
Isbjørnhamna–Hyrneodden	0.268	0.1
Gåshamna–Hyrneodden	0.318	0.1
Isbjørnhamna–Gåshamna	0.07	0.1
Depth		
5–10 m	0.046	0.4

Pairwise analysis of similarities (ANOSIM) did not show any differences in hydroid assemblages between algal host species ($P > 0.05$; Table 3).

The hydroid distribution among the three parts of laminarian thalli differed markedly. A significantly higher number of taxa settled on the rhizoid part of the algae, but there were just two species (*Eudendrium annulatum* and *Opercularella lacerata*) present exclusively on the rhizoid part. Only a few species were present on parts other than the basal (phylloids: *Symplectoscyphus tricuspidatus*, *Calycella syringa*, *Eudendrium* indet. and *Eudendrium* cf. *rameum*; cauloids: *Sertularella rugosa* and *Campanularia integra*).

Hydroid biodiversity was significantly, but very weakly, correlated to the total wet weight of algae ($r = 0.26$, $P = 0.01$). No relationship (at $P > 0.05$) was noted concerning hydroid diversity and the characteristics of the algae, such as age, total length, total volume, rhizoid wet weight, or rhizoid volume.

The algae species *Desmarestia* sp., *Phycodrys rubens* and *Odonthalia* sp. were also colonized by hydroids, with 88% frequency of occurrence on *P. rubens*, 50% frequency of occurrence on *Odonthalia* sp., and 18% frequency of occurrence on *Desmarestia* sp.

Microsubstrate heterogeneity

Hydroids colonized many kinds of microhabitats associated with laminarians. Laminarian thalli often hosted bryozoans or hydroids that created an additional substrate for other hydroids. Bivalves, *Balanus* sp., ascidians, poriferans, and rocks were found connected to algal holdfast, and served as extra ecological niches for epiphytic assemblage. Sample species richness was significantly correlated with the number of microhabitats available ($r^2 = 0.6$, $r = 0.8$, $P < 0.05$).

Three species (*Halecium muricatum*, *Sertularella tenella* and *Eudendrium rameum*) were not noted on any part of algal substrate, but exclusively on secondary substrate attached to seaweed (Table 2). Bryozoans were the preferred secondary substrate for hydroid settlement (Figure 3). As many as 17 species overgrew bryozoans, while hydrozoans and bivalves were also commonly colonized by eight species each (Figure 3). At least six species (of the families Eudendriidae, Haleciidae, Sertulariidae and Lafoeidae) offered a habitat for other hydroid settlement. Ascidians, *Balanus* sp., poriferans, and rocks were occupied by a lower number of hydroid species.




Fig. 3. Total number of hydroid species recorded on different types of kelp associated microsubstrates.

Table 4. Analysis of variance and post hoc analysis testing differences between hydroids species richness at sites.

ANOVA and post hoc test	ANOVA Kruskal–Wallis test <i>P</i> (%)	Mann–Whitney <i>U</i> -test <i>P</i> (%)
Sites		
Isbjørnhamna–Hyrneodden	$P < 0.05$	$P > 0.05$
Gåshamna–Hyrneodden	$P < 0.05$	$P < 0.05$
Isbjørnhamna–Gåshamna	$P < 0.05$	$P < 0.05$
Depth		
5–10 m	$P > 0.05$	–

Environmental factors

Neither sample species richness nor species composition were related to depth, as shown by the Kruskal–Wallis test ($P > 0.05$; Table 4) and the pair-wise ANOSIM test, ($R = 0.046$, $P < 0.05$; Table 3), respectively.

The highest mean sample species richness was in Gåshamna (Figure 4A). This was also the site with the highest frequency of hydroid occurrence and of the most common species (Figure 5). Hydroids were noted in 79% of samples taken at Gåshamna, in 63% of samples taken at Isbjørnhamna, and in 28% samples collected at Hyrneodden.

There were significant differences in hydroid species richness among the studied sites (Kruskal–Wallis test, $P < 0.05$; Table 4). A post hoc pair-wise test revealed significant differences between two pairs of sites: Gåshamna–Isbjørnhamna, and Gåshamna–Hyrneodden (Mann–Whitney *U*-test, $P < 0.05$; Table 4), whereas Isbjørnhamna and Hyrneodden did not differ significantly (Mann–Whitney *U*-test, $P > 0.05$).

The low values of *R* statistic at $P < 0.05$ indicated that the hydroid species composition at three sites did not differ notably (pair-wise ANOSIM test; Table 3), nor was any clear separation of samples coming from the three sites observed on the MDS plot (Figure 6).

Ten species were found in the reproductive stage (as indicated by the presence of gonothecae or gonophores) at the time of the study (Figure 7). All of them were noted at Gåshamna. Furthermore the highest mean sample fertile species richness was in Gåshamna (Figure 4B). Only two fertile species were found at Isbjørnhamna, and one species at Hyrneodden (Figure 7).

There were statistical differences between the number of fertile species found at Gåshamna–Isbjørnhamna, and Gåshamna–Hyrneodden (Kruskal–Wallis test, $P < 0.05$; post hoc Mann–Whitney *U*-test, $P < 0.05$), but again there were no significant differences between Isbjørnhamna and Hyrneodden (Mann–Whitney *U*-test, $P > 0.05$).


DISCUSSION

In the extensive material collected during the present study, 17 species were found growing on the algae. That represents less than 15% of all the species recorded to date in Svalbard waters (117 species are reported by Palerud *et al.*, 2004; Ronowicz, 2007 and Ronowicz & Schuchert, 2007). Other studies of Arctic seaweeds epifauna reported lower species diversity in this specific habitat. Lippert *et al.* (2001) found seven taxa of hydroids on macroalgae collected in Kongsfjord (north-west Spitsbergen), while Różycki & Gruszczyński (1986) recorded only one hydroid species associated with laminarians collected in west Spitsbergen. The studies conducted in lower latitudes reported a similarly low number of species (e.g. 11 species of hydroids were found colonizing *Laminaria digitata* and *L. hyperborea* at Helgoland (Schultze *et al.*, 1990)). However, sampling efforts have never been as comprehensive as during the present study.

In the Mediterranean Sea, 32 epiphytic hydrozoan species were identified on brown algae *Cystoseira amentacea* at a high sampling effort (Frashetti *et al.*, 2006), but they comprise only 7% of the taxa recorded in the Mediterranean area.

Primary substrate characteristics

The pattern of hydroid distribution along the kelp thalli is very pronounced. The most attractive part of the algae for settlement is the holdfast. The basal part of kelp provides protection from disruptive wave action and other disturbances (e.g. iceberg scouring in near-glacial areas). In contrast to rhizoids, phylloids grow seasonally and shed in winter and so they offer a short-lived and unstable substrate (Schultze *et al.*, 1990). A similar pattern of strong preferences for rhizoids by epifauna was observed for example, on *Cystoseira amentacea* (Frashetti *et al.*, 2006) and *Laminaria* species (Schultze *et al.*, 1990; Schmidt & Scheibling, 2006).

**Fig. 4.** Mean (± 0.95 CI) number of species at the three sites; A, all species; B, fertile species (I, Isbjørnhamna; G, Gåshamna; H, Hyrneodden).




Fig. 5. Frequency (%) of hydroid occurrence at sites (I, Isbjørnhamna; G, Gåshamna; H, Hyrneodden).

Biodiversity (number of species) seems not to depend on algal morphology (length and volume of thallus, wet weight) or age. According to studies by Kuklinski *et al.* (2006), on bryozoans the size (surface area) and volume of algal substrate do not correlate with the number of species observed either. The common relation of an increase in epibenthic diversity and abundance with the substrate's size may occur in Hornsund kelp forests for whole benthic communities, but it does not have to be expressed in particular taxa (e.g. Hydrozoa). Different major taxa respond differently to physical factors and biological interactions, and so their distributional patterns may contrast with the patterns observed for the whole benthic community (as shown, for example, by Anderson *et al.*, 2005; Wlodarska-Kowalczyk & Kedra, 2007).

Microsubstrate heterogeneity

The high heterogeneity of microsubstrate increases the hydroid species richness. This relationship has been often recorded both for sessile and mobile fauna: the more heterogeneous habitat the more species it hosts (Abele, 1974; Kuklinski *et al.*, 2006).




Fig. 6. nMDS plot of Bray-Curtis similarities of species composition in samples based on presence/absence data (I, Isbjørnhamna; G, Gåshamna; H, Hyrneodden).

All the noted hydroids are known to be substrate generalists and occur on a variety of substrates without any specificity (Cornelius, 1995a,b; Schuchert, 2001). It may be suggested that the availability of habitat is the main force of settlement. There are studies indicating that competition for space is important in shaping Arctic marine assemblages (Barnes & Kuklinski, 2003). The secondary substrates become important for sessile organisms because they provide a substitute home for inferior competitors. Hydroids are among the first colonizers of vacant substrates, but they often cannot withstand pressure from other colonizers, such as bryozoans, sponges and ascidians (Llobet *et al.*, 1991).

The high species richness occurring on bryozoans, hydroids and bivalves may be a consequence of a lower competition level on these types of substrates. The other reason for favouring bryozoans and hydroids may possibly be the fact that mobile, flexible substrates have an advantage over stable microhabitats by providing protection against burying by sediment and supplying better water flow conditions for suspension feeders, thus better food availability and more efficient waste evacuation (Wahl, 1989; Kuklinski & Bader, 2007).

Physical factors

A total of 83% of species collected in Hornsund belong to Thecata, an order grouping hydroids known to be more resistant to mechanical stress than the Athecata representatives (17%) (Faucci & Boero, 2000). The shallow Arctic sublittoral is an extreme environment for benthic organisms due to heavy icebergs scouring in addition to glacier-derived sedimentation.

Both biodiversity and frequency of occurrence were much lower at the two sites situated in the vicinity of glaciers than at the site not influenced by glacier disturbance.

The distribution pattern of sessile invertebrates is formed by the substrate selection processes occurring during larval settlement (Hayward, 1980). In an environment disturbed by strong sedimentation, the loss of suitable substrata for colonization may inhibit larval recruitment (Irving & Connel, 2002; Airoldi, 2003). Several suspension feeding organisms have been observed to experience reduced survival and mortality as a consequence of burial, scouring, and clogging by




Fig. 7. Frequency of occurrence (%) of fertile specimens at sites.

sediments (Moore, 1977; Smith & Witman, 1999), causing changes in the species composition and diversity of already established communities. The most vulnerable species are excluded, and sediment tolerant species take their place. Naturally, there are many covarying factors that interact together with each other (e.g. sedimentation, hydrodynamics, salinity, turbidity and current velocity). Therefore, it is not clear whether the sedimentation regime is the only factor responsible for lowering hydroid species richness at Isbjørnhamna and Hyrneodden.

On the other hand, species composition is homogeneous at the study area with high small-scale variability. This patchiness might be a consequence of microsubstrate heterogeneity, which play an important role in colonization processes (Ronowicz, 2005).

In the collected material all reproducing species produce fixed gonophores. The loss of the planktonic dispersal stage (as in the case of most of the hydroids cited in the present study) has many advantages. It enables to keep subsequent generations in the vicinity of a favourable habitat, thus reducing the risk of spreading to an environment where the settlement of planulae would be impossible.

An interesting observation is that, at sites influenced by glacier disturbances, the sexual reproduction of hydroids seems to be limited or even stopped. Of ten fertile species found at Gåshamna, representing the 'clearest' site without glacier-induced disturbance, all were noted at other sites, but only two of them were fertile at Isbjørnhamna, and one at Hyrneodden sites which are under the strong influence of such perturbations. Investigations on the impact of environmental stress, such as decreasing temperature and salinity, on sexual reproduction of hydroids have shown a completely different reaction as revealed in the acceleration of gonophore formation (Piraino, 1991). The induction of reproductive processes under unfavourable physical conditions was explained as an adaptive response (Stebbing, 1980; Piraino, 1991). In addition, experiments on the competitive ability of *Hydractinia* [GM] revealed that competition as a disturbing force, also causes an increase in the production of gonozooids (Ferrell, 2004). For organisms lacking the medusa stage, the advantage of sexual over asexual reproduction is in the

better adaptation of new genotypes to unfavourable conditions. However, there are also other studies showing no such response of hydroid colonies in the face of inflicted stress, for instance, the exposure of hydroid colonies to low concentration of copper and mercury, as well as poor conditions of water circulation, does not influence gonozooid production; it neither stops production, nor stimulates it (Piraino, 1991).

Since the hydroids do not reproduce at heavily disturbed sites, the impact of sedimentation may exceed the limit of their adaptability. Another possibility is that the reproduction period is simply shifted at sites with disturbed conditions.

The effect of different levels of mineral suspension concentrations on the growth and reproduction of suspension-feeders has not been experimentally tested yet. Such an experiment is desirable in order to fully understand and explain the environmental pattern.

ACKNOWLEDGEMENTS

We wish to thank Dr Peter Schuchert for kind assistance during the identification of problematic specimens. The authors also acknowledge Dr Marek Zajackowski's help with some physical environmental measurements. This project was supported by the State Committee for Scientific Research grant N304 034 31/1202.

REFERENCES

- Abele L.G. (1974) Species diversity of decapod crustaceans in marine habitats. *Ecology* 55, 156–161.
- Airoldi L. (2003) The effect of sedimentation on rocky coast assemblages. *Oceanography and Marine Biology. Annual Review* 41, 161–236.
- Anderson M.J., Diebel C.E., Blom W.M. and Landers T.J. (2005) Consistency and variation in kelp holdfast assemblages: spatial patterns of biodiversity for the major phyla at different taxonomic resolutions. *Journal of Experimental Marine Biology and Ecology* 320, 35–56.

- Barnes D.K.A. and Kuklinski P.** (2003) High polar spatial competition: extreme hierarchies at extreme latitude. *Marine Ecology Progress Series* 259, 17–28.
- Carlsen B.P., Johnsen G., Berge J. and Kuklinski P.** (2007) Biodiversity patterns of macro-epifauna on different lamina parts of *Laminaria digitata* and *Saccharina latissima* collected during spring and summer 2004 in Kongsfjorden, Svalbard. *Polar Biology* 30, 939–943.
- Cornelius P.F.S.** (1995a) North-west European thecate hydroids and their medusae. Part 1. *Synopses of the British Fauna (New Series)* 50, 1–347.
- Cornelius P.F.S.** (1995b) North-west European thecate hydroids and their medusae. Part 2. *Synopses of the British Fauna (New Series)* 50, 1–347.
- Fauci A. and Boero F.** (2000) Structure of an epiphytic hydroid community on *Cystoseira* at two sites of different wave exposure. *Scientia Marina* 64, 255–264.
- Ferrell D.L.** (2004) Fitness consequences of allorecognition-mediated agonistic interactions in the colonial hydroid *Hydractinia [GM]*. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 206, 173–187.
- Fraschetti S., Giangrande A., Terlizzi A., Miglietta M.P., DellaTommasa L. and Boero F.** (2002) Spatio-temporal variation of hydroids and polychaetes associated with *Cystoseira amentacea* (Fucales: Phaeophyceae). *Marine Biology* 140, 949–957.
- Fraschetti S., Terlizzi A., Bevilacqua S. and Boero F.** (2006) The distribution of hydroids (Cnidaria, Hydrozoa) from micro- to macro-scale: spatial patterns on habitat-forming algae. *Journal of Experimental Marine Biology and Ecology* 339, 148–158.
- Görlich K., Weslawski J.M. and Zajaczkowski M.** (1987) Suspension settling effect on macrobenthos biomass distribution in the Hornsund fjord, Spitsbergen. *Polar Research* 5, 175–192.
- Hansen J. and Haugen I.** (1989) Some observations of intertidal communities on Spitsbergen (79°N), Norwegian Arctic. *Polar Research* 7, 23–27.
- Hayward P.J.** (1980) Invertebrate epiphytes of coastal marine algae. In Price J.H. et al. (eds) *The shore environment, 2: ecosystems*. London and New York: Academic Press, pp. 761–787.
- Hughes R.G., Johnson S. and Smith I.D.** (1991) The growth patterns of some hydroids that are obligate epiphytes of seagrass leaves. *Hydrobiologia* 216/217, 205–210.
- Irving A.D. and Connel S.D.** (2002) Sedimentation and light penetration interact to maintain heterogeneity of subtidal habitats: algal versus invertebrate dominated assemblages. *Marine Ecology Progress Series* 245, 83–91.
- Kain J.M.** (1963) Aspects of the biology of *Laminaria hyperborea*. II. Age, weight and length. *Journal of the Marine Biological Association of the United Kingdom* 43, 129–151.
- Kuklinski P. and Bader B.** (2007) Diversity, structure and interactions of encrusting lithophilic macrofaunal assemblages from Belgica Bank, East Greenland. *Polar Biology* 30, 709–717.
- Kuklinski P., Gulliksen B., Lønne O.J. and Weslawski J.M.** (2006) Substratum as a structuring influence on assemblages of Arctic bryozoans. *Polar Biology* 29, 652–661.
- Lippert H., Iken K., Rachor E. and Wiencke C.** (2001) Macrofauna associated with macroalgae in the Kongsfjord (Spitsbergen). *Polar Biology* 24, 512–522.
- Llobet I., Gili J.M. and Hughes R.G.** (1991) Horizontal, vertical and seasonal distributions of epiphytic hydrozoa on the alga *Halimeda tuna* in the Northwestern Mediterranean Sea. *Marine Biology* 110, 151–159.
- Loeng H.** (1991) Features of the physical oceanographic conditions of the Barents Sea. In *Proceedings of the Pro Mare Symposium on Polar Marine Ecology, Trondheim, 12–16 May 1990*. *Polar Research* 10, 5–18.
- Martin-Smith K.M.** (1993) Abundance of mobile epifauna: the role of habitat complexity and predation by fishes. *Journal of the Experimental Biology and Ecology* 174, 243–260.
- Moore P.G.** (1977) Inorganic particulate suspensions in the sea and their effects on marine animals. *Oceanography and Marine Biology. Annual Review* 15, 225–363.
- Nishihira M.** (1965) The association between Hydrozoa and their attachment substrata with special reference to algal substrata. *Bulletin of the Marine Biological Station of Asamushi* 12, 75–92.
- Palerud R., Gulliksen B., Brattegard T., Sneli J.-A. and Vader W.** (2004) The marine macro-organisms in Svalbard waters. In Prestrud P. et al. (eds) *A catalogue of the terrestrial and marine animals of Svalbard*. Tromsø: Norwegian Polar Institute, pp. 5–56.
- Piraino S.** (1991) The adaptive pattern of growth and reproduction of the colonial hydroid *Clavopsella michaeli*. *Hydrobiologia* 216/217, 229–234.
- Ronowicz M.** (2005) Species diversity of Arctic gravel beach: case study for species poor habitats. *Polish Polar Research* 26, 287–297.
- Ronowicz M.** (2007) Benthic hydroids (Cnidaria: Hydrozoa) from Svalbard waters—biodiversity and distribution. *Journal of the Marine Biological Association of the United Kingdom* 87, 1089–1094.
- Ronowicz M. and Schuchert P.** (2007) *Halecium arcticum*, a new hydroid from Spitsbergen (Cnidaria: Hydrozoa). *Zootaxa* 1549, 55–62.
- Różycki O. and Gruszczynski M.** (1986) Macrofauna associated with laminarians in the coastal waters of West Spitsbergen. *Polish Polar Research* 7, 337–351.
- Schmidt A.L. and Scheibling R.E.** (2006) A comparison of epifauna and epiphytes on native kelps (*Laminaria* species) and an invasive alga (*Codium fragile* ssp. *tomentosoides*) in Nova Scotia, Canada. *Botanica Marina* 49, 315–330.
- Schuchert P.** (2001) Survey of the family Corynidae (Cnidaria, Hydrozoa). *Revue Suisse de Zoologie* 108, 739–878.
- Schultze K., Janke K., Krüß A. and Weidemann W.** (1990) The macrofauna and macroflora associated with *Laminaria digitata* and *L. hyperborea* at the island of Helgoland (German Bight, North Sea). *Helgoländer Meeresunters* 44, 39–51.
- Smith F. and Witman J.D.** (1999) Species diversity in subtidal landscapes: maintenance by physical processes and larval recruitment. *Ecology* 80, 51–69.
- Stebbing A.R.D.** (1980) Increase in gonozooids frequency as an adaptive response to stress in *Campanularia flexuosa*. In Tardent P. and Tardent R. (eds) *Proceedings of the 4th International Coelenterate Conference held in Interlaken, Switzerland, 4–8 September, 1979. Developmental and Cellular Biology of Coelenterates*. Amsterdam and other cities: Elsevier: North-Holland Biomedical Press, pp. 27–32.
- Svendsen J.I., Elverhoi A. and Mangerud J.** (1996) The retreat of the Barents Sea ice sheet on the western Svalbard margin. *Boreas* 25, 244–256.
- Swerpel S.** (1985) The Hornsund Fjord: water masses. *Polish Polar Research* 6, 475–496.
- Wahl M.** (1989) Marine epibiosis. I. Fouling and antifouling: some basic aspects. *Marine Ecology Progress Series* 58, 175–189.
- Watson J.E.** (1992) The hydroid community of *Amphibolis* seagrasses in south-eastern and south-western Australia. *Scientia Marina* 56, 217–227.
- Weslawski J.M., Koszteyn J., Zajaczkowski M., Wiktor J. and Kwaśniewski S.** (1995) Fresh water in Svalbard fjord ecosystems. In Skjoldal H.R. et al. (eds) *Ecology of fjords and coastal waters*. Amsterdam: Elsevier Sci BV, pp. 229–241.

Wiencke C., Clayton M.N., Gómez I., Iken K., Lüder U.H., Amsler C.D., Karsten U., Hanelt D., Bischof K. and Dunton K. (2007) Life strategy, ecophysiology and ecology of seaweeds in polar waters. *Reviews in Environmental Science and Biotechnology* 6, 95–126.

Wlodarska-Kowalczyk M. and Kedra M. (2007) Surrogacy in natural patterns of benthic distribution and diversity: selected taxa versus lower taxonomic resolution. *Marine Ecology Progress Series* 351, 53–63.

and

Wlodarska-Kowalczyk M. and Weslawski J.M. (2001) Impact of climate warming on Arctic benthic biodiversity: a case study of two Arctic glacial bays. *Climate Research* 18, 127–132.

Correspondence should be addressed to:

Marta Ronowicz
Institute of Oceanology
Polish Academy of Sciences ul. Powstancow Warszawy 55
Sopot 81-712, Poland
email: martigor1@o2.pl