

Density of Chromatic Roots in Minor-Closed Graph Families

THOMAS J. PERRETT and CARSTEN THOMASSEN[†]

Department of Applied Mathematics and Computer Science, Technical University of Denmark, DK-2800 Lyngby, Denmark
(e-mail: perrettt02@hotmail.com)

Received 24 March 2016; revised 19 February 2018; first published online 22 April 2018

We prove that the roots of the chromatic polynomials of planar graphs are dense in the interval between $32/27$ and 4, except possibly in a small interval around $\tau + 2$ where τ is the golden ratio. This interval arises due to a classical result of Tutte, which states that the chromatic polynomial of every planar graph takes a positive value at $\tau + 2$. Our results lead us to conjecture that $\tau + 2$ is the only such number less than 4.

2010 *Mathematics subject classification*: Primary 05C31
Secondary 05C15, 05C83

1. Introduction

The chromatic polynomial was introduced in 1912 by Birkhoff [1] in order to study the 4-colour problem which says that 4 is not a root of the chromatic polynomial of any planar graph. Since then, chromatic polynomials and their roots, which we call *chromatic roots*, have also become of interest in statistical physics; see *e.g.* [7], where complex chromatic roots are studied. In the present paper we deal exclusively with real chromatic roots. If \mathcal{G} is a class of graphs, then we let $R(\mathcal{G})$ denote the set of real chromatic roots of the graphs in \mathcal{G} .

It is easy to see that 0 and 1 are the smallest chromatic roots. Jackson [3] proved the surprising result that every other chromatic root is strictly greater than $32/27$ and that this bound is sharp. Thomassen [8] proved that chromatic roots are dense in the interval from $32/27$ to infinity, so that the chromatic roots of all graphs are understood in the sense that if \mathcal{G} denotes the class of all graphs, then $R(\mathcal{G}) = \{0, 1\} \cup [32/27, \infty)$. Here, \bar{A} denotes the closure of a set A .

By contrast, the situation for planar graphs is much less clear. Birkhoff and Lewis [2] proved that all chromatic roots of planar graphs are less than 5, and conjectured that the same should hold with 5 replaced by 4. On the other hand, Thomassen [8] proved that the chromatic roots of

[†] Research supported by ERC Advanced Grant GRACOL, project number 320812.

planar graphs are dense in the interval from $32/27$ to 3 . Thus, for planar graphs, the remaining interval from 3 to 4 is of interest.

Conjecture 1.1 ([8]). *If \mathcal{G} denotes the family of planar graphs, then $[3, 4] \subseteq \overline{R(\mathcal{G})}$.*

Recently, Royle [5] proved that real chromatic roots of planar triangulations can be found in any interval $(4 - \varepsilon, 4)$ for $\varepsilon > 0$. In the present paper we prove that the chromatic roots of planar graphs are dense in most of the interval $[3, 4]$, almost settling Conjecture 1.1.

Theorem 1.2. *The set of chromatic roots of planar graphs contains a dense subset of $(3, t_1) \cup (t_2, 4)$, where t_1 and t_2 are constants with $t_1 \approx 3.618032$ and $t_2 \approx 3.618356$.*

To prove Theorem 1.2, we develop a technique which relates the set $R(\mathcal{G})$ to the numbers at which the chromatic polynomial of some graph in \mathcal{G} is negative. The unsolved interval (t_1, t_2) in Theorem 1.2 arises because our technique breaks down in connection with the following fascinating result of Tutte.

Theorem 1.3 ([9]). *If G is a planar triangulation, then $P(G, \tau + 2) > 0$, where τ is the golden ratio.*

Indeed, note that $\tau + 2 \approx 3.618033$ which falls within the interval (t_1, t_2) . Thus it is possible that Tutte's result [9] can be extended to a small neighbourhood around $\tau + 2$. However, we do not believe this to be the case. In fact, the intermediate results in this paper lead us to a conjecture that $\tau + 2$ is the only number in the interval $(3, 4)$ at which the chromatic polynomials of planar graphs are always positive.

Finally, we remark that one can ask similar questions about the set $R(\mathcal{G})$ for other classes of graphs \mathcal{G} . In [4], the techniques of this paper are used to prove results regarding the classes of $K_{3,3}$ -minor free graphs, K_5 -minor free graphs, and graphs on surfaces. Indeed, when \mathcal{G} denotes the $K_{3,3}$ -minor free graphs, we provide a complete characterization of $R(\mathcal{G})$ unless there are counterexamples to the Birkhoff and Lewis conjecture.

2. Chromatic roots of minor-closed graph families

Let \mathcal{G} be a possibly infinite family of finite graphs. Recall that $R(\mathcal{G})$ denotes the set of all real chromatic roots of graphs in \mathcal{G} . Similarly, we denote by $R_o(\mathcal{G})$ the set of all chromatic roots of odd multiplicity of graphs in \mathcal{G} . Finally, we denote by $N(\mathcal{G})$ the set of all real numbers q such that the chromatic polynomial of some graph in \mathcal{G} is negative at q . It is easy to show that these sets are related.

Lemma 2.1. *If \mathcal{G} is a class of graphs, then $R_o(\mathcal{G}) \subseteq \overline{N(\mathcal{G})}$.*

Proof. Let q_0 be a real number such that $q_0 \in R_o(\mathcal{G})$. Thus, for some graph G in \mathcal{G} , the chromatic polynomial $P(G, q)$ is of the form $(q - q_0)^r S(q)$ where r is an odd natural number and S is a polynomial such that $S(q_0) \neq 0$. By continuity, $S(q)$ is non-zero in an interval $(q_0 - \varepsilon, q_0 + \varepsilon)$ for

some $\varepsilon > 0$. Now, for any ε' such that $0 < \varepsilon' < \varepsilon$, it follows that precisely one of $P(G, q_0 - \varepsilon')$ and $P(G, q_0 + \varepsilon')$ is negative. Thus, q_0 is in the closure of $N(\mathcal{G})$. \square

For two graphs G and H , and an integer $k \geq 2$, a k -clique sum of G and H is any graph formed by identifying a clique of size k in G with a clique of size k in H . Thomassen [8] showed that in certain circumstances, a construction based on a 2-clique sum can produce a graph with a chromatic root in a given interval.

Proposition 2.2 ([8]). *Let q_0 and δ be real numbers with $q_0 > 2$ and $\delta > 0$. Let G be a graph, and $e \in E(G)$. Let $H'_{k,\ell}$ be the graph formed by a 2-clique sum of k copies of G , and ℓ copies of C_4 , where the edge identified is e . Finally, let $H_{k,\ell}$ be the graph formed from $H'_{k,\ell}$ by deleting the edge corresponding to e . If $|P(G, q_0)| < (q_0 - 1)|P(G/e, q_0)|$ and $P(G, q_0)P(G/e, q_0) < 0$, then there exist $k, \ell \in \mathbb{N}$, and a real number $q_1 \in (q_0 - \delta, q_0 + \delta)$ such that $q_1 \in R_o(\{H_{k,\ell}\})$.*

We say that \mathcal{G} is closed under taking k -clique sums if all k -clique sums of graphs $G, H \in \mathcal{G}$ are members of \mathcal{G} . Under certain conditions, we can now prove a stronger relationship between the closures of the sets $R_o(\mathcal{G})$ and $N(\mathcal{G})$.

Theorem 2.3. *Let \mathcal{G} be a class of graphs such that $C_4 \in \mathcal{G}$. If \mathcal{G} is closed under edge deletion and taking 2-clique sums, then $\overline{R_o(\mathcal{G})} \cap [2, \infty) = \overline{N(\mathcal{G})} \cap [2, \infty)$.*

Proof of Theorem 2.3. By Lemma 2.1, it suffices to show that $N(\mathcal{G}) \cap [2, \infty) \subseteq \overline{R_o(\mathcal{G})}$, so suppose that $q_0 \in N(\mathcal{G}) \cap [2, \infty)$. Thus, there is a graph $G \in \mathcal{G}$, such that $P(G, q_0) < 0$. This implies that $q_0 \neq 2$, because the chromatic polynomial of any graph evaluated at a positive integer is non-negative. Assume that G is edge-minimal with this property, so for each edge e we have $P(G - e, q_0) \geq 0$. Note that G is not edgeless, since the chromatic polynomial of a graph with no edges is positive at any $q_0 > 0$. By the deletion–contraction formula, we have

$$P(G - e, q_0) = P(G, q_0) + P(G/e, q_0).$$

Hence $P(G/e, q_0) \geq -P(G, q_0)$ and $P(G/e, q_0) > 0$, and so G and q_0 satisfy the conditions in Proposition 2.2, which implies that for all $\delta > 0$, there is a graph H such that $P(H, q)$ has a root of odd multiplicity in $(q_0 - \delta, q_0 + \delta)$. Furthermore, $H \in \mathcal{G}$ since $C_4 \in \mathcal{G}$ and \mathcal{G} is closed under edge deletion and taking 2-clique sums. Thus $q_0 \in \overline{R_o(\mathcal{G})}$. \square

Let \mathcal{G} be a minor-closed class of graphs, and denote by $\text{Forb}(\mathcal{G})$ the minor-minimal graphs not in \mathcal{G} . We note the following simple fact.

Proposition 2.4. *If \mathcal{G} is a minor-closed class of graphs such that all graphs in $\text{Forb}(\mathcal{G})$ are 3-connected, then \mathcal{G} is closed under taking 2-clique sums.*

Proof. Let G be a graph formed by a 2-clique sum of graphs $G_1, G_2 \in \mathcal{G}$. If G contains a 3-connected minor F , then F must be a minor of G_1 or G_2 , a contradiction. Hence $G \in \mathcal{G}$ as required. \square

Since minor-closed classes of graphs are clearly closed under edge-deletion, we deduce the following from Theorem 2.3 and Proposition 2.4.

Theorem 2.5. *Let \mathcal{G} be a minor-closed class of graphs such that $C_4 \in \mathcal{G}$. If all graphs in $\text{Forb}(\mathcal{G})$ are 3-connected, then $\overline{R_0(\mathcal{G})} \cap [2, \infty) = \overline{N(\mathcal{G})} \cap [2, \infty)$.*

We note one final relationship, which is a simple consequence of the addition–contraction formula. If \mathcal{G} is a class of graphs, let \mathcal{G}_{\max} denote the set of edge-maximal graphs in \mathcal{G} . Also, let $A(\mathcal{G})$ denote the set of real numbers such that $P(G, q) > 0$ for every $G \in \mathcal{G}$.

Lemma 2.6. *If \mathcal{G} is a minor-closed class of graphs, then $A(\mathcal{G}) = A(\mathcal{G}_{\max})$.*

Proof. Clearly $A(\mathcal{G}) \subseteq A(\mathcal{G}_{\max})$ hence it just remains to prove the reverse inclusion. To this end, suppose for a contradiction that there is $q \in A(\mathcal{G}_{\max})$ such that $q \notin A(\mathcal{G})$. Thus, there is a graph $G \in \mathcal{G}$ such that $P(G, q) \leq 0$. Suppose that G has as few vertices as possible and, subject to that, suppose further that G is edge maximal. Since $q \in A(\mathcal{G}_{\max})$, we have that G is not edge-maximal. Thus there are $u, v \in V(G)$ such that $G + uv \in \mathcal{G}$. Now by the addition–contraction formula, we have

$$P(G, q) = \overline{P(G + uv, q)} + P(G/uv, q). \quad (2.1)$$

Since \mathcal{G} is minor-closed, we also have that $G/uv \in \mathcal{G}$. Furthermore, by the vertex-minimality and edge-maximality assumptions on G , we have $P(G + uv, q) > 0$ and $P(G/uv, q) > 0$. It follows from (2.1) that $P(G, q) > 0$, a contradiction. \square

3. Chromatic roots of planar graphs

In this section we apply Theorem 2.5 to deduce Theorem 1.2. However we first discuss the current situation with regard to the chromatic roots of the set \mathcal{G} of planar graphs. Recall the result of Jackson mentioned in the Introduction and the following widely believed conjecture of Birkhoff and Lewis.

Conjecture 3.1 ([2]). *A planar graph has no chromatic root in the interval $[4, \infty)$.*

Collectively, these would imply $R(\mathcal{G}) \subseteq \{0, 1\} \cup (32/27, 4)$. Moreover, the result of Thomassen and the work of this paper suggest that $\overline{R(\mathcal{G})} = \{0, 1\} \cup [32/27, 4]$. However, this simple statement does not capture the richness of behaviour exhibited by the chromatic polynomials of planar graphs.

Indeed, recall Theorem 1.3 which says that $\tau + 2 \in A(\mathcal{G}_{\max})$, where \mathcal{G} is the set of planar graphs. Applying Lemma 2.6, one may deduce that Theorem 1.3 holds not just for planar triangulations but for all planar graphs. We suspect this is known, though we include it here as we have been unable to find a written reference.

Theorem 3.2. *If \mathcal{G} denotes the class of planar graphs, then $\tau + 2 \in A(\mathcal{G})$.*

Table 1. Coefficients of the polynomials p_1 and p_2

	p_1	p_2
q^{16}	1	
q^{15}	-48	1
q^{14}	1098	-45
q^{13}	-15902	961
q^{12}	163358	-12934
q^{11}	-1263582	122848
q^{10}	7622430	-873498
q^9	-36631145	4811140
q^8	141942413	-20939539
q^7	-445623415	72739711
q^6	1131241573	-202024440
q^5	-2299985643	445549685
q^4	3673360947	-766835653
q^3	-4454394269	996884682
q^2	3863934525	-923047942
q^1	-2138431325	543148340
q^0	567438039	-152643002

It follows naturally to ask for a precise description of the set $A(\mathcal{G})$ where \mathcal{G} denotes the class of planar graphs. We believe the answer to be the following.

Conjecture 3.3. *If \mathcal{G} denotes the class of planar graphs, then $A(\mathcal{G}) = \{\tau + 2\} \cup [4, \infty)$.*

Note that Conjecture 3.3 contains Conjecture 3.1, Theorem 3.2 and the Four-Colour Theorem. In addition, it also implies Conjecture 1.1 via an application of Theorem 2.5. In what follows we shall show that $(3, t_1) \cup (t_2, 4) \subseteq N(\mathcal{G})$, where t_1 and t_2 are defined below. Therefore, as in the problem of determining $R(\mathcal{G})$, the only part of Conjecture 3.3 that remains to be proved are the intervals (t_1, t_2) and $(4, 5)$.

Let p_1 and p_2 be the polynomials whose coefficients are listed in Table 1. Let $t_1 \approx 3.618032$ be the largest real root of p_1 , and let $t_2 \approx 3.618356$ be the second largest real root of p_2 .

Lemma 3.4. *If \mathcal{G} denotes the class of planar graphs, then $(3, t_1) \cup (t_2, 3.8) \subseteq N(\mathcal{G})$.*

Proof. Let F and W be the graphs in Figure 1, and let G_1 be the unique graph obtained from the union of F and W by identifying the two distinguished 4-cycles. It may be calculated that G_1 has chromatic polynomial

$$P(G_1, q) = q(q^7 - 18q^6 + 141q^5 - 619q^4 + 1627q^3 - 2525q^2 + 2107q - 714),$$

which is negative in the interval $(3, 3.6)$.

Now let K be the graph in Figure 2. Let G_2 denote the planar triangulation formed from K and two copies of F , say F_1 and F_2 , by identifying the two distinguished 4-cycles in K with the distinguished 4-cycles in F_1 and F_2 respectively. We do this in such a way that the vertices of degree 3 in F_1 and F_2 are at distance 2 in G_2 . It may be computed that this graph has chromatic




Figure 1. The graphs F and W with distinguished 4-cycles in bold.




Figure 2. The graphs K and L with distinguished 4-cycles in bold.

polynomial

$$P(G_2, q) = q(q - 1)(q - 2)(q - 3)^3 p_1(q),$$

where p_1 is the polynomial in Table 1. Among other places, the polynomial $P(G_2, t)$ is negative in the interval $(3.5, t_1)$.

Finally, let G_3 be the unique planar triangulation formed from the disjoint union of F , K and W by identifying the two distinguished 4-cycles in K with those in F and W respectively. It may be computed that the chromatic polynomial of this graph is

$$P(G_3, q) = q(q - 1)(q - 2)(q - 3)^2 p_2(q),$$

where p_2 is the polynomial in Table 1. The polynomial $P(G_3, q)$ is negative in the interval $(t_2, 3.8)$. □

In Section 4, we investigate an infinite family of planar triangulations $T(n)$, $n \in \mathbb{N}$. These graphs are formed from the graphs W , K and L depicted in Figures 1 and 2. We analyse their chromatic polynomials using a transfer matrix approach to obtain the following result.

Lemma 3.5. *For every $q \in (3.7, 4)$, there exists $n \in \mathbb{N}$ such that $P(T(n), q) < 0$.*

We are now able to prove Theorem 1.2.

Proof of Theorem 1.2. Let \mathcal{G} denote the class of planar graphs. Lemmas 3.4 and 3.5 show that $(3, t_1) \cup (t_2, 4) \subseteq N(\mathcal{G})$. By Wagner’s theorem [10], which is equivalent to Kuratowski’s theorem, $\text{Forb}(\mathcal{G}) = \{K_5, K_{3,3}\}$. Since both K_5 and $K_{3,3}$ are 3-connected, Theorem 2.5 now implies the result. □

4. Proof of Lemma 3.5

Let K and L be the graphs in Figure 2. We denote by L^n the graph obtained from n copies of L , say L_1, \dots, L_n , by identifying the inner 4-cycle of L_i with the outer 4-cycle of L_{i+1} for each $i \in \{1, \dots, n-1\}$. The resulting graph is planar and has two distinguished 4-cycles. Also let W denote the wheel on 5 vertices (see Figure 1), and let K' denote the graph formed from K by adding a single edge, triangulating one of the distinguished 4-cycles. In this section, we analyse the family of planar triangulations $T(n)$, $n \in \mathbb{N}$, formed by identifying the distinguished 4-cycles of L^n with the distinguished 4-cycles of W and K' respectively.

The graphs $T(n)$, $n \in \mathbb{N}$ were first studied by Royle [5], who took inspiration from a graph found by Woodall and from graphs studied in the field of statistical mechanics. Indeed, the family of graphs L^n , $n \in \mathbb{N}$ can be viewed as $4 \times n$ strips of the infinite triangular lattice: see [5, Figure 1], where one side of the strip is wrapped around and identified with the other. A standard technique to compute the chromatic polynomial of such graphs is the so-called transfer matrix approach: see [6]. We employ this technique in the form used by Royle [5], and give here a fairly condensed analysis. A more detailed presentation can be found in [5].

Let A be a graph with a distinguished 4-cycle $a_1a_2a_3a_4$. We may partition the colourings ϕ of A into four types.

Type 1: $\phi(a_1) = \phi(a_3)$ and $\phi(a_2) = \phi(a_4)$.

Type 2: $\phi(a_1) = \phi(a_3)$ and $\phi(a_2) \neq \phi(a_4)$.

Type 3: $\phi(a_1) \neq \phi(a_3)$ and $\phi(a_2) = \phi(a_4)$.

Type 4: $\phi(a_1) \neq \phi(a_3)$ and $\phi(a_2) \neq \phi(a_4)$.

Let $P_i(A, q)$ denote the number of q -colourings of A of type i . Note that identifying two vertices a_1 and a_3 , say, gives a graph whose colourings correspond bijectively to the colourings ϕ of A such that $\phi(a_1) = \phi(a_3)$. Alternatively, adding the edge a_1a_3 produces a graph whose colourings correspond bijectively to the colourings ϕ of A such that $\phi(a_1) \neq \phi(a_3)$. Thus, for computational purposes, we have for example that $P_2(A, q) = P(A/a_1a_3 + a_2a_4, q)$. We collect this information in a vector $Q(A, q)$ called the *partitioned chromatic polynomial*:

$$Q(A, q) = \begin{pmatrix} P_1(A, q) \\ P_2(A, q) \\ P_3(A, q) \\ P_4(A, q) \end{pmatrix}.$$

Let $\langle q \rangle_k$ denote the k th falling factorial $q(q-1) \cdots (q-k+1)$ and let q_1, \dots, q_4 be the polynomials whose coefficients are listed in Table 2. The partitioned chromatic polynomials of W and K' are given by

$$Q(W, q) = \begin{pmatrix} \langle q \rangle_3 \\ \langle q \rangle_4 \\ \langle q \rangle_4 \\ \langle q \rangle_5 \end{pmatrix}, \quad Q(K', t) = \begin{pmatrix} \langle q \rangle_5 \cdot q_1(q) \\ \langle q \rangle_4 \cdot q_2(q) \\ \langle q \rangle_4 \cdot q_3(q) \\ \langle q \rangle_4 \cdot (q-3) \cdot q_4(q) \end{pmatrix}$$

Table 2. Coefficients of the polynomials q_1, q_2, q_3 and q_4 : see [5, Table 1]

	q_1	q_2	q_3	q_4
q^{11}		1	1	1
q^{10}		-34	-34	-34
q^9	1	538	538	538
q^8	-28	-5244	-5244	-5246
q^7	362	35078	35078	35148
q^6	-2846	-169490	-169490	-170548
q^5	15036	604806	604806	613920
q^4	-55448	-1595807	-1595807	-1645010
q^3	142716	3051803	3051803	3222645
q^2	-246724	-4024676	-4024676	-4397342
q^1	258889	3286881	3286881	3753346
q^0	-124884	-1255163	-1255163	-1511254

The formula for $Q(W, q)$ is trivial, whereas the formula for $Q(K', q)$ appears in [5] and was verified by the present authors.

In a similar way, for a graph with two distinguished 4-cycles, we may define a square matrix whose entries capture the types of colourings on those two 4-cycles. More precisely, we let the element in the i th row and j th column be the number of colourings of the graph which are of type i on the outer distinguished 4-cycle and type j on the inner distinguished 4-cycle. The matrix M corresponding to the graph L in Figure 2 is given by

$$M = \begin{pmatrix} \langle q \rangle_4 & \langle q \rangle_5 & \langle q \rangle_5 & \langle q \rangle_6 \\ \langle q \rangle_5 & \langle q \rangle_4 + 2\langle q \rangle_5 + \langle q \rangle_6 & \langle q \rangle_4 + 2\langle q \rangle_5 + \langle q \rangle_6 & 4\langle q \rangle_5 + 4\langle q \rangle_6 + \langle q \rangle_7 \\ \langle q \rangle_5 & \langle q \rangle_4 + 2\langle q \rangle_5 + \langle q \rangle_6 & \langle q \rangle_4 + 2\langle q \rangle_5 + \langle q \rangle_6 & 4\langle q \rangle_5 + 4\langle q \rangle_6 + \langle q \rangle_7 \\ \langle q \rangle_6 & 4\langle q \rangle_5 + 4\langle q \rangle_6 + \langle q \rangle_7 & 4\langle q \rangle_5 + 4\langle q \rangle_6 + \langle q \rangle_7 & M_{44} \end{pmatrix},$$

where $M_{44} = 2\langle q \rangle_4 + 16\langle q \rangle_5 + 20\langle q \rangle_6 + 8\langle q \rangle_7 + \langle q \rangle_8$. The expression appears in [5] and was verified by the present authors. We can now state a lemma of Royle.

Lemma 4.1 ([5]). *Let A and B be graphs with distinguished 4-cycles. If $X_{A,B}(n)$ denotes the graph obtained from L^n by identifying its two distinguished 4-cycles with those of A and B respectively, then the chromatic polynomial of $X_{A,B}(n)$ is the sole entry of the 1×1 matrix*

$$Q(A)^T D (MD)^n Q(B),$$

where

$$D = \begin{pmatrix} 1/\langle q \rangle_2 & 0 & 0 & 0 \\ 0 & 1/\langle q \rangle_3 & 0 & 0 \\ 0 & 0 & 1/\langle q \rangle_3 & 0 \\ 0 & 0 & 0 & 1/\langle q \rangle_4 \end{pmatrix}.$$

The matrix MD is called the *transfer matrix* of the graph L , as it describes how colourings transfer from one 4-cycle to the other. As we let n tend to infinity, the matrix $(MD)^n$ is determined by the spectral properties of MD . Indeed let $\lambda_1, \dots, \lambda_4$ and v_1, \dots, v_4 denote the eigenvalues and eigenvectors of MD respectively. Furthermore, let $\|v\|$ denote the norm of the vector v with respect to the inner product $\langle u, v \rangle = u^T Dv$. Royle proves the following.

Lemma 4.2 ([5]). *If A and B are graphs with distinguished 4-cycles and q is a fixed real number in $(\tau + 2, 4)$, then*

$$P(X_{A,B}(n), q) = \alpha_1 \beta_1 \lambda_1^n \|v_1\|^2 + \alpha_2 \beta_2 \lambda_2^n \|v_2\|^2 + \alpha_3 \beta_3 \lambda_3^n \|v_3\|^2 + \alpha_4 \beta_4 \lambda_4^n \|v_4\|^2,$$

where $\alpha_i \|v_i\|^2 = Q(A, q)^T Dv_i$ and $\beta_i \|v_i\|^2 = Q(B, q)^T Dv_i$ for $i \in \{1, 2, 3, 4\}$.

In essence, Lemma 4.2 says that if $\lambda_i > 0$ is the eigenvalue of largest absolute value, then in the limit as n tends to infinity, the chromatic polynomial of $X_{A,B}(n)$ has the sign of $\alpha_i \beta_i$. Moreover, this sign can be determined simply by looking at the partitioned chromatic polynomials of A and B and the eigenvector v_i . With this in mind we first find the eigenvalues and eigenvectors of MD . Indeed, if a is the polynomial $q^4 - 10q^3 + 43q^2 - 106q + 129$, then we have that $\lambda_1 = 2$,

$$\lambda_2 = \frac{1}{2}(q - 3)(q^3 - 9q^2 + 33q - 48 - (q - 4)a(q)^{1/2}),$$

$$\lambda_3 = \frac{1}{2}(q - 3)(q^3 - 9q^2 + 33q - 48 + (q - 4)a(q)^{1/2}),$$

and $\lambda_4 = 0$. The corresponding eigenvectors are $v_1 = (1, -1, -1, 1)^T$,

$$v_2 = \begin{pmatrix} (q^2 - 7q + 15 + a(q)^{1/2}) / (2(q - 2)(q - 3)^2) \\ (q^2 - 9q + 21 + a(q)^{1/2}) / (4(q - 3)^2) \\ (q^2 - 9q + 21 + a(q)^{1/2}) / (4(q - 3)^2) \\ -1 \end{pmatrix},$$

$$v_3 = \begin{pmatrix} (q^2 - 7q + 15 - a(q)^{1/2}) / (2(q - 2)(q - 3)^2) \\ (q^2 - 9q + 21 - a(q)^{1/2}) / (4(q - 3)^2) \\ (q^2 - 9q + 21 - a(q)^{1/2}) / (4(q - 3)^2) \\ -1 \end{pmatrix},$$

and $v_4 = (0, 1, -1, 0)^T$.

We remark that in the notation of this section, the graphs $T(n)$ in Lemma 3.5 can be denoted $X_{K',W}(n)$. We can now prove Lemma 3.5.

Proof of Lemma 3.5. Let t be a fixed real number in $(3.7, 4)$. At t , we have $\lambda_1 > \lambda_2 > \lambda_3 > \lambda_4 = 0$. For $i \in \{1, 2, 3, 4\}$, let α_i be such that $\alpha_i \|v_i\|^2 = Q(K', t)^T Dv_i$, and let β_i be the corresponding value for the graph W . It may be calculated that $\alpha_1 \|v_1\|^2 = 0$ and $\beta_1 \|v_1\|^2 = 0$. Similarly, if b denotes the polynomial $q^3 - 9q^2 + 25q - 24$, and the polynomials c and d are

Table 3. Coefficients of the polynomials c and d

	c	d
q^{13}	1	
q^{12}	-38	
q^{11}	676	1
q^{10}	-7473	-33
q^9	57452	506
q^8	-325572	-4770
q^7	1405168	30784
q^6	-4698525	-143070
q^5	12224002	489214
q^4	-24510275	-1231299
q^3	36830333	2234215
q^2	-39168919	-2778891
q^1	26277660	2126441
q^0	-8337189	-755627

defined as in Table 3, then it may be calculated that

$$\alpha_2 \|v_2\|^2 = \frac{d(q)a(q)^{1/2} - c(q)}{2(q - 3)}$$

and

$$\beta_2 \|v_2\|^2 = \frac{(q - 2)a(q)^{1/2} - b(q)}{2(q - 3)^2}.$$

From these expressions and by Lemma 4.2, the dominant term in the expression for $P(T(n), q)$ is $\alpha_2 \beta_2 \lambda_2^n \|v_2\|^2$. Thus, for large enough n , the sign of $P(T(n), q)$ depends on the sign of $\alpha_2 \beta_2$. A short calculation gives that $\alpha_2 \|v_2\|^2 < 0$ for $q \in (3.7, 4)$ and $\beta_2 \|v_2\|^2 > 0$ for $q \in (3, 4)$, which implies that $\alpha_2 \beta_2$ is negative in $(3.7, 4)$. Therefore, there exists $n \in \mathbb{N}$ such that $P(T(n), q) < 0$. □

4.1. Remarks on the computations

The calculation of chromatic polynomials in this article was done with Maple. The calculation of eigenvalues and eigenvectors of the matrix MD was performed with MATLAB and cross-checked with the series expansion expressions presented in [5].

4.2. Concluding remarks

Theorem 1.2 is close to a proof of Conjecture 1.1. Indeed the missing interval (t_1, t_2) has length less than 0.0004. Because of Tutte’s result [9], it will require an infinite family of planar graphs to shrink this interval to the point $\tau + 2$. It seems conceivable that for some graph S , the family $X_{S,W}(n)$, $n \in \mathbb{N}$ may be useful to dispose of the interval $(\tau + 2, t_2)$. Indeed, by mirroring the proof of Lemma 3.5, one only needs to find a graph S , with a distinguished 4-cycle, such that $Q(S, t)Dv_2 < 0$ for $q \in (\tau + 2, t_2)$. Royle observed the striking fact that the eigenvalues $\lambda_1, \dots, \lambda_4$ are not distinct for $q = \tau + 2$. However, they are distinct for $q \in (3, 4) \setminus \{\tau + 2\}$. Thus

it seems possible that the families $X_{S,W}(n)$, $n \in \mathbb{N}$ may also be useful to dispose of the interval $(t_1, \tau + 2)$.

References

- [1] Birkhoff, G. D. (1912) A determinant formula for the number of ways of coloring a map. *Ann. of Math.* **12** 42–46.
- [2] Birkhoff, G. D. and Lewis, D. C. (1946) Chromatic polynomials. *Trans. Amer. Math. Soc.* **60** 355–451.
- [3] Jackson, B. (1993) A zero-free interval for chromatic polynomials of graphs. *Combin. Probab. Comput.* **2** 325–336.
- [4] Perrett, T. J. (2016) Roots of the chromatic polynomial. PhD thesis, Technical University of Denmark.
- [5] Royle, G. (2008) Planar triangulations with real chromatic roots arbitrarily close to four. *Ann. Combin.* **12** 195–210.
- [6] Salas, J. and Sokal, A. D. (2001) Transfer matrices and partition-function zeros for antiferromagnetic Potts models I: General theory and square-lattice chromatic polynomial. *J. Statist. Phys.* **104** 609–699.
- [7] Sokal, A. D. (2004) Chromatic roots are dense in the whole complex plane. *Combin. Probab. Comput.* **13** 221–261.
- [8] Thomassen, C. (1997) The zero-free intervals for chromatic polynomials of graphs. *Combin. Probab. Comput.* **6** 497–506.
- [9] Tutte, W. T. (1970) The golden ratio in the theory of chromatic polynomials. *Ann. New York Acad. Sci.* **175** 391–402.
- [10] Wagner, K. (1937) Über eine Eigenschaft der ebenen Komplexe. *Math. Ann.* **114** 570–590.