




Neuroimaging Highlight

Progressive Neurological Decline Associated With Intracranial Calcification in Down Syndrome; Fahr Disease Mimic?

Montaha Almudhry^{1,2} , Chitra Prasad^{1,3}, KengYow Tay^{1,4}, Derek B. Debicki^{1,5}  and Asuri N. Prasad^{1,3,5} 

¹London Health Sciences Centre and Western University, London, ON, Canada, ²Department of Neuroscience, King Fahad Specialist Hospital, Dammam, Saudi Arabia, ³Department of Pediatrics, Western University, London, ON, Canada, ⁴Department of Medical Imaging, Western University, London, ON, Canada and ⁵Department of Clinical Neurological Sciences, Western University, London, ON, Canada

Keywords: Down syndrome; calcification; basal ganglia; dentate nucleus; cranial CT scan

(Received 30 September 2023; date of acceptance 19 October 2023; First Published online 7 November 2023)

Intracranial calcification (ICC) has been described commonly in association with congenital infections, head trauma, and endocrine or systemic metabolic derangements. However, it is also noted in patients with pathogenic genetic mutations and defects in inborn error of metabolism.^{1–4} We report the clinical, imaging, and genetic findings in an adult with Down syndrome (DS) referred to the clinic for further investigation for Fahr disease.

A 43-year-old lady with DS, secondary to Robertsonian translocation of chromosomes 13 and 21 and a background history of hypothyroidism who remained highly functional till the age of 38 years, was noted to have progressive neurological decline affecting cognitive, motor, visual, adaptive, and linguistic abilities. At baseline, the patient had a grade 1 level in literacy and writing and was working in a nursing home with minimal supervision. However, over the span of 5 years, she progressively lost executive function, short-term recall, expressive communication, and ambulation capabilities. She also started experiencing epileptic seizures, which were controlled following the introduction of antiseizure medication, levetiracetam, and lacosamide. The patient was adopted at two years of life, details of the family history could not be elicited. A limited examination in the clinic demonstrated no eye contact; she would not interact with the examiner. She was sitting in her wheelchair, and bilateral spasticity was pronounced in the lower extremities with diffuse brisk deep tendon reflexes, however, ankle clonus was not elicited. The patient was rated by her sister for the Dementia Questionnaire for Persons with Mental Retardation⁵ and found to score 30/66, within the range of moderate cognitive difficulties as well as moderately severe behavioral symptoms.

A CT scan requested by her neurologist as part of her investigative work up for neurological decline demonstrated calcifications within the basal ganglia, dentate nuclei, and parasagittal occipital sulci bilaterally (Figs. 1 and 2). The patient has since undergone biochemical testing, showing intact parathyroid and thyroid function; PTH:2.7 pmol/L (1.6–6.9), calcium:2.26



Figure 1: Axial CT illustrates dense calcification of the bilateral lentiform nuclei (white arrowheads) and bilateral parasagittal occipital cortices (thin white arrows).

mmol/L (2.15–2.60), phosphorus:0.98 mmol/L (0.80–1.45), TSH:1.090mIU/L (0.465–4.680). Her karyotype was reported as 46, XX, der (13;21) (q10;q10),+21, consistent with Robertsonian translocation, between the long arms of one copy of each of

Corresponding author: A. N. Prasad; Email: narayan.prasad@lhsc.on.ca

Cite this article: Almudhry M, Prasad C, Tay KY, Debicki DB, and Prasad AN. (2024) Progressive Neurological Decline Associated With Intracranial Calcification in Down Syndrome; Fahr Disease Mimic?. *The Canadian Journal of Neurological Sciences* 51: 683–684, <https://doi.org/10.1017/cjn.2023.309>

© The Author(s), 2023. Published by Cambridge University Press on behalf of Canadian Neurological Sciences Federation. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Figure 2: Coronal CT shows dense calcification of the bilateral cerebellar white matter including the dentate nuclei (thick white arrows).

chromosomes 13 and 21. Molecular genetic testing for Fahr disease was conducted with no pathogenic variant identified.

Patients with DS have a higher prevalence of basal ganglia calcifications with a frequency of 10%–26.2% compared to only 0.3%–0.6% in the general population.^{4,6,7} On scans, globus pallidus and, less commonly, the head of the caudate are the most affected structures, while other areas like the choroid plexus and pineal body are found to be just as frequently calcified in healthy individuals.⁶ Histopathology reveals that the calcification involves the media of small arteries in the perivascular regions and amyloid degeneration.⁷ The exact underlying pathogenesis of calcium deposit in DS remains to be elucidated. Clinically, the identification of ICC has been linked to accelerated aging and premature onset of dementia in DS.⁸

Imaging studies (Figs. 1 and 2) in this individual confirm that the intracranial calcification extends beyond the basal ganglia to

the cerebellum's dentate nuclei, providing additional justification for molecular testing for Fahr disease. Being aware of this phenomenon in DS could help with early diagnosis and prevent unneeded, costly, and time-consuming investigations.

Funding. None.

Competing interests. One of the authors ANP has received payment from the Canadian Medical Protection Association for expert opinion in a medicolegal case that is unrelated to the material or case in the submitted manuscript. A second author, DD, also received payment as a speaker's honorarium at UBC Canada that is unrelated to this manuscript.

Statement of authorship. MA wrote the manuscript; KT reviewed the CT of the patient and provided labeled figures; and CP, ANP, and DD provided consultation and diagnosis for the patient and supervised the manuscript.

References

- Livingston JH, Stivaros S, Van Der Knaap MS, Crow YJ. Recognizable phenotypes associated with intracranial calcification. *Dev Med Child Neurol.* 2013;55:46–57.
- Livingston JH, Stivaros S, Warren D, Crow YJ. Intracranial calcification in childhood: a review of aetiologies and recognizable phenotypes. *Dev Med Child Neurol.* 2014;56:612–26.
- Donzuso G, Mostile G, Nicoletti A, Zappia M. Basal ganglia calcifications (Fahr's syndrome): related conditions and clinical features. *Neurol Sci.* 2019;40:2251–63.
- Eskandary H, Sabba M, Khajehpour F, Eskandari M. Incidental findings in brain computed tomography scans of 3000 head trauma patients. *Surg Neurol.* 2005;63:550–3.
- Prasher VP. Dementia questionnaire for persons with mental retardation (DMR): modified criteria for adults with Down's syndrome. *J Appl Res Intellect Disabil.* 1997;10:54–60.
- Ieshima A, Kisa T, Yoshino K, Takashima S, Takeshita K. A morphometric CT study of down's syndrome showing small posterior fossa and calcification of basal ganglia. *Neuroradiology.* 1984;26:493–8.
- Le Becker S. Basal ganglia calcification in Down's syndrome. *J Neurol Neurosurg Psychiatry.* 1985;48:61–4.
- Wisniewski KE, French JH, Rosen JF, Kozlowski PB, Tenner M, Wisniewski HM. Basal ganglia calcification (BGC) in Down's syndrome (DS)—another manifestation of premature aging. *Ann N Y Acad Sci.* 1982;396:179–89.