

# Fishing down the food web of the Antarctic continental shelf and slope

David G. Ainley

H.T. Harvey & Associates, Los Gatos, CA 95032, USA  
(dainley@penguinscience.com)

Daniel Pauly

Sea Around Us Project, Fisheries Centre, University of British Columbia, Vancouver, Canada, V6T 1Z4

Received November 2012; first published online 27 February 2013

**ABSTRACT.** The history of biotic exploitation for the continental margin (shelf and slope) of the Antarctic Large Marine Ecosystem (LME) is reviewed, with emphasis on the period from 1970 to 2010. In the Antarctic Peninsula portion, marine mammals were decimated by the 1970s and groundfish by the early 1980s. Fishing for Antarctic krill *Euphausia superba* began upon the demise of groundfish and now is the only fishing that remains in this region. Surveys show that cetacean and most groundfish stocks remain severely depressed, harvest of which is now prohibited by the International Whaling Commission and the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR). On the other hand, krill fishing in this region is underway and in recent years has contributed up to 72% of the Southern Ocean catch, depending on fishing conditions and the CCAMLR conservation measures in force. Elsewhere along the Antarctic continental margin, marine mammals were also severely depleted by the 1970s, followed directly by relatively low-level fisheries for krill that continued until the early 1990s. Recently in these areas, where fin-fishing is still allowed, fisheries for Antarctic toothfish *Dissostichus mawsoni* have been initiated, with one of this fish's main prey, grenadiers *Macrourus* spp., being taken significantly as by-catch. Continental margin fishing currently accounts for ~25% of the total toothfish catch of the Southern Ocean. Fishing along the Antarctic continental margin, especially the Antarctic Peninsula region, is a clear case of both the tragedy of the commons and 'fishing down the food web'.

## Introduction

The extraction of fish and other biotic resources from marine systems has had dramatic effects on ocean ecosystems (Pauly and others 1998; Halpern and others 2008; Baum and Worm 2009; Estes and others 2011). The process by which this has been accomplished, described by Pauly and others (1998) as 'fishing down the food web', generally has been to initially target larger, high-trophic level species and then, as these become depleted, to transition to short-lived pelagic fish and low-trophic level invertebrates. These changes result in a reduction of the mean trophic level of the catch, now called the marine trophic index (MTI; Pauly and Watson 2005). An alternative explanation for the change in the average trophic level of species exploited, argued by Essington and others (2006) and called 'fishing through the food web', is that exploitation of upper-trophic level species continues but lower-trophic level ones are added. One problem in disentangling these processes and subsequent interpretation of ecosystem effects is that the most studied ocean areas, where a large portion of fishery extraction has occurred, are coastal, with the ocean there subject to numerous other natural and unnatural, confounding impacts (see, for example, Halpern and others 2008). Most recently, Branch and others (2010) proposed that the catch statistics used in 'fishing down' and 'fishing through' studies are not informative as to trophic changes in fisheries and food

web health, and that direct abundance-survey data are required to gauge current food-web structure and fishing impacts.

Herein an attempt is made to review how the most remote ocean area in the world, the Antarctic continental shelf and slope, has fared in the face of human exploitation. This location, with its daunting environmental conditions, has no resident human population, and only a relatively short history of human exploitation, contrary to the situation in the Arctic (Zeller and others 2011). Thus, the Antarctic continental shelf and slope are destinations that can serve few other purposes than exploitation. Indeed, other than the taking of biotic resources there has been little other anthropogenic influence on the food webs of these waters, except for recent effects of global climate change (Atkinson and others 2004; Halpern and others 2008).

The waters covered in this review are those included in the continental margin portion of the Antarctic large marine ecosystem (LME; Aquarone and Adams 2008; see also Sea Around Us Project 2012). These waters are also contained within Food and Agriculture Organization (FAO) fishery statistical areas, subareas and divisions 48.1; 48.5; 48.6 (in part); and the continental margin portions of 58.4.1, 58.4.2 and 88.1–88.3. They are, obviously, also part of the area covered by the Convention for the Conservation of Antarctic Marine Living Resources (CCAMLR; Fig. 1). Overall, the Antarctic continental shelf and slope in many sectors have received




Fig. 1. (Colour online) Antarctica, its Large Marine Ecosystem (LME, in grey) and the FAO areas, subareas and divisions used in this report (which are also those used by CCAMLR). Note that subarea 48.6 actually extends further north, but was cut here (dotted line), as this contribution deals only with the shelf and slope of the Antarctic continent. The present report covers largely just the grey area: shelf and slope.

relatively little attention by commercial sealers, whalers and fishers, in large part owing to persistent sea ice that deters vessel operations, especially vessels that are not ice

capable. Very few fishing vessels that ply the Southern Ocean are ice capable.

In the present review, details are given of the continued exploitation of Antarctic continental shelf and slope waters only. The fish catch data reported here have been taken from CCAMLR statistical bulletins (CCAMLR 1990–2012).

#### Early history: the marine mammals

The South Shetland and South Orkney islands (FAO subareas 48.1 and 48.2, respectively), continental islands located off the northern coast of the Antarctic Peninsula, were discovered by sealers in 1819–1820, and subsequently the Antarctic fur seal *Arctocephalus gazella* was extirpated (O’Gorman 1961; as were southern elephant seals *Mirounga leonina* in more northern Southern Ocean waters). Thus, human exploitation of Antarctic resources began with a high-trophic level species (Table 1). The last fur seal seen breeding in this area, on Nelson Island, South Shetlands, was in 1904. Not long after, in 1911, the hunt for baleen whales (*Balaenoptera* and *Megaptera* spp.) began among the South Shetlands, and by 1915 well over 6,000 whales had been taken, as well as thousands of tonnes of guano, presumably from penguin colonies (Tønnessen and Johnsen 1982: 190–200). By the late 1920s few whales could be found, but by then the industry had shifted to pelagic whaling in neighbouring waters and elsewhere in the Southern Ocean. In so doing,

Table 1. Scientific name, common name, mean maximum length and trophic levels of Antarctic marine mammals, fish and invertebrates mentioned in this article

Group	Scientific name	Common name	Mean max. length (cm) <sup>a</sup>	Trophic level <sup>b</sup>
<b>Marine mammals</b>	<i>Arctocephalus gazella</i>	Antarctic fur seal	178	3.70
	<i>Balaenoptera musculus</i>	Blue whale	3,200	3.20
	<i>Megaptera novaeangliae</i>	Humpback whale	1,800	3.60
	<i>Balaenoptera acutorostrata</i>	Minke whale	1,000	3.40
	Mysticeti	Baleen whales	2,000	3.30
<b>Fish</b>	<i>Chaenocephalus aceratus</i>	Blackfin icefish	59	3.47
	<i>Chaenodraco wilsoni</i>	Spiny icefish	35	3.32
	<i>Champscephalus gunnari</i>	Mackerel icefish	54	3.23
	<i>Pseudochaenichthys georgianus</i>	South Georgia icefish	53	3.33
	<i>Chionodraco rastrospinosus</i>	Ocellated icefish	43	3.85
	Channichthyidae	Crocodile icefish	59	3.34
	<i>Dissostichus eleginoides</i>	Patagonian toothfish	176	3.96
	<i>Dissostichus mawsoni</i>	Antarctic toothfish	144	4.45
	<i>Notothenia rossii</i>	Marbled rockcod	75	3.61
	<i>Gobionotothen gibberifrons</i>	Humped rockcod	45	3.15
	<i>Lepidonotothen squamifrons</i>	Grey rockcod	45	3.09
	<i>Pleuragramma antarcticum</i>	Antarctic silverfish	22	3.00
	Macrouridae	Rattails, grenadiers	82	3.88
	<i>Micromesistius australis</i>	Southern blue whiting	74	3.79
	Muraenolepididae	Eel cods	30	3.41
Myctophidae	Lanternfish	16	3.30	
Rajidae	Skates	205	3.82	
<b>Invertebrates</b>	<i>Euphausia superba</i>	Antarctic krill	5	2.20

<sup>a</sup> From Trites and Pauly (1998) for marine mammals, FishBase (2012) for fish and Rosenberg and others (1986) for krill.

<sup>b</sup> From Pauly and others (1998) for marine mammals, Fishbase (2012) for fishes and SeaLifeBase (2012) for krill.




Fig. 2. Catch by taxon (A) and by fishing country (B) in the shelf and slope waters of area 48 (48.1, the northern Antarctic Peninsula; 48.5; and shelf/slope waters of 48.6).

the whale catch was shifted to shelf and slope waters to the west, including the Ross Sea and the offshore Balleny Islands. By the early 1930s, blue whales *B. musculus* could no longer be found along the Ross Sea slope (Tønnessen and Johnsen 1982: 346–353; Ainley 2010). By the 1970s, few great whales remained in Southern Hemisphere waters, including those of the Antarctic continental shelf and slope (Hilborn and others 2003: 363–365). Blue whales, for example, which concentrated along the Antarctic continental slope, were reduced to no more than 10% of their historical numbers by the 1950s (Kareiva and others 2006; IUCN 2011). A take of Antarctic minke whales *B. bonaerensis* followed in the late 1960s to early 1980s, mainly in Antarctic continental shelf and slope waters, but this ended when a moratorium was placed on commercial whaling by the International Whaling Commission (IWC) in 1982 (Ainley 2010). A ‘scientific catch’ of minke whales by Japan, 400–1,000 individuals per year, was instituted in 1987 and

has continued in slope waters off East Antarctica to the present day (IWC areas III–VI, FAO areas 58 and 88; see Brown and Brownell 2001).

#### Fishery for groundfish

The early 1960s saw the emergence of fisheries for demersal species in the Southern Ocean, with the early years of this activity being well summarised by Kock (1992: 173–201). Fishing first around South Georgia (FAO subarea 48.3) and the Kerguelen Islands (58.5), exploitation began in waters of the South Orkney Islands (48.2) in 1977 and South Shetland Islands (48.1) in 1978 (a split season, such as 1978/79, is indicated here by the first year, in this case 1978). Target species in continental margin waters included *Notothenia rossii*, *Champrocephalus gunnari*, *Lepidonotothen squamifrons* and *Gobionotothen gibberifrons* (Table 1). Highest catches were made initially, with stocks becoming greatly




Fig. 3. Catch by taxon (A) and by fishing country (B) in the shelf and slope waters of area 88 (88.1, 88.2 (Ross Sea) and 88.3 (southern Antarctic Peninsula), as well as catches in 88 not allocated to one of these subareas).

depleted within 10 years (Fig. 2; Tables A1 and A2). These waters of the southern Scotia Arc and southward were closed to fin-fishing in 1990 by the CCAMLR, an international convention that came into force in 1982. At that time, stocks of almost all groundfish species were estimated to be <10% of pre-fished size (Kock 1992). To this day, fin-fishing in subareas 48.1 and 48.2 is prohibited (CCAMLR 2011), and stocks have yet to recover (Marschoff and others 2012). When the fish stocks disappeared in these Antarctic continental waters, industry turned more intensively to Antarctic krill *Euphausia superba* (see below).

Following closure of subareas 48.1 and 48.2 to fin-fishing in 1990, waters of the Antarctic continental shelf and slope saw no fishing for fin-fish until 1996. In that year, with the fishery for Patagonian toothfish *Dissostichus eleginoides* and other fin-fish fully exploited (or off limits) in insular waters of the northern Southern Ocean (northern portions of areas 48 and 58), a New Zealand vessel conducted exploratory fishing for Antarctic toothfish *D. mawsoni* in the continental shelf and slope

waters of the Ross Sea (subareas 88.1 and 88.2; Fig. 1). Until then, the long distance from home ports and persistent pack ice of these waters had discouraged fishing (Ainley and others 2012). Less than one tonne was caught that first year, but by 1999 the fishers had figured out the technique, finding the most productive areas (slope, Antarctic ridge seamounts) and depths (800–1,400m: slope and seamounts to the north). In 2000, after several years of what CCAMLR terms ‘exploratory fishing’, vessels of other CCAMLR members entered the fishery, though its ‘exploratory fishing’ status remained (owing to insufficient data to conduct a proper assessment). Permits were issued as long as vessels agreed to take observers and abide by CCAMLR rules (such as no fishing in waters <550m deep to protect benthic communities, or location to be changed if the by-catch becomes too high; CCAMLR 2011). By 2003, vessels flagged in 12 countries had fished in area 88 and in that year the quota, based on stock models with the initial stock level unknown, was reached for the first time: the catch was around 3,000–3,500t (Fig. 3A). The catch has remained at about




Fig. 4. The ratio of the by-catch of grenadiers (family Macrouridae) to the catch of toothfish (family Nototheniidae) in areas 58 and 88 (1977–2010). The by-catch is actually much higher in area 58 owing to the large IUU effort in the 58.4.1 and 58.4.2 divisions, for which the level of by-catch is not known.

that level since (Table A3), but the number of countries and vessels fishing has decreased (Table A4). Presumably, the distances to port, conditions and competition encountered have proved too daunting for some. In the last three years about 15 vessels from only five or six countries have fished (for example, four vessels from New Zealand and up to six from the Republic of South Korea), in an Olympic-style scenario in which permitted vessels vie for the same total allowable catch (TAC) (Fig. 3B). Since inception of the fishery, and attracted by the high price that 'Chilean sea bass' commands, several vessels have foundered, caught fire or been holed by collision with ice thus requiring rescue. The Ross Sea (especially subarea 88.1; see Fig. 1) is regularly patrolled by aircraft from New Zealand and this activity is thought to discourage almost all IUU fishing in this area.

In 1998, a cruise was undertaken by Chilean vessels to determine the level of toothfish catch that could be achieved in subarea 48.1, as well as at the base of the Antarctic Peninsula (88.3). Yields were very low, <20g/hook and were equalled or surpassed by by-catch fishes (Arana and Vega 1999). This subarea, characterised by year-round sea-ice cover, subsequently has seen very limited fishery.

The by-catch in the toothfish fishery of non-target, unused species, principally macrourid fish *Macrourus* spp. (Macrouridae) and fish of the genus *Muraenolepis* (Muraenolepidae), and to a lesser extent skates *Raja* spp., has been significant. In the legal fishery, it has ranged to almost 170t, which in some years has been close to 10% of the toothfish catch in the respective subarea (Fig. 4). The ratio of the catches of macrourids to toothfish caught has decreased following the onset of fishing; in other words, the catch is ini-

tially high then gradually declines in subsequent years (Fig. 4).

The waters off East Antarctica (divisions 58.4.1, 58.4.2), the parts of the Antarctic continent facing the Indian Ocean, began to be fished for krill and some bottom fish as early as 1974. Fishing was short-lived and initially discontinued after the early 1990s. However, the success of fishing in area 88 inspired fishing companies to explore the continental margin elsewhere for Antarctic toothfish. Mainly this led to re-opening of the continental margin of subarea 58 to fishing less than 10 years ago (58.4.1, 58.4.2). This 'exploratory' fishery for Antarctic toothfish is conducted by many of the same countries fishing area 88 (Table A5). The TAC has been lower here than elsewhere (Fig. 5), but the IUU take has been huge, often surpassing the legal take. By-catch is sparsely reported (Table A6), the IUU by-catch not at all, and, therefore, the pattern relative to tonnes of toothfish caught in area 88 is difficult to ascertain (Fig. 4).

#### Fishery for forage fish

On occasion, attempts have been made to determine the feasibility of taking some of the mid-water fish in continental waters. Principally, this has involved species of Myctophidae (lanternfish, perhaps mostly *Electrona* spp.) along the slope and the nototheniid *Pleuragramma antarcticum* over the shelf. In both cases these species could be characterised as loosely shoaling, a level of dispersal that is problematic for fuel-efficient trawling (Gjøsaeter and Kawaguchi 1980). Fished during less than 10 seasons in total and not in every area, peak catches for *Pleuragramma* were reported as 653t and 1,517t in subareas 58.4.2 and 88 (subarea not specified), and for




Fig. 5. Catches by taxon (A) and by fishing country (B) in the shelf and slope portions of area 58 (divisions 58.4.1 and 58.4.2).

myctophids 129t in subarea 88, in all cases by USSR vessels (Table A3).

#### Fishery for Antarctic krill

Antarctic krill (Table 1) is the small, shrimp-like crustacean supporting, directly or indirectly, most larger animals in the Antarctic LME, or at least those of the outer shelf and slope (Pikitch and others 2012). Fisheries, since the 1970s, have exploited krill for making fish meal and related products. Japan reported taking 283t in the Indian Ocean sector (58.4.2) in 1974, Chile reported taking 276t and 92t in waters of the South Shetland Islands (48.1) in 1976 and 1977, and the USSR reported taking 3,355t from area 88 (subarea unspecified) in 1977. Subsequently, the fishery for krill in area 88 was largely confined to the western portion, including waters off Wilkes Land and the Balleny Islands (Dolzhnev and others 1990). The fisheries in areas 58 and 88 were discontinued by the 1990s but never reached more than

6,000t or 35,000t, respectively (Fig. 6A). On the other hand, the krill fishery in Antarctic Peninsula waters has been much greater, ranging up to 150,000t, taken by vessels flagged in 14 different countries (Fig. 6B). During the 1980s and early 1990s, the catch in subarea 48.1 was consistently 20–25% of the total Southern Ocean catch. The total has been halved in recent years. However, the subarea 48.1 catch contributed 72% of the total in 2009 (26% in 2008, 5% in 2010). In recent years, vessels from South Korea, Japan (less so than initially) and Norway have replaced those from Eastern European countries as those taking the largest proportion of the annual catch in continental margin waters (Fig. 6B).

Krill trawlers seek to fish on swarms that are as pure as possible, to avoid contamination by other organisms, especially gelatinous ones (Kelleher 2005). At times, however, the by-catch can be significant, but it is not routinely reported. Observers take samples for analysis, but not every trawl is observed. The low level of reporting is related to difficulties in separating out




Fig. 6. Catches by taxon (A) and by fishing country (B) on the shelf and slope of the Antarctic continent (= portion of the Antarctica large marine ecosystem (LME) corresponding to the grey area of Figure 1).

by-catch owing to the method of catch and initial processing of krill. In most operations, krill are smashed together in fine-mesh trawls and the catch then has to be processed (oil removed, meat freeze- or air-dried) within a few hours to avoid the rapid enzymatic breakdown releasing toxic fluorides in the stored product (FAO 2005). Specific studies of by-catch have shown that lanternfish (Myctophidae) and the larvae of other fin-fish can be caught in significant amounts (Pakhomov and Pankratov 1994; Watters 1996) to the point that it is a concern for CCAMLR (CCAMLR 2006). Management thus far seeks to avoid secondary effects of an elevated by-catch of fin-fish juveniles, and competition with land-based predators, by spreading the krill fishing as broadly as possible among spatial management units (CCAMLR 2011). The degree to which the catch of fin-fish larvae and juveniles affects the stocks of by-catch species is open to question (Marschoff and others 2012). Some researchers posit that krill exploitation by fishing where predators concentrate

is limiting the recovery of some top trophic-level species, such as cetaceans (Leaper and Miller 2011).

### Discussion

To date, owing to severe conditions and destinations a long way from well-equipped ports, there has been no reason for humans to be active along the Antarctic continental margin except for scientific research or biotic exploitation. Coordination of research was instituted in 1959 by the signing of the Antarctic Treaty (Fogg 1992). Subsequently, treaty nations established a number of logistics hubs along the coast of FAO areas 48, 58 and 88 (Tin and others 2009). Treaty parties excluded future minerals development by an agreement that came into force in 1998. On the other hand, biotic exploitation along the continental margin (and elsewhere in the Southern Ocean), which had been underway for almost 200 years, was allowed to continue. Populations of certain




Fig. 7. The extraction of biotic resources from the Antarctic Peninsula (subarea 48.1) best illustrates the 'fishing down' process that occurred in Antarctic waters as a whole, starting with Antarctic fur seals, then on to baleen whales, fishes and, finally, krill (based on trophic levels in Table 1 and catches in Table A1). The open dots for 1976 and 1977 refer to miniscule catches (276t and 92t, respectively) taken by Chile, which were probably exploratory trawls.

seals and whales were decimated. To prevent a repetition of the fur (and elephant) seal decimation, a treaty banning the take of fur seals and controlling commercial take of other species (pack ice 'true' seals) entered into force in 1978 (Convention for the Conservation of Antarctic Seals). A few very small areas (two in subarea 88.1) were closed entirely to sealing to protect scientifically important populations. The International Whaling Commission, recognising that the stocks of large whales had been decimated, banned commercial whaling of all cetacean species in 1986. Fin-fishing then increased but quickly decimated the targeted demersal fishes. One of the first conservation measures adopted by CCAMLR, which the members brought into force in 1982, was to close subareas 48.1 and 48.2 to fin-fishing (by 1990). Fishing for Antarctic krill became regulated during the 1980s, and the management of this has been continually refined since then (CCAMLR 2011). To this day, krill fishing occurs in areas closed to fin-fishing as well as other areas.

Fur seals have recovered in the South Orkneys and South Shetlands (Boyd 2002), but baleen whales have not (IUCN 2011), although one species, humpback whale *M. novaeangliae*, is beginning to do so (Branch 2011). Some fin-fish species in area 48.1 are showing signs of recovery (for example, *N. rossii*), while others are not (Marschoff and others 2012). Fishing for Antarctic krill continues in this region, with the continental margin fishery being important, its contribution (5–72% in recent years) depending on conditions and conservation measures in force (CCAMLR 2011; see Figures 2, 3, 5 and 6).

The history of fishing in Antarctic continental waters is a good example of 'fishing down the food web'. Figure 7 summarises 'fishing down' as it occurred in the vicinity of the Antarctic Peninsula (subarea 48.1) and, by extension, around the Antarctica margin as a whole.

Clearly, what occurred was not 'fishing through', as described by Essington and others (2006): the mean trophic level of the fishery (or previously the hunt for marine mammals) declined not (only) because new, low trophic-level species came under exploitation, but because the high trophic-level animals that were targeted had been decimated and their exploitation therefore had to cease for economic reasons. Thus, the MTI, in this case, does its job as an indicator of high-trophic level biodiversity (Pauly and Watson 2005) and points to a vastly altered ecosystem (see also Ballance and others 2006; Ainley and Blight 2009; Trivelpiece and others 2011), a victim of the 'tragedy of the commons' (Hardin 1968). Ironically, the MTI in recent decades may be increasing, not so much owing to the recovery of fur seals and baleen whales (which is still in its infancy), but because krill abundance is decreasing due to climate change (Atkinson and others 2004), perhaps exacerbated by decimation of the top trophic levels (Nicol and others 2010).

In the southern portions of areas 58 and 88, fur seals were never present but large baleen whales, once plentiful, were severely reduced. The take of smaller minke whales was greatly curtailed before that stock was depleted, and since appears to have recovered; stocks of humpback whales are beginning to recover in spectacular fashion in slope waters of western area 88 (Balleny Islands) and eastern area 58, but stocks of blue whales remain greatly depressed (Ainley 2010; Branch 2011). Nowadays, fishing in these areas, which once included krill, is confined to bottom fishes (many demersal fishes in the northern parts of area 58 were depleted by the late 1970s; Kock 1992: 191–192). The target species in waters of the Antarctic margin is Antarctic toothfish, but it appears that grenadiers (Macrouridae), caught as by-catch, are significantly taken as well. Ironically,



these macrourids are an important prey for the target species in the main fishing grounds along the continental slope (Fenaughty and others 2003). Impacts to the toothfish have not been studied, but the principles of ecosystem-based fishery management are probably being challenged.

The total catch of Antarctic toothfish in areas 58 and 88 accounts for almost all the catch for this species in the Southern Ocean or 26% (in 2010) of the total toothfish catch (~15,000t). The remainder of the total is contributed by *D. eleginoides* taken from northern areas of the Southern Ocean. Although krill were taken in relatively small amounts from areas 58 and 88 early, in recent years the fishery continues only in the SW Atlantic sector (area 48), probably owing to the greater densities of krill there as well as easier access by fishing vessels (as it is closer to ports and population centres).

In summary, then, exploitation of biotic resources along the continental margin of Antarctica has played a role in humans' progressive exploitation of global marine resources: ever farther away, deeper and more species (see, for example, Pauly and others 2005; Swartz and others 2010), and, with successive depletions, has not been something of which human society can be proud. During the last few decades, efforts have been made to try to recover this reputation in these Antarctic continental waters, though whether continued severe reduction in toothfish stocks can be offered as an example of enlightened management, that is, fishing to 50% of grossly estimated pre-fished spawning biomass (Constable and others 2000), is debatable (Ainley and Brooks 2013). To what degree the affected food webs will respond or can recover from this form of exploitation remains to be seen, despite the fact that the CCAMLR articles require that exploited populations are not so badly affected that they cannot recover in 20–30 years. With rapid climate change now one of fishery management's challenges (Cheung and others 2010, 2012), the latter requirement may have become especially problematic.

### Acknowledgements

This summary was prepared under NSF grant ANT-0944411; views expressed here do not necessarily represent those of NSF. The authors acknowledge support from *The Sea Around Us*, a research collaboration between the University of British Columbia and the Pew Environment Group. They thank A. Ulman for her assistance in preparing figures and tables, and P. Penhale, P. Trathan, T. Branch and an anonymous reviewer for very helpful comments that improved the article greatly.

### References

- Ainley, D.G. 2010. A history of the exploitation of the Ross Sea, Antarctica. *Polar Record* 46: 233–243.
- Ainley, D.G., and L.K. Blight. 2009. Ecological repercussions of historical fish extraction from the Southern Ocean. *Fish and Fisheries* 10: 13–38.
- Ainley, D.G., and C.M. Brooks. 2013. Exploiting the Southern Ocean: rational use or reversion to tragedy of the commons? In: Liggett, D., and A.D. Hemmings (editors). *Exploring Antarctic values*. Christchurch: University of Canterbury, Gateway Antarctica Special Publication Series, in press.
- Ainley, D.G., N. Nur, J.T. Eastman, G. Ballard, C.L. Parkinson, C.W. Evans and A.L. DeVries. 2012. Decadal trends in abundance, size and condition of Antarctic toothfish in McMurdo Sound, Antarctica, 1972–2011. *Fish and Fisheries* (doi:10.1111/j.1467–2979.2012.00474.x).
- Aquarone, M.C., and S. Adams. 2008. In: Sherman, K., and G. Hempel (editors). The UNEP large marine ecosystem report: a perspective on changing conditions in LMEs of the world's regional seas. *UNEP Regional Seas Reports and Studies* No. 182: 765–773.
- Arana, P.M., and R. Vega. 1999. Exploratory fishing for *Dissostichus* spp. in the Antarctic region (subareas 48.1, 48.2 and 88.3). *CCAMLR Science* 6: 1–17.
- Atkinson, A., V. Siegel, E. Pakhomov and P. Rothery. 2004. Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature* 432: 100–103.
- Ballance, L., R.L. Pitman, R.P. Hewitt, D.B. Siniff, W.Z. Trivelpiece, P.J. Clapham and R.L. Brownell, Jr. 2006. The removal of large whales from the Southern Ocean: evidence for long-term ecosystem effects? In: Estes, J.A., D.P. Demaster, D.F. Doak, T.E. Williams and R.L. Brownell, Jr. (editors). *Whales, whaling and ocean ecosystems*. Berkeley, CA: University of California Press: 215–230.
- Baum, J.K., and B. Worm. 2009. Cascading top-down effects of changing oceanic predator abundances. *Journal of Animal Ecology* 78: 699–714.
- Boyd, I.L. 2002. Estimating food consumption of marine predators: Antarctic fur seals and Macaroni penguins. *Journal of Applied Ecology* 39: 103–119.
- Branch, T.A. 2011. Humpback abundance south of 60°S from three complete circumpolar sets of surveys. *Journal of Cetacean Research and Management* (special issue) 3: 53–69.
- Branch, T.A., R. Watson, E.A. Fulton, S. Jennings, C.R. McGilliard, G.T. Pablico, D. Ricard and S.R. Tracey. 2010. The trophic fingerprint of marine fisheries. *Nature* 468: 431–435.
- Brown, M.R., and R.L. Brownell. 2001. Review of catches of great whales taken in the proposed South Pacific sanctuary region. *Paper SC52/033*. Cambridge: International Whaling Commission.
- CCAMLR (Convention for the Conservation of Antarctic Marine Living Resources). 2006. Report of the twenty-fifth meeting of the Scientific Committee of the Convention for the Conservation of Antarctic Marine Living Resources. (SC-CCAMLR-XXV). Hobart: CCAMLR.
- CCAMLR. 2011. Conservation measures in force 2011/12. Hobart: CCAMLR. URL: [http://www.ccamlr.org/pu/e/e\\_pubs/cm/11–12/toc.htm](http://www.ccamlr.org/pu/e/e_pubs/cm/11–12/toc.htm) (accessed October 2012).
- CCAMLR. 1990–2012. CCAMLR Statistical Bulletin, Vols 1, 2, 5, 13, 21 and 24. Hobart: CCAMLR. URL: [http://archive.ccamlr.org/pu/e/e\\_pubs/sb/intro.htm](http://archive.ccamlr.org/pu/e/e_pubs/sb/intro.htm) (accessed October 2012).
- Cheung, W.W.L., V.W.Y. Lam, J.L. Sarmiento, K. Kearney, R. Watson, D. Zeller and D. Pauly. 2010. Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Global Change Biology* 16: 24–35.
- Cheung, W.W.L., J.L. Sarmiento, J. Dunne, T.L. Frölicher, V. Lam, M.L.D. Palomares, R. Watson and D. Pauly. 2012. Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nature Climate Change* (doi:10.1038/nclimate1691).

- Constable, A.J., W.K. de la Mare, D.J. Agnew, I. Everson and D. Miller. 2000. Managing fisheries to conserve the Antarctic marine ecosystem: practical implementation of the Convention on the Conservation of Antarctic Marine Living Resources (CCAMLR). *ICES Journal of Marine Science* 57: 778–791.
- Dolzhenkov, V.N., I.A. Zhigalov, E.A. Kovalev, V.P. Timonin and V.A. Spiridonov. 1990. The distribution pattern and fishery for the Antarctic krill (*Euphausia superba*) off the Wilkes Land and Balleny Islands (with notes on the application of CPUE data as indices of krill abundance). CCAMLR document WG-KRILL-90/16. Hobart: CCAMLR. URL: <http://www.ccamlr.org/en/wg-krill-90/16> (accessed October 2012).
- Essington, T.E., A.H. Beaudreau and J. Wiedenmann. 2006. Fishing through marine food webs. *Proceeding of the National Academy of Sciences* 103: 3171–3175.
- Estes, J.A., J. Terborgh, J.S. Brashares, M.E. Power, J. Berger, W.J. Bond, S.R. Carpenter, T.E. Essington, R.D. Holt, J.B.C. Jackson, R.J. Marquis, L. Oksanen, T. Oksanen, R.T. Paine, E.K. Pikitch, W. Ripple, S.A. Sandin, M. Scheffer, T.W. Schoener, J.B. Shurin, A.R.E. Sinclair, M.E. Soulé, R. Virtanen and D.A. Wardle. 2011. Trophic downgrading of planet Earth. *Science* 333: 301–306.
- FAO (Food and Agriculture Organization). 2005. Species fact sheet *Euphausia superba*. Rome: Food and Agriculture Organization of the United Nations.
- Fenaughty, J.M., D.W. Stevens and S.L. Hanchet. 2003. Diet of the Antarctic toothfish (*Dissostichus mawsoni*) from the Ross Sea, Antarctica, CCAMLR statistical subarea 88.1). *CCAMLR Science* 10: 113–123.
- FishBase 2012. <http://www.fishbase.org> (accessed 15 Oct 2012).
- Fogg, G.E. 1992. *The history of Antarctic science*. Cambridge: Cambridge University Press.
- Gjøsaeter, J., and K. Kawaguchi. 1980. A review of the world resources of mesopelagic fish. *Fisheries Technical Paper 193*. Rome: Food and Agriculture Organization of the United Nations.
- Halpern, B.S., S. Walbridge, K.A. Selkoe, C.B. Kappel, F. Micheli, C. D'Agrosa, J.F. Bruno, K.S. Casey, C. Ebert, H.E. Fox, R. Fujita, D. Heinemann, H.S. Lenihan, E.M.P. Madin, M.T. Perry, E.R. Selig, M. Spalding, R. Steneck and R. Watson. 2008. A global map of human impact on marine ecosystems. *Science* 319: 948–951.
- Hardin, G. 1968. The tragedy of the commons. *Science* 162: 1243–1248.
- Hilborn, R., T.A. Branch, B. Ernst, A. Magnusson, C.V. Minte-Vera, M.D. Scheuerell and J.L. Valero. 2003. State of the world's fisheries. *Annual Review of Environmental Resources* 28: 359–399.
- IUCN (International Union for Conservation of Nature and Natural Resources). 2011. Red list of threatened species. Version 2011.2. URL: <http://www.iucnredlist.org> (accessed 20 February 2012).
- Kareiva, P., C. Yuan-Farrell and C. O'Conner. 2006. Whales are big and it matters. In: J.A. Estes, D.P. DeMaster, D.F. Doak, T.M. Williams and R.L. Brownell, Jr. (editors). *Whales, whaling, and ocean ecosystems*. Berkeley: University of California Press: 379–387.
- Kelleher, K. 2005. Discards in the world's marine fisheries: an update. *FAO Fisheries Technical Paper 470*. Rome: Food and Agriculture Organization of the United Nations.
- Kock, K.H. 1992. *Antarctic fish and fisheries*. New York: Cambridge University Press.
- Leaper, R., and C. Miller. 2011. Management of Antarctic baleen whales amid past exploitation, current threats and complex marine ecosystems. *Antarctic Science* 23: 503–529.
- Marschoff, E.R., E.R. Barrera-Oro, N.S. Aleccio and D.G. Ainley. 2012. Slow recovery of previously depleted demersal fish at the South Shetland Islands, 1983–2010. *Fisheries Research* 125/126: 206–213.
- Nicol, S., A. Bowie, S. Jarman, D. Lannuzel, K.M. Meiners and P. van der Merwe. 2010. Southern Ocean iron fertilization by baleen whales and Antarctic krill. *Fish and Fisheries* 11: 203–209.
- O'Gorman, F.A. 1961. Fur seals breeding in the Falkland Islands Dependencies. *Nature* 192: 914–916.
- Pakhomov, E.A., and S.A. Pankratov. 1994. By-catch, growth and feeding of Antarctic juvenile fish taken in krill (*Euphausia superba* Dana) fisheries in the South Georgia area, in 1992. *CCAMLR Science* 1: 129–142.
- Pauly, D., V. Christiansen, J. Dalsgaard, R. Froese and F. Torres, Jr. 1998. Fishing down marine food webs. *Science* 279: 860–863.
- Pauly, D., and R. Watson. 2005. Background and interpretation of the 'Marine Trophic Index' as a measure of biodiversity. *Philosophical Transactions of the Royal Society B* 360: 415–423.
- Pauly, D., R. Watson and J. Alder. 2005. Global trends in world fisheries: impacts on marine ecosystems and food security. *Philosophical Transactions of the Royal Society B* 360: 5–12.
- Pikitch, E., P.D. Boersma, I.L. Boyd, D.O. Conover, P. Cury, T. Essington, S.S. Heppell, E.D. Houde, M. Mangel, D. Pauly, É. Plagányi, K. Sainsbury and R.S. Steneck. 2012. *Little fish, big impact: managing a crucial link in ocean food webs*. Washington, D.C.: Lenfest Ocean Program.
- Rosenberg, A.A., J.R. Beddington and M. Basson. 1986. Growth and longevity of krill during the first decade of pelagic whaling. *Nature* 324: 152–154.
- Sea Around US Project. 2012. <http://www.seaaroundus.org> (accessed 15 Oct 2012).
- SeaLifeBase. 2012. <http://www.sealifebase.org> (accessed 15 Oct 2012).
- Swartz, W., E. Sala, S. Tracey, R. Watson and D.L. Pauly. 2010. The spatial expansion and ecological footprint of fisheries (1950 to present). *PLoS ONE* 5 (12): e15143 (doi:10.1371/journal.pone.0015143).
- Tin, T., Z.L. Fleming, K.A. Hughes, D.G. Ainley, P. Convey, C.A. Moreno, S. Pfeiffer, J. Scott and I. Snape. 2009. Impacts of local human activities on the Antarctic environment. *Antarctic Science* 21: 3–33.
- Tønnessen, J.N., and A.O. Johnsen. 1982. *The history of modern whaling*. London: C. Hurst and Co.
- Trites, A.W., and D. Pauly. 1998. Estimating mean body masses of marine mammals from maximum body lengths. *Canadian Journal of Zoology* 76: 886–896.
- Trivelpiece, W.Z., J.T. Hinke, A.K. Miller, C.S. Reiss, S.G. Trivelpiece and G.M. Watters. 2011. Variability in krill biomass links harvesting and climate warming to penguin population changes in Antarctica. *Proceedings of the National Academy of Sciences* 108 (18): 7625–7628.
- Watters, G. 1996. By-catch of fishes captured by the krill fishing vessel Chiyo Maru No. 2 in statistical area 58 (January to March 1995). *CCAMLR Science* 3: 111–123.
- Zeller, D., S. Booth, E. Pakhomov, W. Swartz and D. Pauly. 2011. Arctic fisheries catches in Russia, USA and Canada: baselines for neglected ecosystems. *Polar Biology* 34: 955–973.

## Appendices

Table A1. Catches (t) by taxon in the parts of FAO area 48 covered in this article (that is, subareas 48.1, 48.5 and the southern portion of 48.6; note that until 1990, all of these catches were from 48.1). The 'Other icefish' category includes blackfin icefish, ocellated icefish, spiny icefish, South Georgia icefish and other crocodile icefish categories

Year	Antarctic silverfish	Grey rockcod	Humped rockcod	Antarctic krill	Lanternfish	Marbled rockcod	Other icefish	Marine fishes n.e.i.
1976	-	-	-	276	-	-	0	-
1977	-	-	-	92	-	-	397	-
1978	-	-	-	-	-	-	-	-
1979	-	-	3,280	-	-	470	-	79
1980	-	-	765	49,439	48	18,763	47,733	551
1981	-	36	50	92,859	-	-	2,586	4,230
1982	-	-	-	69,240	-	-	1,700	-
1983	-	110	-	531	-	-	-	16
1984	-	-	-	32,863	-	-	2,604	-
1985	-	-	-	11,592	-	-	1	-
1986	-	-	-	42,113	-	-	37	1
1987	4	-	54	70,544	-	-	-	-
1988	-	-	11	78,923	304	-	75	2
1989	-	-	667	64,671	17	-	2	-
1990	-	-	-	42,477	-	-	142	-
1991	-	-	-	64,671	-	-	-	-
1992	-	-	-	78,385	50	-	-	-
1993	-	-	-	45,093	-	-	-	-
1994	-	-	-	35,025	-	-	-	-
1995	-	-	-	62,384	-	-	-	-
1996	-	-	-	51,285	-	-	-	-
1997	-	-	-	49,837	-	-	-	-
1998	-	-	-	38,897	-	-	-	-
1999	-	-	-	62,588	-	-	-	-
2000	1	-	-	52,559	-	-	-	-
2001	-	-	-	10,646	-	-	-	-
2002	-	-	-	35,377	-	-	-	-
2003	-	-	-	13,881	-	-	-	-
2004	-	-	-	7,096	-	-	-	-
2005	14	-	-	88,876	-	-	-	-
2006	-	-	-	18,419	-	-	17	-
2007	-	-	-	2,884	-	-	-	-
2008	-	-	-	33,971	-	-	-	-
2009	-	-	-	153,682	-	-	-	-
2010	-	-	-	9,209	-	-	-	-

Table A2. Total catches (t) by country in the parts of FAO area 48 covered in this article (that is, subareas 48.1, 48.5 and 48.6, catches from the shelf area only)

Year	Chile	Japan	South Korea	Norway	Poland	Ukraine	Former USSR	USA
1976	276	-	-	-	-	-	-	-
1977	92	-	-	-	-	-	-	-
1979 <sup>ab</sup>	-	-	-	-	20,596	28,319	15	-
1980 <sup>a</sup>	-	-	-	-	2,675	717	68,722	-
1981	-	3,751	-	-	-	1,700	93,641	-
1982	-	4,978	-	-	-	-	64,045	-
1983	396	96	-	-	-	2,604	900	-
1984	1,649	30,479	-	-	-	-	-	-
1985	2,598	9,032	-	-	-	-	-	-
1986	3,264	36,874	-	-	1,976	-	-	-
1987 <sup>c</sup>	4,063	64,483	1,503	-	-	60	373	-
1988 <sup>b</sup>	5,938	71,814	1,111	-	57	-	408	-
1989	5,329	75,912	1,615	-	1,827	140	21,557	-
1990	4,501	33,936	4,040	-	-	-	-	-
1991	3,679	54,720	1,211	-	310	-	4,751	-
1992 <sup>d</sup>	6,066	61,631	519	-	641	636	8,925	-
1993	3,834	41,251	-	-	-	-	-	-
1994	-	29,070	-	-	1,278	4,677	-	-
1995	-	45,719	-	-	14,927	1,738	-	-
1996 <sup>e</sup>	-	37,479	-	-	13,498	-	-	-
1997 <sup>e</sup>	-	34,430	890	-	13,883	-	-	-
1998 <sup>f</sup>	-	26,106	-	-	8,151	-	-	-
1999	-	39,952	4,677	-	17,959	-	-	-
2000	-	39,553	5,781	-	2,302	3,362	-	1,562
2001	-	9,207	-	-	-	-	-	1,439
2002	-	27,094	2,360	-	2,783	324	-	2,816
2003 <sup>g</sup>	-	5,069	1,608	-	1,148	-	-	903
2004	-	6,090	142	-	436	413	-	15
2005 <sup>bd</sup>	-	19,756	42,386	8,360	5,496	12,878	-	-
2006	-	1,608	11,636	2,868	2,307	-	-	-
2007	-	435	-	2,449	-	-	-	-
2008	-	8,897	19,526	2,198	3,350	-	-	-
2009 <sup>h</sup>	-	28,924	41,863	75,803	6,605	-	-	-
2010 <sup>h</sup>	-	222	4,999	1,417	489	-	-	-

<sup>a</sup>Bulgaria had 25t of catches in 1979 and 112t in 1980.

<sup>b</sup>Germany had 3,032t of catches in 1979, 13t in 1988 and 1t in 2005.

<sup>c</sup>Spain had 200t of catches in 1987.

<sup>d</sup>The Russian Federation had 50t of catches in 1992 and 30t in 2005.

<sup>e</sup>The United Kingdom had 308t of catches in 1996 and 634t in 1997.

<sup>f</sup>Argentina had 4,640t of catches in 1998.

<sup>g</sup>Vanuatu had 5,153t of catches in 2003.

<sup>h</sup>China had 67t of catches in 2009 and 2,082t in 2010.

Table A3. Catches (t) by taxon in the parts of FAO area 88 covered in this article (that is, catches in subareas 88.1–88.3 and catches in area 88 not allocated to any subarea)

Year	Antarctic krill	Antarctic silverfish	Antarctic toothfish	Crocodile icefish	Eel cods	Mackerel icefish	Rattails	Rays, skates	Misc. marine fish
1977	3,355	-	-	-	-	-	-	-	-
1978	554	21	-	-	-	-	-	-	2
1979	2,862	-	-	-	-	-	-	-	200
1980	3,181	-	-	-	-	-	-	-	-
1981	4,234	1,517	-	-	-	-	-	-	-
1982	6,637	90	-	-	-	15	-	-	-
1983	10,637	-	-	-	-	-	-	-	-
1984 <sup>a</sup>	641	-	-	-	-	-	-	-	2
1985	4,721	-	-	-	-	-	-	-	-
1986	3,892	-	-	-	-	-	-	-	-
1987	394	-	-	-	-	-	-	-	-
1989 <sup>a</sup>	658	-	-	-	-	-	-	-	-
1990	2,150	-	-	-	-	-	-	-	-
1991	3	-	-	-	-	-	-	-	-
1992	50	-	-	-	-	-	-	-	-
1997	-	-	41	-	-	-	9	5	-
1998	-	-	296	-	5	-	22	18	-
1999	-	-	751	-	7	-	70	41	-
2000	-	-	624	2	3	-	48	9	-
2001	-	-	1,446	2	6	-	158	25	-
2002	-	-	1,912	9	4	-	75	17	-
2003	-	-	2,797	3	14	-	339	23	-
2004	-	-	3,534	3	5	-	361	63	-
2005	-	-	3,481	3	2	-	132	4	-
2006	-	-	3,425	11	21	-	200	35	-
2007	-	1	2,851	5	8	-	113	7	-
2008	-	-	2,915	0	3	-	130	7	-
2009	-	-	3,182	3	4	-	128	6	-
2010	-	-	3,460	1	4	-	133	6	-

<sup>a</sup>Lanternfish was reported for 129t of catches in 1984 and 1,110t in 1989.

Table A4. Total catches (t) by country in the parts of FAO area 88 covered in this article (that is, catches in subareas 88.1–88.3 and catches in which the subarea was not recorded)

Year	Argentina	Japan	South Korea	New Zealand	Norway	Russian Federation	South Africa	Spain	Former USSR	United Kingdom	Uruguay	IUU <sup>a</sup>
1977	-	-	-	-	-	-	-	-	3,355	-	-	-
1978 <sup>b</sup>	-	518	-	-	-	-	-	-	21	-	-	-
1979	-	2,262	-	-	-	-	-	-	800	-	-	-
1980	-	3,134	-	-	-	-	-	-	47	-	-	-
1981	-	1,154	-	-	-	-	-	-	4,597	-	-	-
1982	-	2,544	-	-	-	-	-	-	4,198	-	-	-
1983	-	4,718	-	-	-	-	-	-	5,919	-	-	-
1984	-	626	-	-	-	-	-	-	146	-	-	-
1985	-	4,721	-	-	-	-	-	-	-	-	-	-
1986	-	2,008	-	-	-	-	-	-	1,884	-	-	-
1987	-	106	-	-	-	-	-	-	288	-	-	-
1989–1999	-	<sup>c</sup>	-	<sup>d</sup>	-	-	-	-	<sup>e</sup>	-	-	-
2000	-	-	-	642	-	-	21	-	-	-	23	-
2001	-	-	-	1,545	-	-	-	-	-	-	-	92
2002	-	-	-	1,155	-	712	150	-	-	-	-	-
2003 <sup>f,g</sup>	225	-	108	1,306	179	264	125	118	-	31	189	240
2004	284	-	-	1,971	292	669	-	1	-	318	407	23
2005	261	-	-	1,480	385	718	-	-	-	408	370	-
2006	256	-	465	1,220	326	605	51	-	-	511	258	-
2007	-	-	425	1,135	-	288	121	44	-	711	75	186
2008 <sup>h</sup>	-	-	657	910	-	-	140	385	-	762	99	-
2009	38	-	1,025	1,396	-	1	-	363	-	500	-	-
2010	-	-	800	1,172	1	459	-	427	-	722	23	-

<sup>a</sup>IUU not reported to country level.

<sup>b</sup>Poland had a catch of 38t in 1978.

<sup>c</sup>Japan had 3t of catch in 1991 and 50t in 1992.

<sup>d</sup>New Zealand had 55t of catch in 1997, 341t in 1998 and 869t in 1999.

<sup>e</sup>The former USSR had 1,768t of catch in 1989 and 2,150t in 1990.

<sup>f</sup>Ukraine had catches of 158t in 2003 and 1t in 2004.

<sup>g</sup>The USA had a catch of 233t in 2003.

<sup>h</sup>Chile had a catch of 102t in 2008.

Table A5. Catches (t) by taxon in the parts of FAO area 58 covered in this article (that is, subareas/divisions 58.4.1 and 58.4.2)

Year	Antarctic toothfish	Antarctic krill	Rattailsa	Antarctic silverfish	Rays, skatesa
1974	-	283	-	-	-
1975	-	2,642	-	-	-
1976	-	4,399	-	-	-
1977	-	11,991	-	-	-
1978	-	24,685	-	-	-
1979	-	34,699	-	-	-
1980	-	33,046	-	-	-
1981	-	22,793	-	-	-
1982	-	27,168	-	-	-
1983	-	32,066	-	307	-
1984	-	8,195	-	-	-
1985	-	2,249	-	384	-
1986	-	5,262	-	653	-
1987	-	3,974	-	30	-
1988	-	6,490	-	67	-
1989	-	217	-	44	-
1990	-	29,753	-	-	-
1991	-	1,329	-	-	-
1992	-	5,762	-	-	-
1993	-	899	-	-	-
1994	-	1,266	-	-	-
2002	215	-	12	-	-
2003	217	-	1	-	-
2004	690	-	32	-	3
2005	1,372	-	20	-	-
2006	1,577	-	49	-	1
2007	762	-	45	-	1
2008	616	-	9	-	-
2009	1,629	-	5	-	-
2010	350	-	8	-	-

<sup>a</sup>Rattails, grenadiers and rays are by-catch.

Table A6. Catches (t) by country in the parts of FAO area 58 covered in this article (that is, subareas/divisions 58.4.1 and 58.4.2)

Year	Japan	South Korea	Spain	Former USSR	Uruguay	IUU <sup>h</sup>
1974–1984	a	-	-	b	-	-
1985	2,249	-	-	384	-	-
1986	5,262	-	-	653	-	-
1987	3,974	-	-	30	-	-
1988	-	-	-	6,557	-	-
1989	-	-	-	261	-	-
1990 <sup>c</sup>	28,250	-	-	-	-	-
1991	1,329	-	-	-	-	-
1992	5,762	-	-	-	-	-
1993	899	-	-	-	-	-
1994	1,266	-	-	-	-	-
2002 <sup>d</sup>	-	-	-	-	-	98
2003 <sup>d</sup>	-	-	-	-	-	197
2004 <sup>ef</sup>	-	226	154	-	-	86
2005 <sup>e</sup>	-	294	235	-	3	789
2006 <sup>g</sup>	-	343	243	-	37	914
2007 <sup>g</sup>	-	389	73	-	10	136
2008	20	226	-	-	51	328
2009	86	206	-	-	-	1,342
2010	5	277	76	-	-	-

<sup>a</sup>In 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983 and 1984, Japan had 283t, 2,642t, 4,399t, 11,991t, 24,685t, 34,699t, 33,046t, 22,793t, 27,168t, 32,066t and 8,195t of catches, respectively.

<sup>b</sup>In 1983, the former USSR had 307t of catches.

<sup>c</sup>In 1990, Ukraine had 1,503t of catches.

<sup>d</sup>In 2002 and 2003, Australia had 129t and 21t of catches, respectively.

<sup>e</sup>In 2004 and 2005, Chile had 196t and 71t of catches, respectively.

<sup>f</sup>In 2004, New Zealand had 63t of catches.

<sup>g</sup>In 2006 and 2007, Namibia had 90t and 200t of catches, respectively.

<sup>h</sup>Toothfish.