

PARITY BIAS IN FUNDAMENTAL UNITS OF REAL QUADRATIC FIELDS

FLORIAN BREUER  and CAMERON SHAW-CARMODY 

(Received 6 September 2024; accepted 27 October 2024)

Abstract

We compute primes $p \equiv 5 \pmod{8}$ up to 10^{11} for which the Pellian equation $x^2 - py^2 = -4$ has no solutions in odd integers; these are the members of sequence A130229 in the Online Encyclopedia of Integer Sequences. We find that the number of such primes $p \leq x$ is well approximated by

$$\frac{1}{12}\pi(x) - 0.037 \int_2^x \frac{dt}{t^{1/6} \log t},$$

where $\pi(x)$ is the usual prime counting function. The second term shows a surprising bias away from membership of this sequence.

2020 *Mathematics subject classification*: primary 11D09; secondary 11R27, 11Y65.

Keywords and phrases: Pellian equation, fundamental unit, prime, OEIS A130229.

1. Introduction

For a prime $p \equiv 5 \pmod{8}$, consider the real quadratic field $K = \mathbb{Q}(\sqrt{p})$, with ring of integers $\mathcal{O}_K = \mathbb{Z}[\frac{1}{2}(1 + \sqrt{p})]$ and fundamental unit $\varepsilon_p = \frac{1}{2}(x_0 + y_0\sqrt{p}) > 1$. Then, (x_0, y_0) is a fundamental solution to the Pellian equation

$$x^2 - py^2 = -4. \tag{1.1}$$

The prime 2 is inert in K/\mathbb{Q} , and $\varepsilon_p \equiv 1 \pmod{2\mathcal{O}_K}$ if and only if (1.1) has no odd integer solutions. Primes $p \equiv 5 \pmod{8}$ satisfying the above equivalent conditions define sequence A130229 in [5]. They also appear in [1, 2, 7].

Since $\varepsilon_p \pmod{2\mathcal{O}_K}$ can take any of three nonzero values in $\mathcal{O}_K/2\mathcal{O}_K \cong \mathbb{F}_4$, it is reasonable to expect roughly one third of all primes $p \equiv 5 \pmod{8}$ to be members of this sequence.

The first author was supported by the Alexander von Humboldt Foundation. The second author was supported by a 2020–2021 Vacation Research Scholarship from the Australian Mathematical Sciences Institute (AMSI).

© The Author(s), 2025. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.

Define

$$\chi(p) := \begin{cases} 1 & \text{if } p \equiv 5 \pmod 8 \text{ and } \varepsilon_p \equiv 1 \pmod{2O_K}, \\ -\frac{1}{2} & \text{if } p \equiv 5 \pmod 8 \text{ and } \varepsilon_p \not\equiv 1 \pmod{2O_K}, \\ 0 & \text{if } p \not\equiv 5 \pmod 8, \end{cases}$$

and define the modified counting function

$$\theta_\chi(x) := \sum_{p \leq x} \chi(p) \log p.$$

Then, the above heuristic leads us to expect $\theta_\chi(x) = o(x)$ as $x \rightarrow \infty$.

In this note, we report on computations of $\theta_\chi(x)$ for $x \leq 10^{11}$, which show a surprising bias away from the $\varepsilon_p \equiv 1 \pmod{2O_K}$ case, hinted at in related computations reported in [1, Section 4]. We thus pose a conjecture.

CONJECTURE 1.1. There exists a constant $c \approx -0.066$ for which

$$\theta_\chi(x) \sim cx^{5/6}$$

as $x \rightarrow \infty$.

2. Results

We computed ε_p using the continued fraction method in [3, Section 3.3], with the modification that B_i and G_i are only computed modulo 2, since we only need to know the parity of ε_p . This significantly reduces the memory requirements of the calculation.

We implemented the algorithm to run on a GPU using the PYTHON Numba library [4]. The final computation for all $p < 10^{11}$ took approximately 17 hours on an entry-level gaming laptop with an Nvidia RTX 3050 GPU. The source code and data are available at <https://github.com/florianbreuer/A130229>.

Table 1 lists some values for the naive counting function

$$\pi_1(x) = \sum_{p \leq x, \chi(p)=1} 1.$$

However, it is advantageous to study the ‘smoothed’ counting function $\theta_\chi(x) = \sum_{p \leq x} \chi(p) \log p$. Figure 1 plots $-\theta_\chi(x)$ for $x \leq 10^{11}$ on logarithmic axes. The plot approximates a straight line with slope 5/6. The least squares best fit of the form $f(x) = cx^{5/6}$ is found to have $c \approx -0.06626$, computed using the `find_fit` method in SAGEMATH v9.3 [6]. The error term $\theta_\chi(x) - cx^{5/6}$ is shown in Figure 2. This provides evidence for Conjecture 1.1. Moreover, it appears likely that the error is of the order $O(x^{1/2+\varepsilon})$.

From this, we may also deduce a good approximation for $\pi_1(x)$. Define

$$\pi_{-1/2}(x) = \sum_{p \leq x, \chi(p)=-1/2} 1 \quad \text{and} \quad \pi_\chi(x) = \sum_{p \leq x} \chi(p) = \pi_1(x) - \frac{1}{2}\pi_{-1/2}(x).$$

TABLE 1. Some values of the counting function $\pi_1(x)$ for sequence A130229.

x	$\pi_1(x)$	Approximation	x	$\pi_1(x)$	Approximation
10^2	1	1	2×10^{10}	72770931	72761719
10^3	15	11	3×10^{10}	107298975	107293481
10^4	98	90	4×10^{10}	141363308	141357259
10^5	741	735	5×10^{10}	175085540	175080418
10^6	6200	6187	6×10^{10}	208542967	208537579
10^7	53382	53348	7×10^{10}	241775700	241776120
10^8	468223	468144	8×10^{10}	274823028	274829667
10^9	4164936	4165422	9×10^{10}	307723656	307723171
10^{10}	37490293	37483463	10^{11}	340472393	340476359




FIGURE 1. Log-log plot of $-\theta_\chi(x)$ for $x \leq 10^{11}$.




FIGURE 2. Plot of the error term $\theta_\chi(x) - cx^{5/6}$ for $c \approx -0.06626$.

Then, $\theta_\chi(x) \sim cx^{5/6} \approx c \cdot \frac{5}{6} \int_2^x t^{-1/6} dt$ suggests

$$\pi_\chi(x) \approx c \cdot \frac{5}{6} \int_2^x \frac{t^{-1/6}}{\log t} dt \sim c \frac{x^{5/6}}{\log x}.$$

Then, from $\pi_1(x) + \pi_{-1/2}(x) \approx \frac{1}{4}\pi(x)$, where $\pi(x)$ is the usual prime counting function, we arrive at

$$\pi_1(x) \approx \frac{1}{12}\pi(x) + \frac{2}{3}\pi_\chi(x) \approx \frac{1}{12}\pi(x) + c \cdot \frac{5}{9} \int_2^x \frac{t^{-1/6}}{\log t} dt.$$

These approximations are compared with the computed values of $\pi_1(x)$ in Table 1.

Acknowledgements

The first author would like to thank James Punch for useful discussions, Frank Calegari for suggesting that the correct exponent is likely to be $5/6$ and GPT-4 for help with porting the algorithm to a GPU.

References

- [1] F. Breuer, ‘Periods of Ducci sequences and odd solutions to a Pellian equation’, *Bull. Aust. Math. Soc.* **100** (2019), 201–205.
- [2] F. Breuer, ‘Multiplicative orders of Gauss periods and the arithmetic of real quadratic fields’, *Finite Fields Appl.* **73** (2021), Article no. 101848.
- [3] M. J. Jacobson and H. C. Williams, *Solving the Pell Equation*, CMS Books in Mathematics (Springer, New York, 2009).
- [4] S. T. Lam, A. Pitrou and S. Seibert, ‘Numba: a LLVM-based Python JIT compiler’, in: *LLVM’15: Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC*, ACM Digital Library (ed. H. Finkel) (Association for Computing Machinery, New York, 2015), Article no. 7, 6 pages.
- [5] *The On-line Encyclopedia of Integer Sequences*, entry #A130229, <https://oeis.org/A130229>.
- [6] The Sage Developers, SageMath, the Sage Mathematics Software System (Version 9.3), 2021. <https://www.sagemath.org>.
- [7] J. Xue, T.-C. Yang and C.-F. Yu, ‘Numerical invariants of totally imaginary quadratic $\mathbb{Z}[\sqrt{p}]$ -orders’, *Taiwanese J. Math.* **20**(4) (2016), 723–741.

FLORIAN BREUER, School of Information and Physical Sciences,
University of Newcastle, Callaghan, New South Wales 2308, Australia
e-mail: florian.breuer@newcastle.edu.au

CAMERON SHAW-CARMODY, School of Information and Physical Sciences,
University of Newcastle, Callaghan, New South Wales 2308, Australia
e-mail: cameron.shaw-carmody@newcastle.edu.au