

TRADE-OFFS BETWEEN BIOMASS USE AND SOIL COVER. THE CASE OF RICE-BASED CROPPING SYSTEMS IN THE LAKE ALAOTRA REGION OF MADAGASCAR

By K. NAUDIN^{†,‡}, E. SCOPEL[†], A. L. H. ANDRIAMANDROSO[§],
M. RAKOTOSOLOFO[§], N. R. S. ANDRIAMAROSOA RATSIMBAZAFY[§],
J. N. RAKOTOZANDRINY[§], P. SALGADO[¶] and K. E. GILLER^{††}

[†]CIRAD, UPR Systèmes de Culture Annuels, F- 34398 Montpellier, France, [§]Université
d'Antananarivo, ESSA, Département « Élevage », 101, Antananarivo, Madagascar, [¶]CIRAD,
UMR Systèmes d'Élevage Méditerranéens et Tropicaux, F-97410 Saint-Pierre, La Réunion,
France and ^{††}Plant Production Systems, Wageningen University, P.O. Box 430, 6700 AK
Wageningen, The Netherlands

(Accepted 14 September 2011; First published online 20 October 2011)

SUMMARY

Farmers in the Lake Alaotra region of Madagascar are currently evaluating a range of conservation agriculture (CA) cropping systems. Most of the expected agroecological functions of CA (weed control, erosion control and water retention) are related to the degree of soil cover. Under farmers' conditions, the grain and biomass productivity of these systems is highly variable and the biomass is used for several purposes. In this study, we measured biomass production of cover crops and crops in farmers' fields. Further, we derived relationships to predict the soil cover that can be generated for a particular quantity of mulch. We used these relationships to explore the variability of soil cover that can be generated in farmers' fields, and to estimate how much of the biomass can be removed for use as livestock feed, while retaining sufficient soil cover. Three different kinds of cropping systems were investigated in 91 farmers' fields. The first two cropping sequences were on the hillsides: (i) maize + pulse (*Vigna unguiculata* or *Dolichos lablab*) in year 1, followed by upland rice in year 2; (ii) the second crop sequence included several years of *Stylosanthes guianensis* followed by upland rice; (iii) the third crop sequence was in lowland paddy fields: *Vicia villosa* or *D. lablab*, which was followed by rice within the same year and repeated every year. The biomass available prior to rice sowing varied from 3.6 t ha⁻¹ with *S. guianensis* to 7.3 t ha⁻¹ with *V. villosa*. The relationship between the mulch quantity (M) and soil cover (C) was measured using digital imaging and was well described by the following equation: $C = 1 - \exp(-A_m \times M)$, where A_m is an area-to-mass ratio with $R^2 > 0.99$ in all cases. The calculated average soil cover varied from 56 to 97% for maize + *V. unguiculata* and *V. villosa*, respectively. In order to maintain 90% soil cover at rice sowing, the average amount of biomass of *V. villosa* that could be removed was at least 3 t ha⁻¹ for three quarters of the fields. This quantity was less for other annual or biennial cropping systems. On average the *V. villosa* aboveground biomass contained 236 kg N ha⁻¹. The study showed that for the conditions of farmers of Malagasy, the production and conservation of biomass is not always sufficient to fulfil all the above-cited agroecological functions of mulch. Inventory of the soil cover capacity for different types of mulch may help farmers to decide how much biomass they can remove from the field.

INTRODUCTION

Conservation agriculture (CA) is defined by three principles: minimum soil disturbance, permanent organic mulch covering the soil and diversified crop rotations

[‡]Corresponding author. Email: krishna.naudin@cirad.fr

and associations (Food and Agriculture Organization (FAO), 2010a; Hobbs, 2007; Reicosky, 2008). Mulch plays an important role in CA benefits. In particular, soil cover acts on (i) weeds control, (ii) erosion control and (iii) improvement of crop–water balance. Weed control, besides allelopathic effects, results from physical effects of mulch on temperature, light extinction and physical obstruction of weed seedling emergence (Bilalis *et al.*, 2003; Teasdale and Mohler, 1993, 2000). The percentage of ground cover has more direct influence than the quantity of biomass on weed emergence (Teasdale and Mohler, 2000), on erosion control (Smets *et al.*, 2008) and on improvements in the crop–water balance (Scopel *et al.*, 2004). By contrast, other benefits of mulch, such as contributions to increase soil carbon contents (Neto *et al.*, 2010) or provision of nitrogen for subsequent crop growth (Maltas *et al.*, 2009), are directly proportional to the amount of mulch and its content of each element.

In the Lake Alaotra region of Madagascar, farmers face different constraints in different fields within their farming systems. In upland fields, soil's organic matter stocks are declining because of reduced fallow time. On these types of fields, dry spells can have a strong impact on crops' yields. In most of paddy fields, rice transplanting is delayed and because of poor water control, weeds threaten rice yield. The average rice yield for conventional fields is around 1 t ha⁻¹ for upland rice and 2.5 t ha⁻¹ in paddy fields with poor water control (Penot *et al.*, 2010).

As part of a long-term research and development programme exploring options for enhancing the productivity and sustainability of farming systems in Madagascar, considerable emphasis has been devoted to identify suitable CA systems (Husson *et al.*, 2010). In the Lake Alaotra region, in 2009, 1420 farmers have implemented CA cropping systems on a total of 1000 ha, i.e. on average 0.7 ha per adopting farmer (Rakotondramanana *et al.*, 2010). Extension agents regularly monitor grain production, but biomass production is measured neither in terms of quantity nor in terms of soil cover.

Various authors have stressed on the lack of biomass for use as mulch in smallholder farming systems in Africa (Erenstein, 2003; Giller *et al.*, 2009; Wezel and Rath, 2002). First it is difficult to produce enough biomass without external inputs, and second, once biomass is produced, it is difficult to retain it as mulch because of competing uses, especially for livestock feed. Surprisingly, few quantitative data are available concerning mulch availability under smallholder conditions, or on the amount of mulch required to fulfil different ecological functions. To the best of our knowledge, previous research has not addressed the question to what quantity of the available biomass can be removed from field while maintaining a degree of soil cover required for specific agronomic functions. Some authors propose thresholds for biomass exportation but provide few justifications. For example, Govaerts *et al.* (2007) suggest that it is possible to remove 50–70% of the residue while keeping adequate benefits to the soil only considering cereal yields.

The hypotheses of this study are that when CA systems are implemented by smallholder farmers, in some cases the production and/or conservation of biomass lead to a partial soil cover; consequently, we can assume that not all cover functions will be effective. We can support farmers and technicians' decisions, in terms of biomass




Figure 1. Rainfall in the Ambongabe village ($48^{\circ}28'11.2''S$, $17^{\circ}51'50.6''E$), lower and upper quartile for the 2000–2010 time period and 2008–2009 season rainfall.

management, by establishing curves taking into relation soil cover and mulch quantity by mulch type.

MATERIALS AND METHODS

Location

All fields investigated were located in the Lake Alaotra region, Madagascar, between $17^{\circ}28.0'S$ and $17^{\circ}53.0'S$, $48^{\circ}08.0'E$ and $48^{\circ}38.0'E$, and 760–950 m above sea level. The mid-altitude tropical climate has a mean annual temperature of $22^{\circ}C$. Average rainfall near Ambatondrazaka was 994 mm from 2000 to 2010 and 1553 mm from October 2008 to September 2009 (Figure 1) (Bas Rhône Languedoc (BRL), 2010). The hillside soils are Cambisols (texture 20% clay, 38% silt and 42% sand). Lowland paddy fields are Ferralsols (texture 39% clay, 29% silt and 32% sand) (A. Albrecht, personal communication 2010; FAO, 2010b; Razafimbelo *et al.*, 2010). The hillside soils *C* stocks (0–20-cm layer) are smaller (15.6 to 19.7 t ha $^{-1}$) than the paddy soils (23.6 to 29.0 t ha $^{-1}$) (Razafimbelo *et al.*, 2010).

Experimental design

The study was conducted in 91 farmers' fields in 2008 and 2009. The study was done in one crop cycle but the aim was to stress the intra-annual variability coming from farmers' management. Cropping systems differ according to their location in the landscape. On the hillsides, locally called *tanety*, all the crops are rain-fed. In the




Figure 2. (Colour online available at journals.cambridge.org/EAG) Examples of crop and cover crop sequences in CA cropping systems in the Lake Alaotra region, Madagascar. (a) A two-year rotation on hillsides with maize + *D. lablab* in year n , and upland rice in year $n + 1$; (b) a multi-annual succession on hillsides with a crop + *S. guianensis* in year n , *S. guianensis* alone in year $n + 1/2/3$, upland rice the last year; (c) a double crop sequence within a year in lowland fields with *V. villosa* in the off-season and rice in the main season. Modified from Séguy *et al.* (2009).

lowland, paddy field crops are irrigated but with poor water control, as the irrigation network is not fully functional, i.e. the farmers largely depend on rainfall and natural drainage in and out of fields. Two cropping sequences were studied on the hillsides. The first sequence was maize + pulse in year n , followed by upland rice in year $n + 1$. Pulses were cowpea (*Vigna unguiculata* (L.) Walp) or dolichos (*Dolichos lablab* L.), (Figure 2a). The second crop sequence included one year of the forage legume *Stylosanthes guianensis* Aubl., ‘CIAT 184’. In year n , *S. guianensis* was sown alone or intercropped with main crops such as Bambara nut (*Vigna subterranea* (L.) Verdc.), groundnut (*Arachis hypogaea* L.), maize (*Zea mays* L.), and cassava (*Manihot esculenta* Crantz). In year $n + 1/+2/+3$, *S. guianensis* was grown alone for as long as the farmer wished. The last year of rotation, i.e. years $n + 3$ or $n + 4$, *S. guianensis* was killed mechanically by cutting the crown. After 2–3 weeks, when the mulch had been flattened, rice was sown (Figure 2b). The third sequence studied was in lowland paddy fields with poor water control, where a cover crop was sown during off-season and rice

Table 1. Number of fields investigated for each of the crop–cover crop combinations. In four fields, measurements were taken in both 2008 and 2009, giving a total of 95 samples.

Types of fields and cover crop–crop combinations	Number of fields
Tanety (hillside)	
<i>S. guianensis</i>	17
Maize + <i>V. unguiculata</i>	16
Maize + <i>D. lablab</i>	22
Paddy fields	
<i>D. lablab</i>	15
<i>V. villosa</i>	21
Total	91

was sown into the mulch of the cover crop at the beginning of the rainy season. The cover crop was hairy vetch (*Vicia villosa* Roth) or *D. lablab* (Figure 2c). In all cropping systems rice was directly seeded without tillage. Less than one-fifth of the maize + pulse fields received nitrogen, phosphorus and potassium (NPK) (in ratio of 11:22:16) or urea fertilizer, and in each case less than 50 kg ha⁻¹ of fertilizer was used. The season before *V. villosa* and *D. lablab* were grown less than one-fifth of the paddy crops were fertilized. These fields were fertilized with less than 50 kg ha⁻¹ urea.

Sizes of the fields were diverse but relatively small, ranging from 100 m² to 5000 m². Farmers conducted all cultural operations. Table 1 shows the distribution of fields regarding the crop sequence and their locations.

Aboveground biomass measurement

The available biomass was estimated from October to the first week of December, when rice is usually sown. Where biomass was still living (e.g. *S. guianensis* on hillside and *V. villosa* and *D. lablab* in paddy fields) it was cut close at 5 to 15 cm above the soil surface. Where the plants had already senesced (e.g. maize + *D. lablab* or maize + *V. unguiculata*), the dead material was removed from the soil to be weighed. Five sub-samples of 1 m² were taken in each field, one in the centre of the field and others at the middle of each diagonal linking the centre and the corners of the field. Each sub-sample was weighed separately and a composite sample prepared. The composite sample was weighed in the field, air-dried and finally reweighed. Samples of the biomass (200 g) were oven-dried at 55 °C for 48 h to allow correction for moisture content and stored for near-infrared reflectance spectroscopy (NIRS) predictions. All biomass values are expressed on a dry matter basis.

For some fields of maize + *V. unguiculata* and maize + *D. Lablab*, the aboveground biomass was also measured at the end of the growing season (March–April). At this date, five plots of 2.5 m² were sampled from each field using the above-described pattern. As the maize rows were spaced 1 m apart, each sample included 2.5-m length of one row of maize.

Soil cover measurement

The relationship between mulch mass and soil cover was determined by measuring soil cover of the known mass of plant residue. Residues of *D. lablab*, *V. villosa*, maize + *D. lablab* mixture and *S. guianensis* were collected from farmers' fields. In order to give uniform background, quantities of residues equivalent to 1, 3, 6, 9, 12 and 15 t ha⁻¹ were spread on a 1 m² blue plastic tarpauline. A nadir view photograph of the residue was taken. Digital images were processed using the Photoshop[®] software to determine the visible area of the blue background. From this we inferred the proportion of the area covered by plant residues. For each quantity of residue, two replicate pictures were taken with a different random arrangement of residues. For randomly distributed mulch elements, the fraction of the soil covered by mulch (C) can be related to the mulch mass (M) by

$$C = 1 - \exp(-A_m \times M), \quad (1)$$

where A_m is an area-to-mass ratio depending on mulch type (Gregory, 1982; Scopel *et al.*, 1999; Smets *et al.*, 2008). The coefficient A_m has physical dimension of area covered by one average straw per mass of one average straw. We determined A_m by adjusting a non-linear regression to observed data using the 'non-linear regression' function of the XLStat 2010.1.01 software.

Nitrogen content

An NIRS prediction was used to determine the nitrogen content of samples. This method has proved to be an efficient tool to screen the quality of organic resources (Shepherd *et al.*, 2003). Dried samples were finely grounded (1 mm) and scanned twice at 2-nm intervals over the 1100–2500-nm wavelengths on a monochromator (FOSS—NIR Systems 5000, Silver Spring, MD, USA). Mathematical analysis of the spectral data was performed with WinISI III Version 1.63 software (Infrasoft International, Port Matilda, PA, USA). The NIRS prediction referential used in the present study consisted of a large tropical and temperate forage database pairing reflectance values and reference analyses for concentrations of nitrogen (Tran *et al.*, 2009). The nitrogen content is reported here only for *S. guianensis* on hillside and *V. villosa*, *D. lablab* on paddy fields, as the analysis has been made on biomass sample just before the seeding of rice. Thus, part of the nitrogen content of this biomass is available to following rice through mulch decomposition.

RESULTS

Production of biomass and amount of mulch available

The mulch available at the end of the dry season (October 2009) compared to the biomass produced at the beginning of the dry season (April–May 2009) was higher for maize + *D. lablab* fields and lower for maize + *V. unguiculata* ones (Figure 3). The mean quantity of mulch available prior to sowing of rice on hillsides was 3.6 t ha⁻¹ for fields of *S. guianensis*, 4.0 t ha⁻¹ for maize + *V. unguiculata* fields and 5.4 t ha⁻¹ for maize + *D. lablab* fields. In paddy fields, the mean mulch available was 6.8 t ha⁻¹ with *V. villosa*




Figure 3. Relationship between the amount of vegetative biomass (dry matter) produced by cereal (+ cover crop) in April–May 2009 and the amount of mulch left in October 2009. Data from 17 farmers' fields.

and 7.3 t ha^{-1} with *D. lablab* (Figure 4a). For all types of mulch, there was considerable variability between the hillside fields, but less variability in the paddy fields.

Soil cover

The digital picture analysis allowed relationships between the quantity of mulch and soil cover to be derived for four types of mulch (*S. guianensis*, maize + *D. lablab*, *D. lablab* alone and *V. villosa*). Equation (1) proved to be a good descriptor of this relationship, as the coefficient of determination between observed soil cover and curve fit was greater than 0.99 in all cases (Figure 5a). A_m for maize + *D. lablab*, *D. lablab*, *S. guianensis* and *V. villosa* are presented in Table 2. The capacity of plant residues to cover the soil varied strongly between different residues. For example, 3 t ha^{-1} of maize + *D. lablab* covered around 50% of the soil surface, 3 t ha^{-1} of *D. lablab* covered 60%, whereas a similar quantity of *V. villosa* biomass covered nearly 90% of the soil surface. Ninety-five percent of soil cover was obtained with less than 5 t ha^{-1} of *V. villosa*, but the same cover rate required 10 t ha^{-1} of *D. lablab*.

The range of biomass quantity (Figure 4a) was then converted to soil cover (Figure 4b) using Equation (1) and the A_m values given in Table 2. The calculated average soil cover (lower and upper quartile between commas) for *S. guianensis*, maize + *V. unguiculata*, maize + *D. lablab*, *V. villosa* and *D. lablab* was 66% (58–79%), 56% (30–74%), 70% (62–84%), 97% (99–100%) and 87% (84–94%), respectively. The range of variability observed for mulch quantity was different from those of soil cover. For example, CV of the average quantity of mulch of *V. villosa* was 34%, but the CV for soil cover was only 8%. For maize + *D. lablab* cover the CV varied from 27 to 41% (Figures 4a, b).




Figure 4. (a) Aboveground dry biomass available as mulch prior to sowing of rice. (b) Soil cover calculated from the amount of biomass measured in the field. Measured in the hillside fields and paddy fields in the Lake Alaotra region, 2008–2009. Number of fields (n): *S. guianensis* = 19; maize + *V. unguiculata* = 17; maize + *D. lablab* = 23; *V. villosa* = 21; *D. lablab* = 15. Box plot: median (horizontal continuous line), mean (cross).

Impact of biomass removal on soil cover

Using the biomass production of *V. villosa*, *D. lablab*, *S. guianensis*, and maize + *V. unguiculata* measured in the field (Figure 4a) and the soil cover curves derived from this data (Figure 5b), estimates were made of the effects of biomass removal on soil cover (Figure 6). This was done using the upper and the lower quartiles of biomass production among farmers' fields. For *V. villosa*, points A, B, C and D mark the maximum quantity of biomass that can be removed before reaching 90% of soil cover (A, B) or 30% (C, D), for three quarters of fields (A, C) or one-quarter of fields (B, D). For three quarters of the *V. villosa* fields, 3 t ha⁻¹ can be removed while maintaining 90% soil cover, and 5.6 t ha⁻¹ can be removed from one-quarter of the fields (Figure 6). If the target is 30% of soil cover, then the removable




Figure 5. (a) Soil cover (%) as a function of the amount of mulch for different crop–cover crop combinations. *Data from this study, [†]data from Teasdale and Mohler, (2000), [‡]data from Scopel *et al.* (1999). Equation: $C = 1 - \exp(-A_m \times M)$, where C is the fraction of the soil covered by mulch, M is the mulch mass in t ha⁻¹ and A_m is an area-to-mass ratio depending on mulch type. The R^2 for the fitted curves for *S. guianensis*, maize + *D. lablab*, *D. lablab* and *V. villosa* are, 0.991, 0.990, 0.998 and 0.998, respectively. (b) Soil cover calculated from the quantities of biomass measured in the field. The relationship between mulch quantity and soil cover for maize + *V. unguiculata* has been inferred from the relationship for maize + *D. lablab*. Number of fields (n): *S. guianensis* = 19; maize + *V. unguiculata* = 17; maize + *D. lablab* = 23; *V. villosa* = 21; *D. lablab* = 15.

biomass will be 5.6 and 7.9 t ha⁻¹ for three quarters or one-quarter of the fields, respectively.

Nitrogen content

The average nitrogen content of samples was respectively 2.7% of dry matter for *S. guianensis*, 3.4% for *V. villosa* and 1.8% for *D. lablab*. Combining with total biomass available, this gave 82 (± 21) kg N ha⁻¹ in the mulch for *S. guianensis*, 236 (± 97) kg N ha⁻¹ for *V. villosa*, and 123 (± 46) kg N ha⁻¹ for *D. lablab* (Figure 7).

Table 2. Area-to-mass ratio values (A_m) from this study and from the literature for different crops and cover crops.

Crop and cover crop	Type of residue	Area-to-mass ratio (ha t^{-1})	Source
<i>Avena sativa</i>	Not decomposed	1.370	(Steiner <i>et al.</i> , 2000)
	Unknown	1.400	(Gregory, 1982)
<i>Dolichos lablab</i>	Not decomposed	0.320	This study
<i>Glycine max</i>	Unknown	0.720	(Gregory, 1982)
<i>Hordeum vulgare</i>	Not decomposed	1.170	(Steiner <i>et al.</i> , 2000)
<i>Secale cereale</i>	Unknown	0.420	(Teasdale and Mohler, 2000)
<i>Stylosanthes guianensis</i>	Not decomposed	0.377	This study
<i>Triticum aestivum</i>	Unknown	0.540	(Gregory, 1982)
	Unknown	0.450	(Gregory, 1982)
<i>Triticum aestivum</i> spring	Not decomposed	1.830	(Steiner <i>et al.</i> , 2000)
<i>Triticum aestivum</i> winter	Not decomposed	1.380	(Steiner <i>et al.</i> , 2000)
<i>Vicia villosa</i>	Not decomposed	0.690	(Teasdale and Mohler, 2000)
	Not decomposed	0.742	This study
<i>Zea mays</i>	Not decomposed	0.114	(Gilley <i>et al.</i> , 1986)
	Unknown	0.190	(Teasdale and Mohler, 2000)
	Unknown	0.400	(Gregory, 1982)
	Not decomposed	0.367	(Scopel <i>et al.</i> , 1999)
	Partially decomposed	0.271	(Scopel <i>et al.</i> , 1999)
	Partially decomposed, stem without leaves	0.092	(Scopel <i>et al.</i> , 1999)
	<i>Zea mays</i> + <i>Dolichos lablab</i>	Partially decomposed	0.251




Figure 6. Effect of biomass removal on soil cover for five different cover crops, *V. villosa*, *D. lablab*, maize + *D. lablab*, *S. guianensis* and maize + *V. unguiculata* and quartile values from farmers' fields. Points A, B, C and D mark the maximum quantity of dry biomass, which can be removed while maintaining 90% soil cover (A, B) or 30% (C, D), for 3/4 of the fields (A, C) or 1/4 of the fields (B, D). These quantities for A, B, C and D are 3.0, 5.3, 5.6 and 7.9 t ha⁻¹, respectively.




Figure 7. Amount of nitrogen (kg ha^{-1}) contained in the aboveground dry biomass of different cover crops. Number of fields (n): *S. guianensis*, $n = 5$; *V. villosa*, $n = 21$; *D. lablab*, $n = 15$. Box plot: median (continuous line), mean (cross).

DISCUSSION

Production and conservation of biomass

Although maize + *D. lablab* fields had more biomass at the end of the dry season than at the beginning, less biomass remained in almost all maize + *V. unguiculata* fields. Three reasons can explain the difference between these two cover crops. First, *V. unguiculata* had ceased to grow before the end of the rainy season, whereas *D. lablab* continued to grow into the dry season. Second, cattle herders tend not to graze their cattle in fields of maize + *D. lablab* fields, as they see *D. lablab* is still growing there. As all the standing biomass dries *in situ* in maize + *V. unguiculata* fields, herders consider it to be a 'normal' field available for grazing. Third, farmers grew *D. lablab* only to produce biomass for the next crop, and not for edible grain. By contrast, farmers grew *V. unguiculata* for grain with the additional benefit of biomass for use as mulch. Nevertheless, the amount of biomass remaining at the end of the dry season in the Alaotra region of Madagascar is large compared with CA systems in other countries of sub-Saharan Africa, e.g. 3.5 t ha^{-1} (Naudin *et al.*, 2010) or 2 t ha^{-1} (Wezel and Rath, 2002).

S. guianensis can be cut and killed at the beginning of the third year after sowing to produce mulch where rice can be sown (Husson *et al.*, 2010). All *S. guianensis* fields investigated were in the third, fourth or fifth year but the average biomass available at the beginning of the subsequent rainy season was 3.6 t ha^{-1} , a small amount compared with the other cover crops, and much less than reported elsewhere (e.g. Saito *et al.* 2010 reported 7.4 t ha^{-1} for a two-year stand in Benin). Under controlled conditions, *S. guianensis* produced from 5 to 20 t ha^{-1} (Husson *et al.*, 2008), but under real farmers' conditions most of these fields had been partially grazed during the dry seasons, which explained the relatively small amount of remaining biomass. *S. guianensis* is well known to support multiple cuts during the growing season to provide fresh forage for animal feed, and is resistant to grazing (Roberge and Toutain, 1999). Nevertheless, this reduces its final growth and biomass available. Furthermore, *S. guianensis* is usually grown on the worst fields where farmers intend to improve soil fertility and can afford to leave

the field uncropped. The 2008–2009 cropping season was rainy season (1553 mm) compared with the average rains (994 mm), thus the biomass obtained on hillsides was close to the optimum attainable in this region. Biomass production on paddy fields should be less sensitive to this climatic condition, as the water is not limiting in this kind of fields.

In the lowland paddy fields, biomass production of *D. lablab* and *V. villosa* was similar at around 7 t ha⁻¹, and greater than reported earlier in the literature, e.g. 2.44 to 5.16 t ha⁻¹ (Sainju *et al.*, 2006). None of these *V. villosa* or *D. lablab* fields have been grazed. Farmers prefer to grow *V. villosa* in this kind of field, as it can be intercropped with vegetables. *V. villosa* requires more water than *D. lablab*, so it is found only in lower lying fields with fine soil texture that allow capillary rise. When water is more limiting, *D. lablab* is selected.

In the Lake Alaotra region of Madagascar where no basal fertiliser is applied, large amount of legume biomass was achieved in the lowland fields, but less biomass was produced in the upland fields probably due to poorer soil fertility. In particular, this poor production can be linked with low phosphorus availability. In many parts of the tropics basal fertilisation with phosphorus and other nutrients is required to get good legume growth and nitrogen fixation (Giller and Cadisch, 1995). In paddy fields, the use of adapted legumes (*D. lablab* and *V. villosa*) on relatively fertile soils allowed production of a large amount of biomass each year without competing with other crops. The paddy fields are usually under exploited during the off-season, as vegetables are the only crops grown where manual irrigation is possible. The area covered by vegetables is small due to the labour required, leaving a large area where cover crops could be grown.

Relationships between biomass and soil cover

The capacity of plant residues to cover soil varied strongly between different residues. The presence of small leaves in *V. villosa*, *S. guianensis* and *D. lablab* gives the higher A_m value compared with cereal residues alone so that much less biomass is needed to obtain the same percentage of soil cover. The digital picture analysis proved to be a useful tool for generating predictive equations to relate biomass with soil cover for different residue mixtures (Figure 5a). This method is relatively easy to use even with low resources. It should be used more in order to better characterise mulch characteristics and thus to allow a better explanation for CA cropping systems impacts. As we can see in Figure 4a, the variability in terms of biomass production is relatively high, as is commonly found in smallholder cropping systems in developing countries (Naudin *et al.*, 2010; Tittonell *et al.*, 2008). This variability results in a wide range of soil cover (Figure 4b) and nitrogen input (Figure 7). These examples demonstrate the wide variability in biomass yield found under farmer's conditions, even for one type of cropping system, so that the agronomic benefits expected from CA are not necessarily fulfilled. Further, the agronomic benefits are not linearly linked with the quantity of mulch and therefore thresholds should be defined for specific combinations of environmental conditions, cover crop and expected function.

Maintaining sufficient mulch

We can infer from Smets *et al.* (2008) that a minimum of 30% soil cover is required to reduce inter-rill soil erosion substantially, whereas a target of 90% is the minimum required to obtain a good weed control (Bilalis *et al.*, 2003; Teasdale and Mohler, 2000). The amount of mulch required to achieve these rates of soil cover can be readily derived from Figure 5a. On the hillside fields where the biomass production was less than in the lowland paddy fields, the amount of biomass that could be removed was substantially less. For example, for *S. guianensis*, 90% of soil cover was reached in less than a quarter of the fields. With a target of 30% of soil cover, the removable biomass was between 1.4 t ha⁻¹ for three quarters of the fields and 3.4 t ha⁻¹ for a quarter of the fields. Thus, the amount of biomass that can be removed for livestock, or grazed *in situ* varies strongly between the hillside and lowland paddy fields and between different legumes or residue mixtures. Govaerts *et al.* (2005) stressed the need to establish critical amount of residue required for maintaining soil productivity while using part of the biomass as fodder. These authors also mentioned that zero tillage with residue retention give better cereal yield results than without residue. But they did not specified the quantity of mulch retained and even less the percentage of corresponding soil cover.

Knowing the relationship between potential removable biomass and impact in terms of soil cover rate can help farmers to take decisions regarding the possibility to use part of the biomass produced in field. It also helps to compare the management flexibility of different cropping systems. In fact, in no-till cropping systems, the lack of mulch, less or equal to 30% of soil cover, can lead to increased erosion (Volk *et al.*, 2004) and weed competition (Bilalis *et al.*, 2003) compared with tilled cropping.

Nitrogen availability and role on short-term productivity and long-term fertility

Beyond the quantity of biomass produced, the quality also varies among cover crops and fields. Again, for the same types of field (paddy field) and cropping system (annual rotation with rice), the quantity of nitrogen available in the residues can double with the type of cover crop, e.g. 123 kg N ha⁻¹ for *D. lablab* against 236 kg N ha⁻¹ for *V. villosa*. Values for *V. villosa* are higher than those observed by Sainju *et al.* (2006), which varied from 76 to 167 kg N ha⁻¹ depending on the year. These authors showed that even with the smaller amount of biomass added, the available inorganic nitrogen content increased in the soil when *V. villosa* was killed resulting in increased grain and biomass yields of the subsequent sorghum crop.

The biomass nitrogen can be partially returned to soil to benefit the following rice crop, or be fed to cattle to improve animal productivity. As stressed by Rufino *et al.* (2006), the direct application of plant materials to soil results in more efficient cycling of nitrogen, with fewer losses from the system than from materials fed to livestock and then returned to the soil through manure. However, livestock provide many other benefits, and animal manure can contain large amount of available nitrogen, which can promote crop growth in short term (Rufino *et al.*, 2006). The partial allocation of the biomass to cattle or to mulch is driven by the goals of the farmer; especially by

trade-offs between expected benefits from rice yield improvement, reduction in labour required for weeding and enhanced cattle production.

The short-term effects of mulch, such as water balance improvement (Scopel *et al.*, 2004; Thierfelder and Wall, 2010), are more easily perceived by farmers than long-term effects on soil fertility. Although after eight years of implementation of CA in the Lake Alaotra region, the C stock was consistently greater in CA plots (between 1.1 t ha⁻¹ and 3.5 t ha⁻¹) than in ploughed plots, but the difference was not statistically significant (Razafimbelo *et al.*, 2010). Furthermore, these results were obtained when all of the plant residues were returned as mulch in the CA plots (rarely achieved in farmers' fields) compared with complete removal of crop residues in the ploughed plots. These results reinforce the conclusion that the fulfilment of agroecological functions by CA will depend on the amount of biomass returned to soil and length of time the system is implemented.

CONCLUSION

Our results showed that it is possible to produce and keep sufficient biomass in the field for CA systems even under smallholder farming conditions where livestock graze freely during dry season. However, the quantity of biomass produced varies strongly between hillsides and valleys, and between cover crops and farmers' management. Soil cover is not linearly related to mulch quantity. Thus, for a given quantity of biomass exported to feed cattle, the impact is different depending on the cover crop, the initial amount of biomass and the agroecological functions of mulch searched by farmers. When comparing benefits of different types of CA cropping systems, it is important to report the amount and quality of biomass produced, and the corresponding rate of soil cover. In terms of the agroecological functions of soil cover, such as weed control, erosion control and water retention, different amount of mulch is required with different cover crops. The relationships between biomass export for cattle feed and these agroecological functions require more systematic study. The decision on how much biomass can be removed from the field will depend on the local biophysical conditions, the biomass characteristics and the farmer's goals for his/her whole farm system.

Acknowledgements. This study was conducted in collaboration with the BV Lac project (Bassins versants–Lac Alaotra) of the Malagasy Ministry of Agriculture implemented by Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD) and funded by the Agence Française de Développement (AFD). The farmers were advised by Agronomes et Vétérinaires Sans Frontières (AVSF) and Bas Rhône Languedoc-Madagascar (BRL-Madagascar). We are grateful to all of them for their technical support. This study was funded by (i) PEPITES project from Agence Nationale de la Recherche (ANR), (ii) PAMPA project from AFD, Fond Français pour l'Environnement Mondial (FFEM) and Ministère des Affaires Étrangères (MAE) and (iii) CIRAD.

REFERENCES

- Bas Rhône Languedoc (BRL) (2010). Cuvette du lac Alaotra, réseau pluviométrique. Annuaire 2000 à 2010. Ambatondrazaka, Madagascar: BRL-Madagascar, CIRAD, 154 pp.
- Bilalis, D., Sidiras, N., Economou, G. and Vakali, C. (2003). Effect of different levels of wheat straw soil surface coverage on weed flora in Vicia faba crops. *Journal of Agronomy and Crop Science* 189:233–241.
- Erenstein, O. (2003). Smallholder conservation farming in the tropics and sub-tropics: a guide to the development and dissemination of mulching with crop residues and cover crops. *Agriculture, Ecosystems & Environment* 100:17–37.
- Food and Agriculture Organization (FAO) (2010a). <http://www.fao.org/ag/ca/> (Accessed 28 June 2011).
- Food and Agriculture Organization (FAO) (2010b). <http://www.fao.org/nr/land/soils/soil/wrb-soil-maps/classification-key/en/#c25142> (Accessed 10 November 2010).
- Giller, K. E. and Cadisch, G. (1995). Future benefits from biological nitrogen fixation. An ecological approach to agriculture. *Plant and Soil* 174:255–277.
- Giller, K. E., Witter, E., Corbeels, M. and Tittonell, P. (2009). Conservation agriculture and smallholder farming in Africa: the heretics' view. *Field Crops Research* 114:23–34.
- Gilley, J., Finkner, S., Spomer, R. and Mielke, L. (1986). Runoff and erosion as affected by corn residue: Part I. Total losses. *Transactions of the ASAE* 29:157–160.
- Govaerts, B., Sayre, K. D. and Deckers, J. (2005). Stable high yields with zero tillage and permanent bed planting? *Field Crops Research* 94:33–42.
- Govaerts, B., Sayre, K. D., Lichter, K., Dendooven, L. and Deckers, J. (2007). Influence of permanent raised bed planting and residue management on physical and chemical soil quality in rain fed maize/wheat systems. *Plant and Soil* 291:39–54.
- Gregory, J. (1982). Soil cover prediction with various amounts and types of crop residue. *Transactions of the ASAE* 25:1333–1337.
- Hobbs, P. R. (2007). Conservation agriculture: what is it and why is it important for future sustainable food production? *Journal of Agricultural Science* 145:127–137.
- Husson, O., Charpentier, H., Raharison, T., Razanamparany, C., Moussa, N., Rasolomanjaka, J., Michellon, R., Naudin, K., Rakotoarivivo, C., Rakotondramanana, Enjalric, F. and Séguy, L. (2010). Les systèmes SCV adaptés aux différentes zones agroécologiques de Madagascar. In *Manuel Pratique du Semis Direct à Madagascar* (Ed CIRAD). Antananarivo, Madagascar: GSDM, CIRAD.
- Husson, O., Charpentier, H., Razanamparany, C., Moussa, N., Michellon, R., Naudin, K., Rakotoarivivo, C., Rakotondramanana and Séguy, L. (2008). *Stylosanthes guianensis*. In *Manuel Pratique du Semis Direct à Madagascar. Vol. III. Chap. 3.2.1. Fiches Techniques Plantes de Couverture: Légumineuses Pérennes*, 12 (Eds CIRAD, TAFA, GSDM, AFD and MAEP). Antananarivo, Madagascar: CIRAD.
- Maltas, A., Corbeels, M., Scopel, E., Wery, J. and da Silva, F. A. M. (2009). Cover crop and nitrogen effects on maize productivity in no-tillage systems of the Brazilian cerrados. *Agronomy Journal* 101:1036–1046.
- Naudin, K., Gozè, E., Balarabe, O., Giller, K. E. and Scopel, E. (2010). Impact of no tillage and mulching practices on cotton production in North Cameroon: a multi-locational on-farm assessment. *Soil and Tillage Research* 108:68–67.
- Neto, M. S., Scopel, E., Corbeels, M., Cardoso, A. N., Douzet, J. M., Feller, C., Piccolo, M. D., Cerri, C. C. and Bernoux, M. (2010). Soil carbon stocks under no-tillage mulch-based cropping systems in the Brazilian Cerrado: an on-farm synchronic assessment. *Soil and Tillage Research* 110:187–195.
- Penot, E., Scopel, E., Domas, R. and Naudin, K. (2010). La durabilité est elle soluble dans le développement ? L'adoption des techniques de conservation de l'agriculture dans un contexte d'incertitudes multiples au lac Alaotra, Madagascar. In *Colloque "Agir en Situation D'incertitude,"* 10 (Eds IFSA and CIRAD). Montpellier, Madagascar: IFSA-CIRAD.
- Rakotondramanana, Enjalric, F. and Husson, O. (2010). *Documentation et synthèse de l'Agriculture de Conservation à Madagascar (FAO)*. Antananarivo, Madagascar: GSDM, 96 pp.
- Razafimbelo, T., Albrecht, A., Feller, C., Ravelojaona, H., Moussa, N., Razanamparany, C., Rakotoarivivo, C., Razafintsalama, H., Michellon, R., Naudin, K. and Rabeharisoa, L. (2010). Stockage de carbone dans les sols sous systèmes de culture en semis direct sous couvert végétal (SCV) dans différents contextes pédoclimatiques à Madagascar. *Etude et Gestion des Sols* 17:143–158.
- Reicosky, D. (2008). Carbon sequestration and environmental benefits from no-till systems. In *No-Till Farming Systems, Vol. 3*, 43–58 (Ed T. E. A. Goddard). Bangkok, Thailand: World Association of Soil and Water Conservation.
- Roberge, G. and Toutain, B. (1999). *Cultures Fourragères Tropicales*. Montpellier, Madagascar: CIRAD.

- Rufino, M. C., Rowe, E. C., Delve, R. J. and Giller, K. E. (2006). Nitrogen cycling efficiencies through resource-poor African crop-livestock systems. *Agriculture Ecosystems & Environment* 112:261–282.
- Sainju, U. M., Whitehead, W. F., Singh, B. P. and Wang, S. (2006). Tillage, cover crops, and nitrogen fertilization effects on soil nitrogen and cotton and sorghum yields. *European Journal of Agronomy* 25:372–382.
- Saito, K., Azoma, K. and Oikeh, S. O. (2010). Combined effects of *Stylosanthes guianensis* fallow and tillage management on upland rice yield, weeds and soils in southern Benin. *Soil and Tillage Research* 107:57–63.
- Scopel, E., Chavez Guerra, E. and Arreola-Tostado, J. (1999). Le semis direct avec paillis de résidus dans l’ouest mexicain: une histoire d’eau? *Agriculture et Développement* 21:71–86.
- Scopel, E., Da Silva, F. A. M., Corbeels, M., Affholder, F. O. and Maraux, F. (2004). Modelling crop residue mulching effects on water use and production of maize under semi-arid and humid tropical conditions. *Agronomie* 24:383–395.
- Séguy, L., Husson, O., Charpentier, H., Bouzinac, S., Michellon, R., Chabanne, A., Boulakia, S., Tivet, F., Naudin, K., Enjalric, F., Chabierski, S., Rakotondralambo, P. and Rakotondramanana (2009). La gestion des écosystèmes cultivés en semis direct sur couverture végétale permanente. In *Manuel du semis direct à Madagascar*, Vol I, Chap. 2., 32 (Eds GSDM and CIRAD). Antananarivo, Madagascar: CIRAD.
- Shepherd, K. D., Palm, C. A., Gachengo, C. N. and Vanlauwe, B. (2003). Rapid characterization of organic resource quality for soil and livestock management in tropical agroecosystems using near-infrared spectroscopy. *Agronomy Journal* 95:1314–1322.
- Smets, T., Poesen, J. and Knapen, A. (2008). Spatial scale effects on the effectiveness of organic mulches in reducing soil erosion by water. *Earth-Science Reviews* 89:1–12.
- Steiner, J. L., Schomberg, H. H., Unger, P. W. and Cresap, J. (2000). Biomass and residue cover relationships of fresh and decomposing small grain residue. *Soil Science Society of America Journal* 64:2109–2114.
- Teasdale, J. R. and Mohler, C. L. (1993). Light transmittance, soil-temperature, and soil-moisture under residue of hairy vetch and rye. *Agronomy Journal* 85:673–680.
- Teasdale, J. R. and Mohler, C. L. (2000). The quantitative relationship between weed emergence and the physical properties of mulches. *Weed Science* 48:385–392.
- Thierfelder, C. and Wall, P. C. (2010). Rotation in conservation agriculture systems of zambia: effects on soil quality and water relations. *Experimental Agriculture* 46:309–325.
- Tittonell, P., Vanlauwe, B., Corbeels, M. and Giller, K. E. (2008). Yield gaps, nutrient use efficiencies and response to fertilisers by maize across heterogeneous smallholder farms of western Kenya. *Plant and Soil* 313:19–37.
- Tran, H., Salgado, P. and Lecomte, P. (2009). Species, climate and fertilizer effects on grass fibre and protein in tropical environments. *The Journal of Agricultural Science* 147:555–568.
- Volk, L. B. S., Cogo, N. P. and Streck, E. V. (2004). Water erosion influenced by surface and subsurface soil physical conditions resulting from its management in the absence of vegetal cover. *Revista Brasileira De Ciencia Do Solo* 28:763–774.
- Wezel, A. and Rath, T. (2002). Resource conservation strategies in agro-ecosystems of semi-arid West Africa. *Journal of Arid Environments* 51:383–400.