

Early Vertebrate Evolution

Palaeogeography of tropical seasonal coastal wetlands in northern Britain during the early Mississippian Romer's Gap

David MILLWARD^{1*}, Sarah J. DAVIES², Peter J. BRAND¹,
Michael A. E. BROWNE¹, Carys E. BENNETT², Timothy I. KEARSEY¹,
Janet E. SHERWIN² and John E. A. MARSHALL³

¹ British Geological Survey, The Lyell Centre, Edinburgh EH14 4AP, UK.

Email: dmill@bgs.ac.uk

² School of Geography, Geology and the Environment, University of Leicester, Leicester LE1 7RH, UK.

³ Ocean and Earth Sciences, University of Southampton, National Oceanography Centre, European Way, Southampton SO14 3ZH, UK.

*Corresponding author

ABSTRACT: The lower Mississippian Ballagan Formation of northern Britain is one of only two successions worldwide to yield the earliest known tetrapods with terrestrial capability following the end-Devonian mass extinction event. Studies of the sedimentary environments and habitats in which these beasts lived have been an integral part of a major research project into how, why and under what circumstances this profound step in the evolution of life on Earth occurred. Here, a new palaeogeographic map is constructed from outcrop data integrated with new and archived borehole material. The map shows the extent of a very low-relief coastal wetland developed along the tropical southern continental margin of Laurussia. Coastal floodplains in the Midland Valley and Tweed basins were separated from the marginal marine seaway of the Northumberland–Solway Basin to the south by an archipelago of more elevated areas. A complex mosaic of sedimentary environments was juxtaposed, and included fresh and brackish to saline and hypersaline lakes, a diverse suite of floodplain palaeosols and a persistent fluvial system in the east of the region. The strongly seasonal climate led to the formation of evaporite deposits alternating with flooding events, both meteoric and marine. Storm surges drove marine floods from the SW into both the western Midland Valley and Northumberland–Solway Basin; marine water also flooded into the Tweed Basin and Tayside in the east. The Ballagan Formation is a rare example in the geological record of a tropical, seasonal coastal wetland that contains abundant, small-scale evaporite deposits. The diverse sedimentary environments and palaeosol types indicate a network of different terrestrial and aquatic habitats in which the tetrapods lived.



KEY WORDS: Ballagan Formation, coastal wetlands, Midland Valley of Scotland, palaeoenvironment, Tournaisian, Tweed Basin.

The Mississippian (Tournaisian) Ballagan Formation of northern Britain hosts a diverse assemblage of vertebrate fossils, including the earliest known tetrapods with terrestrial capability (Clack 2002; Clack & Finney 2005; Smithson *et al.* 2012; Clack *et al.* 2016; Smithson & Clack 2018). This is one of only two successions in the world where such fossils have been found within the period of about 15 million years at the beginning of the Carboniferous and has become known as Romer's Gap. The other is the Horton Bluff Formation in Nova Scotia, Canada (Anderson *et al.* 2015), which at this time was relatively close along the southern margin of Laurussia. The emergence of tetrapods on land and their subsequent diversification follows the mass extinction events in late Devonian times (Kaiser *et al.* 2015). Understanding how, why and under what conditions this step change in the evolution of life on Earth occurred was part of the remit of the Tweed Project led by Jenny Clack from 2012 until 2016. So, what were the landscape, palaeoenvironments and habitats like at this time?

Francis *et al.* (1970) and later studies have all inferred a coastal floodplain setting, although earlier Belt *et al.* (1967) had proposed a marginal marine to restricted marine environment for dolostone deposition in the Ballagan Formation of the Midland Valley of Scotland. The Anderton (1985) conceptual facies model for the important fish-bearing succession at Foulden in the Tweed Basin was based on detailed sedimentary logging. This type of model has been developed further by Bennett *et al.* (2016, 2017), Kearsey *et al.* (2016) and Millward *et al.* (2018) from detailed investigations of sandy siltstones, ichnofacies, palaeosols and evaporite rocks displayed in the successions through the formation exposed at Burnmouth and in the cores from the Norham West Mains Farm Borehole near Berwick upon Tweed (Fig. 1). All of these models show the rapidly changing character of the sedimentary environments in the rock record, suggesting the juxtaposition of a mosaic of distinctive habitats within a strongly seasonal, tropical coastal wetland. These studies are applicable particularly to the Tweed




Figure 1 Outcrop and subsurface map of Tournaisian rocks in the Midland Valley of Scotland, Tweed and Northumberland-Solway basins. These strata include the Ballagan Formation and part of the Lyne Formation in the last basin. Major boreholes and thickness of the formation are shown. Outcrop data from the British Geological Survey DiGMapGB© NERC 2015; subsurface extent in the Northumberland-Solway basin from Chadwick *et al.* (1995).

Basin (Fig. 1), but how do they extend across the broader region of northern Britain? This is important because the palaeogeography of the region and its control on the distribution of palaeoenvironments and habitats may have played a role in shaping vertebrate evolution.

The Tournaisian palaeogeography of the Midland Valley, and the Tweed and Northumberland-Solway basins presented here has its origins in the seminal atlases of the UK by Wills (1951) and Cope *et al.* (1992). The Tweed Project studies have provided greater clarity of detail and connections across the region between the sedimentary environments, habitats and the tetrapods that lived in them (Clack *et al.* 2016). The new map combines data from field sections and borehole records through the Ballagan Formation in the Midland Valley and Tweed Basin and includes reflections on published interpretations of the Northumberland-Solway Basin. These investigations provide new insights into this rare example in the geological record of an equatorial seasonal coastal wetland.

1. Regional setting and stratigraphy

During the early Mississippian, interconnected strike-slip fault-controlled sedimentary basins opened across northern Britain as a result of the break-up of the equatorial, southern part of the Laurussian continental margin (Coward 1993; Domeier & Torsvik 2014). The northern basins include those in the Midland Valley, Tweed and Northumberland-Solway

regions. Volcanism accompanied opening of the Tweed and Northumberland-Solway basins (Leeder 1974a; Chadwick *et al.* 1995). The early sedimentary fill to the basins comprises the Ballagan Formation, which was preceded by the upper Devonian Kinnesswood Formation and succeeded in various parts of the region by the Clyde Sandstone, Garleton Hills Volcanic, Lyne and Fell Sandstone formations (Figs 1, 2).

The Ballagan Formation is preserved at outcrop and in the subsurface across the Midland Valley and Tweed Basin, and this passes southward and stratigraphically upward into the Lyne Formation in the Northumberland-Solway Basin (Fig. 2). The Ballagan Formation consists of grey siltstone and thin beds of argillaceous ferroan dolostone (historically referred to as ‘cementstone’), with varying proportions of sandstone, flood-deposited sandy siltstone, palaeosols and evaporite rock (Anderton 1985; Scott 1986; Andrews *et al.* 1991; Andrews & Nabi 1994, 1998; Bennett *et al.* 2016; Kearsley *et al.* 2016; Millward *et al.* 2018). The formation is best exposed in the 520 m thick, vertically dipping, coastal section at Burnmouth (Fig. 1). Though a late Tournaisian, CM Miospore Biozone age (e.g., Neves *et al.* 1973; Neves & Ioannides 1974; Stephenson *et al.* 2002) has been determined for the formation in central Scotland, Smithson *et al.* (2012) recorded earliest Tournaisian, VI Biozone miospores from the base of the formation at Burnmouth. Most recently, Marshall *et al.* (2019) provide evidence that in the Tweed Basin the formation spans the Tournaisian. In East Lothian, the upper part of the formation has yielded early Visean Pu Biozone miospores (Neves *et al.* 1973; Neves




Figure 2 Stratigraphy chart for the region (after Waters *et al.* 2011). Ballagan Formation (Browne 1980) subsumes the Tynninghame Formation (Chisholm *et al.* 1989), Cementstone Group (Miller 1887; Greig 1988) and part of the Lower Border Group (Lumsden *et al.* 1967; Day 1970; Leeder 1974b). The Kinnesswood, Ballagan and Clyde Sandstone formations comprise the Inverclyde Group (Browne *et al.* 1999). The miospore biozones of the Ballagan Formation are from Stephenson *et al.* (2002, 2004a, b), Williams *et al.* (2005) and Smithson *et al.* (2012). Abbreviations: KCBV = Kelso, Cottonshope and Birrenswark volcanic formations; W = Whita Sandstone Member.

& Ioannides 1974); these upper beds are therefore laterally equivalent to the Clyde Sandstone Formation of the west of the Midland Valley which is inferred to be lower Viséan (Waters *et al.* 2011).

The thickness of the Ballagan Formation ranges from a few tens of metres on the Isle of Bute, to 80 m in north Arran (Young & Caldwell 2012), 150 m in the west of the Midland Valley and in Fife, up to 355 m in Ayrshire (Stephenson *et al.* 2002), and possibly up to 900 m west of Edinburgh (Fig. 1; Mitchell & Mykura 1962). At least 418 m are present in East Lothian, but only 200 m at Birnieknowes at the NE margin of the Southern Upland massif. Approximately 520 m occur in the Tweed Basin whereas in the Northumberland–Solway Basin Tournaisian and Chadian strata may be up to 3400 m thick in the centre of this asymmetric graben (Chadwick *et al.* 1995). The formation is absent from the central part of the Midland Valley, having been eroded during mid-Viséan (Arundian) uplift prior to the Clyde Plateau volcanism (Millward & Stephenson 2011).

In stark contrast to the Ballagan Formation, the Kinnesswood and Clyde Sandstone formations comprise braided fluvial sandstone in upward-fining cycles, interbedded with overbank red and purple mudstone, multiple intervals with carbonate nodules and thin beds of concretionary limestone ('cornstones'; Eyles *et al.* 1949; Read & Johnson 1967; Francis *et al.* 1970; Leeder 1976; Leeder & Bridges 1978; Young & Caldwell 2011, 2012). The 'cornstones' are palaeosols with pedogenic carbonate nodules (Burgess 1961; Leeder 1976; Andrews *et al.* 1991; Wright *et al.* 1993).

In the Northumberland–Solway Basin the Lyne Formation (formerly Lower Border Group of, e.g., Lumsden *et al.* 1967; Leeder 1975a, b) is Tournaisian and early Viséan in age, and contains significant marine clastic and carbonate sedimentary rocks and fossils (Leeder 1974b; Brand 2018). The Lyne and Ballagan formations are overlain unconformably by the Fell Sandstone Formation in the Tweed Basin and eastern part of the Northumberland–Solway Basin. The Fell Sandstone is composed of a stacked succession of mainly coarse-grained

sandstone deposited from the first major Carboniferous fluvio-deltaic system to cross the region (Kearsey *et al.* 2018).

2. Methods

The palaeogeography is reconstructed using data from field exposures across the Midland Valley and Tweed basins, and from about 40 borehole logs from the British Geological Survey (BGS) archive that proved significant occurrences of Tournaisian strata. A sub-set of 12 boreholes, commissioned by BGS as part of the stratigraphical investigation of Upper Devonian and Mississippian rocks of the region from 1967 to 1981, provides descriptions and consistent information on the spatial and temporal occurrence, context and general form of the different lithofacies. Cores have been retained for only one of these, the Glenrothes Borehole, and visual inspection of these provided additional information.

It is to be expected that the archived logs vary in amount and quality of information they contain because they were made over more than 100 years. The BGS sub-set provides the greatest volume of data, though even here variations in data quantity and quality arise because they were logged by different geologists. Information is particularly sparse on palaeosols and sandy siltstones. Palaeosols are under-described in UK geology generally, though rooted 'seat-earths' have long been recognised by Geological Survey staff because of extensive work in the coalfields. Sandy siltstone is a facies only recently identified from the Ballagan Formation and so is not recorded (Bennett *et al.* 2016). Therefore, interpretation of the facies from these boreholes (Table 1) is informed and benchmarked against the facies analysis of the Norham West Mains Farm Borehole undertaken by Bennett *et al.* (2016; Table 1). This borehole is located in the Tweed Basin, about 10 km SW of Berwick upon Tweed, and fully cored 491 m of the Ballagan Formation (Fig. 1; Millward *et al.* 2013).

The BGS collections contain macrofossils from the Ballagan and Lyne formations from 235 localities, including many of

Table 1 Facies, facies associations and environments in the Ballagan Formation of the Midland Valley of Scotland and Tweed Basin. The marginal marine facies of the Lyne Formation Northumberland–Solway Basin are omitted. Facies descriptions are from the archived borehole logs in the BGS collections and from key exposures. Facies association after Bennett *et al.* (2016).

Facies	Facies description	Where seen	Depositional environment	Facies association
Dolostone	Parallel beds of ferroan dolostone ('cementstone') and dolomitised siltstone; grey but some reddish brown; hard, compact, splintery; sporadic records of ostracods, some with shell fragments and <i>Chondrites</i> ; some beds nodular or with bulbous base; varieties include laminated dolostone, others with diffuse top and base; tops may be cracked or brecciated; commonly hosts evaporite nodules	In all boreholes and exposures, varying abundance	Saline–hypersaline flood-plain lakes, cryptic marine flooding	Freshwater, saline and hypersaline lake
Reddened pseudobrecciated dolostone	Dolomicrite, bluish to greenish grey, pseudo-brecciated, cracked and reddened profile; Fe-rich claystone fills to cracks; low-angle, concave upwards fractures in upper part of profile resemble pedogenic slickensides	Mains of Errol Member, Glenrothes, East Dron	Desiccated hypersaline, ?alkaline lakes	
Limestone	Grey limestone, porcellaneous, in part pseudo-brecciated; or sandy, cross-bedded; with ostracods, fish scales and teeth, Spirorbids	Spilmersford, East Linton (top part), Tantallan	Flood-plain lakes	
Evaporite	Diverse suite of subfacies of gypsum and anhydrite nodules, chicken-wire masses, evaporite rock and laminae, hosted by dolostone and laminated grey siltstone facies. Pseudomorphs after halite in siltstone facies	Widespread, except in East Lothian	Hypersaline lakes, brine pans and sabka	
Laminated grey siltstone	Planar laminated grey and dark grey siltstone and mudstone; disseminated pyrite; ostracods, bivalves, fish scales, conchostracans and burrows; sporadic marine taxa; desiccation cracks and roots generally absent; typically interbedded with dolostone	All boreholes and exposures	Flood-plain lakes, cryptic marine flooding	
Heterogeneous siltstone	Grey, purple-grey, mottled grey and brown, red-brown or red and purple; massive to laminated, some rippled or disturbed bedding; some rooting and <i>Stigmaria</i> . Interbedded with laminated or ripple-laminated sandstone	In all boreholes and exposures	Overbank flooding	Overbank
Ripple-laminated sandstone	Ripple-laminated or cross-laminated, fine-grained with siltstone laminae; plant, shell and fish debris; bioturbation in some beds, sporadic records of ? <i>Monocraterion</i> and <i>Planolites</i> ; interbedded with siltstone; units typically <3 m	In all boreholes and exposures	Overbank and storm flooding into lakes; small stream deposits	
Palaeosols	Few identified in logs; best indicator is record of carbonised root traces in grey siltstone and 'seatclay' or 'seatearth' (?gleyed inceptisols) but depth of rooting and thickness not given; also rooted fine-sandstone (?entisols); unbedded mottled red-brown siltstone may represent vertisols	Most boreholes; Bute, Burnmouth	Vegetated flood-plain	
'Mudstone breccia' (sandy siltstone?)	Greenish grey dolomitised siltstone with irregular dark grey siltstone clasts up to 10 mm. Beds 10–84 cm. (described similarly as 'Clast-rock' in Ascog)	Glenrothes, East Dron, Ascog	Unconfined debris-flow deposit	
Conglomerate	Round dolostone clasts 1–20 mm diameter with calcareous mudstone matrix; plant fragments; beds 18 cm, siltstone above and below	East Linton, Everton	Unconfined debris-flow deposit	
Cross-bedded sandstone	Fine, medium and coarse-grained sandstone, trace siltstone; conglomerate at base in some units; carbonate and micaceous coatings to laminae; massive or cross-bedded, some convolute bedded; some units ripple laminated towards top; sporadic beds with mudstone flakes. Units 3–15 m thick. Unit base sharp, irregular, erosive. Minor bioturbation, rooting at top. Plant debris, ' <i>Stigmaria</i> ' pieces.	Spilmersford, East Linton, Birnieknowes, Marshall Meadows, Blairmulloch; Edington Mill	Fluvial channels	Fluvial
Volcaniclastic sandstone	'Tuff', 'agglomerate', and tuffaceous siltstone and sandstone; grain size from medium to coarse sand, granules and pebbles, coarsest up to 15 cm; clasts of green amygdaloidal 'lava'; dolostone, siltstone; bedded units up to 2.5 m thick, occurring sporadically and in packages up to 8 m.	East Linton, Spilmersford	Reworked older volcanic deposits and ripped-up substrate, probably fluvial	

the boreholes, collected over a period of more than 100 years. The fossils have been re-examined for this study to ensure taxonomic consistency (Brand 2018). The biogeographical distribution of marine taxa and '*Estheria*' are included herein as they provide additional insight alongside the sedimentary data. Raymond (1946, p. 219) noted that '*Estheria*' is an invalid name for a conchostracan genus as the name was pre-occupied, and he erected a number of genera in which to allocate the known species. Unfortunately, he did not allocate all the Scottish described species to his genera, and preservation of the material from the Ballagan Formation often precludes even a generic allocation.

3. Sedimentary facies

Facies identified in selected exposures and the archived borehole logs are described in Table 1, and examples illustrated in Figure 3. Summary statistics for selected features in the borehole logs are given in Table 2. The stratigraphical variation of facies across the Midland Valley is shown in Figures 4 and 5.

The succession in the west and north of the Midland Valley mostly comprises small-scale intercalations of packages of dolostone and grey siltstone, along with heterogeneous siltstone and fine-grained sandstone, and locally, evaporite rock; these abruptly interdigitated facies constitute the saline and hypersaline lake and overbank facies associations (Figs 3, 4). Laterally continuous distinctive units that might be suitable for correlation across the region are not present, though the uppermost part of the formation is sand-rich in parts. At the base of the formation in Fife and Tayside, a unit of distinctive dolostone beds, 32 m thick, comprises the Mains of Errol Member (Browne 1980). By contrast, in the east, the succession is more heterogeneous with thick distinctive packages of, for example, fluvial sandstone and floodplain-lake siltstones within a mixed assemblage of fluvial and overbank facies associations (Figs 3, 5). The position of the base of the Pu Miospore Biozone suggests that some of the packages in East Lothian may be correlated, at least over a distance of about 15 km.

3.1. Siltstone, dolostone and evaporite facies

Thin beds of micritic ferroan dolostone interbedded with planar laminated, grey siltstone are the defining characteristic of the Ballagan Formation. Siltstone occurs in both the saline and hypersaline lake facies association, and the overbank facies association (Fig. 3). Laminated grey siltstone comprises a high proportion of the fines component in some of the boreholes, for example 99% in Blairmulloch and 89% in Loch Humphrey (Table 2), but is generally lower in the East Lothian boreholes. Thick packages of laminated grey siltstone in the boreholes with high proportions of fine sediment are likely to represent floodplain-lake deposits, though these may include misidentified thin beds of sandy siltstone. Laminated grey siltstone associated with the upper part of massive and cross-bedded sandstone may fill fluvial channels. The overbank facies association includes grey, yellow, red and purple siltstone, only some of which is described as having traces of bedding; these may be palaeosols, but the descriptions are insufficient for a diagnosis.

After removing the sandstone component from each borehole, dolostones comprise 3–21% of the thickness of the remainder of the succession. The number of dolostone beds varies from nine in the Ascog Borehole to 277 in Norham, with the average thickness ranging from 9 cm in Ascog to 34 cm in East Linton (Table 2). Thus, in the boreholes of the western and northern part of the region, dolostone beds occur on average every 0.9 m (Glenrothes) to 1.7 m (Knocknairshill, Blairmulloch; Fig. 3). By contrast, dolostone beds are, on average, less common in

the East Lothian boreholes, but there the concentration of dolostone beds within packages of laminated grey siltstone up to 35 m thick is strongly evident, with significant packages of overbank siltstone and fine-grained sandstone devoid of dolostone beds (Fig. 5).

Apart from the thickness, few descriptive details of the dolostone beds are typically given, though some are described as ironstone (Davies *et al.* 1986; Greig 1988). Belt *et al.* (1967), Francis *et al.* (1970) and Scott (1986) identified dolostone nodules, laminated beds (Fig. 8d), evaporite-bearing beds and massive beds; the last of these become more argillaceous at the base and top. Identification of these differing facies is aided by slicing the cores, but information from the borehole logs is insufficient to make the distinctions in all but the evaporite-bearing dolostones.

Small-scale evaporite deposits are a characteristic of the Ballagan and other Tournaisian formations in the British Isles (Millward *et al.* 2018). They are distributed spatially and temporally throughout the succession in the Midland Valley and Tweed Basin, though they are absent from the Lothians and from the area bordering the Highlands (Figs 4, 5). Nodules of gypsum occur in siltstone and dolostone; calcite and dolomite pseudomorphs occur in the latter at Burnmouth and in some of the boreholes (Scott 1986; Millward *et al.* 2018). Siltstone pseudomorphs after hopper crystals of halite are recorded locally in boreholes (Fig. 4) and in exposures, for example in the Langholm area, Chattlehope Burn, Northumberland and Willie's Hole, Scottish Borders (Leeder 1974b; Cater *et al.* 1989).

3.2. Mains of Errol Member

The Mains of Errol Member (Fig. 4; Browne 1980) comprises beds of hard dolomitic, with a characteristic reddened, cracked and 'pseudobrecciated' profile (Fig. 6). The top of each profile is a sharp planar contact. Roots and burrows are absent; ostracods and fish fragments are rare and other fossils absent. Dolostone beds, 0.22–1.70 m thick, are either stacked on top of each other, or they are intercalated with units of laminated siltstone or sandy siltstone. The dolostone is composed of planar-textured subhedral dolomite with crystal size up to 10 μm . A pseudobrecciated texture shows 'clasts' of finer grained dolomite set in coarser grained dolomite. Quartz grains scattered throughout the rock are etched, suggesting corrosion in alkaline fluids. A few rhombohedra composed of dolomite aggregates are probably pseudomorphs after gypsum. Scattered clasts up to 500 μm composed of chlorite and feldspar may be volcanic in origin.

From just above each dolostone unit base, isolated, narrow sinuous cracks widen upwards as they unite with adjacent ones (Fig. 6). At their upper termination against subhorizontal planar dislocations these cracks are polygonal and up to 20 mm across on the bedding surface. The cracks are filled with iron (Fe)-rich claystone. Above the cracked zone, 20–40 cm of dolostone are intensely brecciated, with poorly sorted, angular clasts of dolostone from 1 to 20 mm across, set in a Fe-rich claystone. In this zone and the top part of the underlying cracked zone, low-angle, concave-upwards dislocations resemble pedogenic slickensides. In the most fully preserved profiles the uppermost 5–10 cm are composed mainly of red-brown, Fe-rich claystone, with sporadic dolostone clasts, 1–5 mm across. Similar beds have been reported from a borehole in the Stirling area (Francis *et al.* 1970, plate 4, fig. 7).

3.3. Sandstone

Sandstone facies occur within the fluvial and overbank facies associations (Bennett *et al.* 2016). All sandstone contains




Figure 3 Sedimentary logs of sections drawn up from the archived BGS borehole logs to illustrate features of the facies associations. Glenrothes Borehole: flood-plain lake facies, saline and hypersaline lake facies association intercalated with overbank facies association; original written log by D. N. Halley, A. A. McMillan and M. A. E. Browne. East Linton Borehole: overbank facies association, with subordinate saline and hypersaline lake facies association; original written log by A. D. McAdam. Birnieknowes Borehole: fluvial facies associations, with overbank facies association and floodplain lakes; author of original written log not recorded.

Table 2 Summary statistics for Ballagan Formation in selected BGS boreholes. Abbreviations: BGN = Ballagan Formation; CYD = Clyde Sandstone Formation; FESD = Fell Sandstone Formation; GHV = Garleton Hills Volcanic Formation; GUL = Gullane Formation; KNW = Kinnesswood Formation; LLAGS = Lower Limestone Formation.

Borehole name	BGS Registered number	BGN thickness metres	Unit above	Unit below	Sandstone % of formation	No of beds	As % of fines	Dolostones (cementstones)					Number rooted horizons/palaeosols	Number evaporite units	% fines grey and bedded
								average bed thickness centimetres	bed frequency, metres, sand free	Number desiccation crack horizons	Number beds burrowed/bioturbated	Number rooted horizons/palaeosols			
Ascog	NS06SE 8	33.09	CYD	KNW	27.8	9	3.4	9.1	2.65	0	0	0	0	0	n.d. ¹
Knocknairshill	NS37SW 10	70.87	CYD	KNW	20.6	33	7.5	12.9	1.70	8	0	0	0	17	49
Barnhill	NS47NW 2	123.08	CYD	KNW	8.0	82	11.4	15.7	1.38	2	6	0	0	30	85
Loch Humhrey	NS47NE 1	162.95	CYD	Not reached	6.7	126	12.4	15.1	1.21	6	2	0	0	64	89
East Dron	NO11NW 24	209.54	Not seen	KNW	9.7	127	10.9	16.3	1.49	110	45	10	10	114	57
Glenrothes	NO20SE 385	157.43	PDB	KNW	7.5	159	18.0	16.5	0.92	137	21	2	2	47	83
Deaconhill	NS43SE 81	150.03	LLGS	KNW	1.9	94	16.0	25.1	1.57	4	9	2	2	12	94
Blairmulloch	NS52NE 21	166.58	CYD	Not reached	14.8	83	11.5	19.65	1.71	3	21	1	1	34	99
Spilmersford	NT46NE 73	258.24	GHV	Fault	31.9	63	8.3	23.1	2.82	7	24	2	2	0	44
East Linton	NT57NE 2	418.17	GHV	Not reached	23.8	59	6.6	34.1	5.41	15	62	26	26	0	42
Birnknowes	NT77SE 9	200.21	GUL	KNW	46.5	33	5.0	16.3	3.24	2	20	1	1	0	59
Norham	NT94NW 20	490.50	Not seen	Not reached	35.5	277	21.5	24.5	1.14	131	128	216	216	50	16

¹ Log descriptions of mudstone and siltstone units in the Ascog Borehole contain insufficient detail for analysis.




Figure 4 Summary lithological logs and environmental interpretations for selected boreholes in the west and north of the Midland Valley of Scotland. Borehole locations are shown in Figure 1. The black curves within the stratigraphical columns represent the proportional thickness of grey, laminated siltstone/mudstone per 10 m interval through the succession. The key to the lithology colours is given in Figure 5.




Figure 6 Sedimentary logs through selected dolostone units in the Mains of Errol Member in the Glenrothes cores.

abundant partings of mica and carbonaceous detritus. The proportion of sandstone in the succession varies across the region (2–46%; Table 2), with the higher amounts in East Lothian and the Tweed Basin (24% in East Linton Borehole, 32% in Spilmersford and 46% in Birnieknowes) (Fig. 7). By contrast, in boreholes in the north of the Midland Valley, sandstone is most commonly seen at the base and top to the formation, though in the Ascog Borehole it occurs throughout (Fig. 4). Units up to 3 m thick occur in all the boreholes, with the median occurrence less than 50 cm thick (Fig. 7). Typically, these thin units of sandstone are described as fine-grained and ripple-laminated or cross-laminated. Many are interbedded with siltstone and are interpreted as deposited by overbank flooding and crevasse splays. A few units in this interval are recorded as having a sharp, erosive base and massive or cross-bedded and may represent channel sandstone bodies.

By contrast, the Blairmulloch and East Lothian boreholes also contain sandstone units more than 3 m thick (Fig. 7). Though the number of such units is small, they make up a high proportion of the total thickness of the sandstone component in the succession; for example, three units in Spilmersford constitute 31% of the sandstone thickness and in Birnieknowes 63% (Table 2). These sandstone bodies are described as sharp-based, medium and coarse grained, massive or cross-bedded (Fig. 3) and may include mudstone rip-up clasts and conglomerate layers in the lower part. A thin upper part typically is finer grained and planar laminated to ripple laminated. Thicknesses vary from just a few metres up to 15 m in the Spilmersford, East Linton and Birnieknowes boreholes. The 11.4 m thick sandstone unit illustrated in Figure 3 is typical of many, comprising stacked subunits separated by a thin bed of siltstone, suggesting that the thicker sandstone bodies represent multiple-storey channel fills. Lenticular beds of conglomerate at the base of sandstone units at Burnmouth host disarticulated vertebrate bones.

The geometry of one of these sandstone units is seen in the river cliff of the Whiteadder River at Edington Mill [NT 894

548] (Fig. 8a, c). The lower part of the channel cuts sharply down about 4 m into the underlying siltstone-dolostone succession. Rotated blocks of siltstone-dolostone suggest that the channel margin collapsed during formation (Fig. 8c). The channel fill shows lenticular units of cross-bedded sandstone with lateral accretion at the channel base. Clasts of dolostone in the thalweg can be traced to a bed cut by the channel. Beds in the upper part of the channel thicken to the right in Figure 8a. Another unit, more than 20 m thick, near the top of the formation, and exposed along the SE bank of the River Tweed at Norham is strongly trough cross-bedded with sediment transport consistently towards the SW.

At the coastal locality of Burnmouth in the Tweed Basin, a number of sandstone fluvial bodies are exposed in the almost vertical strata of the wave-cut platform and cliff. Depicted on a geological excursion map, Scrutton & Turner (1995) numbered the large sandstone bodies 1 (youngest) to 14 (oldest). Analysis of these bodies indicates that a number are multistorey and multilateral, and can be traced across the wave-cut platform for up to a kilometre. Lateral accretion deposits (2–3 m thick) are recognised in several bodies. Whereas individual channels may only reach 3 m in thickness, one or two of the multistorey systems comprise a total in excess of 30 m. In general, the erosive bases to the channels cut into the underlying floodplain sedimentary rocks, but some systems cut into the dolostones, representing lake deposits.

Poorly bedded volcanoclastic rocks within the Ballagan Formation are described here for the first time from the Spilmersford and East Linton boreholes (Fig. 5). They occur both within, and at the top of, the formation, interbedded with sandstone and siltstone. A unit in the lower part of the East Linton Borehole is 7.9 m thick with beds ranging from 24 cm to 1.5 m thick. They are composed of multicoloured fragments of volcanic rock of medium sand through gravel grade, with some dolostone, siltstone and sporadic chert clasts. These rock types have not been reported from elsewhere in the formation.

3.4. Emergent surfaces and palaeosols

Desiccation cracks and brecciated, upper parts of beds, both at outcrop and in boreholes, represent short-lived episodes of emergence and drying out of the sediment. At least five horizons are seen, for example, in the upper part of the shallowly dipping succession in the well exposed coastal section at Belhaven Bay, west of Dunbar (Fig. 8), but there is great variation in abundance across the Midland Valley (Fig. 9; Table 2). Glenrothes and East Dron borehole logs contain 138 and 110 horizons respectively, similar to 131 in the Norham core of the Tweed Basin (Figs 3, 9). By contrast, the very low numbers of desiccated surfaces in the other boreholes may inversely relate to the high proportion of laminated grey mudstone present, representing extended periods of sedimentation in perennial bodies of standing water. Alternatively, the low numbers may reflect under-recording of detail.

The preservation of plant roots and palaeosols indicates established episodes of emergence and vegetation cover. Carpenter *et al.* (2015) interpreted the mottled red mudstone beds containing carbonate nodules and vertic cracks within the thin succession on the Isle of Bute as well drained, seasonally dry palaeosols. Near Foulden, Retallack & Dilcher (1988) inferred that gleyed inceptisols were populated by small shrubby ferns such as *Lyrasperma scotica* and vertisols by tall forest trees such as the reconstructed *Stammotoma huttonense*. Kearsy *et al.* (2016) described 216 palaeosols from the Norham core and 64 from the exposed section at Burnmouth in the Tweed Basin. Nineteen vertisols occur in the Norham core and 15 at




Figure 7 Histograms for key Midland Valley boreholes showing the frequency of sandstone bodies of a given thickness interval. The percentages shown for the thicker units are of the total sandstone thickness in the borehole succession.

Burnmouth, representing extended periods of stabilised floodplain. Some 60 % of the vertisols have a gleyed top, indicating that the soils were finally flooded by surface waters (Kearsey *et al.* 2016).

Few palaeosols are described in the archived borehole logs (Table 2). In the Glasgow and Ayrshire logs there are sporadic mentions of ‘seatearths’ and ‘seat clay’, typically associated with carbonised root traces; the absence of coals capping these suggest that they probably equate to the gleyed inceptisols of Kearsey *et al.* (2016). Roots in the top of sandstone beds are likely to be the entisols described by Kearsey *et al.* (2016). However, the depth and form of rooting is not given. Massive, red-, brown- and yellow-mottled siltstone, particularly in the East Lothian boreholes and sporadically elsewhere, might suggest that vertisols are also present there, but indicative evidence of vertic cracks and pedogenic slickensides is missing. However, in many of the logs from the west and north of the Midland Valley, the high proportion of laminated grey siltstone present would seem to preclude the presence of well-

developed palaeosols from those parts of the succession. This is also the case with the Glenrothes core, visual inspection of which showed the presence of very few palaeosols of any type.

3.5. Sandy siltstone

Matrix-supported sandy siltstone with millimetre-sized rock clasts and bioclasts was first described from the Ballagan Formation from the Norham core and at Burnmouth by Bennett *et al.* (2016). The 146 and 71 beds respectively from the two sections are mostly less than 1 m thick; sandy siltstone makes up about 6 % of the succession and typically overlies palaeosols or desiccated horizons. The deposits have lateral extents of up to several tens of metres at Burnmouth. The beds are inferred to have been generated by episodes of high rainfall and deposited as unconfined, cohesive flows across the floodplain, forming temporary pools and small lakes. Bennett *et al.* (2016) suggested that because of the small size of the clasts, the facies could be misinterpreted as a massive siltstone, with colour varying from




Figure 8 Images illustrating features of the Ballagan Formation. (a) Fluvial channel cut into interbedded siltstone, fine-grained sandstone and dolostone alongside the Whiteadder Water at Edington Mill [NT 894 548]. Detail in white rectangle in (c). Fisherman, bottom centre, for scale. (b) Desiccation cracks, Dunbar shore. (c) Detail of channel side, showing multiple dislocations probably caused by collapse of the channel bank. (d) Laminated dolostone, Glenrothes borehole, depth 305.65 m; core 95 mm wide. (e) Sandy siltstone in Mains of Errol Member – dolomitised siltstone containing small dark fragments of siltstone; Glenrothes borehole, depth 358.21 m; core 78 mm diameter. (f) Scan of thin section of oncolite-bearing limestone. Located at top of the formation, just beneath the Fell Sandstone, Coomsdon Burn [NY 710 038].




Figure 9 Occurrence of sedimentary features and fossils recorded in selected boreholes across the east of the Midland Valley.

black, grey and red to light green. Thus, it is likely to be that the sandy siltstone facies is more widespread in the Ballagan Formation than hitherto reported. This facies is not described in many of the borehole logs throughout the Midland Valley, but inspection of the Glenrothes core revealed at least three beds in the Mains of Errol Member. Described in the logs as ‘mudstone breccia’, these are 10–44 cm thick and composed of greenish grey (?) dolomitic siltstone with elongate irregular clasts of dark grey siltstone (Fig. 8e).

3.6. Fauna and flora

The Ballagan Formation has a generally sparse and low-diversity invertebrate fauna dominated by bivalves (particularly *Modiolus latus*), ostracods, shrimps and other arthropods (Cater *et al.* 1989), along with rarer occurrences of *Spirorbis*, *Serpula* and scolecodonts (Bennett *et al.* 2017). In a few beds invertebrates may be locally common (Brand 2018), and *Modiolus* and ostracods can occur in large numbers (Williams *et al.* 2005, 2006). By contrast, the vertebrate fauna appears to have been diverse with actinopterygians, dipnoans, chondrichthyans, acanthodians and tetrapods (e.g., Andrews 1985; Gardiner 1985; Clack 2002; Clack & Finney 2005; Smithson *et al.* 2012, 2016; Carpenter *et al.* 2015; Clack *et al.* 2016; Richards *et al.* 2018; Smithson & Clack 2018). A varied flora has been reported from plant elements and palynomorphs (Scott & Meyer-Berthaud 1985; Retallack & Dilcher 1988; Bateman & Scott 1990; Stephenson *et al.* 2004a, b); furthermore, indeterminate plant debris is a ubiquitous component of many lithofacies.

In the BGS fossil collections (Brand 2018), most of the invertebrate species from the Ballagan Formation are considered to be either non-marine or euryhaline. Important indicators of terrestrial lakes are the sporadic conchostracans which occur in the Midland Valley and Tweed Basin, but which are absent from the Northumberland–Solway Basin (Fig. 10). Recorded from several levels in most of the boreholes (Figs 4, 5, 9), they are particularly abundant in the East Linton Borehole (340.66–347.50 m depth), where specimens were recorded from eight levels within 8 m of dark purple and grey, bedded or poorly bedded, mudstone and siltstone.

There is no sedimentary evidence for fully established open marine conditions across the Midland Valley and Tweed Basin: for example, carbonate rocks of this type are absent, as are successions indicative of the upward transition from storm wave-base to shore face. Beach and tidal deposits have also not been reported. However, marine fossils are recorded sporadically. Typically, the occurrences are restricted to a single taxon per interval and include orthocone nautiloids, the bivalves *Schizodus pentlandicus?*, *Phestia traquairi*, *Sanguinolites* cf. *clavatus*, *S.* aff. *striatolamellosus* and *S.* cf. *roxburgensis*, *Cardiomorpha*, very rare rhynchonellid brachiopods and *Lingula mytilloides*. The shells are disarticulated or fragmentary, except at one level (496.17 m depth) from the Norham Borehole where an articulated *Phestia traquairi* was found vertical to bedding, in a life position. *Sanguinolites*, *Cardiomorpha*, orthocones and brachiopods occur in the west of the Midland Valley, notably in Ayrshire but also near Glasgow (Fig. 10). Also, four marine intervals are present in the lower




Figure 10 Map showing the locations of fossil records from the Ballagan Formation in the BGS biostratigraphy collections. The distribution of marine taxa (see Section 3.6) and ‘*Estheria*’ and *Leaia* (conchostracan genera) are highlighted.

part of the Blairmulloch Farm Borehole. Similarly, an orthocone and *Sanguinolites* occur near the top of succession in the East Dron Borehole. Larger marine macrofossils occur at five intervals in the Norham core. In stark contrast, at outcrop along the northern flank of the Northumberland Basin, records of marine species are more widespread and assemblages from individual localities are more varied (Fig. 10; Brand 2018).

A significantly greater cryptic marine influence has been demonstrated from 128 thin beds in the Norham core of a *Chondrites* ichnofacies, particularly within dolostones of the saline–hypersaline lake facies association (Bennett *et al.* 2017). These occurrences are described as single-tier colonisations, with a high ichnofabric index. By contrast, very few bioturbated cementstone beds were recorded in the Midland Valley borehole logs, where burrows and bioturbation are most commonly recorded in mudstone and siltstone (Table 2; Figs 3–5). ‘Flat’ burrows were noted in some finer grained rocks and simple vertical burrows in sandstone, but there are few records of the ichnotaxa present. Exceptions are in the East Linton Borehole where ?*Monocraterion* and *Planolites* were recorded from ripple-bedded sandstone. Eighteen beds containing scolecodont (marine polychaete) jaws were identified from grey siltstones in the Norham core (Bennett *et al.* 2017) providing more evidence of cryptic marine incursions.

4. Palaeogeography

The new map (Fig. 11) combines elements of the palaeogeography from throughout the Tournaisian, a period of

some 12 million years. The map shows areas of more elevated land with a coastal floodplain occupying a broad swath through the Midland Valley into the Tweed Basin, and connected east of the Southern Uplands and Cheviot massifs to a shallow seaway in the Northumberland–Solway Basin. The western Midland Valley is connected through Northern Ireland to full marine conditions in the south of Ireland (Clayton & Higgs 1979; Clayton *et al.* 1986). At times, shallow saline lagoons were established in the Belfast Lough area, as evidenced by the presence of stromatolites at Cultra (Clayton 1986), and the thick succession of dolostone and evaporite containing marine invertebrates proved in the Belfast Harbour Borehole suggest restricted marine conditions (Smith 1986). The coastlines migrated from time to time in the Northumberland–Solway Basin (Sherwin 2018), but storm-driven marine flooding (Bennett *et al.* 2017) reached much greater extents into the western Midland Valley and in the east.

Over most of the region, floodplain lakes and overbank facies are juxtaposed. Three floodplain-lake successions hosting significant fish communities are indicated on the Isle of Bute (Carpenter *et al.* 2015) in the Midland Valley, at Foulden in the Tweed Basin (Clarkson 1985; Wood & Rolfe 1985) and in Coomsdon Burn on the flanks of the Northumberland–Solway Basin (Fig. 11; Moy-Thomas 1938). Detailed distributions are given where known for small-scale temporary inland ponds (see below), sabkhas and perennial hypersaline lakes; the early Tournaisian Mains of Errol alkaline lakes in Fife and Tayside are also shown. Forested areas are known from the Tweed Basin and on the Isle of Bute, but were probably more extensive, particularly adjacent to upland massifs. The approximate




Figure 11 Palaeogeographic map of the Ballagan Formation.

likely extent of the belt of fluvial sandbodies that dominated the east of the Midland Valley and Tweed Basin in the upper part of the succession is indicated (Fig. 11). Alluvial fans along the margin of the Southern Uplands and Cheviot massifs are early features. The environments of the Northumberland–Solway Basin are described only briefly for context.

4.1. Massifs

Tournaisian strata are absent from the isolated outcrops of Carboniferous rocks in the western part of the Highlands of Scotland, which has long been regarded as terrestrial in the Tournaisian. This upland may have extended south to the Ochil Fault (Figs 1, 11), though this is difficult to prove. The Lower Devonian Ochil Volcanic Formation, which today forms the eponymous hills, comprises 2400 m of largely sub-aerial volcanic rocks that interdigitate north-eastwards with fluvial and lacustrine rocks of the Arbutnott Group (Stephenson *et al.* 1999). Though the sedimentary strata may have been buried by up to about 3 km of later strata by the late Carboniferous (Marshall *et al.* 1994), it is possible that the core of the volcanic massif in what is now Fife and Tayside remained elevated through early Carboniferous times.

South of the Midland Valley, an upland massif of Lower Palaeozoic rocks is inferred to approximate to the area of the modern-day Southern Uplands, extending SW to include the Longford–Down massif in Northern Ireland. The Sanquhar and Thornhill coalfields crop out in a northerly trending half graben within the Southern Uplands. The oldest Carboniferous rocks are Pendleian in the former coalfield and upper Viséan in the latter (Davies 1970; McMillan 2002). Furthermore, on the northern margin of the Northumberland–Solway Basin northerly derived palaeocurrent directions in the Whita Sandstone at the base of the Ballagan Formation suggest that the massif was a source of sediment and probably remained significantly elevated for much of the Tournaisian (Nairn 1958; Leeder 1974b).

Though the lower Tournaisian basalts of the Northumberland–Solway Basin margin were rapidly buried by the Ballagan Formation (Leeder 1974a), the eroded remains of the Middle Devonian Cheviot volcano probably stood above the floodplain. Relict outcrops of Ballagan Formation in the intervening ground suggest that the Cheviot massif was separated from the Southern Uplands. On the flanks of the Cheviot massif, alluvial fan conglomerates at Roddam Dene, Ramshope Burn and Windy Gyle reworked the volcanic rocks and granite (Purnell & Cossey 2004). These rocks are thought to be coeval with the Kinnesswood Formation (Stone *et al.* 2010) and are therefore latest Devonian in age (Marshall *et al.* 2019). Seismic mapping and borehole interpretation across the Mid North Sea High suggest that several small elevated areas also persisted eastwards across the coastal plain (Arsenikos *et al.* 2018; Kearsey *et al.* 2018), like an archipelago separating the Tweed and Northumberland–Solway basins. To the south of the latter basin, the Manx–Lake District–Alston Block was also terrestrial at this time, with fluvial (possibly as valley fills) and alluvial fan deposits of the Marsett Formation on the northern flank of the Lake District and on the present-day Pennine escarpment, suggesting that that block had some relief (Burgess & Holliday 1979).

Pointers to the location of the margins of upland areas are indicated in the uppermost part of the underlying Kinnesswood Formation, by local occurrences of massive carbonate rock, several metres thick and typically associated with stringers, nodules and irregular masses of chert; these have been interpreted as super-mature calcretes (Wright *et al.* 1993). Examples occur in Bute (Young & Caldwell 2011), Ayr (Burgess 1961), at Pease Bay [NT 795 712], east of Dunbar (Clough *et al.* 1910; Andrews *et al.* 1991) and in the Tweed valley, about 5 km SW of Coldstream, where the Carham Limestone is at least 7.5 m thick (Carruthers *et al.* 1932). These are highly significant as they may represent disconformity with a significant time span (Ettensohn *et al.* 1988; Wright *et al.* 1993;

Alonso-Zarza 2003). The calccrete-bearing palaeosols may have formed in areas elevated above the floodplain for a long time, perhaps up to several million years, for example as terraces adjacent to upland areas or through tectonic uplift, or on terraces formed by uplift associated with volcanism (Leeder 1976; Leeder & Bridges 1978).

4.2. Coastal floodplain of the Midland Valley and Tweed basin

Over the western and northern parts of the Midland Valley boreholes show abrupt intercalations of facies belonging to the saline–hypersaline and overbank facies associations. Long core runs of grey laminated siltstone suggest perennial floodplain lakes dominated much of the area through the Tournaisian. The common fauna of these rocks are ostracods, which were locally monospecific and in great abundance (Stephenson *et al.* 2004a; Williams *et al.* 2006). Stable isotope analysis of the ostracod shells from different communities indicates that salinity of the lakes varied from brackish to saline and hypersaline (Williams *et al.* 2006), though later work indicated that the ostracods' calcite carapace had been recrystallised as a result of diagenetic alteration (Bennett *et al.* 2011).

It is difficult to estimate the size of the floodplain lakes. Unpublished studies of the laminated siltstone beds and dolostones at Burnmouth and in the Norham core by the authors suggest deposition in very shallow lakes up to a few kilometres across, based on the outcrops of planar beds of dolostone and their thickness on the foreshore at Burnmouth. Dolostones can be traced some 500 m or so, though some are less extensive, but no beds can be correlated between Burnmouth and Norham, a distance of about 10 km. None of the lakes was thought to have been long-lived, with mud-cracks at the top of many of the thin units indicating drying out of the surface. However, the area west and north of Glasgow was repeatedly occupied by lakes throughout most of the formation. Similar lacustrine-dominated successions are recorded in the middle of the formation in the area between Spilmersford and East Linton in East Lothian, some 15 km apart, suggesting a substantial area that was persistently occupied by lakes.

Thin intervals of overbank flood and crevasse-splay deposits become dominant sedimentary environments in the upper part of the formation around the Isle of Bute and west of Glasgow. Few palaeosols are recorded from this area, though in the East Dron and Glenrothes boreholes in the NE dried out surfaces are common (Table 2; Fig. 3). Vegetated land seems to have been sparse as indicated by the dominance of floodplain lakes. These features are well demonstrated by Figures 3 (Glenrothes Borehole) and 4. Correlation between boreholes is not possible, suggesting a mosaic of juxtaposed and ever-changing sedimentary environments. No aeolian deposits are known, though differentiating these in fine-grained successions may be difficult.

A more complex picture occurs in the SE of the Midland Valley and in the Tweed Basin where overbank facies dominate and fluvial channel systems developed (Figs 3, 5). Brackish to saline floodplain lakes continue throughout, but are particularly concentrated in two 30–40 m thick intervals of strata in the upper Tournaisian, as indicated by the base of the Pu Miospore Biozone (Neves & Ioannides 1974; McAdam & Tulloch 1985). Waterlogged palaeosols appear to be more common in this region. Recognition of these is important because these are the likely habitat of the creeping lycopod *Oxroadia*, the spores of which are abundant where tetrapods have been located in the Burnmouth section (Clack *et al.* 2016). In the upper part of the succession in the Tweed Basin (Kearsey *et al.* 2016) and on the Isle of Bute (Carpenter *et al.* 2015), well-developed reddened vertisols indicative of forests

demonstrate areas of greater floodplain stability. Reddened siltstone and mudstone constitute a significant proportion of boreholes in the SE of the Midland Valley, suggesting that vertisols may have developed more widely in this region, but descriptions are insufficient to be secure in the interpretation.

4.2.1. Perennial floodplain lakes with vertebrate communities.

Fragmentary vertebrate material including teeth, scales and spines is a common component of many of the floodplain lake successions. However, at least three of these successions have revealed more abundant and complete specimens within diverse communities that included bivalves, malacostracan crustacea and *Spirorbis* and these sites are located in Figure 11. The best studied of these is the Foulden Fish Bed, which occurs at the top of a lacustrine succession that began development as a lycopod wetland, became a semi-permanent floodplain lake and finally was silted-up to return the area to a floodplain environment (Clarkson 1985; Wood & Rolfe 1985). The abundant complete fish remains and beds of shrimps are likely to represent mass mortality events. Kills of this sort are well known in the seasonal tropics today, for example in northern Australia, where they result from sudden changes in water quality caused by the influx of sediment at the onset of the wet season (e.g., Townsend *et al.* 1992). The precise position of the Foulden Fish Bed within the Ballagan Formation is not known, though its outcrop location and its CM Miospore Biozone age (Clayton 1986) suggest the upper part.

Moy-Thomas (1938) described six species of fish from Coomsdon Burn [NY 710 038], in north Northumberland that resemble those seen at Foulden. Exposure in the stream banks today is poor and it is difficult to interpret the sedimentary context of these fossils from the two contrasting published sedimentary logs of the succession by Day (1970, p. 280) and Cater *et al.* (1989, fig. 6). However, the presence of nearly complete specimens of both fish and shrimps suggests similar palaeoenvironments in the upper part of the formation. Carpenter *et al.* (2015) described two fish beds, possibly from two different stratigraphical levels, on the Isle of Bute. Though the material is mostly fragmentary, it contains possible juveniles and the community occupied shallow fresh- to brackish water lakes.

4.2.2. Temporary inland ponds. The presence of abundant, small temporary lakes is revealed from the widespread occurrence of conchostracans across the Midland Valley and Tweed Basin (Fig. 10). Tasch (1969, pp. 146–49) described the main habitat of modern conchostracans as small temporary inland ponds, generally no larger than about 0.4 hectares (1 acre) in extent. Modern forms have also been recorded from spring water, from the margins of large lakes in Africa and from coastal salt flats in Brazil. Further, he found that fossil forms may have inhabited temporary pools close to coastal areas subject to fluctuating tides, possibly in lagoons or at shorelines subject to marine inundations. Gueriau *et al.* (2016) also suggested that conchostracans typically inhabit temporary freshwater environments. Their discovery of resting eggs associated with conchostracans in a late Famennian deposit at Strud in Belgium suggests a mechanism of dispersal from such temporary environments and offers a capability of surviving seasonal desiccation.

4.2.3. Sabkhas and hypersaline lakes. In the Tweed Basin evaporite rocks are most abundant in the lower part of the succession, but elsewhere these rocks occur throughout the formation (Fig. 4). In contrast, evaporite rocks are notably absent from the SE part of the Midland Valley. Most of the evaporite occurrences are located within areas where there is evidence for marine flooding (see below). Millward *et al.* (2018) have described this unusual feature of seasonal coastal wetlands that developed in evaporative closed hypersaline

lakes and perennial brine pans. Sabkhas, marsh and microbial mats surrounded some of these. Floodplain lake–evaporite cycles were illustrated from the Tweed Basin by Millward *et al.* (2018, fig. 6) whereby flooding events over a palaeosol surface established a floodplain lake with a low-diversity ichnofauna and invertebrate assemblage. Subsequent evaporation led to deposition of dolomite and hypersalinity ultimately to gypsum; saline groundwater and marine flooding events recharged the lakes.

4.2.4. Mains of Errol alkaline lakes. The reddened and pseudobrecciated dolostone beds that comprise most of the Mains of Errol Member in the NE of the Midland Valley probably represent similar lakes that were hypersaline and alkaline: repeated episodes of dolomite precipitation in shallow evaporating lakes, which dried out and desiccated. The very sparse, low-diversity fauna and lack of bioturbation suggest that these lakes were not conducive to life. The absence of roots suggests that the dried out lake beds were barren.

4.2.5. Fluvial systems. The occurrence of thick packages of fluvial sandstone and dominance of overbank siltstones over floodplain lakes suggests that a major fluvial system persisted for much of the Tournaisian in the east of the Midland Valley. Palaeocurrent indicators show sediment transport to the SW in the Tweed Basin, presumably contributing to the sediment budget of the Northumberland–Solway Basin. The fluvial systems appear to be integral components of the palaeo-environment (Bennett *et al.* 2016), and no valley systems are recognised at outcrop. Lateral accretion deposits characterise some of the sandstone bodies, suggesting that meandering systems dominated, though braided systems were also present. At times, sand appears to have also been diverted by the NE extent of the Southern Uplands massif along its Midland Valley margin. However, the SW extent of these sandstone bodies is unclear, as very little sand characterises the Ayrshire succession. Though the source of the sand is not known because no heavy mineral studies have been published yet, derivation is likely from somewhere northwards along the North Sea to the Greenland or Norway Caledonides mountains (Cliff *et al.* 1991; Coward 1993).

In the rest of the Midland Valley fine sand appears to have been distributed mostly by overbank flooding and small streams. In the area north of Glasgow, the small proportion of fine sand suggests that there was no significant volume of sediment supplied from the north. Late in the Tournaisian in Fife, overbank flooding and small streams supplied sediment, either from the north, or more likely from the major river system. Some alluvial fan deposits were likely localised along the northern flank of the Southern Uplands massif in Ayrshire for example (Monaghan 2004).

The intercalations of volcanoclastic rocks are localised to a small area adjacent to the NE extent of the Southern Upland massif. In the written logs, use of the terms tuff and agglomerate implies that these rocks represent primary pyroclastic deposits rather than contemporaneously re-sedimented ash or eroded and reworked older volcanic rocks. Such a distinction is difficult to make from the descriptions, particularly as other sedimentary lithologies form a component of some beds and could have been included as pyroclasts or through erosion of substrate material. However, the volcanoclastic beds intercalated with Ballagan facies at the top of the formation, beneath the Garleton Hills Volcanic Formation (Fig. 5), probably represents the initial explosive eruptions, or contemporaneous fluvial reworking of such deposits.

The derivation of the volcanoclastic rocks at lower levels in the formation in the East Linton and Spilmersford boreholes (Fig. 5) is enigmatic. Though basaltic volcanism preceded the Ballagan Formation in the Northumberland–Solway Basin,

the only record of contemporary volcanism elsewhere in northern Britain during the Tournaisian is at Heads of Ayr (Stephenson *et al.* 2002). Thus, it seems likely that the volcanoclastic rocks in the east of the Midland Valley were eroded from older outcrops, transported and re-deposited in river systems. The closest source is the Lower Devonian volcanoes near Eyemouth and these may have had a more extensive outcrop at the time. However, no palaeocurrent or geochemical data are available to support this inference.

4.2.6. Marine flooding. The sporadic occurrences in the succession of this area yielding low-diversity marine taxa suggest cryptic marine events. Bennett *et al.* (2017) clearly demonstrated that such intervals were abundant in the Tweed Basin from the presence of 128 beds containing single tier colonisations of *Chondrites* in the 500 m thick succession of the Norham Borehole. They attributed emplacement of the burrowing fauna to storm surges.

The marine taxa seen in the western part of the Midland Valley include *Cardiomorpha* and brachiopods such as *Lingula*, which are not seen in the Tweed Basin and Fife. This suggests that their marine connections were from different directions; their emplacement may represent true marine transgressions, rather than surges. A pathway from Ayrshire to the SW is likely, on the grounds that brachiopods are recorded from the coeval and sedimentologically similar Ballycultra Formation in Northern Ireland (Griffith & Wilson 1982). The Tweed Basin is, by contrast, much closer to the more marine Northumberland–Solway Basin and here the storm surges could have been funnelled inland along drowned river systems. The presence of two marine intervals near the top of the East Dron succession is more difficult to understand, though here again access could have been via the same route and these two events probably represent the most energetic storms seen.

4.3. Marginal marine Northumberland–Solway Basin

A narrow coastal floodplain existed along the southern margin of the Southern Uplands massif (Leeder 1974b). The basalt lavas of the Birrenswark Volcanic Formation were buried beneath fluvial sands of the Whita Sandstone Member at the base of the Ballagan Formation. The sandstone unit comprises up to 500 m of massive and cross-bedded, fine to coarse-grained sandstone in typically fining upwards units up to 6 m thick (Lumsden *et al.* 1967; Leeder 1974b). It is clean, carbonate cemented and without the characteristic micaceous and carbonaceous partings of most other Ballagan sandstones. Leeder (1974b) interpreted these as fluvial channel deposits derived from the Southern Uplands massif. Floodplain siltstones and thin dolostones are interbedded with the Whita sandstone units, but there are very few palaeosols. Surface exposures reveal little evidence of evaporite deposits in this belt, with the exception of the Hoddum Borehole, near Ecclefechan, which proved abundant layers and nodules of gypsum deposited in a perennial brine pan (Millward *et al.* 2018).

To the SW, bordering the Solway Firth, Maguire *et al.* (1996) interpreted early Mississippian environments. Coarsely clastic alluvial fan deposits built up adjacent to the active syn-depositional Solway–Gillnockie faults marking the southern margin of the Southern Uplands massif and were bordered by low-relief coastal plains. Prograding deltas, supplied with sediment along the axis of the basin, along with lime muds and storm-emplaced sheet sands were deposited in a shallow offshore region. Biostratigraphical control on some of the sections in this area is poor and only some of these rocks may be Tournaisian in age.

Sherwin (2018) described mixed marginal marine and vegetated floodplain environments with vertebrate remains from three sites along the northern edge of the basin that further

illustrate the proximity and migration of contemporary coastlines. The succession in Whitrope Burn [NY 507 965], near Langholm comprises carbonate rocks, deposited by high-density turbidity currents into a sheltered lagoon (Richards *et al.* 2018). Some beds contain an abundant and diverse assemblage of shark teeth. At Coldstream (Fig. 1), and in the River Coquet, south of the Cheviot massif, shallow marine carbonate, coarsening-upward bay-fill and near-shore facies assemblages are intercalated with a vegetated floodplain facies assemblage.

Further southward towards the basin depocentre, there is increasing evidence of marine conditions, which are characteristic of the Lyne Formation. There are coarsening upward siltstone to fine sandstone units, interpreted as bay fills (Leeder 1974b). The dolostones are replaced by carbonate rocks, some including stromatolitic bioherms and oncolite beds, whereas others contain an abundant and diverse marine fauna including crinoids, foraminifera and brachiopods (Fig. 10; Leeder 1975a, b; Brand 2018).

In the centre of the Northumberland–Solway Basin, 1153 m of anhydrite-bearing carbonate rocks and mudstone within the Lyne Formation were proved in the Easton No 1 well, north of Carlisle (Fig. 1; Ward 1997). The uppermost part of the evaporite succession is upper Chadian–lower Arundian (Ward 1997), but the age of the lower part is not well-constrained biostratigraphically and may be of Tournaisian age. The succession contains about 120 anhydrite beds ranging from 30 cm to 7.9 m thick with some of the thickest evaporite units in the lower part. The subsurface extent of these evaporites mapped by Ward (1997) may extend eastwards to the Newcastle area. Sulphate deposits cementing Permian breccias along the Ninety-Fathom Fault and from brines encountered in the local collieries (Younger *et al.* 2016) may have been derived from this deep source. Ward (1997) suggested that the evaporites were deposited in salinas, but the thickness and association with carbonate rocks suggests a restricted marine environment.

5. Discussion

The facies change at the base of the Ballagan Formation signals a profound change in the paleoenvironments along this part of the southern margin of Laurussia. The upper Devonian Kinnesswood Formation represents braided or meandering river channels filling several small depocentres in the Midland Valley (Read *et al.* 2002), while in southern Scotland rivers fed sediment from the catchments into an elongate, north-eastwards orientated endorheic basin (Leeder 1976). Throughout, calcrete-bearing palaeosols developed on stable alluvial terraces in a semi-arid seasonal climate (Burgess 1961; Leeder 1976; Wright *et al.* 1993). In late Famennian times (Marshall *et al.* 2019), basalt lava and pyroclastic rocks were erupted from volcanoes during rapid extension along the northern margin of the Northumberland–Solway Basin. The change to a wetland regime was likely to have been brought about by a combination of events related to the juxtaposition of Gondwana and Laurussia causing fragmentation of the continental margin and opening of strike-slip basins through the region, and to climate change (Coward 1993; Falcon-Lang 1999; Domeier & Torsvik 2014). Contemporary faulting played a significant role in the particularly thick accumulation of Tournaisian to Chadian strata in the Northumberland–Solway Basin (Chadwick *et al.* 1995). Subsidence related to the major faults bounding the Southern Uplands is inferred from the presence of alluvial fan deposits, and at Dunbar, for

example, by local thickening of the flood-plain sedimentary rocks toward the fault.

The analysis presented here of the Ballagan Formation extends the detailed picture that has emerged from recent studies of the Norham cores and the coastal section at Burnmouth by Bennett *et al.* (2016, 2017), Kearsy *et al.* (2016) and Millward *et al.* (2018) across the Tweed and Midland Valley basins. The spatially and temporally diverse set of palaeoenvironments defines a tropical, strongly seasonal, coastal wetland with very low-relief landscape, developed in the hinterland of an archipelago of more elevated landmasses. Abundant fresh to brackish water, saline and hypersaline floodplain lakes co-existed on the floodplain. Evaporation of some of the closed saline and hypersaline lakes during the dry season or drier intervals formed sabkhas. Microbial mats were associated with these areas (Millward *et al.* 2018). A persistent, dominantly meandering, fluvial system in the east of the region hosts the greatest diversity of terrestrial habitats with the riparian strip extensively vegetated as indicated by a diverse suite of palaeosols ranging from ephemeral plant colonies to marsh and forest (Retallack & Dilcher 1988; Kearsy *et al.* 2016). Forest habitats were also present at times along the northern margin in the Isle of Bute (Carpenter *et al.* 2015), but the true extent of such habitats is poorly known because of low confidence in the interpretation of palaeosols in the archived borehole data. Other areas were dominated by perennial floodplain lakes. By contrast, rainfall, overbank and marine flooding distributed sediment across the floodplain during the wet season, forming and recharging lakes (Bennett *et al.* 2016, 2017). The large lateral extents reached during the storm surges suggest that the floodplain had very low relief and was only just above sea level.

High rainfall during tropical storms is likely to have formed many of the floodplain lakes. However, storm surges caused marine flooding from the SW and northward into the east of the region, reaching significant parts of the Midland Valley and at times as far north as Fife and Tayside, some 140 km due north of the nearest likely coastline. Are such distances supported from modern examples? Williams (2009) and Goodbred & Hine (1995) documented the transport by storm surges of sediment containing sporadic marine fossils inland from the Louisiana and Florida coasts respectively for distances up to 6 km. However, far greater distances have been recorded occasionally. The storm surge associated with the category 1 hurricane Isaac in Louisiana in 2012 caused the Mississippi river to flow upstream for a distance of 480 river kilometres and at about two-thirds of this distance the river level was still 2.6 m above normal (Berg 2013). Substantial areas were flooded to depths up to 5 m, though the deepest water was recorded adjacent to man-made levees.

Another factor may have contributed to the extensive existence of saline and hypersaline lakes. Evaporation in the coastal lagoons and marshes during the dry season may have initiated the upstream flow of saline water when there is low fresh water runoff. This was demonstrated for example by Barousseau *et al.* (1985) to occur in the Salum and Gambia rivers of West Africa. There the effects of a ‘salinity tide’ and increasing salinity inland are seen up-river for up to 250 km.

Many equatorial coastal zones, for example around the Gulf of Mexico, in Florida and Ghana, are barred with barrier islands protecting lagoons. Breaching of the barriers during tropical storms can increase the flooded area hugely and this could have been a further mechanism of flooding the Ballagan wetlands. However, barrier island deposits have low preservation potential and their recognition in the geological record is

difficult. Sherwin (2018) considered the possibility of such an origin for a succession of planar bedded, wave-rippled sandstones in Coquetdale, though these rocks overlie floodplain strata and were more likely to have represented transgression, establishing more open bay conditions.

Though the Ballagan floodplain was dynamic in that local sedimentary environments changed from time to time, probably instigated by river flooding events, the coastal wetland regime remained remarkably constant for the duration of the Tournaisian, a period of at least 12 million years. The succession is aggradational, in which a delicate balance was maintained between subsidence, compaction and sedimentation rates to keep the floodplain just above sea level. Few hiatuses in the succession are seen and are mostly evident in erosive channels at the base of fluvial sandstone units. Another exception is the floodplain terraces represented in the succession at Eastern Hole near Pease Bay on the NE margin of the Southern Uplands massif where the formation is substantially thinner than in nearby areas (Andrews *et al.* 1991). Preservation of delicate desiccated and cracked surfaces (Bennett *et al.* 2016; Kearsy *et al.* 2016) and abundant palaeosols testify to continued 'passive' burial of the floodplain. Time gaps of centuries to thousands of years are mostly evident in the substantial number of vertisols that are seen in the upper part of the succession in the Norham Borehole and at Burnmouth (Kearsy *et al.* 2016).

Johnson (1982) likened the landscape of northern England during Mississippian times to that of the Gulf Coast plain of SE USA today. However, the picture that has emerged during Tournaisian times is not of a floodplain associated with a major delta system. That scenario was initiated in Arundian times with the onset of the Fell Sandstone fluvial distributive system and the overlying Yoredale facies (Kearsy *et al.* 2018). Modern analogues for the Ballagan Formation appear to be few, despite an extensive search.

The Everglades of southern Florida demonstrate some aspects in common. This area has a seasonal climate and is frequently inundated by tropical storms. Freshwater and brackish lakes and marshes are abundant; palustrine carbonates and peats are accumulating; and desiccation features and pedogenic overprinting are widespread (Platt & Wright 1992). These give way at the coast to marginal marine conditions, and offshore in the Bahamas, gypsum precipitates within the carbonate system (Glunk *et al.* 2011).

The seasonal wetlands of the Pantanal of Brazil in South America have a precipitation rate of about 1100 mm per year, similar to that estimated for the Ballagan Formation by Kearsy *et al.* (2016); during the wet season the wetland is flooded through the major rivers (Costa *et al.* 2015). The landscape is gently undulating with variations in elevation of up to 5 m, but the overall gradient across the region is less than 1%. A major feature of the southern part of the region is the juxtaposition on the floodplain of round to elongate freshwater and saline lakes; the latter are locally concentrated. Typically, the saline lakes have surface areas of less than 0.15 km² and are at a slightly lower elevation than the freshwater lakes covering areas of up to 0.8 km² (Costa *et al.* 2015). The pH of the saline lakes is greater than 9 and, though carbonate minerals are precipitated, the lakes never dry out and gypsum or halite is unknown. Seasonal drainage channels provide connections between the lakes and rivers and the intervening areas are covered in forest or open, mainly herbaceous, cover vegetation.

The vast expanse of the northern Great Plains of western Canada is characterised by undulating topography and hosts many hyposaline to hypersaline and alkaline lakes with surface areas that vary from a few tens of metres across to the immense Lake Manitoba (Last & Slezak 1988). Some of the hypersaline lakes are meromictic; carbonate sediment accumu-

lation is common with gypsum and other evaporitic salts in some. Large rivers traverse the region and there are many regions of endorheic drainage. The climate is cold and semi-arid.

Comparative examples in the geological record are sparse too. The lower Cretaceous Leza Formation of the Cameros Basin in NE Spain contains evidence for a coastal wetland setting, with a low-diversity biota, terrestrial sediments with a range of edaphic features, lacustrine environments from freshwater to hypersaline, and leading into a transition to the marine environment; alluvial fans fed clastic detritus into some lakes (Suarez-Gonzalez *et al.* 2015).

The Late Miocene Solimoes Formation in the Amazon Basin comprises a thick vertebrate-bearing succession of typically red or blue-grey, fine-grained sandstones and mudstones (Latrubesse *et al.* 2010). The sandstones are stacked river-channel deposits of avulsive Andean megafans and floodplain distributaries. The mudstones are lacustrine and floodplain deposits with pedogenic layers, abundant desiccation cracks and some gypsum veins. Latrubesse *et al.* (2010) interpreted the succession as an extensive wetland that lay between the Andes and the Purus Arch, prior to development of the current Amazon River system.

Fluvial and lacustrine environments are intercalated in the Miocene Siwalik Group in northern Pakistan. Laminated siltstones, decimetres to several metres thick, were deposited in lakes estimated to have water depths of no more than 14 m and many less than 10 m; the lakes were from a few hundred metres to 25 km across (Zaleha 1997). Desiccation cracks are common and siltstone beds typically overlie palaeosols. Units of very fine- to medium-grained sandstone are generally 5–10 m thick, though some exceed 100 m and comprise up to ten storeys. The sedimentary characteristics indicate these to be river-channel deposits. Thin sandstone beds intercalated with mudstone are thought to have been deposited from non-channelised and channelised flows during flood episodes (Zaleha 1997).

None of these examples is a particularly good fit for the early Mississippian Ballagan palaeoenvironments. The Northern Great Plains, Pantanal, Siwalik Group and Solimoes Formation are all located in tectonic settings different from that of the Ballagan Formation. A coastal setting with transition to marginal marine is present only in the Everglades and the Leza Formation. Tropical seasonal climates are present in the Siwalik Group, Pantanal and the Everglades. However, none of these successions hosts evaporites in abundance comparable with the Ballagan Formation. Primary dolostones are absent from all, though palustrine carbonates are a characteristic of the Everglades. However, the Great Plains, Siwalik Group and Solimoes Formation are examples of the association of lacustrine and fluvial systems. Furthermore, lakes of various salinities characterise both the Great Plains and Pantanal regions. The Leza Formation has perhaps the best range of comparable palaeoenvironments, though the wetland was small, estimated to have been about 10 × 30 km, was not equatorial and evaporites are a minor component; though alluvial fans fed sediment into lakes, established fluvial systems were absent. These examples serve to illustrate the unusual nature of the Ballagan wetlands.

6. Conclusions

The Ballagan Formation is an unusual example in the geological record of a succession with abundant lacustrine deposits of various scales and salinities, interspersed with fluvial and evaporitic facies within a tropical, seasonal, coastal wetland palaeoenvironmental setting. It has few parallels in the literature, particularly with the abundance of small-scale evaporite deposits. An extensive, very low-relief coastal wetland developed

along the equatorial southern continental margin of Laurussia and lasted for the 12 million years of the Tournaisian. The Midland Valley and Tweed basins were separated from the marginal marine seaway of the Northumberland–Solway Basin to the south by an archipelago of upland areas.

The new palaeogeographical map constructed from borehole and outcrop data across the Midland Valley and Tweed Basin shows the juxtaposition of a mosaic of sedimentary environments from fresh and brackish to saline and hypersaline lakes, a diverse suite of floodplain palaeosols and a persistent fluvial system in the east of the region. The strongly seasonal climate led to episodic flooding, both meteoric and marine, the latter driven from the SW into both the western Midland Valley and Northumberland–Solway Basin; marine flooding also reached the Tweed Basin and Tayside in the east via the river floodplain. The sedimentary environments and palaeosol types indicate a broad mosaic of different terrestrial and aquatic habitats that hosted a diverse vertebrate fauna that included tetrapods which had, for the first time, terrestrial capability. The areal size of the region is comparable to that of coastal wetlands today, such as the Everglades of southern Florida.

7. Acknowledgements

This research formed part of the TW:eed Project (Tetrapod World: early evolution and diversification), led by Jenny Clack (University Museum of Zoology, Cambridge) and funded by N.E.R.C. consortium grants to the British Geological Survey (NE/J021067/1) and the Universities of Leicester (NE/J020729/1) and Southampton (NE/J021091/1). The authors are grateful for the helpful comments of two anonymous referees. D.M., P.J.B., T.I.K. and M.A.E.B. publish with the permission of the Executive Director, British Geological Survey (N.E.R.C.).

8. References

- Alonso-Zarza, A. M. 2003. Palaeoenvironmental significance of palustrine carbonates and calcretes in the geological record. *Earth Science Reviews* **60**, 261–98.
- Anderson, J. S., Smithson, T., Mansky, C. F., Meyer, T. & Clack, J. 2015. A diverse tetrapod fauna at the base of ‘Romer’s Gap’. *PLoS ONE* **10**, 1–27.
- Anderton, R. 1985. Sedimentology of the Dinantian of Foulden, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **76**, 7–12.
- Andrews, J. E., Turner, M. S., Nabi, G. & Spiro, B. 1991. The anatomy of an early Dinantian terraced floodplain: palaeo-environment and early diagenesis. *Sedimentology* **38**, 271–87.
- Andrews, J. E. & Nabi, G. 1994. Lithostratigraphy of the Dinantian Inverclyde and Strathclyde Groups, Cockburnspath Outlier, East Lothian–North Berwickshire. *Scottish Journal of Geology* **30**, 105–19.
- Andrews, J. E. & Nabi, G. 1998. Palaeoclimatic significance of calcretes in the Dinantian of the Cockburnspath Outlier (East Lothian–North Berwickshire). *Scottish Journal of Geology* **34**, 153–64.
- Andrews, S. M. 1985. Rhizodont crossopterygian fish from the Dinantian of Foulden, Berwickshire, Scotland, with a re-evaluation of this group. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **76**, 67–95.
- Arsenikos, S., Quinn, M. F., Kimbell, G., Williamson, P., Pharaoh, T., Leslie, G. & Monaghan, A. A. 2018. Structural development of the Devonian–Carboniferous plays of the UK North Sea. In Monaghan, A. A., Underhill, J. R., Hewett, A. J. & Marshall, J. E. A. (eds) *Paleozoic Plays of NW Europe*. Special Publication. London: Geological Society of London.
- Barousseau, J. P., Diop, E. H. S. & Saos, J. L. 1985. Evidence of dynamics reversal in tropical estuaries, geomorphological and sedimentological consequences (Salum and Casamance Rivers, Senegal). *Sedimentology* **32**, 543–52.
- Bateman, R. M. & Scott, A. C. 1990. A reappraisal of the Dinantian floras at Oxroad Bay, East Lothian, Scotland. 2. Volcanicity, palaeoenvironments and palaeoecology. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **81**, 161–94.
- Belt, E. S., Freshney, E. C. & Read, W. A. 1967. Sedimentology of Carboniferous cementstone facies, British Isles and Eastern Canada. *The Journal of Geology* **75**, 711–21.
- Bennett, C. E., Williams, M., Leng, M. J., Siveter, D. J., Davies, S. J., Sloane, H. J. & Wilkinson, I. P. 2011. Diagenesis of fossil ostracods: implications for stable isotope based palaeoenvironmental reconstruction. *Palaeogeography, Palaeoclimatology, Palaeoecology* **305**, 150–61.
- Bennett, C. E., Kearsley, T. I., Davies, S. J., Millward, D., Clack, J. A., Smithson, T. R. & Marshall, J. E. A. 2016. Early Mississippian sandy siltstones preserve rare vertebrate fossils in seasonal flooding episodes. *Sedimentology* **63**, 1677–700.
- Bennett, C. E., Howard, A. S., Davies, S. J., Kearsley, T. I., Millward, D., Brand, P. J., Browne, M. A. E., Reeves, E. J. & Marshall, J. E. A. 2017. Ichnofauna record cryptic marine incursions onto a coastal floodplain at a key early Mississippian tetrapod site. *Palaeogeography, Palaeoclimatology, Palaeoecology* **468**, 287–300.
- Berg, R. 2013. *Tropical cyclone report: Hurricane Isaac (AL092012)*. Miami, FL: National Hurricane Center, National Oceanic and Atmospheric Administration. https://www.nhc.noaa.gov/data/tcr/AL092012_Isaac.pdf
- Brand, P. J. 2018. A list of fossil specimens in the BGS biostratigraphy collections from the Ballagan Formation in Scotland and from the former Lower Border Group of the Northumberland–Solway Basin. British Geological Survey Internal Report IR/18/03.
- Browne, M. A. E. 1980. The Upper Devonian and Lower Carboniferous (Dinantian) of the Firth of Tay, Scotland. Institute of Geological Sciences Report series 80/9.
- Browne, M. A. E., Dean, M. T., Hall, I. H. S., McAdam, A. D., Monro, S. K. & Chisholm, J. I. 1999. A lithostratigraphical framework for the Carboniferous rocks of the Midland Valley of Scotland. British Geological Survey, Research Report RR/99/07.
- Burgess, I. C. 1961. The fossil soils of the Upper Old Red Sandstone of South Ayrshire. *Transactions of the Geological Society of Glasgow* **24**, 138–53.
- Burgess, I. C. & Holliday, D. W. 1979. *Geology of the country around Brough-Under-Stainmore*. Memoir of the Geological Survey of Great Britain, sheets 31 and parts of sheets 25 and 30, England and Wales. London: HMSO.
- Carpenter, D. K., Falcon-Lang, H. J., Benton, M. J. & Henderson, E. 2015. Carboniferous (Tournaisian) fish assemblages from the Isle of Bute, Scotland: systematics and palaeoecology. *Palaeontology* **57**, 1215–40.
- Carruthers, R. G., Burnett, G. A. & Anderson, W. 1932. *The geology of the Cheviot Hills*. Memoir of the Geological Survey England & Wales, explanation of sheets 3 & 5. London: HMSO.
- Cater, J. M. L., Briggs, D. E. G. & Clarkson, E. N. K. 1989. Shrimp-bearing sedimentary successions in the Lower Carboniferous (Dinantian) Cementstone and Oil Shale groups of northern Britain. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **80**, 5–15.
- Chadwick, R. A., Holliday, D. W., Holloway, S. & Hulbert, A. G. 1995. *The structure and evolution of the Northumberland–Solway Basin and adjacent areas*. Subsurface memoir of the British Geological Survey. London: HMSO.
- Chisholm, J. I., McAdam, A. D. & Brand, P. J. 1989. Lithostratigraphical classification of Upper Devonian and Lower Carboniferous rocks in the Lothians. British Geological Survey, Technical Report WA/89/26.
- Clack, J. A. 2002. An early tetrapod from ‘Romer’s Gap’. *Nature* **418**, 72–76.
- Clack, J. A., Bennett, C. E., Carpenter, D. K., Davies, S. J., Fraser, N. C., Kearsley, T. I., Marshall, J. E. A., Millward, D., Otoo, B. K. A., Reeves, E. J., Ross, A. J., Ruta, M., Smithson, K. Z., Smithson, T. R. & Walsh, S. A. 2016. Phylogenetic and environmental context of a Tournaisian tetrapod fauna. *Nature Ecology and Evolution* **1**, 11.
- Clack, J. A. & Finney, S. M. 2005. *Pederpes finneyae*, an articulated tetrapod from the Tournaisian of western Scotland. *Journal of Systematic Palaeontology* **2**, 311–46.
- Clarkson, E. N. K. 1985. Palaeoecology of the Dinantian of Foulden, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **76**, 97–100.
- Clayton, G. 1986. Late Tournaisian miospores from the Ballycultra Formation at Cultra, County Down, Northern Ireland. *Irish Journal of Earth Sciences* **8**, 73–79.
- Clayton, G., Graham, J. R., Higgs, K., Sevastopulo, G. D. & Welsh, A. 1986. Late Devonian and early Carboniferous palaeogeography of southern Ireland and southwest Britain. *Annales de la Société géologique de Belgique* **109**, 103–11.

- Clayton, G. & Higgs, K. 1979. The Tournaisian marine transgression in Ireland. *Journal of Earth Sciences, Royal Dublin Society* **2**, 1–10.
- Cliff, R. A., Drewery, S. E. & Leeder, M. R. 1991. Sourcelands for the Carboniferous Pennine river system: constraints from sedimentary evidence and U-Pb geochronology using zircon and monazite. In Morton, A. C., Todd, S. P. & Haughton, P. D. W. (eds) *Developments in sedimentary provenance studies*, **57**, 137–59. Special Publication. London: Geological Society of London.
- Clough, C. T., Barrow, G., Crampton, C. B., Maufe, H. B., Bailey, E. B. & Anderson, E. M. 1910. *The geology of East Lothian, including the counties of Edinburgh and Berwick*. Memoir of the Geological Survey of Scotland, Explanation of sheet 33 with parts of 34 and 41. 2nd edn. Edinburgh: HMSO.
- Cope, J. C. W., Guion, P. D., Sevastopulo, G. D. & Swan, A. R. H. 1992. Carboniferous. In Cope, J. C. W., Ingham, J. K. & Rawson, P. F. (eds) *Atlas of palaeogeography and lithofacies*, **13**, 67–86. London: Geological Society.
- Costa, M., Telmer, K. H., Evans, T. L., Almeida, T. I. R. & Diakun, M. T. 2015. The lakes of the Pantanal: inventory, distribution, geochemistry, and surrounding landscape. *Wetlands Ecology and Management* **23**, 19–39.
- Coward, M. P. 1993. The effect of Late Caledonian and Variscan continental escape tectonics on basement structure, Paleozoic basin kinematics and subsequent Mesozoic basin development in NW Europe. *Geological Society of London Petroleum Geology Conference* **4**, 1095–108.
- Davies, A. 1970. Carboniferous rocks of the Sanquhar outlier. *Geological Survey of Great Britain Bulletin* **31**, 37–87.
- Davies, A., McAdam, A. D. & Cameron, I. B. 1986. *Geology of the Dumbar District*. Memoir of the British Geological Survey, Sheet 33E and part of 41 (Scotland). London: HMSO.
- Day, J. B. W. 1970. *Geology of the country around Bewcastle*. Memoir Geological Survey, Great Britain, England and Wales Sheet 12. London: HMSO.
- Domeier, M. & Torsvik, T. H. 2014. Plate tectonics in the late Paleozoic. *Geoscience Frontiers* **5**, 303–50.
- Ettensohn, F. R., Dever G. R. Jr. & Grow, J. S. 1988. A paleosol interpretation for profiles exhibiting subaerial exposure 'crusts' from the Mississippian of the Appalachian Basin. In Reinhart, J. & Sigleo, W. R. (eds) *Paleosols and weathering through geologic time*, **216**, 49–79. Special Paper. Boulder, CO: Geological Society of America.
- Eyles, V. A., Simpson, J. B. & MacGregor A. G. 1949. *Geology of Central Ayrshire*. Memoir Geological Survey, Scotland (sheet 14). 2nd edn. Edinburgh: HMSO.
- Falcon-Lang, H. J. 1999. The early Carboniferous (Courceyan–Arundian) monsoonal climate of the British Isles: evidence from growth rings in fossil woods. *Geological Magazine* **136**, 177–87.
- Francis, E. H., Forsyth, I. H., Read, W. A. & Armstrong, M. 1970. *The Geology of the Stirling District*. Memoir Geological Survey, Scotland (sheet 39). Edinburgh: HMSO.
- Gardiner, B. G. 1985. Actinopterygian fish from the Dinantian of Foulden, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **76**, 61–66.
- Glunk, C., Dupraz, C., Braissant, O., Gallagher, K. L., Verrecchia, E. P. & Visscher, P. T. 2011. Microbially mediated carbonate precipitation in a hypersaline lake, Big Pond (Eleuthera, Bahamas). *Sedimentology* **58**, 720–38.
- Goodbred, S. L. & Hine, A. C. 1995. Coastal storm deposition: salt-marsh response to a severe extratropical storm, March 1993, west-Central Florida. *Geology* **23**, 679–82.
- Greig, D. C. 1988. *Geology of the Eyemouth District*. Memoir British Geological Survey, Scotland sheet 34. London: HMSO.
- Griffith, A. E. & Wilson, H. E. (1982) *Geology of the country around Carrickfergus and Bangor*. Memoir Geological Survey Northern Ireland (sheet 29). Belfast: HMSO.
- Gueriau, P., Rabet, N., Clement, G., Lagebro, L., Vannier, J., Briggs, D. E. G., Charbonnier, S., Olive, S. & Berthou, O. 2016. A 365-million-year-old freshwater community reveals morphological and ecological stasis in Branchiopod crustaceans. *Current Biology* **26**, 383–90.
- Johnson, G. A. L. 1982. Geographical change in Britain during the Carboniferous period. *Proceedings of the Yorkshire Geological Society* **44**, 181–203.
- Kaiser, S. I., Aretz, M. & Becker, R. T. 2015. The global Hangenberg Crisis (Devonian–Carboniferous transition): review of a first-order mass extinction. In Becker, R. T., Königshof, P. & Brett, C. E. (eds) *Devonian climate, sea level and evolutionary events*, **423**, 423–29. Special Publication. London: Geological Society of London.
- Kearsey, T. I., Bennett, C. E., Millward, D., Davies, S. J., Gowing, C. J. B., Kemp, S. J., Leng, M. J., Marshall, J. E. A. & Browne, M. A. E. 2016. The terrestrial landscapes of tetrapod evolution in earliest Carboniferous seasonal wetlands of S.E. Scotland. *Palaeogeography, Palaeoclimatology, Palaeoecology* **457**, 52–69.
- Kearsey, T., Millward, D., Ellen, R., Whitbread, K. & Monaghan, A. A. 2018. Revised stratigraphic framework of pre-Westphalian Carboniferous petroleum system elements from the Outer Moray Firth to the Silverpit Basin, North Sea, UK. In Monaghan, A. A., Underhill, J. R., Hewett, A. J. & Marshall, J. E. A. (eds) *Paleozoic plays of NW Europe*. Special Publication. London: Geological Society of London.
- Last, W. M. & Slezak, L. A. 1988. The salt lakes of western Canada: a paleolimnological overview. *Hydrobiologia* **158**, 301–16.
- Latrubesse, E. M., Cozzuol, M., da Silva-Caminha, S. A. F., Rigsby, C. A., Absy, M. L. & Jaramillo, C. 2010. The Late Miocene paleogeography of the Amazon Basin and the evolution of the Amazon River system. *Earth-Science Reviews* **99**, 99–124.
- Leeder, M. R. 1974a. Origin of the Northumberland Basin. *Scottish Journal of Geology* **10**, 283–96.
- Leeder M. R. 1974b. Lower Border Group (Tournaisian) fluvio-deltaic sedimentation and palaeogeography of the Northumberland Basin. *Proceedings of the Yorkshire Geological Society* **40**, 129–80.
- Leeder, M. R. 1975a. Lower Border Group (Tournaisian) limestones from the Northumberland Basin. *Scottish Journal of Geology* **11**, 151–67.
- Leeder, M. R. 1975b. Lower Border Group (Tournaisian) stromatolites from the Northumberland Basin. *Scottish Journal of Geology* **11**, 207–26.
- Leeder, M. R. 1976. Palaeogeographic significance of pedogenic carbonates in the topmost Upper Old Red Sandstone of the Scottish Border Basin. *Journal of Geology* **11**, 21–28.
- Leeder, M. R. & Bridges P. H. 1978. Upper Old Red Sandstone near Kirkbean, Dumfries and Galloway. *Scottish Journal of Geology* **14**, 267–72.
- Lumsden, G. I., Tulloch, W., Howells, M. F. & Davies, A. 1967. *The geology of the neighbourhood of Langholm*. Memoir of the Geological Survey, Scotland (Sheet 11). Edinburgh: HMSO.
- Maguire, K., Thompson, J. & Gowland, S. 1996. Dinantian depositional environments along the northern margin of the Solway Basin, UK. In Strogon, P., Somerville, I. D. & Jones, G. L. (eds) *Recent advances in Lower Carboniferous geology*, **107**, 163–82. Special Publication. London: Geological Society of London.
- Marshall, J. E. A., Haughton, P. D. W. & Hillier, S. J. 1994. Vitrimite reflectivity and the structure and burial history of the Old Red Sandstone of the Midland Valley of Scotland. *Journal of the Geological Society, London* **151**, 425–38.
- Marshall, J. E. A., Reeves, E. J., Bennett, C. E., Davies, S. J., Kearsey, T. I., Millward, D., Smithson, T. R. & Browne, M. A. E. 2019. Reinterpreting the age of the uppermost 'Old Red Sandstone' and Early Carboniferous in Scotland. *Earth and Environmental Transactions of the Royal Society of Edinburgh*, **109**.
- McAdam, A. D. & Tulloch, W. 1985. *Geology of the Haddington District*. Memoir Geological Survey of Scotland, sheet 33W and part of Sheet 41. London: HMSO.
- McMillan, A. A. 2002. *Geology of the New Galloway and Thornhill District*. Memoir of the British Geological Survey, Sheets 9W and 9E (Scotland). Norwich, UK: The Stationary Office.
- Miller, H. 1887. *The Geology of the Country Around Otterburn and Elsdon*. Memoir Geological Survey, England and Wales, sheet 8. London: HMSO.
- Millward, D., Kearsey, T. I. & Browne, M. A. E. 2013. Norham West Mains Farm Borehole: operations report. British Geological Survey Internal Report IR/13/033.
- Millward, D., Davies, S. J., Williamson, F., Curtis, R., Kearsey, T. I., Bennett, C. E., Marshall, J. E. A. & Browne, M. A. E. 2018. Early Mississippian evaporites of coastal tropical wetlands. *Sedimentology*. <https://doi.org/10.1111/sed.12465>.
- Millward, D. & Stephenson, D. 2011. Bedrock GSI3D models from interpreted data in geologically complex Carboniferous terrains: a work in progress from the Clyde catchment area, Midland Valley of Scotland. British Geological Survey Internal Report, IR/11/052.
- Mitchell, G. H. & Mykura, W. 1962. *The geology of the neighbourhood of Edinburgh*. Memoir of the Geological Survey of Great Britain, Sheet 32, 3rd edn. Scotland.
- Monaghan, A. 2004. *Geology of the Daily area, Ayrshire*. British Geological Survey Internal Report, IR/04/035.
- Moy-Thomas, J. A. 1938. Carboniferous Palaeoniscids from Northumberland and Berwickshire. *Geological Magazine* **75**, 308–18.
- Nairn, A. E. M. 1958. Petrology of the Whita Sandstone, Southern Scotland. *Journal of Sedimentary Petrology* **28**, 57–64.
- Neves, R., Gueinn, K. J., Clayton, G., Ioannides, N. S., Neville, R. S. W. & Kruszewska, K. 1973. Palynological correlations within the

- Lower Carboniferous of Scotland and Northern England. *Transactions of the Royal Society of Edinburgh* **69**, 23–70.
- Neves, R. & Ioannides, N. 1974. Palynology of the Lower Carboniferous (Dinantian) of the Spilmersford Borehole, East Lothian, Scotland. *Geological Survey of Great Britain Bulletin* **45**, 73–98.
- Platt, N. H. & Wright, V. P. 1992. Palustrine carbonates and the Florida Everglades: towards an exposure index for the freshwater environment. *Journal of Sedimentary Petrology* **62**, 1058–71.
- Purnell M. A. & Cossey P. J. 2004. Northumberland trough. Chapter 3. In Cossey, P. J., Adams, A. E., Purnell, M. A., Whiteley, M. J., Whyte, M. A. & Wright, V. P. (eds) *British Lower Carboniferous stratigraphy*, 109–66. Geological Conservation Review Series 29. Peterborough: Joint Nature Conservation Committee.
- Raymond P. E., 1946. The Genera of fossil Conchostraca, an order of bivalve Crustacea. *Harvard University of Comparative Zoology Bulletin* **96**, 218–307.
- Read, W. A., Browne, M. A. E., Stephenson, D. & Upton, B. G. J. 2002. Carboniferous. In Trewin, N. H. (ed.) *The geology of Scotland*. 4th edn. 251–99. London: The Geological Society.
- Read, W. A. & Johnson, S. R. H. 1967. The sedimentology of sandstone formations within the Upper Old Red Sandstone and lowest Calciferous Sandstone Measures west of Stirling, Scotland. *Scottish Journal of Geology* **3**, 242–67.
- Retallack, G. J. & Dilcher, D. L. 1988. Reconstructions of selected seed ferns. *Annals of the Missouri Botanic Garden* **75**, 1010–57.
- Richards, K. R., Sherwin, J. E., Smithson, T. R., Bennion, R. F., Davies, S. J., Marshall, J. E. A. & Clack, J. A. 2018. Diverse and durophagous: early Carboniferous chondrichthyans from the Scottish Borders. *Earth and Environmental Transactions of the Royal Society of Edinburgh* **108**, 67–87.
- Scott, A. C. & Meyer-Berthaud, B. 1985. Plants from the Dinantian of Foulden, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **76**, 13–20.
- Scott, W. B. 1986. Nodular carbonates in the Lower Carboniferous, Cementstone Group of the Tweed Embayment, Berwickshire: evidence for a former sulphate evaporite facies. *Scottish Journal of Geology* **22**, 325–45.
- Scrutton, C. & Turner, B. 1995. The geology of Eyemouth and Burnmouth. Excursion 2. In Scrutton, C. (ed.) *Northumbrian rocks and landscape: a field guide*, 31–41. Maryport: Yorkshire Geological Society, Ellenbank Press.
- Sherwin, J. E. 2018. In an alternating marine and non-marine depositional setting, where and how are early Carboniferous tetrapods preserved? M. Phil Thesis, University of Leicester, UK.
- Smith, R. A. 1986. Permo-Triassic and Dinantian rocks of the Belfast harbour Borehole. British Geological Survey Report 18, No 6.
- Smithson, T. R., Wood, S. P., Marshall, J. E. A. & Clack, J. A. 2012. Earliest Carboniferous tetrapod and arthropod faunas from Scotland populate Romer's Gap. *Proceedings of the National Academy of Science of America* **109**, 4532–37.
- Smithson, T. R., Richards, K. R. & Clack, J. A. 2016. Lungfish diversity in Romer's Gap: reaction to the end-Devonian mass extinction. *Palaeontology* **59**, 29–44.
- Smithson, T. R. & Clack, J. A. 2018. A new tetrapod from Romer's Gap reveals an adaptation for walking. *Earth and Environmental Transactions of the Royal Society of Edinburgh* **108**, 89–97.
- Stephenson, D., Bevins, R. E., Millward, D., Highton, A. J., Parsons, I., Stone, P. & Wadsworth, W. J. 1999. *Caledonian igneous rocks of Great Britain*. Geological Conservation Review, **17**. Peterborough, UK: Joint Nature Conservation Committee.
- Stephenson, M. H., Williams, M., Monaghan, A. A., Arkley, S. & Smith, R. A. 2002. Biostratigraphy and palaeoenvironments of the Ballagan Formation (Lower Carboniferous) in Ayrshire. *Scottish Journal of Geology* **38**, 93–111.
- Stephenson, M. H., Williams, M., Leng, M. J. & Monaghan, A. A. 2004a. Aquatic plant microfossils of probable non-vascular origin from the Ballagan Formation (Lower Carboniferous), Midland Valley, Scotland. *Proceedings of the Yorkshire Geological Society* **55**, 145–58.
- Stephenson, M. H., Williams, M., Monaghan, A. A., Arkley, S., Smith, R. A., Dean, M., Browne, M. A. E. & Leng, M. 2004b. Palynomorph and ostracod biostratigraphy of the Ballagan Formation, Midland Valley of Scotland, and elucidation of intra-Dinantian unconformities. *Proceedings of the Yorkshire Geological Society* **55**, 131–43.
- Stone, P., Millward, D., Young, B., Merritt, J. W., Clarke, S. M., McCormac, M. & Lawrence, D. J. 2010. *British regional geology: Northern England*, 5th edn. Keyworth, Nottingham: British Geological Survey.
- Suarez-Gonzalez, P., Quijada, I. E., Benito, M. I. & Mas, R. 2015. Sedimentology of ancient coastal wetlands: insights from a Cretaceous multifaceted depositional system. *Journal of Sedimentary Research* **85**, 95–117.
- Tasch, P. 1969. Branchiopoda. In Moore, R. C. (ed.) *Treatise on invertebrate palaeontology, Conchostraca*. Part R Arthropoda Vol. 4(1), R128–91. Lawrence, Kansas: University of Kansas and Geological Society of America.
- Townsend, S. A., Boland, K. T. & Wrigley, T. J. 1992. Factors contributing to a fish kill in the Australian wet/dry tropics. *Water Research* **26**, 1039–44.
- Ward, J. 1997. Early Dinantian evaporites of the Easton-1 well, Solway Basin, onshore, Cumbria, England. In Meadows, N. S., Trueblood, S. P., Hardman, M. & Cowan, G. (eds) *Petroleum geology of the Irish Sea and adjacent areas*, **124**, 277–96. Special Publication. London: Geological Society of London.
- Waters, C. N., Somerville, I. D., Jones, N. S., Cleal, C. J., Collinson, J. D., Waters, R. A., Besly, B. M., Dean, M. T., Stephenson, M. H., Davies, J. R., Freshney, E. C., Jackson, D. I., Mitchell, W. I., Powell, J. H., Barclay, W. J., Browne, M. A. E., Leveridge, B. E., Long, S. L. & McLean, D. (eds) 2011. *A revised correlation of Carboniferous rocks in the British Isles*. Special Report 26. London: Geological Society of London.
- Williams, H. F. L. 2009. Stratigraphy, sedimentology, and microfossil content of hurricane Rita storm surge deposits in Southwest Louisiana. *Journal of Coastal Research* **25**, 1041–51.
- Williams, M., Stephenson, M. H., Wilkinson, I. P., Leng, M. J. & Miller, C. G. 2005. Early Carboniferous (Late Tournaisian-Early Viséan) ostracods from the Ballagan Formation, central Scotland, UK. *Journal of Micropalaeontology* **24**, 77–94.
- Williams, M., Leng, M. J., Stephenson, M. H., Andrews, J. E., Wilkinson, I. P., Siveter, D. J., Horne, D. J. & Vannier, J. M. C. 2006. Evidence that Early Carboniferous ostracods colonised coastal flood plain brackish water environments. *Palaeogeography, Palaeoclimatology, Palaeoecology* **230**, 299–318.
- Wills, L. J. 1951. *A palaeogeographical atlas of the British Isles and adjacent parts of Europe*. London: Blackie and Son Ltd.
- Wood, S. P. & Rolfe, W. D. I. 1985. Introduction to the palaeontology of the Dinantian of Foulden, Berwickshire, Scotland. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **76**, 1–6.
- Wright, V. P., Turner, M. S., Andrews, J. E. & Spiro, B. 1993. Morphology and significance of super-mature calcretes from the Upper Old Red Sandstone of Scotland. *Journal of the Geological Society of London* **150**, 871–83.
- Young, G. M. & Caldwell, W. G. E. 2011. Early Carboniferous stratigraphy in the Firth of Clyde area: new information from the Isle of Bute. *Scottish Journal of Geology* **47**, 143–56.
- Young, G. M. & Caldwell, W. G. E. 2012. The Northeast Arran Trough, the Corrie conundrum and the Highland Boundary Fault in the Firth of Clyde, SW Scotland. *Geological Magazine* **149**, 578–89.
- Younger, P. L., Manning, D. A. C., Millward, D., Busby, J. P., Jones, C. R. C. & Gluyas, J. G. 2016. Geothermal exploration in the Fell Sandstone Formation (Mississippian) beneath the city centre of Newcastle upon Tyne, UK: the Newcastle Science Central Deep Geothermal Borehole. *Quarterly Journal Engineering Geology and Hydrogeology* **49**, 350–63.
- Zaleha, M. J. 1997. Fluvial and lacustrine palaeoenvironments of the Miocene Siwalik Group, Khaur are, northern Pakistan. *Sedimentology* **44**, 349–68.