

# Stressful life events and leukocyte telomere attrition in adulthood: a prospective population-based cohort study

S. L. van Ockenburg<sup>1</sup>, E. H. Bos<sup>1</sup>, P. de Jonge<sup>1</sup>, P. van der Harst<sup>2</sup>, R. O. B. Gans<sup>3</sup> and J. G. M. Rosmalen<sup>1\*</sup>

<sup>1</sup>Interdisciplinary Center Psychopathology and Emotion regulation, University of Groningen, University Medical Center Groningen, The Netherlands

<sup>2</sup>Department of Cardiology, University of Groningen, University Medical Center Groningen, The Netherlands

<sup>3</sup>Department of Internal Medicine, University of Groningen, University Medical Center Groningen, The Netherlands

**Background.** Telomere attrition might be one of the mechanisms through which psychosocial stress leads to somatic disease. To date it is unknown if exposure to adverse life events in adulthood is associated with telomere shortening prospectively. In the current study we investigated whether life events are associated with shortening of telomere length (TL).

**Method.** Participants were 1094 adults (mean age 53.1, range 33–79 years) from the PREVENT cohort. Data were collected at baseline (T1) and at two follow-up visits after 4 years (T2) and 6 years (T3). Life events were assessed with an adjusted version of the List of Threatening Events (LTE). TL was measured by monochrome multiplex quantitative PCR at T1, T2, and T3. A linear mixed model was used to assess the effect of recent life events on TL prospectively. Multivariable regression analyses were performed to assess whether the lifetime life events score or the score of life events experienced before the age of 12 predicted TL cross-sectionally. All final models were adjusted for age, sex, body mass index, presence of chronic diseases, frequency of sports, smoking status, and level of education.

**Results.** Recent life events significantly predicted telomere attrition prospectively ( $B = -0.031$ ,  $p = 0.007$ ). We were not able to demonstrate a significant cross-sectional relationship between the lifetime LTE score and TL. Nor did we find exposure to adverse life events before the age of 12 to be associated with TL in adulthood.

**Conclusions.** Exposure to recent adverse life events in adulthood is associated with telomere attrition prospectively.

Received 16 December 2014; Revised 16 April 2015; Accepted 23 April 2015; First published online 29 July 2015

**Key words:** Adversity, life events, observational cohort study, prospective, stress, telomere.

## Introduction

Psychosocial stress is a well-known risk factor for various somatic disorders, including cardiovascular disease (CVD) (Ohlin *et al.* 2004; Rosengren *et al.* 2004). In attempts to unravel the relationship between psychosocial stress and adverse health, much attention has been given to the functioning of three important stress-responsive systems; the hypothalamic-pituitary-adrenal axis (Miller *et al.* 2007; Chida & Hamer, 2008), the autonomic nervous system (Chida & Hamer, 2008; Chida & Steptoe, 2010), and the immune system (Segerstrom & Miller, 2004; Steptoe *et al.* 2007; Miller *et al.* 2009).

Stress-responsive system activation might lead to telomere shortening (Epel *et al.* 2006). Telomere length (TL) might thus reflect cumulative damage due to exposure to psychosocial stress particularly well. Indeed, since Epel and colleagues reported for the first time that psychosocial stress is associated with shorter TL in 2004 (Epel *et al.* 2004), many studies have focused on telomere shortening as a potential downstream consequence of exposure to psychosocial stress.

Telomeres are TTAGGG nucleotide tandem repeats at the ends of chromosomes in eukaryotic cells. As DNA polymerases cannot copy the end of the DNA strand, telomeres progressively shorten with each cell division. When telomeres become critically short this causes chromosomal instability (Chan & Blackburn, 2002) and cellular senescence (Flores *et al.* 2005). Consequently, TL is considered a biological marker of ageing (Harley *et al.* 1990). Although a special enzyme telomerase exists which is capable of lengthening

\* Address for correspondence: Professor J. G. M. Rosmalen, Interdisciplinary Center Psychopathology and Emotion regulation, University of Groningen, University Medical Center Groningen, P.O. Box 30.001, 9700 RB, CC72, Groningen, The Netherlands (Email: j.g.m.rosmalen@umcg.nl)

telomeres, in most somatic cells only limited amounts of this enzyme are present (Chan & Blackburn, 2002). Telomere shortening is predictive of increased mortality rate (Cawthon *et al.* 2003; Honig *et al.* 2006) and various age-related diseases, such as cancer (Willeit *et al.* 2010) and Alzheimer's disease (Honig *et al.* 2006).

According to the recent literature review by Price *et al.* (2012) most studies investigating the relationship between psychosocial stress and TL found a significant negative association, although some studies also failed to demonstrate any relationship. Furthermore, the authors point out that except for one study (Shalev *et al.* 2012) all studies investigating the relationship between psychosocial stress and TL have been cross-sectional (Price *et al.* 2012). This is problematic because of the high variability in TL between individuals which is present at birth, and gender differences in rate of change in TL, limiting the power of studies utilizing only one time point (Price *et al.* 2012; Takubo *et al.* 2002). Although longitudinal studies usually have not measured TL at birth either, they do have the possibility to statistically adjust for 'baseline TL', a composite measure of TL at birth plus environmental effects over life. Moreover, when studying processes such as the effects of psychosocial stress on biological systems the real interest lies in *change* over time.

The only two longitudinal studies on the effects of psychosocial stress on TL until now have been performed in children (Shalev *et al.* 2012), and in post-menopausal, non-smoking, disease-free women (Puterman *et al.* 2015). We know from life-course epidemiology that for certain change effects 'critical periods' or 'sensitive periods' exist for some biological processes (Ben-Shlomo & Kuh, 2002). Thus, it is currently unknown if psychosocial stress in adulthood has the same damaging effect on telomere attrition as it has in childhood. Although the results of Puterman's recent study in post-menopausal women certainly are pointing in this direction, as they showed that major life stressors across a 1-year period predicted telomere attrition in that same year (Puterman *et al.* 2015). Their study was, however, conducted in a highly selected group and is therefore not generalizable to the general population. Furthermore, it might be that repair mechanisms, such as telomere elongation by the enzyme telomerase (Harley *et al.* 1990), prevent short-term stressors from leaving a permanent mark on TL in the long run. In this case it might be that cumulative exposure to life events over life is needed to create an adverse outcome (Ben-Shlomo & Kuh, 2002). In the current study we wished to investigate the multiple possibilities mentioned above and have formulated four hypotheses. First, we postulate that recent adverse life events in adulthood are associated with telomere attrition over

time prospectively. Second, we hypothesize that the rate of change in TL associated with recent adverse life events is moderated by cumulative exposure to adverse life events over life. Third, we hypothesize that life events experienced in childhood have long-lasting negative effects on TL that can still be detected in adulthood. Finally, we hypothesize that the cumulative exposure to stressful life events over life is associated with shorter telomeres cross-sectionally.

The statistical models were adjusted for the following covariates because of their known association with life events or TL: gender (Barrett & Richardson, 2011), age (Chen *et al.* 2011), body mass index (BMI) (Valdes *et al.* 2005), the presence of chronic diseases, smoking (Valdes *et al.* 2005), frequency of sports (Du *et al.* 2012), and level of education (Stephoe *et al.* 2011). It has previously been reported in the cohort of the current study that the presence of a generalized anxiety disorder, but not depression, is associated with TL (Hoen *et al.* 2013). To exclude the possibility that anxiety was mediating the association between stressful life events and telomere attrition we performed *post-hoc* analyses where we additionally adjusted for the presence of generalized anxiety disorder.

## Method

### Study population

Our study has been performed in a cohort derived from Prevention of Renal and Vascular ENd stage Disease (PREVEND), a population cohort study originally designed to investigate microalbuminuria as a risk factor for renal disease and CVD. The recruitment of participants for PREVEND has been extensively described elsewhere (Pinto-Sietsma *et al.* 2000). The PREVEND baseline sample consisted of 8592 subjects randomly selected from the population of the city of Groningen, The Netherlands, with oversampling for albuminuria (T1). Selection of subjects for the present study was aimed at recruiting a representative sample from the general population of Groningen, while simultaneously rectifying PREVEND's oversampling for albuminuria. Research assistants approached participants in the PREVEND study during their visit to the outpatient clinic during follow-up (T2) (2554 participants). Questionnaires were completed by a total of 1094 participants (43%), forming the population cohort of the present study. There was no significant difference in gender, age, or scores on a 12-item neuroticism scale between PREVEND participants who were invited to participate in the present study but declined and PREVEND participants who agreed to participate. The sample consisted of 588 (53.7%) females and 506 (46.3%) males. Their mean age was 53.1 (s.d. = 11.4,




Fig. 1. Flow chart of the recruitment and selection process of PREVEND.

range 33–79) years and their ethnicity was predominantly Caucasian. Follow-up measurements took place between January 2002 and November 2003 (T2) and approximately 2 years later, between April 2004 and November 2006 (T3). Average time between T1 and T2 was 4.1 years; average time between T2 and T3 was 2.4 years. Follow-up measurements were completed by a

total of 976 (89%) participants at T3. A flow chart of the recruitment process of the PREVEND study and the current sub-study can be found in Fig. 1. The study was approved by the local Medical Ethical Committee for human research of the University Medical Center Groningen. All participants provided written informed consent for participation in this study.

### Adverse life events

Stressful life events were assessed at T2 and T3 by means of the Dutch version of the List of Threatening Events (LTE), a 12-item self-report questionnaire (Brugha & Cragg, 1990). The LTE comprises 12 major categories of stressful life events that were selected for their established long-term consequences. It assesses the occurrence of events such as the death of a loved one, losing one's job, or being hospitalized for a physical disease. The original LTE covers the last 6 months, but we used an adjusted version of the LTE with response categories for the previous year and for five age groups (0–12, 13–18, 19–39, 40–60, >60 years) (Rosmalen *et al.* 2012). Respondents were instructed not to include the last year when scoring the age groups since this period was covered by a separate response category 'previous year'.

We used these responses to define three exposure variables: 'childhood LTE score' (number of different adverse life events before age 12), 'recent LTE score' (number of different adverse life events in previous year), and 'lifetime LTE score' (number of different adverse life events in all completed age groups). Underreporting of adverse life events is an established bias of retrospective questionnaires (Hardt & Rutter, 2004). To correct for underreporting of life events in the childhood LTE score and the lifetime LTE score, we scored a life event as 1 (took place) if it was reported at either T2 or T3, and as 0 (did not take place) if it was not reported at any of these two measurement occasions. The current version of the LTE has been extensively validated as described elsewhere (Rosmalen *et al.* 2012). In brief, participants completed the LTE at two occasions; once at T2 and once at T3. The stability of the retrospective reporting of life events was satisfactory. The test–retest correlation for the lifetime LTE score was large: 0.606. The construct validity of the list is indicated by its positive associations with psychological distress, mental health problems, and neuroticism.

### Telomere length

Fasting blood samples were collected from all participants by a nurse during a visit to the research facilities at T1, T2, and T3. In case of influenza or a febrile temperature, blood collection was postponed to a later time. DNA was extracted from leukocytes. In order to prevent batch effects, the samples of the three different time points (T1, T2, T3) were randomly assigned for DNA extraction. The same extraction method was used for all samples with a standard kit according to the manufacturer's instructions (QIamp 96 DNA blood kit, catalogue no. 51162; Qiagen, The Netherlands). TL was measured by a monochrome

multiplex quantitative polymerase chain reaction (PCR) method, whereby the telomere and single copy gene are amplified in the same tube (Cawthon, 2009). Samples were run in triplicate measured in the same well position on different plates. The intra-assay coefficient of variation (CV) was 2% (T), 1.9% (S) and 4.5% (T/S ratio). Reproducibility data was obtained for 216 subjects from PREVENT and good agreement between T/S ratios was observed ( $R^2 = 0.99$ ,  $p < 0.0001$ , inter-run CV 3.9%) (Huzen *et al.* 2011). The samples of the three time points were equally divided over the PCR schedule to reduce potential time or seasonal effects. The calibrator sample used was made up of a mixture of DNAs from young adult individuals (around 25 years). There was a highly significant decline in T/S ratio with age in PREVENT ( $B = -0.0047$ ,  $s.e. = 0.0004$ ,  $p < 0.001$ ) confirming the internal validity of the assay. TL in the present study was available for 1019 people at T1 (93.1%) and for 982 people at T3 (89.8%). At T2 for a large part of the cohort DNA was not available and TL could thus be determined only for a subset of the population ( $n = 445$ , 40.7%). Likelihood-based methods can still provide reliable estimates if there is missingness in the outcome variable (Kenward & Molenberghs, 1998). Thus, we used maximum likelihood estimation in all models involving the T2 data as is explained in more detail below.

### Statistical analyses

#### Covariates

We had information about the presence of a chronic disease in the following categories: coronary heart disease (CHD), cerebrovascular accident (CVA), diabetes mellitus, chronic liver disease, chronic kidney disease, malignancy, rheumatoid arthritis, chronic obstructive pulmonary disease or asthma, severe skin disease, severe bowel disease lasting >3 months. All somatic diseases, except for diabetes, CHD, and CVA were self-reported diseases that were present in the previous year. Diabetes was defined as the use of antidiabetic treatment according to self-report or pharmacy data. CHD and CVA were defined as self-report of CHD/CVA upon inclusion in the study and/or confirmed occurrence of CHD/CVA as registered Dutch national registry of hospital discharge diagnoses between inclusion and date of visit to the research facilities at T2. Smoking was divided into six categories (0, 1–5, 6–10, 11–15, 16–20, >20 cigarettes/day). Frequency of sports was defined as: I don't exercise, I exercise once per week, or I exercise twice or more per week. Educational level was categorized as: none, lower secondary education or less, higher secondary education, or tertiary education. In the statistical models smoking, frequency of sports, and level of education were



entered as dummy variables with the lowest category serving as a reference. The presence of a generalized anxiety disorder in the past 12 months was assessed by means of the Composite International Diagnostic Interview 2.1 (CIDI 2.1) at T2 and T3.

#### *Adverse life events in adulthood and telomere attrition*

Our data had a hierarchical structure, that is, repeated measurements were nested within individuals. Therefore we chose a statistical model that takes into account the non-independence of observations. We used a linear mixed model to test the hypotheses that recent adverse life events in adulthood are associated with telomere attrition over time prospectively, and secondly that this effect is moderated by cumulative exposure to adverse life events. All models contained TL (T2 and T3) as the dependent variable and had the 'recent LTE score' (the year before T2 and the year before T3) as the predictor variable. Moreover, all models were adjusted for sex, age, BMI, presence of chronic diseases, smoking, frequency of sports, level of education, TL at baseline (T1), and time in years between measurement occasions. Our model can thus be viewed as an analysis of covariance, adjusting TL at follow-up for baseline TL (Vickers & Altman, 2001). All predictors were time-varying except sex, TL at baseline, level of education, and number of chronic diseases at T2. A random intercept was used to account for nesting of observations within individuals. In the first model we tested the hypothesis that life events in the previous year were associated with telomere attrition. In the second model, effect modification analysis was used to examine whether associations of recent life events with TL were dependent on the level of lifetime exposure to stressful life events. This model was identical to the model described above, except for the addition of the 'recent LTE score' by 'lifetime LTE score' cross-product term. In *post-hoc* analyses we updated the first model and added the presence of a generalized anxiety disorder at T2 and T3 as a time-varying predictor. The maximum likelihood method was used for model estimation. The outcome variable was checked for normality and was natural log-transformed to meet the assumptions of a normal distribution. Results were considered statistically significant for a two-sided  $p$  value  $<0.05$ . All models were analysed using the nlme package (Pinheiro *et al.* 2012) in R, version 2.15.2 (R Foundation for Statistical Computing, 2012).

#### *Childhood and lifetime adversity and TL in adulthood*

To test our hypotheses that exposure to adverse life events during childhood or cumulative exposure to stressful life events over life is associated with shorter

TL in adulthood, we performed multivariable regression analysis with as a predictor variable either the 'childhood LTE score' or the 'lifetime LTE score', respectively. All models were adjusted for gender and age. The outcome variable was TL at T3. The outcome variable was checked for normality and was natural log-transformed to meet the assumptions of a normal distribution. The final models were also adjusted for BMI, presence of chronic diseases, smoking, frequency of sports, and level of education. To illustrate the influence of the covariates, we also present results of models that are only adjusted for gender and age. Results were considered statistically significant for a two-sided  $p$  value  $<0.05$ . All models were analysed using R, version 2.15.2 (R Foundation for Statistical Computing, 2012).

#### *Exploratory analyses*

Previous longitudinal studies investigating TL have reported both the possibility of telomere attrition and telomere lengthening (Aviv *et al.* 2009; Nordfjall *et al.* 2009; Chen *et al.* 2011; Svenson *et al.* 2011). Most studies defined attrition as a decrease in TL  $>15\%$ , and lengthening as an increase in TL  $>15\%$  between baseline and follow-up measures. We investigated the dynamics of TL in our cohort using the same definitions.

## Results

### *Descriptive statistics*

General characteristics of the study population can be found in Table 1. As mentioned above we used a decrease in TL  $>15\%$  and an increase in TL  $>15\%$  between baseline and follow-up measures as a definition of telomere attrition and lengthening respectively. Over an average time of 6.0 years between baseline (T1) and follow-up (T3) 63.0% of participants showed decrease in TL, 6.3% remained stable, and 30.7% showed lengthening of telomeres.

### *Adverse life events in adulthood and telomere attrition*

The results of the analysis of the model testing the hypothesis that recent adverse life events in adulthood are associated with telomere attrition adjusted for confounders and mediators can be found in Table 2. The recent LTE score was a significant predictor of telomere attrition ( $B = -0.028$ ,  $s.e. = 0.011$ ,  $p = 0.012$ ) in a random intercept model adjusted only for baseline TL and gender and age. Likewise, in a random intercept model that was additionally adjusted for lifestyle variables, the recent LTE score predicted a significant decrease in TL (Table 2). Age,

**Table 1.** General characteristics of the study population at T2

	N = 1094
Gender (female, %)	53.7
Age, mean (s.d.)	53.1 (11.4)
Race (%)	
Caucasian	97.6
Negroid	0.6
Asian	0.7
Other	1.0
Lifetime LTE score, mean (s.d.)	7.2 (3.8)
Recent LTE score* (%)	
0 event	75.9
1 event	14.6
2 events	5.9
3 events	2.8
≥4 events	0.9
Childhood LTE score* (%)	
0 events	60.7
1 event	26.7
2 events	8.0
3 events	3.0
≥4 events	1.6
Telomere length at T1, mean (s.d.)	0.417 (0.276)
Level of education (%)	
None	4.7
Low	26.5
Middle	27.1
High	41.7
Smoking, yes (%)	23.9
Smoking, cigarettes/day (%)	
None	76.1
1–5	4.3
6–10	3.9
11–15	6.4
16–20	5.4
>20	3.8
Exercise (%)	
I don't exercise	51.9
I exercise once per week	27.7
I exercise twice or more times per week	20.4
Body mass index (kg/m <sup>2</sup> ), mean (s.d.)	26.5 (4.1)
Chronic diseases (%)	
Healthy	80.9
1 chronic disease	15.6
2 chronic diseases	2.6
3 chronic diseases	0.6
4 chronic diseases	0.2
5 chronic diseases	0.1

s.d., standard deviation; LTE, List of Threatening Events.

\*Data not normally distributed thus for descriptive purposes only we created categories.

BMI, and smoking 16–20 cigarettes a day compared to none were also significant predictors of telomere attrition. The other categories of smoking status, gender,

**Table 2.** Three mixed models predicting telomere length at T2 and T3 by recent life events adjusted for various mediators and confounders

Beta	Model 1	Model 2	Model 3
Telomere length T1	0.209*	0.195*	0.199*
Age	−0.003*	−0.003*	−0.003*
Recent LTE score	−0.028*	−0.031*	−0.032*
Gender (female)	0.019	0.024	0.029
Time	0.006	0.006	0.004
Frequency of sports			
Once per week		−0.030	−0.024
Twice or more/week		0.027	0.029
Smoking (no. of cigarettes/day)			
1–5		−0.002	−0.008
6–10		−0.010	−0.046
11–15		0.005	0.005
16–20		−0.094*	−0.115*
>20		−0.057	−0.062
Body mass index		−0.007*	−0.006*
Chronic diseases		−0.006	−0.008
Education			
Low		−0.024	−0.019
Middle		−0.042	−0.058
High		−0.015	−0.011
Generalized anxiety disorder			−0.040

LTE, List of Threatening Events.

The recent LTE score = the sum score of adverse life events that took place in the year before T2 and the sum score of life events in the year that took place before T3.

\*Significant at  $\alpha < 0.05$ .

and level of education, however, did not predict changes in TL (Table 2). In a *post-hoc* analysis we also assessed if the effects of recent adverse life events were mediated by the presence of a generalized anxiety disorder by adding generalized anxiety disorder as a time-varying predictor to our final model. The recent LTE score remained a significant predictor of telomere attrition and the presence of a generalized anxiety disorder could not significantly explain the variance in telomere attrition (Table 2). We also tested the hypothesis that the effect of life events in the previous year (recent LTE score) was dependent on the lifetime LTE score by adding an interaction term between these two variables to the model. There was no significant interaction between the recent LTE score and the lifetime LTE score ( $p = 0.716$ ).

### Childhood and lifetime adversity and TL in adulthood

To assess whether exposure to adverse life events between the ages of 0 and 12 years has a long-lasting

**Table 3.** Multivariable regression analyses predicting telomere length at T3 by the childhood LTE score, adjusted for confounders and mediators

N = 898	Coefficient	S.E.	p value
Age	-0.003	0.001	0.003
Childhood LTE score	-0.001	0.013	0.936
Gender (female)	0.032	0.023	0.171
Frequency of sports			
Once per week	-0.007	0.028	0.793
Twice or more/week	0.044	0.029	0.122
Smoking (no. of cigarettes/day)			
1-5	-0.031	0.065	0.637
6-10	-0.038	0.056	0.504
11-15	0.023	0.049	0.637
16-20	-0.083	0.055	0.129
>20	-0.042	0.082	0.608
Body mass index	-0.007	0.003	0.019
Chronic diseases	-0.012	0.023	0.602
Education			
Low	0.043	0.058	0.463
Middle	0.037	0.059	0.535
High	0.051	0.058	0.378

S.E., Standard error.

Childhood LTE score = the sum score of adverse life events that took place in the 0-12 years age group.

effect on TL that can still be detected in adulthood, we investigated whether the childhood LTE score was negatively associated with TL in adulthood at T3. Multivariable regression analysis showed no relationship between the childhood LTE score and TL at T3 in model adjusted for confounders and mediators (Table 3), nor in a model adjusted only for gender and age ( $B = -0.001$ , S.E. = 0.012,  $p = 0.880$ ).

Furthermore, we tested if the cumulative exposure to stressful life events over life was associated with TL. The results of multivariable regression analysis showed that the lifetime LTE score was not significantly associated with TL at T3 in model adjusted for confounders and mediators (Table 4), nor in a model adjusted only for gender and age ( $B = 0.000$ , S.E. = 0.003,  $p = 0.959$ ). Only age and BMI were significant negative predictors of TL at T3.

## Discussion

To our knowledge, this is the first longitudinal population-based study demonstrating that exposure to adverse life events in adulthood is associated with telomere attrition. The effect of recent life events on telomere attrition was not moderated by the cumulative exposure to stressful life events over life, and not mediated by the presence of a generalized anxiety

**Table 4.** Multivariable regression analyses predicting telomere length at T3 by the lifetime LTE score, adjusted for confounders and mediators

N = 889	Coefficient	S.E.	p value
Age	-0.003	0.001	0.003
Lifetime LTE score	0.001	0.003	0.851
Gender (female)	0.031	0.023	0.181
Frequency of sports			
Once per week	-0.007	0.028	0.797
Twice or more/week	0.044	0.029	0.121
Smoking (no. of cigarettes/day)			
1-5	-0.031	0.065	0.634
6-10	-0.038	0.056	0.502
11-15	0.022	0.049	0.656
16-20	-0.083	0.055	0.127
>20	-0.044	0.082	0.595
Body mass index	-0.007	0.003	0.019
Chronic diseases	-0.013	0.024	0.573
Education			
Low	0.042	0.058	0.466
Middle	0.036	0.059	0.548
High	0.050	0.059	0.391

S.E., Standard error.

Lifetime LTE score = the sum score of adverse life events experienced over the entire life.

disorder. In addition to the effect of life events on telomere attrition, we have also found that age, BMI, and smoking between 16 and 20 cigarettes a day compared to none, are associated with increased telomere attrition. Strangely enough we did not find that smoking >20 cigarettes a day was associated with telomere attrition. At T3 the group that smoked >20 cigarettes a day (2.3%) was only half the size of the group that smoked between 16 and 20 cigarettes per day (4.9%). Perhaps we were underpowered to find an effect of smoking >20 cigarettes a day. We did not find a cross-sectional relationship between the lifetime LTE score or the childhood LTE score and TL.

There are several strengths and limitations of the current study that need to be taken into consideration when interpreting our results. The first major strength of this study is that it was conducted in a large population-representative cohort, which increases the generalizability of our findings. The second strength is the prospective nature of the design, allowing us to adjust for TL at baseline (T1) and therefore estimate change in TL over time.

Our finding that recent life events are prospectively associated with telomere attrition is in agreement with the results from the only two other prospective studies in this area of research, which were performed in children (Shalev *et al.* 2012) and post-menopausal,

non-smoking, disease-free women (Puterman *et al.* 2015). It is interesting that, although the life events in our study were of a very different nature than the ones of Shalev's study on child abuse, and took place in very different age groups, the effects on TL are the same. As opposed to the results of our longitudinal analysis, we did not find a cross-sectional relationship between the lifetime LTE score or the childhood LTE score and TL. Several explanations for this difference in findings might be offered. First, at birth there is already a high variability in TL between individuals (Takubo *et al.* 2002). This limits the power to detect environmental effects and might explain the often small effects reported in cross-sectional study psychophysiological research. A longitudinal study may focus on within-individual change and is thus less affected by large natural differences in TL between individuals. Some authors even claim that a cross-sectional study on TL would require five times the sample size a longitudinal study would need (Aviv *et al.* 2006). Nonetheless, also for the cross-sectional part of our study the sample size was fairly large.

Second, the lifetime LTE is a retrospective questionnaire that asks participants to report events that happened sometimes decades ago. Results from retrospective questionnaires suffer from recall bias. The literature shows that retrospective questionnaires mainly carry the risk of underreporting of events (Hardt & Rutter, 2004). This would lead to a loss of power to demonstrate a true effect of lifetime LTE score or the childhood LTE score on TL. We have tried to obviate this drawback by administering the same questionnaire twice and by scoring any event that was reported either at the first or second time the questionnaire was completed. Another limitation of the lifetime LTE score is that every event could only be scored once in every age group. Especially at old age it is not unthinkable that some events happen more than once. This would lead to an underestimation of the true number of life events. These limitations might in part explain the discrepancy in our findings between the cross-sectional and longitudinal part of our study, as the longitudinal part suffers less from recall bias.

Thirdly, TL is said to be dynamic and might depend upon both eroding (e.g. oxidative stress) and protective factors (e.g. antioxidants and telomerase). This is illustrated by the fact that telomere attrition per month is larger in the short term (after 6 months follow-up) than in the long term (after 10 years follow-up) (Svenson *et al.* 2011), indicating that repair mechanisms might play a role. In rats and humans there is evidence that exposure to uncontrollable stress (Beery *et al.* 2012) or depression (Wolkowitz *et al.* 2012) leads to higher blood cell telomerase activity than in healthy unstressed controls. In contrast, there is also a study in

which telomerase activity was reduced in caregivers *v.* non-caregiving controls (Epel *et al.* 2004). It might be that the body is able to counteract telomere erosion for a certain amount of time, such as during a short-term depression or after a stressful life event, but that repair systems become exhausted as stress exposure becomes more chronic. Although upregulation of the telomerase enzyme under certain conditions is a possibility, there is great controversy of whether telomere lengthening as reported in many studies (Aviv *et al.* 2009; Nordfjall *et al.* 2009; Chen *et al.* 2011; Svenson *et al.* 2011), including ours (van Ockenburg *et al.* 2014) really exists. A recently published study demonstrated that the reported telomere lengthening in longitudinal studies is most likely an artifact of laboratory measurement error, which is exacerbated by a short follow-up time (Steenstrup *et al.* 2013). In our study we used monochrome multiplex quantitative PCR and had an average follow-up time of 6 years. Although our measurement error was not that large and we demonstrated good reproducibility of results (Huzen *et al.* 2011), our method has a larger measurement error than for instance southern blot (Aviv *et al.* 2011). From the graphs in the paper of Steenstrup and colleagues it becomes clear that if we had run our samples in duplicate we would have misclassified about 30% of the population as telomere lengtheners. However, we ran our samples in triplicate, therefore reducing measurement error further than is exemplified in their paper. Furthermore, Steenstrup *et al.* classify any increase above zero as lengthening. We classify only those people that have at least a 15% increase in TL from baseline as lengtheners, which should reduce the chance of misclassification. Nonetheless, we acknowledge that measurement error is a problem in our field that deserves much more attention as a possible explanation for the reported results than is currently the case. We propose that future prospective studies should measure both TL and telomerase activity simultaneously. If telomerase activity is indeed higher in telomere lengtheners, this would give support for a biological explanation for telomere lengthening in addition to the role of measurement error.

In conclusion, this is the first study to demonstrate that exposure to recent adverse life events in adulthood is associated with decreased TL prospectively. Telomere attrition might thus be one of the mechanisms by which psychosocial stress exerts negative effects on health.

#### Declaration of Interest

None.



## Acknowledgements

This study was financially supported by a grant of the Innovational Research Incentives Scheme Program of The Netherlands (NWO, VENI, P. van der Harst, grant no. 916.76.170). The PREVENT was funded by the Dutch Kidney Foundation (Grant E033). The authors thank those that participated in the PREVENT study.

## References

- Aviv A, Chen W, Gardner JP, Kimura M, Brimacombe M, Cao X, Srinivasan SR, Berenson GS (2009). Leukocyte telomere dynamics: longitudinal findings among young adults in the Bogalusa Heart Study. *American Journal of Epidemiology* **169**, 323–329.
- Aviv A, Hunt SC, Lin J, Cao X, Kimura M, Blackburn E (2011). Impartial comparative analysis of measurement of leukocyte telomere length/DNA content by Southern blots and qPCR. *Nucleic Acids Research* **39**, e134.
- Aviv A, Valdes AM, Spector TD (2006). Human telomere biology: pitfalls of moving from the laboratory to epidemiology. *International Journal of Epidemiology* **35**, 1424–1429.
- Barrett EL, Richardson DS (2011). Sex differences in telomeres and lifespan. *Aging Cell* **10**, 913–921.
- Beery AK, Lin J, Biddle JS, Francis DD, Blackburn EH, Epel ES (2012). Chronic stress elevates telomerase activity in rats. *Biology Letters* **8**, 1063–1066.
- Ben-Shlomo Y, Kuh D (2002). A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *International Journal of Epidemiology* **31**, 285–293.
- Brugha TS, Cragg D (1990). The List of Threatening Experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatrica Scandinavica* **82**, 77–81.
- Cawthon RM (2009). Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Research* **37**, e21.
- Cawthon RM, Smith KR, O'Brien E, Sivatchenko A, Kerber RA (2003). Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet* **361**, 393–395.
- Chan SW, Blackburn EH (2002). New ways not to make ends meet: telomerase, DNA damage proteins and heterochromatin. *Oncogene* **21**, 553–563.
- Chen W, Kimura M, Kim S, Cao X, Srinivasan SR, Berenson GS, Kark JD, Aviv A (2011). Longitudinal versus cross-sectional evaluations of leukocyte telomere length dynamics: age-dependent telomere shortening is the rule. *Journal of Gerontology, Series A, Biological Sciences and Medical Sciences* **66**, 312–319.
- Chida Y, Hamer M (2008). Chronic psychosocial factors and acute physiological responses to laboratory-induced stress in healthy populations: a quantitative review of 30 years of investigations. *Psychological Bulletin* **134**, 829–885.
- Chida Y, Steptoe A (2010). Greater cardiovascular responses to laboratory mental stress are associated with poor subsequent cardiovascular risk status: a meta-analysis of prospective evidence. *Hypertension* **55**, 1026–1032.
- Du M, Prescott J, Kraft P, Han J, Giovannucci E, Hankinson SE, De Vivo I (2012). Physical activity, sedentary behavior, and leukocyte telomere length in women. *American Journal of Epidemiology* **175**, 414–422.
- Epel ES, Blackburn EH, Lin J, Dhabhar FS, Adler NE, Morrow JD, Cawthon RM (2004). Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of Sciences USA* **101**, 17312–17315.
- Epel ES, Lin J, Wilhelm FH, Wolkowitz OM, Cawthon R, Adler NE, Dolbier C, Mendes WB, Blackburn EH (2006). Cell aging in relation to stress arousal and cardiovascular disease risk factors. *Psychoneuroendocrinology* **31**, 277–287.
- Flores I, Cayuela ML, Blasco MA (2005). Effects of telomerase and telomere length on epidermal stem cell behavior. *Science* **309**, 1253–1256.
- Hardt J, Rutter M (2004). Validity of adult retrospective reports of adverse childhood experiences: review of the evidence. *Journal of Child Psychology and Psychiatry, and Allied Disciplines* **45**, 260–273.
- Harley CB, Futcher AB, Greider CW (1990). Telomeres shorten during ageing of human fibroblasts. *Nature* **345**, 458–460.
- Hoehn PW, Rosmalen JG, Schoevers RA, Huzen J, van der Harst P, de Jonge P (2013). Association between anxiety but not depressive disorders and leukocyte telomere length after 2 years of follow-up in a population-based sample. *Psychological Medicine* **43**, 689–697.
- Honig LS, Schupf N, Lee JH, Tang MX, Mayeux R (2006). Shorter telomeres are associated with mortality in those with APOE epsilon4 and dementia. *Annals of Neurology* **60**, 181–187.
- Hoven J, Peeters W, de Boer RA, Moll FL, Wong LS, Codd V, de Kleijn DP, de Smet BJ, van Veldhuisen DJ, Samani NJ, van Gilst WH, Pasterkamp G, van der Harst P (2011). Circulating leukocyte and carotid atherosclerotic plaque telomere length: interrelation, association with plaque characteristics, and restenosis after endarterectomy. *Arteriosclerosis, Thrombosis, and Vascular Biology* **31**, 1219–1225.
- Kenward MG, Molenberghs G (1998). Likelihood based frequentist inference when data are missing at random. *Statistical Science* **13**, 236–247.
- Miller AH, Maletic V, Raison CL (2009). Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biological Psychiatry* **65**, 732–741.
- Miller GE, Chen E, Zhou ES (2007). If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychological Bulletin* **133**, 25–45.
- Nordfjall K, Svenson U, Norrback KF, Adolfsson R, Lenner P, Roos G (2009). The individual blood cell telomere attrition rate is telomere length dependent. *PLoS Genetics* **5**, e1000375.
- Ohlin B, Nilsson PM, Nilsson JA, Berglund G (2004). Chronic psychosocial stress predicts long-term cardiovascular morbidity and mortality in middle-aged men. *European Heart Journal* **25**, 867–873.

- Pinheiro J, Bates D, DebRoy S, Sarkar D** (2012). nlme: Linear and Nonlinear Mixed Effects Models R package version 3.1–105.
- Pinto-Sietsma SJ, Janssen WM, Hillege HL, Navis G, de ZD, de Jong PE** (2000). Urinary albumin excretion is associated with renal functional abnormalities in a nondiabetic population. *Journal of the American Society of Nephrology* **11**, 1882–1888.
- Price LH, Kao HT, Burgers DE, Carpenter LL, Tyrka AR** (2012). Telomeres and early-life stress: an overview. *Biological Psychiatry* **73**, 15–23.
- Puterman E, Lin J, Krauss J, Blackburn EH, Epel ES** (2015). Determinants of telomere attrition over 1 year in healthy older women: stress and health behaviors matter. *Molecular Psychiatry* **20**, 529–535.
- R Foundation for Statistical Computing** (2012). **R: A Language and Environment for Statistical Computing.**
- Rosengren A, Hawken S, Ounpuu S, Sliwa K, Zubaid M, Almahmeed WA, Blackett KN, Sitthi-Amorn C, Sato H, Yusuf S** (2004). Association of psychosocial risk factors with risk of acute myocardial infarction in 11119 cases and 13648 controls from 52 countries (the INTERHEART study): case-control study. *Lancet* **364**, 953–962.
- Rosmalen JGM, Bos EH, de Jonge P** (2012). Validation of the Long-term Difficulties Inventory (LDI) and the List of Threatening Experiences (LTE) as measures of stress in epidemiological population-based cohort studies. *Psychological Medicine* **42**, 2599–2608.
- Segerstrom SC, Miller GE** (2004). Psychological stress and the human immune system: a meta-analytic study of 30 years of inquiry. *Psychological Bulletin* **130**, 601–630.
- Shalev I, Moffitt TE, Sugden K, Williams B, Houts RM, Danese A, Mill J, Arseneault L, Caspi A** (2012). Exposure to violence during childhood is associated with telomere erosion from 5 to 10 years of age: a longitudinal study. *Molecular Psychiatry* **18**, 576–581.
- Steenstrup T, Hjelmborg JV, Kark JD, Christensen K, Aviv A** (2013). The telomere lengthening conundrum—artifact or biology? *Nucleic Acids Research* **41**, e131.
- Stephoe A, Hamer M, Butcher L, Lin J, Brydon L, Kivimaki M, Marmot M, Blackburn E, Erusalimsky JD** (2011). Educational attainment but not measures of current socioeconomic circumstances are associated with leukocyte telomere length in healthy older men and women. *Brain, Behavior, and Immunity* **25**, 1292–1298.
- Stephoe A, Hamer M, Chida Y** (2007). The effects of acute psychological stress on circulating inflammatory factors in humans: a review and meta-analysis. *Brain, Behavior, and Immunity* **21**, 901–912.
- Svenson U, Nordfjall K, Baird D, Roger L, Osterman P, Hellenius ML, Roos G** (2011). Blood cell telomere length is a dynamic feature. *PLoS ONE* **6**, e21485.
- Takubo K, Izumiyama-Shimomura N, Honma N, Sawabe M, Arai T, Kato M, Oshimura M, Nakamura K** (2002). Telomere lengths are characteristic in each human individual. *Experimental Gerontology* **37**, 523–531.
- Valdes AM, Andrew T, Gardner JP, Kimura M, Oelsner E, Cherkas LF, Aviv A, Spector TD** (2005). Obesity, cigarette smoking, and telomere length in women. *Lancet* **366**, 662–664.
- van Ockenburg SL, de Jonge P, van der Harst P, Ormel J, Rosmalen JG** (2014). Does neuroticism make you old? Prospective associations between neuroticism and leukocyte telomere length. *Psychological Medicine*, **44**, 723–729.
- Vickers AJ, Altman DG** (2001). Statistics notes: analysing controlled trials with baseline and follow up measurements. *BMJ (Clinical Research ed.)* **323**, 1123–1124.
- Willeit P, Willeit J, Mayr A, Weger S, Oberhollenzer F, Brandstatter A, Kronenberg F, Kiechl S** (2010). Telomere length and risk of incident cancer and cancer mortality. *Journal of the American Medical Association* **304**, 69–75.
- Wolkowitz OM, Mellon SH, Epel ES, Lin J, Reus VI, Rosser R, Burke H, Compagnone M, Nelson JC, Dhabhar FS, Blackburn EH** (2012). Resting leukocyte telomerase activity is elevated in major depression and predicts treatment response. *Molecular Psychiatry* **17**, 164–172.