

THE COMPLEXITY OF HOMEOMORPHISM RELATIONS ON SOME CLASSES OF COMPACTA*

PAWEŁ KRUPSKI AND BENJAMIN VEJNAR

Abstract. We prove that the homeomorphism relation between compact spaces can be continuously reduced to the homeomorphism equivalence relation between absolute retracts, which strengthens and simplifies recent results of Chang and Gao, and Cieřla. It follows then that the homeomorphism relation of absolute retracts is Borel bireducible with the universal orbit equivalence relation. We also prove that the homeomorphism relation between regular continua is classifiable by countable structures and hence it is Borel bireducible with the universal orbit equivalence relation of the permutation group on a countable set. On the other hand we prove that the homeomorphism relation between rim-finite metrizable compacta is not classifiable by countable structures.

§1. Introduction. Measuring the complexity of structures is a very general task. Usually we study the complexity in a relative way by comparing with other structures. One possible approach for this by using embeddings of categories was elaborated by Pultr and Trnkova in [25]. However, in this paper we use the notions of *Borel reducibility* and *Borel bireducibility* to relate the complexities of equivalence relations on standard Borel spaces or Polish spaces in order to compare the complexities of classification problems. For more details we refer to the book of Gao [10].

Several equivalence relations became milestones in the theory of Borel reductions, we will mention four of them:

- the equality on an uncountable Polish space,
- the S_∞ -universal orbit equivalence relation,
- the universal orbit equivalence relation,
- the universal analytic equivalence relation.

Let us give several examples to make the above relations more familiar. A classical example is a result of Gromov (see e.g., [10, Theorem 14.2.1]) who proved that the isometry equivalence relation of compact metric spaces is a *smooth* equivalence relation, which means that it is Borel reducible to the equality of real numbers (or equivalently of an uncountable Polish space). The isomorphism relation of countable graphs or the isomorphism relation of countable linear orders are Borel bireducible with the S_∞ -universal orbit equivalence. The isometry equivalence relation of Polish metric spaces and the isometry relation of separable Banach

Received July 1, 2019.

2020 *Mathematics Subject Classification.* 03E15, 54H05, 54F15.

Key words and phrases. Borel reduction, countable structures, universal orbit equivalence relation, absolute retract, LC^n -compactum, regular continuum.

*Dedicated to the memory of Věra Trnkova.

© 2020, Association for Symbolic Logic
0022-4812/20/8502-0013
DOI:10.1017/jsl.2020.3

spaces were proved by Gao and Kechris in [11] and by Melleray in [22] to be Borel bireducible with the universal orbit equivalence relation (see a survey paper by Motto Ros [23]). Ferenczi, Louveau, and Rosendal proved in [9] that the isomorphism equivalence relation of separable Banach spaces is Borel bireducible with the universal analytic equivalence relation.

In order to capture all the structures in one space we need some sort of *coding*. This can be done by considering some universal space (e.g., the Hilbert cube, the Urysohn space or the Gurarii space) and all its subspaces with some natural Polish topology or Borel structure (e.g., the hyperspace topology or the Effros Borel structure). Sometimes there are other natural ways to encode a given structure. For example the class of Polish metric spaces can be coded by the set of all metrics on \mathbb{N} where two metrics are defined to be equivalent if the completions of the respective spaces are isometric. Fortunately by [10, Theorem 14.1.3] it does not matter which coding we choose. It is generally believed that this independence on a natural coding is common to other structures and thus the statements are usually formulated for all structures without mentioning the current coding.

In this paper we are dealing solely with the classification problem of compact metrizable spaces up to homeomorphism. Unless otherwise stated we assume that the coding of compact metrizable spaces is given by the hyperspace of the Hilbert cube. Based on coding techniques developed by Sabok [27], Zielinski proved in [30] that the homeomorphism equivalence relation of metrizable compacta is Borel bireducible with the universal orbit equivalence relation. Soon after Chang and Gao proved in [4] that a similar result is true if we restrict the relation only to continua. Recently Cieřla proved in [5] that a similar result holds true also for locally connected continua. We prove in Theorem 6 that the homeomorphism relation restricted to absolute retracts still has the same complexity. By a similar idea we prove in Theorem 9 that at most n -dimensional continua are Borel reducible to at most n -dimensional LC^{n-1} -continua, for every $n \in \mathbb{N}$. However, the exact complexity of at most n -dimensional compacta remains unknown.

Camerlo, Darji, and Marcone proved in [2] that the homeomorphism relation of dendrites is Borel bireducible with the S_∞ -universal orbit equivalence relation, especially dendrites are classifiable by countable structures. A similar result was proved by Camerlo and Gao in [3] for zero-dimensional compacta instead of dendrites. We prove in Theorem 16 that the homeomorphism equivalence relation of regular continua has the complexity of countable structures. On the other hand it is surprising that the homeomorphism relation of rim-finite compacta, which is a natural small class containing all dendrites and zero-dimensional compacta, is not classifiable by countable structures; see Theorem 18.

§2. Definitions and notations. Recall that a *standard Borel space* is a measurable space (X, \mathcal{S}) such that there is a Polish topology τ on X for which the family $\text{Borel}(X, \tau)$ of Borel subsets of (X, τ) is equal to \mathcal{S} . In order to compare the complexities of equivalence relations we use the notion of Borel reducibility.

DEFINITION 1. Suppose that X and Y are standard Borel spaces and let E, F be (usually analytic) equivalence relations on X and Y respectively. We say that E is *Borel reducible* to F , and we denote this by $E \leq_B F$, if there exists a Borel mapping

$f: X \rightarrow Y$ such that

$$x E x' \iff f(x) F f(x'),$$

for every $x, x' \in X$. The function f is called a *Borel reduction*. We say that E is *Borel bireducible* with F , and we write $E \sim_B F$, if E is Borel reducible to F and F is Borel reducible to E .

In a similar fashion we can define being *continuously reducible* if we suppose that X and Y are Polish spaces and f is continuous.

In the whole paper we denote $I = [0, 1]$ and $Q = I^{\mathbb{N}}$. For a separable metric space X we denote by 2^X the hyperspace of all compacta in X with the Hausdorff distance d_H . The hyperspace $C(X)$ stands for all continua in X . Since we are dealing mainly with the *homeomorphism equivalence relation* between elements of some class \mathcal{C} of metric compacta, we restrict the relation to $\mathcal{C} \cap 2^Q$. For example, we say that the class of continua is continuously reducible to the class of compacta instead of the longer expression that the homeomorphism relation of the class of all continua in the Hilbert cube is continuously reducible to the homeomorphism relation of all compacta in the Hilbert cube.

We say that an equivalence relation E defined on a standard Borel space X is *classifiable by countable structures* if there is a countable relational language \mathcal{L} such that E is Borel reducible to the isomorphism relation of \mathcal{L} -structures whose underlying set is \mathbb{N} . An equivalence relation E on a standard Borel space X is said to be an *orbit equivalence relation* if there is a Borel action of a Polish group G on X such that $x E x'$ if and only if there is some $g \in G$ for which $gx = x'$.

Let \mathcal{C} be a class of equivalence relations on standard Borel spaces. An element $E \in \mathcal{C}$ is called *universal* for \mathcal{C} if $F \leq_B E$ for every $F \in \mathcal{C}$. It is known that for every Polish group G there is an equivalence relation (denoted by E_G) on a standard Borel space that is universal for all orbit equivalence relations given by G -actions. We are particularly interested in the *universal S_∞ -equivalence relation* E_{S_∞} . It is known that an equivalence relation is classifiable by countable structures if and only if it is Borel reducible to E_{S_∞} . Also there exists *universal orbit equivalence relation* which is denoted by E_{G_∞} . We should also note that all the mentioned equivalence relations are analytic.

A space X is an *absolute retract* (*absolute neighborhood retract*) if it is a retract of any space Y (of an open neighborhood of X in Y) containing X as a closed subset. A space X is C^n (*connected in dimension n*) if for each $0 \leq m \leq n$ every continuous map from the unit m -sphere S^m into X has a continuous extension over the unit $(m+1)$ -ball B^{m+1} . X is LC^n (*locally connected in dimension n*) if for each $x \in X$, for each $0 \leq m \leq n$ and each neighborhood U of x there is a neighborhood $V \subset U$ of x such that every continuous map $S^m \rightarrow V$ extends to a continuous map $B^{m+1} \rightarrow U$. Locally connected metric continua (which are also called Peano continua since they are continuous images of I) are known to be both C^0 and LC^0 . An *arc* is a space homeomorphic to I and an *n -od* is a space homeomorphic to the cone over the set $\{1, \dots, n\}$.

A continuum is called *arc-like* if it is an inverse limit of arcs with continuous and onto bonding mappings. A *dendrite* is a locally connected continuum which does not contain a simple closed curve. It is worth mentioning that a nondegenerate dendrite

is fully characterized as an absolute retract of dimension one. A point x of a space X is of order $\leq n$ in X ($\text{ord}_x X \leq n$) if there is a neighborhood base of x whose members have boundaries of cardinality $\leq n$. If $\text{ord}_x X \leq n$ and $\text{ord}_x X \not\leq n-1$ then we write $\text{ord}_x X = n$ and say that x is of order n in X . A compact space is said to be *rim-finite* if it has an open base formed by sets with finite boundaries. A rim-finite continuum is usually called a *regular continuum*. Recall here that if X is a regular continuum then X is hereditarily locally connected, thus a closed set $A \subseteq X$ separates X between two distinct points $x, y \in X \setminus A$ if and only if A cuts X between x and y if and only if every arc from x to y intersects A .

In the next section we are going to use properties of Z -sets in the Hilbert cube or in the Menger universal continuum M_n ([28] and [1] are good references). The crucial facts are:

- the Z -set unknotting theorem: each homeomorphism between Z -sets in Q or in M_n extends to a homeomorphism of Q or M_n , respectively, onto itself [28, Theorem 5.3.7] and [1, Corollary 3.1.5],
- Z -approximation theorem: if B is a compactum such that $\dim B \leq n$, then any continuous map $f: B \rightarrow M_n$ can be approximated by a Z -embedding (i.e., by an embedding such that $f(B)$ is a Z -set in M_n) [1, Theorem 2.3.8],
- each closed subset of a Z -set is a Z -set,
- the Hilbert cube as well as the Menger universal continua contain their respective copies as Z -sets. In fact $Q \times \{0\}$ is a Z -set in $Q \times I$.

For the definition and basic properties of the topological dimension and continuum theory we refer the reader to [8, Chapter 7] and to [24], [28] and [29].

§3. Main results. We present three classification results in theorems 6, 9, and 16. Also there is one nonclassification result in Theorem 18. The following proposition was first proved by Knaster and Reichbach [17, Théorème 5]. It is also a consequence of [14, Proposition 8.8] and the first sentence of its proof. It can be proved using the back and forth argument.

PROPOSITION 2. *Let $K \subseteq A, L \subseteq B$ be four nonempty compact metrizable spaces such that $A \setminus K$ and $B \setminus L$ are dense sets of isolated points in A and B respectively. Then every homeomorphism of K onto L can be extended to a homeomorphism of A onto B .*

The following lemma is a consequence of [7]; see [12, Theorem 2.6] for more details. It will be extremely useful in the proof of Theorem 6.

LEMMA 3. *If X is a nondegenerate Peano continuum then there exists a homotopy $H: 2^X \times I \rightarrow 2^X$ for which*

- $H(A, 0) = A$ for every $A \in 2^X$,
- $H(A, t)$ is finite for every $t > 0$ and $A \in 2^X$.

Recall that if $Y \subseteq X$ and $\varepsilon > 0$ we say that X is ε -deformable into Y if there exists a continuous mapping $\varphi: X \times [0, 1] \rightarrow X$ such that $\varphi(x, 0) = x$, $\varphi(x, 1) \in Y$ and the

diameter of $\varphi(\{x\} \times [0, 1])$ is at most ε for every $x \in X$. The following proposition was proved in [18, 1.1 and 1.3].

PROPOSITION 4. *Let X be a compact space such that for every $\varepsilon > 0$ there exists an absolute neighborhood retract (absolute retract) $Y \subseteq X$ for which X is ε -deformable into Y . Then X is an absolute neighborhood retract (absolute retract, resp.).*

The proof of Proposition 4 is based on the Lefschetz characterization of absolute neighborhood retracts via extending partial realizations of finite polyhedra to their full realizations. By restricting polyhedra of arbitrary dimensions to polyhedra of dimension $\leq n$, one gets a similar characterization of LC^{n-1} -compacta (see [28, Proposition 4.2.29]) and the proof works for LC^{n-1} compacta. So we get the following analogue of Proposition 4.

PROPOSITION 5. *Let X be a compact space such that for every $\varepsilon > 0$ there exists an LC^{n-1} -compactum $Y \subseteq X$ for which X is ε -deformable into Y . Then X is an LC^{n-1} -compactum.*

THEOREM 6. *Compacta are continuously bireducible with absolute retracts.*

PROOF. Let us denote

$$\begin{aligned} Q_0 &= Q \times \{(0, 0, 0)\}, \\ Q_1 &= Q \times I \times \{(0, 0)\}, \\ Q_2 &= Q \times I \times I \times \{0\}, \\ Q_3 &= Q \times I \times I \times I. \end{aligned}$$

It follows that Q_i are Hilbert cubes such that Q_i is a Z -set in Q_{i+1} for $i = 0, 1, 2$. We construct a reduction $\varphi: 2^Q \rightarrow 2^{Q_3}$. Suppose that $H: 2^Q \times I \rightarrow 2^Q$ is given by Lemma 3 and let

$$\psi(K) = (K \times \{(0, 0, 0)\}) \cup \bigcup_{n=0}^{\infty} H(K, 2^{-n}) \times \{(2^{-n}, 0, 0)\}.$$

Hence $\psi(K)$ is a compact subspace of Q_1 containing a homeomorphic copy of K and a countable set of isolated points which is open and dense in $\psi(K)$. Denote

$$\varphi(K) = Q_2 \cup \bigcup_{n=0}^{\infty} H(K, 2^{-n}) \times \{(2^{-n}, 0)\} \times [0, 2^{-n}].$$

Roughly speaking, $\varphi(K)$ consists of the Hilbert cube Q_2 to which a null sequence (i.e., a sequence whose diameters converge to zero) of mutually disjoint segments is attached. Therefore, by Proposition 4, $\varphi(K)$ is an absolute retract. Continuity of the mapping φ is straightforward so it only remains to prove that φ is a reduction.

Suppose first that $K, L \in 2^Q$ and $h: K \rightarrow L$ is a homeomorphism. By Proposition 2 there is a homeomorphism $h_1: \psi(K) \rightarrow \psi(L)$ extending the homeomorphism $h \times \{(0, 0, 0)\}$. Since Q_1 is a Z -set in Q_2 , there is a homeomorphism

$h_2: Q_2 \rightarrow Q_2$ which extends h_1 . Now, extend h_2 to $h_3: \varphi(K) \rightarrow \varphi(L)$ on the remaining arcs linearly. Hence $\varphi(K)$ is homeomorphic to $\varphi(L)$.

On the other hand suppose that $\varphi(K)$ is homeomorphic to $\varphi(L)$ by some homeomorphism g . Since g as well as g^{-1} needs to map points of order two (local cut points) to points of order two (local cut points, resp.) it follows that g is sending the following countable set

$$\bigcup_{n=0}^{\infty} H(K, 2^{-n}) \times \{(2^{-n}, 0, 0)\}$$

onto the set

$$\bigcup_{n=0}^{\infty} H(L, 2^{-n}) \times \{(2^{-n}, 0, 0)\},$$

because these sets are formed exactly by local cut points of $\varphi(K)$ and $\varphi(L)$, respectively, which are not of order two. Since $K \times \{(0, 0, 0)\}$ forms the set of cluster points of the first set and $L \times \{(0, 0, 0)\}$ forms the set of cluster points of the second one, it means that $g(K \times \{(0, 0, 0)\}) = L \times \{(0, 0, 0)\}$. Hence K and L are homeomorphic. \dashv

An n -dimensional analogue of the Hilbert cube is the universal n -dimensional Menger continuum M_n which is characterized as an n -dimensional LC^{n-1} , C^{n-1} continuum that satisfies the disjoint n -cells property [1]. Also, LC^{n-1} compacta are natural analogues of compact absolute neighborhood retracts among compacta of dimension $\leq n$ and LC^{n-1} , C^{n-1} -continua of such dimensions correspond to absolute retracts. Therefore a question arises whether the class \mathcal{C}_n of all compacta of dimension $\leq n$ is continuously bireducible with the class $\mathcal{C}_n \cap LC^{n-1} \cap C^{n-1}$ of at most n -dimensional LC^{n-1} , C^{n-1} continua. Notice that \mathcal{C}_n is a G_δ -subset of 2^Q and, by a Kuratowski's result [20, Théorème A], the family LC^{n-1} is a one-to-one continuous image of a complete space, so it is Borel. Unfortunately, it is unknown if the family $\mathcal{C}_n \cap LC^{n-1} \cap C^{n-1}$ is Borel for any $n > 1$ (for $n = 1$ it is known to be $F_{\sigma\delta}$). We are able to show the Borel bireducibility of \mathcal{C}_n to $\mathcal{C}_n \cap LC^{n-1}$ continua. The strategy is first to Borel reduce \mathcal{C}_n to 2^{M_n} (here, we do not know if this can be done continuously) and next, to proceed similarly as in the proof of Theorem 6.

LEMMA 7. *At most n -dimensional compacta in the Hilbert cube are Borel reducible to compacta in the universal n -dimensional Menger continuum for each $n \in \mathbb{N}$.*

PROOF. Let us denote by $H_p(Q, M_n)$ the set of all partial embeddings $f: K \rightarrow M_n$ of compact subsets $K \subseteq Q$. It is shown in [19] to be a Polish space with the topology induced by the convergence

$$f_k \rightarrow f \Leftrightarrow d_H(\text{dom } f_k, \text{dom } f) \rightarrow 0 \text{ and } f_k(x_k) \rightarrow f(x)$$

for each sequence $x_k \rightarrow x$, $x_k \in \text{dom } f_k$, $x \in \text{dom } f$, where dom denotes the domain of the respective mapping. Consider the multivalued mapping Φ from \mathcal{C}_n given by

$$\Phi(K) = \{f \in H_p(Q, M_n) : \text{dom } f = K\}.$$

CLAIM 8. Φ is lower semicontinuous.

Indeed, let U be an open subset of $H_p(Q, M_n)$. We are going to show that the set

$$\{K \in \mathcal{C}_n : \Phi(K) \cap U \neq \emptyset\}$$

is open in \mathcal{C}_n . Suppose it is not. Then there is an embedding $f: K \rightarrow M_n$ in U such that a sequence $K_i \in \mathcal{C}_n$ exists satisfying $K_i \rightarrow K$ and $\Phi(K_i) \cap U = \emptyset$. By the Z -approximation theorem for M_n [1, Theorem 2.3.8], we can assume that f is a Z -embedding. Since $K \cup \bigcup_i K_i \in \mathcal{C}_n$ and M_n is an absolute extensor for the class \mathcal{C}_n , there is a continuous extension $\bar{f}: K \cup \bigcup_i K_i \rightarrow M_n$ of f . Again, by the Z -approximation theorem, \bar{f} can be approximated by Z -embeddings $g: K \cup \bigcup_i K_i \rightarrow M_n$ which agree with f on K . It follows that mappings $g_i = g|_{K_i}$ converge to f in $H_p(Q, M_n)$, so $g_i \in U$ for sufficiently large i . This contradicts the equality $\Phi(K_i) \cap U = \emptyset$.

By applying the Kuratowski and Ryll-Nardzewski selection theorem [21] to the mapping Φ we get a Borel selection map $\varphi: \mathcal{C}_n \rightarrow H_p(Q, M_n)$. The required reduction is the function $\mathcal{C}_n \ni K \mapsto \pi\varphi(K) \in 2^{M_n}$, where π is the projection $Q \times M_n \rightarrow M_n$. \dashv

THEOREM 9. At most n -dimensional compacta are Borel bireducible with at most n -dimensional LC^{n-1} -continua, for every $n \in \mathbb{N}$.

PROOF. By Lemma 7, it remains to continuously reduce the family 2^{M_n} to the family of at most n -dimensional LC^{n-1} continua in Q . Fix a pair $M_n \subseteq M'_n$ of topological copies of the universal n -dimensional Menger continua in Q such that M_n is a Z -set in M'_n (clearly, this is possible by the Z -approximation theorem). Thus, for each $k \in \mathbb{N}$, there is a continuous mapping $f_k: M'_n \rightarrow M'_n \setminus M_n$ such that

$$d_{sup}(f_k, id_{M'_n}) < \frac{1}{k} \quad \text{and} \quad f_{k+1}(M_n) \cap f_i(M_n) = \emptyset \quad \text{for} \quad i < k + 1.$$

Put $f_0 = id_{M'_n}$. There is also a homotopy $H: 2^{M_n} \times I \rightarrow 2^{M_n}$ through finite sets by Lemma 3. Given $K \in 2^{M_n}$, let

$$\psi(K) = (K \times \{0\}) \cup \bigcup_{k=0}^{\infty} f_k(H(K, 2^{-k})) \times \{0\}$$

and define a continuous mapping

$$\varphi(K) = (M'_n \times \{0\}) \cup \bigcup_{k=0}^{\infty} f_k(H(K, 2^{-k})) \times [0, 2^{-k}].$$

Hence, $\varphi(K)$ is the Menger continuum M'_n with a null sequence of mutually disjoint segments attached, so it is n -dimensional. Moreover, observe that attaching to an LC^{n-1} -compactum finitely many disjoint arcs yields an LC^{n-1} -compactum and $\varphi(K)$ is ε -deformable into such a compactum for every $\varepsilon > 0$. Thus, by Proposition 5, $\varphi(K)$ is an LC^{n-1} -continuum.

In order to show that φ is a reduction, suppose $h: K \rightarrow L$ is a homeomorphism between $K, L \in 2^{M_n}$. Extend $\bar{h}(x, 0) := (h(x), 0)$, $x \in K$, to a homeomorphism $h_1: \psi(K) \rightarrow \psi(L)$ by Proposition 2. Since $\psi(K)$ and $\psi(L)$ are Z -sets in $M'_n \times \{0\}$, there is a homeomorphism $h_2: M'_n \times \{0\} \rightarrow M'_n \times \{0\}$ that extends h_1

[1, Corollary 3.1.5]. Finally, extend h_2 to a homeomorphism $h_3: \varphi(K) \rightarrow \varphi(L)$ linearly over the attached segments.

Conversely, if $g: \varphi(K) \rightarrow \varphi(L)$ is a homeomorphism then, as in the proof of Theorem 6, we conclude that

$$g\left(\bigcup_{k=0}^{\infty} f_k(H(K, 2^{-k})) \times \{0\}\right) = \bigcup_{k=0}^{\infty} f_k(H(L, 2^{-k})) \times \{0\}$$

and, consequently, $g(K \times \{0\}) = L \times \{0\}$, hence K and L are homeomorphic. \dashv

PROPOSITION 10. *Let X be a standard Borel space, Y be a Polish space and $A \subseteq X \times Y$ be a Borel set whose vertical sections are σ -compact. Then $\pi_X(A)$ is Borel and there exist Borel functions $f_n: \pi_X(A) \rightarrow Y$ whose graphs are subsets of A and such that for every $x \in \pi_X(A)$ the set $\{f_n(x): n \in \mathbb{N}\}$ is dense in the vertical section A_x .*

PROOF. Let $\{B_n: n \in \mathbb{N}\}$ be an open countable base of Y with $B_1 = Y$. Notice that $A \cap (X \times B_n)$ has σ -compact vertical sections. Hence, for every $n \in \mathbb{N}$ we can apply the Arsenin-Kunugui Uniformization Theorem [16, Theorem 35.46] to the Borel set $A \cap (X \times B_n) \subseteq X \times Y$ to get a Borel uniformization $g_n: X_n \rightarrow B_n$ where $X_n = \pi_X(A \cap (X \times B_n))$ is Borel in X . Now, it is enough to define

$$f_n(x) = \begin{cases} g_n(x), & x \in X_n, \\ g_1(x), & x \in \pi_X(A) \setminus X_n. \end{cases}$$

Clearly all the functions f_n are Borel and the set $\{f_n(x): n \in \mathbb{N}\}$ is dense in A_x for every $x \in X$. \dashv

If $A \subseteq X$ is an arc with end points a and b such that $A \setminus \{a, b\}$ is an open subset of X then $A \setminus \{a, b\}$ is called a *free arc* in X . If $A \setminus \{a, b\}$ is a free arc, then the union of all free arcs in X containing $A \setminus \{a, b\}$ will be called a maximal free arc in X . Let A_X be the union of all free arcs in X .

Let us denote by \mathcal{R} the set of all regular subcontinua of the Hilbert cube Q and for $X \in \mathcal{R}$, denote by B_X the set of all *branch points* (i.e., points of order ≥ 3) which are also *local cut points* of X (i.e., cut points of some open connected neighborhoods).

LEMMA 11. *Let X be a regular continuum and let $x, y \in X$ be distinct. Then there is a finite set $H \subseteq A_X \cup B_X$ which separates x from y . In particular, $A_X \cup B_X$ is dense in X .*

PROOF. Let $F \subseteq X$ be a minimal finite set separating x and y . Each point of F is a local cut point of X [29, Corollary 9.42]. Also, for every $s \in F$ there is an arc from x to y containing s (otherwise, F would not be a minimal separating set). Let $T \subseteq F$ be the set of points with order two. For every $t \in T$ choose an arc A_t with end points x and y containing t and an open connected neighborhood U_t of t whose closure is disjoint from $F \setminus \{t\}$ and such that $A_t \cap U_t$ is an arc without end points. We can assume that t cuts U_t .

Let $\text{bd}(A_t \cap \overline{U_t}) = \{a, b\}$ and

$$M = \{p \in U_t \cap A_t : p \text{ separates } a \text{ and } b \text{ in } \overline{U_t}\}.$$

Notice that $t \in M$ since F separates x and y . It is enough to prove that at least one of the following holds:

- (1) there is a branch point $z \in A_t \cap U_t$ which separates a and b in $\overline{U_t}$,
- (2) there is a point $z \in A_t \cap U_t$ lying on a free arc which separates a and b in $\overline{U_t}$.

Indeed, in both cases point t can be replaced by z , i.e., the set $(F \setminus \{t\}) \cup \{z\}$ separates $\{x\}$ and $\{y\}$; after finitely many such replacements we get desired set H .

So, suppose condition (1) is not satisfied. We prove that condition (2) holds true. By our assumption every point of M is a point of order two. One can easily prove that

- CLAIM 12. a) *Every point in M is approximated by points in M from both sides with respect to a natural order of $A_t \cap U_t$. Another way of stating this is that no point of M is accessible from $(A_t \cap U_t) \setminus M$.*
 b) *The set M is closed in $A_t \cap U_t$.*

Proof of a) If $z \in M$ then z is of order two. We consider two-point boundaries of small neighborhoods of z contained in U_t . The boundary points separate a from b in $\overline{U_t}$ and hence they are in M .

Proof of b) The limit of a sequence from M is either outside U_t or it separates a from b in $\overline{U_t}$. Hence the limit must be in M as well.

Since $M \neq \emptyset$, it follows by a) and b) that $M = A_t \cap U_t$. Hence every point of U_t is of order two and thus U_t is a free arc and $t \in U_t$ separates a and b in $\overline{U_t}$. Hence, condition (2) is proved. □

DEFINITION 13. Suppose that X and Y are regular continua and let $C \subseteq X$ and $D \subseteq Y$. A bijection $f : C \rightarrow D$ is called *separation preserving* if, for every pair $x, y \in C$ and a finite set $F \subseteq C$, x and y are separated by F if and only if $f(x)$ and $f(y)$ are separated by $f(F)$.

LEMMA 14. *Let X and Y be regular continua and let C and D be dense subsets of X and Y respectively, for which $B_X \subseteq C \subseteq B_X \cup A_X$ and $B_Y \subseteq D \subseteq B_Y \cup A_Y$. Then every separation preserving bijection $f : C \rightarrow D$ can be extended to a homeomorphism of X onto Y .*

PROOF. Since X and Y are compact there is only one uniformity on X and Y respectively. Let us prove that f is uniformly continuous. Suppose for contradiction that this is not the case. Then there exist $\varepsilon > 0$ and two sequences $(x_n), (y_n)$ such that the distances of x_n and y_n converge to zero, but the distance of $f(x_n)$ and $f(y_n)$ is bigger than ε . Since X and Y are compact, we can assume that $f(x_n)$ converges to some $a \in Y$, $f(y_n)$ converges to some $b \in Y$ and both the sequence x_n and y_n converge to the same point $z \in X$.

By Lemma 11 there is a finite set $H_1 \subseteq B_Y \cup A_Y$ separating a and b . By a small adjustment of points in $H_1 \cap A_Y$ we can suppose that $H_1 \subseteq D$. Similarly, there is H_2 with the same properties which is disjoint from H_1 . This follows, e.g., from [24,

Proposition 10.18] where the “additive-hereditary system” is the family of finite subsets of $B_Y \cup A_Y$ and H_2 is the boundary of a neighborhood of a disjoint from H_1 , slightly adjusted to be in D . There is i such that $f^{-1}(H_i)$ does not contain the point z . Since X is locally connected, there is a connected neighborhood U of z which is disjoint from $f^{-1}(H_i)$. For n sufficiently large, $x_n, y_n \in U$ and hence x_n and y_n are not separated by $f^{-1}(H_i)$.

Since Y is locally connected there are connected neighborhoods of a and b disjoint from H_i , hence it follows that for n sufficiently large, $f(x_n)$ and $f(y_n)$ are separated by H_i . We get a contradiction, since the function f was supposed to be separation preserving.

Since f is uniformly continuous it can be extended to a continuous mapping of X to Y [8, Theorem 4.3.17]. Since, moreover, we can do the same with f^{-1} , the extended mapping is a homeomorphism. ⊖

For the purpose of the following lemma and Theorem 16 let us fix the sets

$$\mathcal{A} = \{(X, a) : X \in \mathcal{R}, a \in A_X\}, \quad \mathcal{B} = \{(X, b) : X \in \mathcal{R}, b \in B_X\}.$$

LEMMA 15. *The sets \mathcal{A} and \mathcal{B} are Borel subsets of $C(Q) \times Q$.*

PROOF. We will mostly follow a corresponding idea for dendrites from the proof of [2, Lemma 6.4]. So, in order to show that the sets are Borel, we check that they are both analytic and coanalytic. The proof for coanalyticity of \mathcal{B} is exactly the same for regular continua as for dendrites. To show that \mathcal{B} is analytic we have to modify the respective formula as follows:

$(X, b) \in \mathcal{B}$ iff $b \in X$, $X \in \mathcal{R}$ and there exist arcs $A_1, A_2, A_3 \subseteq X$ such that $b \in A_1 \cap A_2 \cap A_3$ and for every $x \in Q$ and every distinct $i, j \in \{1, 2, 3\}$ if $x \in A_i \cap A_j$ then $x = b$.

Before considering \mathcal{A} , let us first check that the set $\mathcal{E} := \{(X, e) : X \in \mathcal{R}, e \in E_X\}$, where E_X is the set of all points $e \in X$ of order $\text{ord}_X e = 1$, is Borel in $C(Q) \times Q$. Indeed, its analyticity can be shown as for the set \mathcal{E}^D in [2, Lemma 6.4]. To see that it is coanalytic, one can characterize \mathcal{E} in the following way:

$(X, e) \in \mathcal{E}$ iff $e \in X$, $X \in \mathcal{R}$ and, for every two arcs $A_1, A_2 \in C(Q)$, if $e \in A_1 \cap A_2$ and $A_1, A_2 \subseteq X$ then there is $x \in Q$ such that $x \in A_1 \cap A_2$ and $x \neq e$.

It follows that the set $\mathcal{O} := \{(X, x) : X \in \mathcal{R}, \text{ord}_X x = 2\} = \{(X, x) : X \in \mathcal{R}, x \in X\} \setminus (\mathcal{B} \cup \mathcal{E})$ is Borel.

Now, we are ready to characterize \mathcal{A} :

$(X, a) \in \mathcal{A}$ iff $(X, a) \in \mathcal{O}$ and there exists $k \in \mathbb{N}$ such that, for each $x \in Q$, if $d(a, x) < \frac{1}{k}$ and $x \in X$ then $(X, x) \in \mathcal{O}$.

This shows coanalyticity of \mathcal{A} . On the other hand, the following description gives its analyticity. Let $\{B_i : i \in \mathbb{N}\}$ be an open countable base in Q . Then

$(X, a) \in \mathcal{A}$ iff $a \in X$, $X \in \mathcal{R}$ and there is an arc $A \in C(Q)$ such that $a \in A$, $A \subseteq X$ and there is i such that $a \in B_i$, $B_i \cap X \subseteq A$. ⊖

THEOREM 16. *Regular continua are classifiable by countable structures.*

PROOF. We will assign to every regular continuum X a sequence $(R_n^X)_{n=1}^\infty$ of relations, such that $R_n^X \subseteq X^{n+2}$ for which $(x, y, x_1, x_2, \dots, x_n) \in R_n^X$ iff $\{x_1, \dots, x_n\}$

separates x and y . Note that the sets A_X are open in X and hence σ -compact, and, by [29, p. 606], the sets B_X are countable for $X \in \mathcal{R}$. Notice that $A_X \cap B_X = \emptyset$ and that $A_X \cup B_X$ is dense in X by Lemma 11. Let $\mathcal{R}_A, \mathcal{R}_B$ denote the sets of continua $X \in \mathcal{R}$ such that $A_X \neq \emptyset, B_X \neq \emptyset$, respectively.

Clearly

$$\mathcal{R} = (\mathcal{R} \setminus \mathcal{R}_B) \cup (\mathcal{R}_B \setminus \mathcal{R}_A) \cup (\mathcal{R}_B \cap \mathcal{R}_A).$$

If we prove that each of the summands is Borel and it is classifiable by countable structures, it will follow that \mathcal{R} is classifiable by countable structures as well. The class $\mathcal{R} \setminus \mathcal{R}_B$ contains only three topological types of continua: a point, an arc and a simple closed curve. By [26, Theorem 1] it follows that they are Borel. Hence the set $\mathcal{R} \setminus \mathcal{R}_B$ is easily classifiable by countable structures (this is even a *smooth* equivalence relation). The case of $\mathcal{R}_B \setminus \mathcal{R}_A$ can be handled similarly as the remaining one, so we will skip it. In the rest we prove that $\mathcal{R}_B \cap \mathcal{R}_A$ is classifiable by countable structures.

The vertical sections of the set \mathcal{A} are σ -compact and \mathcal{A} is Borel by Lemma 15. Hence, by Proposition 10 the set \mathcal{R}_A is Borel and there are Borel mappings $b_{2n-1}: \mathcal{R}_A \rightarrow \mathcal{Q}$ such that $\{b_{2n-1}(X): n \in \mathbb{N}\}$ is dense in A_X . We can suppose that every point of the dense set occurs infinitely many times.

The vertical sections of \mathcal{B} are countable and \mathcal{B} is Borel by Lemma 15. It follows by the Lusin-Novikov Uniformization Theorem [16, Theorem 18.10] that \mathcal{R}_B is Borel and there are countably many Borel mappings $b_{2n}: \mathcal{R}_B \rightarrow \mathcal{Q}$ such that $\{b_{2n}(X): n \in \mathbb{N}\} = B_X$. We can easily assume that every point occurs infinitely many times.

We define an $(n + 2)$ -ary relation S_n^X on \mathbb{N} . For $k, l, m_1, \dots, m_n \in \mathbb{N}$ let

$$S_n^X(k, l, m_1, \dots, m_n) \text{ iff } R_n^X(b_k(X), b_l(X), b_{m_1}(X), \dots, b_{m_n}(X)),$$

and let

$$\Phi(X) = (\mathbb{N}, (S_n^X)_{n \in \mathbb{N}}).$$

Roughly speaking, the mappings b_n allow us to code relations R_n^X by relations on \mathbb{N} in a Borel way.

Let us check that Φ is a Borel reduction of $\mathcal{R}_A \cap \mathcal{R}_B$ to countable structures. Suppose that $h: X \rightarrow Y$ is a homeomorphism and $X, Y \in \mathcal{R}_A \cap \mathcal{R}_B$. Clearly, $h(A_X) = A_Y$ and $h(B_X) = B_Y$. For any $x \in B_X$, the sets $\{n \in \mathbb{N}: b_{2n}(X) = x\}$ and $\{n \in \mathbb{N}: b_{2n}(Y) = h(x)\}$ are infinite. So let φ_x be an arbitrary bijection between them.

Let A be a maximal free arc in X . Note that \overline{A} is either an arc or a simple closed curve. The set $E_X = \{b_{2n-1}(X): n \in \mathbb{N}\} \cap A$ is countable and dense in A . The same is true for the set $E_Y = \{b_{2n-1}(Y): n \in \mathbb{N}\} \cap h(A)$ in $h(A)$. Since the real line is countable dense homogeneous, there is a homeomorphism $\psi_A: \overline{A} \rightarrow h(\overline{A})$ such that $\psi_A|_{(\overline{A} \setminus A)} = h|_{(\overline{A} \setminus A)}$ and $\psi_A(E_X) = E_Y$. Now, for every $x \in E_X$, the sets $\{n \in \mathbb{N}: b_{2n-1}(X) = x\}$ and $\{n \in \mathbb{N}: b_{2n-1}(Y) = \psi_A(x)\}$ are infinite, hence there is a bijection φ_x between these sets. Let $\varphi = \bigcup \{\varphi_x: x \in b_{\mathbb{N}}(X)\}$. It is straightforward to check that $\varphi: \Phi(X) \rightarrow \Phi(Y)$ is an isomorphism.

On the other hand, assume that $\varphi: \Phi(X) \rightarrow \Phi(Y)$ is an isomorphism. Let us define $h(b_k(X)) = b_{\varphi(k)}(Y)$, which is a correct definition (not depending on k but only on $b_k(X)$). Indeed if $b_{\varphi(k)}(Y) \neq b_{\varphi(l)}(Y)$, then using Lemma 11 we can find a finite set $\{y_1, \dots, y_n\} \subseteq b_{\mathbb{N}}(Y)$ separating $b_{\varphi(k)}(Y)$ and $b_{\varphi(l)}(Y)$. Suppose that $y_i = b_{m_i}(Y)$.

Then $S_n^Y(\varphi(k), \varphi(l), m_1, \dots, m_n)$, thus $S_n^X(k, l, \varphi^{-1}(m_1), \dots, \varphi^{-1}(m_n))$, since φ is an isomorphism. Hence the points $b_k(X)$ and $b_l(X)$ are separated and so they are different points. Similarly, if $b_k(X) \neq b_l(X)$, then $b_{\varphi(k)}(Y) \neq b_{\varphi(l)}(Y)$.

Let $C = b_{\mathbb{N}}(X)$ and let $D = b_{\mathbb{N}}(Y)$. Clearly, $h: C \rightarrow D$ is a separation preserving bijection and hence, by Lemma 14, it can be extended to a homeomorphism of X onto Y . ←

Let $c_0 = \{x \in \mathbb{R}^{\mathbb{N}} : x_n \rightarrow 0\}$. For a set X of sequences of real numbers we denote by X/c_0 the equivalence relation for which two sequences are equivalent if and only if their difference converges to zero.

LEMMA 17. *The equivalence relation $I^{\mathbb{N}}/c_0$ is not classifiable by countable structures.*

PROOF. By the proof of [15, Lemma 6.2.2] it follows that $\mathbb{R}^{\mathbb{N}}/c_0$ is Borel reducible to $I^{\mathbb{N}}/c_0$. The equivalence relation $\mathbb{R}^{\mathbb{N}}/c_0$ is known to be *turbulent* and hence by [13, Example 3.23] it is not classifiable by countable structures. Hence $I^{\mathbb{N}}/c_0$ is not classifiable by countable structures too. ←

THEOREM 18. *Rim-finite compacta are not classifiable by countable structures.*

PROOF. By Lemma 17, it is enough to prove that $I^{\mathbb{N}}/c_0$ is Borel reducible to rim-finite compacta. Let $J = I \times \{0\}$. For $n \geq 3$ denote by $T_n \subseteq \mathbb{R} \times (-\infty, 0) \cup \{(0, 0)\}$ a simple n -od with diameter less than 2^{-n-1} whose vertex is located at the point $(0, 0)$. For $a, b \in \mathbb{R}$ denote $T_n(a, b) = (a, b) + T_n$. Let $\{q_n : n \in \mathbb{N}\}$ be a dense subset of $(0, 1)$ enumerated without repetitions and let

$$L = J \cup \bigcup_{n \in \mathbb{N}} T_{k_n}(q_n, 0),$$

where $k_n \in \mathbb{N}$ are odd, $3 < k_1 < k_2 < \dots$ and $T_{k_n}(q_n, 0)$ is disjoint from $T_{k_m}(q_m, 0)$ for $m \neq n$. For $x \in I^{\mathbb{N}}$ define a compact subset of the plane

$$\Phi(x) = L \cup \bigcup_{n \in \mathbb{N}} T_{2n+2}(x_n, 2^{-n}).$$

Let us verify that Φ is a reduction. Suppose that $x, y \in I^{\mathbb{N}}$ and $x - y \in c_0$. There are homeomorphisms $h_n : T_{2n+2}(x_n, 2^{-n}) \rightarrow T_{2n+2}(y_n, 2^{-n})$, simply let $h_n(a, b) = (a - x_n + y_n, b)$. It follows that we can define $h : \Phi(x) \rightarrow \Phi(y)$ in such a way that h is the identity on L and $h(z) = h_n(z)$ if $z \in T_{2n+2}(x_n, 2^{-n})$. One can easily check that h is one-to-one continuous and thus it is a homeomorphism.

On the other hand, suppose that h is a homeomorphism of $\Phi(x)$ and $\Phi(y)$ for $x, y \in I^{\mathbb{N}}$. In $\Phi(x)$ as well as in $\Phi(y)$, there is at most one point of order n for every $n \geq 4$. Hence h needs to preserve these points, especially $h|_J$ equals the identity on a dense subset of J and, by continuity, $h|_J$ is the identity. Moreover, we get $h(x_n, 2^{-n}) = (y_n, 2^{-n})$. Since

$$|x_n - y_n| = \|(x_n, 0) - (y_n, 0)\| \leq \|(x_n, 0) - (y_n, 2^{-n})\| = \|h(x_n, 0) - h(x_n, 2^{-n})\|,$$

we get $x - y \in c_0$, by the uniform continuity of h .

The reduction Φ is not only Borel, but even continuous. This easy verification is left to the reader. \dashv

We should remark that in the preceding proof we used simple n -ods as countably many markers in order to distinguish some points. We can even prove that compacta of order at most two are not classifiable by countable structures. As in the preceding proof we find a Borel reduction of $I^{\mathbb{N}}/c_0$ to such compacta. For a sequence $(x_n) \in I^{\mathbb{N}}$ we attach to I some compact countable spaces whose Cantor-Bendixson rank is some countable ordinal number instead of simple n -ods (we can use e.g., the ordinal $\omega^n + 1$ with the order topology instead of the simple n -od T_n). So we get a compact space $\Phi(x)$ with only one nondegenerate component all of whose points are of order at most two. Hence we get the following result.

THEOREM 19. *Compacta of order at most two are not classifiable by countable structures.*

§4. Auxiliary results and questions. In this section we provide two simple ways how to prove that continua are Borel bireducible with universal orbit equivalence, once we know that compacta have this property. This simplifies the result of [4] substantially. Also we prove that arc-like continua and hereditarily decomposable continua are not classifiable by countable structures.

FACT 20. [6] *There exists a nondegenerate continuum C such that for every subcontinuum $D \subseteq C$ any continuous map $f: D \rightarrow C$ is either the identity or a constant map. Any continuum with these properties is called a Cook continuum.*

A construction of the Cook continuum can be also found in [25, pp. 319–341].


NOTE 21. *Below we present a simple construction of a continuous reduction of compacta to continua.*

Let $C \subseteq Q$ be a Cook continuum. Fix a point $c \in C$. Let $Q' = Q \times C / Q \times \{c\}$ be the quotient space of $Q \times C$ where $Q \times \{c\}$ is shrunk to a point and denote by $q: Q \times C \rightarrow Q'$ the corresponding quotient mapping. To every $K \in 2^Q$ assign $\varphi(K) = q(K \times C)$. Hence, $\varphi(K)$ can be viewed as a cone over K with the Cook continuum instead of an arc as its fiber. Let us verify that $\varphi: 2^Q \rightarrow 2^{Q'}$ is a reduction of the homeomorphism relation of compacta to the homeomorphism relation of continua. It is obvious that if K and L are homeomorphic, then $\varphi(K)$ and $\varphi(L)$ are homeomorphic. On the other hand, assume that $h: \varphi(K) \rightarrow \varphi(L)$ is a homeomorphism. Let $q(x, a) \in \varphi(K)$ and $h(q(x, a)) = q(y, b) \in \varphi(L \times C)$. Suppose that $a \neq b$. Denote by $i_x: C \rightarrow q(\{x\} \times C)$ the embedding given by $i_x(d) = q(x, d)$. Let $\tilde{\pi}_2: Q' \rightarrow C$ be such that $\tilde{\pi}_2 \circ q = \pi_2$, where π_2 is the projection of $Q \times C$ onto C . Consider the mapping $\tilde{\pi}_2 \circ h \circ i_x: C \rightarrow C$ which maps the point a to b . Since these points are distinct, the preceding mapping needs to be a constant map by the properties of the Cook continuum mentioned in Fact 20. It follows that h maps the set $i_x(C)$ into the set $q(L \times \{b\})$. In particular, the point $q(x, c)$ is mapped to $q(x, b)$. Since $q(x, c) = q(x', c)$ for every $x' \in K$, it follows analogously that h maps every set $i_{x'}(C)$ into the set $q(L \times \{b\})$. Thus, h is not an onto mapping, a contradiction. Consequently, h maps $q(K \times \{a\})$ onto $q(L \times \{a\})$ for every $a \in C$. Hence, K and L are homeomorphic. It is also easy to see that φ is continuous.

NOTE 22. *There are more simple ways of obtaining such a reduction. Let us sketch the one which avoids a Cook continuum, but uses a nondegenerate arc-less continuum B . Fix a point $b \in B$. To a compactum X assign its cone, $\text{cone}(X) = X \times I / X \times \{1\}$, and attach $X \times B$ to $X \times \{0\} \subseteq \text{cone}(X)$ along $X \times \{b\}$. Thus obtained continuum $\varphi(X)$ has only one arc component with nonempty interior (namely the cone) and its boundary is $X \times \{0\}$. It follows easily that φ is a continuous reduction of all compacta to continua.*

PROPOSITION 23. *Neither arc-like continua nor hereditarily decomposable continua are classifiable by countable structures.*

PROOF. Similarly to the proof of Theorem 18 it suffices to prove that $I^{\mathbb{N}}/c_0$ is Borel reducible to arc-like hereditarily decomposable continua. We describe the idea just very informally. For a sequence $(x_n) \in I^{\mathbb{N}}$, consider a continuum in the plane consisting of an arc I with a ray L approximating I from the left which makes the arc rigid for self-homeomorphisms. From the other side take another ray R approximating I with small two-sided $\sin(1/x)$ -curves inserted into small neighborhoods of the points A_1, A_2, \dots which encode sequence (x_n) . See the figure below. ⊣



NOTE 24. *Dendrites can be naturally coded either as a subset of the hyperspace of the Hilbert cube or as the hyperspace of all subcontinua of the universal dendrite. Fortunately both these codings are equivalent in the sense that the corresponding homeomorphism equivalence relations are Borel bireducible one with the other. This follows actually by the proof of [2]. Alternatively, one can Borel reduce countable linear orders to dendrites in the universal dendrite. Since countable linear orders are Borel bireducible with the universal S_∞ orbit equivalence relation we obtain the same for dendrites in the universal dendrite.*

QUESTION 25. *Are there some Borel reductions decreasing the topological dimension?*

QUESTION 26. *Is the universal orbit equivalence Borel reducible to rim-finite compacta? Is it reducible to compacta of order at most two?*

Acknowledgment. Partially supported by the Faculty of Fundamental Problems of Technology, Wrocław University of Science and Technology, grant 0401/0017/17.

The author was supported by the grant GAČR 17-04197Y. This work has been supported by Charles University Research Centre program No. UNCE/SCI/022.

REFERENCES

- [1] M. BESTVINA, *Characterizing k -dimensional universal Menger compacta*. *Memoirs of the American Mathematical Society*, vol. 71 (1988), no. 380.
- [2] R. CAMERLO, U. B. DARJI, and A. MARCONE, *Classification problems in continuum theory*. *Transactions of the American Mathematical Society*, vol. 357 (2005), no. 11, pp. 4301–4328.
- [3] R. CAMERLO and S. GAO, *The completeness of the isomorphism relation for countable Boolean algebras*. *Transactions of the American Mathematical Society*, vol. 353 (2001), no. 2, pp. 491–518.
- [4] C. CHANG and S. GAO, *The complexity of the classification problem of continua*. *Proceedings of the American Mathematical Society*, vol. 145 (2017), no. 3, pp. 1329–1342.
- [5] T. CIEŚLA, *Completeness of the homeomorphism relation of locally connected continua*. *Proceedings of the American Mathematical Society*, vol. 147 (2019), no. 3, pp. 1269–1276.
- [6] H. COOK, *Continua which admit only the identity mapping onto non-degenerate subcontinua*. *Fundamenta Mathematicae*, vol. 60 (1967), pp. 241–249.
- [7] D. W. CURTIS, *Hyperspaces of finite subsets as boundary sets*. *Topology and Its Applications*, vol. 22 (1986), no. 1, pp. 97–107.
- [8] R. ENGELKING, *General Topology*, Sigma Series in Pure Mathematics, vol. 6, second ed., Heldermann Verlag, Berlin, 1989.
- [9] V. FERENCZI, A. LOUVEAU, and C. ROSENDAL, *The complexity of classifying separable Banach spaces up to isomorphism*. *Journal of the London Mathematical Society*, vol. 79 (2009), no. 2, pp. 323–345.
- [10] S. GAO, *Invariant Descriptive Set Theory*, Pure and Applied Mathematics (Boca Raton), vol. 293, CRC Press, Boca Raton, FL, 2009.
- [11] S. GAO and A. S. KECHRIS, *On the classification of Polish metric spaces up to isometry*. *Memoirs of the American Mathematical Society*, vol. 161 (2003), no. 766, p. viii+78.
- [12] H. GLADDINES and J. VAN MILL, *Hyperspaces of Peano continua of Euclidean spaces*. *Fundamenta Mathematicae*, vol. 142 (1993), no. 2, pp. 173–188.
- [13] G. HJORTH, *Classification and Orbit Equivalence Relations*, Mathematical Surveys and Monographs, vol. 75, American Mathematical Society, Providence, RI, 2000.
- [14] A. ILLANES and S. B. NADLER, JR., *Hyperspaces: Fundamentals and Recent Advances*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 216, Marcel Dekker, Inc., New York, 1999.
- [15] V. KANOVEI, *Borel Equivalence Relations: Structure and Classification*, University Lecture Series, vol. 44, American Mathematical Society, Providence, RI, 2008.
- [16] A. S. KECHRIS, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995.
- [17] B. KNASTER and M. REICHBACH, *Notion d'homogénéité et prolongements des homéomorphismes*. *Fundamenta Mathematicae*, vol. 40 (1953), pp. 180–193.
- [18] J. KRASINKIEWICZ, *On a method of constructing ANR-sets. An application of inverse limits*. *Fundamenta Mathematicae*, vol. 92 (1976), no. 2, pp. 95–112.
- [19] K. KURATOWSKI, *Sur l'espace des fonctions partielles*. *Annali di Matematica Pura ed Applicata*, vol. 40 (1955), no. 4, pp. 61–67.
- [20] ———, *Sur une méthode de métrisation complète de certains espaces d'ensembles compact*. *Fundamenta Mathematicae*, vol. 43 (1956), pp. 114–138.
- [21] K. KURATOWSKI and C. RYLL-NARDZEWSKI, *A general theorem on selectors*. *Bulletin L'Académie Polonaise des Science, Série des Sciences Mathématiques, Astronomiques et Physiques*, vol. 13 (1965), pp. 397–403.
- [22] J. MELLERAY, *Computing the complexity of the relation of isometry between separable Banach spaces*. *Mathematical Logic Quarterly*, vol. 53 (2007), no. 2, pp. 128–131.
- [23] L. MOTTO ROS, *Can we classify complete metric spaces up to isometry?*. *Bollettino dell'Unione Matematica Italiana*, vol. 10 (2017), no. 3, pp. 369–410.

- [24] S. B. NADLER, JR., *An Introduction: Continuum Theory*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 158, Marcel Dekker, Inc., New York, 1992.
- [25] A. PULTR and V. TRNKOVÁ, *Combinatorial, Algebraic and Topological Representations of Groups, Semigroups and Categories*, North-Holland Mathematical Library, vol. 22, North-Holland Publishing Co., Amsterdam/New York, 1980.
- [26] C. RYLL-NARDZEWSKI, *On a Freedman's problem*. *Fundamenta Mathematicae*, vol. 57 (1965), pp. 273–274.
- [27] M. SABOK, *Completeness of the isomorphism problem for separable C^* -algebras*. *Inventiones Mathematicae*, vol. 204 (2016), no. 3, pp. 833–868.
- [28] J. VAN MILL, *The Infinite-Dimensional Topology of Function Spaces*, North-Holland Mathematical Library, vol. 64, North-Holland Publishing Co., Amsterdam, 2001.
- [29] G. THOMAS WHYBURN, *Analytic topology*, American Mathematical Society Colloquium Publications, vol. 28, American Mathematical Society, New York, 1942.
- [30] J. ZIELINSKI, *The complexity of the homeomorphism relation between compact metric spaces*. *Advances in Mathematics*, vol. 291 (2016), pp. 635–645.

DEPARTMENT OF COMPUTER SCIENCE
FACULTY OF FUNDAMENTAL PROBLEMS OF TECHNOLOGY
WROCIĄW UNIVERSITY OF SCIENCE AND TECHNOLOGY
WROCIĄW, POLAND

E-mail: pawel.krupski@pwr.edu.pl

DEPARTMENT OF MATHEMATICAL ANALYSIS
FACULTY OF MATHEMATICS AND PHYSICS
CHARLES UNIVERSITY
PRAGUE, CZECHIA

E-mail: vejnar@karlin.mff.cuni.cz