

Perverse Leray filtration and specialisation with applications to the Hitchin morphism

BY MARK ANDREA A. DE CATALDO

*Department of Mathematics, Stony Brook University, Stony Brook,
NY, 11794-3651, U.S.A*

e-mail: mark.decataldo@stonybrook.edu

(Received 4 April 2020; revised 4 February 2021; accepted 3 February 2021)

Abstract

We initiate and develop a framework to handle the specialisation morphism as a filtered morphism for the perverse, and for the perverse Leray filtration, on the cohomology with constructible coefficients of varieties and morphisms parameterised by a curve. As an application, we use this framework to carry out a detailed study of filtered specialisation for the Hitchin morphisms associated with the compactification of Dolbeault moduli spaces in [de 2018].

2020 Mathematics Subject Classification: 14A10, 14D06, 14D20, 14H60 (Primary).

1. Introduction, notation and preliminaries

Let $v : Y \rightarrow S$ be a morphism into a connected nonsingular curve, let $s \in S$ be a point, let $G \in D_c^b(Y)$ be a bounded constructible complex, let $t \in S$ be a suitably general point (in a more algebraic set-up, the geometric generic point of the curve, or of an Henselian trait), let Y_s and Y_t be the corresponding fibers.

It is natural, and of fundamental importance, to compare the cohomology groups $H^j(Y_s, G|_{Y_s})$ and $H^j(Y_t, G|_{Y_t})$. One classical example, is the study of Lefschetz pencils. Similarly, if $X \rightarrow Y$ is an S -morphism, then it is equally important to compare, for $F \in D_c^b(X)$, the cohomology groups $H^j(X_s, F|_{X_s}) \rightarrow H^j(X_t, F|_{X_t})$.

When it is defined, the specialisation morphism $H^j(Y_s, G|_{Y_s}) \rightarrow H^j(Y_t, G|_{Y_t})$, and similarly for X_s and X_t , is a key tool for this comparison.

The paper [de-Ma 2018] studies the Hitchin S -morphism $f : X \rightarrow Y$ for a smooth and projective family \mathcal{X}/S and establishes that the resulting specialisation morphisms in intersection cohomology exist, and that they are filtered isomorphisms for the respective perverse Leray filtrations. The two authors realised that there seems to be no available discussion in the literature of the specialisation morphism as a filtered morphism, so that they developed a criterion for having such a filtered isomorphism that worked in the context of the Hitchin morphism associated with a family of smooth varieties over a base.

In this paper, we initiate and develop a general framework to study the specialisation morphism as a filtered morphism for the perverse (Leray) filtration. We then apply such a framework to a more detailed study of the Hitchin morphism.

1.1. *Motivation and outline of the results*

Let us consider the toy model situation in Remark 3.5.7 based on the Set-up 3.4.1, where we start with a smooth morphism $f : X^o \rightarrow Y^o$ over a smooth curve S , and we compactify f by adding divisors Z and W , to get $f : X^o \cup Z = X \rightarrow Y = Y^o \cup W$, so that all resulting morphisms are smooth and proper, except for X^o and Y^o over S . Then the morphisms of long exact sequences (69) for the relative singular cohomology of the pairs $(X_{t,s}, X_{t,s}^o)$ and $(Y_{t,s}, Y_{t,s}^o)$ relating any two point $s, t \in S$, is a filtered isomorphisms for the Leray spectral sequences for the morphisms $f_t : X_t \rightarrow Y_t$ and $f_s : X_s \rightarrow Y_s$.

What happens when the morphisms are not proper, the varieties are singular, we take coefficients in an arbitrary bounded constructible complex of sheaves in $D_c^b(X)$, and we consider the perverse Leray filtration? This is the question addressed in this paper.

First of all, why the perverse Leray filtration? The middle perversity t -structure is more suitable to study singularities, so we focus on the perverse (Leray) filtration, instead of the Grothendieck (Leray) filtration. In principle the Leray filtration can be studied with the methods of this paper, but the results I can think of are much weaker, essentially due to the fact that the vanishing/nearby cycle functors have several important perverse t -exactness properties, without having a counterpart for the usual standard t -structure.

Let us focus on an arbitrary situation $X/Y/S$ with $K \in D_c^b(X)$, with $s \in S$ a point and $t \in S$ a suitably general point (the geometric generic point of the curve/Henselian trait, if the reader prefers that language). In this general context, the specialisation morphism (“from a special point to a general point”) $H^*(X_s, K|_{X_s}) \rightarrow H^*(X_t, K|_{X_t})$ is not even defined. This is due to the failure of the relevant base-change morphisms to be isomorphisms. Even when the specialisation morphism is defined, its perverse filtered counterpart may fail to be well-defined.

The purpose of this paper is to develop a framework where these questions can be studied systematically. What follows is a list of some of the outcomes of this study, presented in a weaker and less complete form with respect to what is found in the body of the paper, so that the reader may get an idea of the techniques introduced and of the results that are proved.

Let us start with some of the preparatory results in Section 2. In the set up of morphisms $X \rightarrow Y \rightarrow S$ and of the specialisation to a point s on the curve S , the fibers X_s and Y_s over the special point are Cartier divisor. A key point is to study, in the more general set-up of a Cartier divisor T' inside a variety T , the failure of commutativity of the perverse truncation functors ${}^p\tau_{\leq k}$ with the restriction i^* to T' , when applied to a complex $G \in D(T)$. This is codified in the failure to be isomorphisms of a certain morphisms that we introduce and denote by δ (22). Proposition 2.1.5 says that the δ are isomorphisms when G has no constituents (e.g. no non-trivial direct summands) supported on the divisor T' . Lemma 2.2.1, which is placed in the context of specialisation, gives a criterion for the δ to be isomorphisms in terms of the vanishing $\phi G = 0$ of the vanishing cycles of the complex G . Proposition 2.3.2 is another criterion for the δ being isomorphisms for a complex f_*F , when we are in a projective morphism $f : X \rightarrow Y$ situation (not necessarily over a curve S), where a Cartier divisor on X , pulled back from Y , has suitable transversality conditions with respect to the

complex F on X : essentially, one requires the complex F on X to be semisimple and to stay semisimple after restriction to the Cartier divisor. Corollary 2.4.2 shows that this kind of transversality can be achieved in a simple normal crossing singularities situation, where the restriction does not stay semisimple.

As it may be clear by now, the main goal of this paper is to identify conditions that ensure that specialisation morphisms are defined, and, when they are defined, that they are isomorphisms, and, in the filtered context, that they are filtered isomorphisms.

Following the preparation in Section 2, concerning the morphisms δ , we zero in on the problem by defining the perverse filtered version of the specialisation morphism (when it exists), and by offering some criteria in the main section, Section 3 of this paper.

Let me now state a version of the main Theorem 3.3.7 which is a list of different criteria for having well defined specialisation filtered isomorphisms. Let us stress that when v is not proper, without additional constraints on the situation, the specialisation morphism may fail to be well-defined. We are in the situation $f : X \rightarrow Y$, $v : Y \rightarrow S$, $F \in D_c^b(X)$. In order to try to convey the flavor of the theorem, we offer two different sets of conditions, each of which is a sufficient set of conditions: f and v are proper, and $\phi(F) = 0$; f proper, F semisimple and $\phi(F) = 0$.

We apply these results to a compactification of the morphism f as above, i.e. as in Set-up 3.4.1. In this case, we prove Theorem 3.5.4, i.e. that, under suitable sets of hypotheses, the situation of the toy model discussed at the beginning of this section can be reproduced in its entirety: the specialisation morphisms for X_s , X_t , X_s^o , X_t^o and Z_s , Z_t are filtered isomorphisms compatibly with the restriction and Gysin morphisms (which one needs to show are well-defined) stemming from the inclusions.

Finally, Theorem 4.4.2 is our application of the methods of this paper, and especially of Theorem 3.5.4, to the compactification of Dolbeault moduli spaces constructed in [de 2018], thus showing that the various criteria developed in this paper can actually be implemented in a highly non-trivial and geometrically interesting situation. Perhaps surprisingly, this is true whether we consider intersection cohomology, or singular cohomology.

1.2. More precise outline of the contents of the paper

Section 1 includes this introduction Section 1.1, lists the general notation Section 1.3 and then discusses the formalism of the vanishing/nearby cycles functors; more precisely: Section 1.4 summarises the parts of the formalism of vanishing/nearby cycles functors that we need in this paper; in particular, Fact 1.4.6, and (13) are of key importance; Section 1.5 explains in some detail the geometric intuition behind the nearby cycle functor.

Section 2 is preparatory in nature: it introduces and discusses the morphisms of type δ ; these are key to this paper since their being isomorphisms is necessary to the specialisation morphisms being filtered isomorphisms for the perverse (Leray) filtration. The heart of this paper, i.e. the discussion of the specialisation morphism as a filtered morphism for the perverse Leray filtration, is Section 3. In order to carry out that discussion we need to measure the failure of the restriction-to-the-special-fiber functor to commute with perverse truncation. This failure is measured by the morphisms of type δ , which are defined in Lemma 2.1.2 and Remark 2.1.3, in the more general context of effective Cartier divisors (the special fiber being one such). Proposition 2.1.5 gives a sufficient condition for the morphisms of type δ to be isomorphisms. Lemma 2.2.1 establishes the key facts we need when dealing with the morphisms of type δ together with the vanishing/nearby cycle functors; in particular,

the vanishing of the vanishing cycle functor gives a sufficient condition for the morphisms of type δ being isomorphisms. Proposition 2.3.2 provides another such criterion under the assumption that certain relative hard Lefschetz symmetries are in place; Corollary 2.4.2 ensures that if we are in a simple normal crossing divisor situation, then such symmetries are in place.

The aforementioned framework is developed in Section 3. We refer to the beginning of that section for a more detailed account of its contents. Here we simply list the main points. The set-up is the one of an S -morphism $X \rightarrow Y$, where S is a nonsingular connected curve, and $s \in S$ is a point, the “special” point. The definition of the specialisation morphism as a filtered morphism for the perverse Leray filtration is contained in Definition 3.3.3. The main Theorem of this paper is Theorem 3.3.7, which establishes various criteria for this morphisms to exist and to be a filtered isomorphism. One of the main themes here is to work with non proper structural morphisms $X \rightarrow S$ and $Y \rightarrow S$, for in this case, the base change morphisms are not isomorphisms in general. Compactifying the situation is one traditional way to circumvent this issue; this introduces additional base change issues, and Proposition 3.4.2 provides criteria to resolve them. Once a compactification is in place, one has the long exact sequence of the resulting triple (boundary, compactification, original space), and Theorem 3.5.4 provides three criteria ensuring that the resulting three specialisation morphisms gives rise to an isomorphism of filtered long exact sequences associated with the special and general triples.

Section 4 applies the abstract framework developed in Section 3 in the case of the Hitchin morphisms arising from the compactification of Dolbeault moduli spaces introduced in [de 2018]. We refer to the beginning of that section for a detailed outline of the contents of this section. The main result is Theorem 4.4.2, to the effect that we get the desired isomorphism of filtered long exact sequences associated with the special and general triples stemming from the particular compactification [de 2018]. Certain preliminary results concerning descending certain properties along \mathbb{G}_m -quotients, which could be of general independent interest, are established along the way.

Remark 1.2.1. (Other algebraically closed fields, generic geometric points vs general points). We have chosen to work with vanishing/nearby cycles with respect to a morphism into a nonsingular curve, over the field of complex numbers, with the classical topology, and with finite algebraic Whitney stratifications (which play a role only in the background, by merely existing and having the usual properties). While this is mostly a matter of expository style, we also work in a more global (not over a curve/Henselian trait) context in parts of Section 2. By complementing the references given in Section 1.4 with [II 1994, Section 4], which also deals with vanishing/nearby cycles and the perverse t-structure, the exposition and the results of this paper remain valid, mutatis mutandis, for varieties over algebraically closed fields of characteristic zero and for the étale cohomology with coefficients in $\overline{\mathbb{Q}}_\ell$, for any prime ℓ ; for example, the role played in this paper by a suitably general point t on a nonsingular curve S is now played by the geometric generic point of S , or by the geometric generic point of an Henselian trait. Similarly, except for Section 4, where in the application to the compactification of the Hitchin morphism we need to consider quotients by finite groups on which we have little control, the results remain valid over an algebraically closed field of positive characteristic and $\overline{\mathbb{Q}}_\ell$ -coefficients, with ℓ not dividing the characteristic of the field.

1.3. Notation

By variety, we mean a separated scheme of finite type over the field of complex numbers \mathbb{C} . By point, we mean a closed point. See [de-Mi 2009] for a quick introduction and for standard references concerning the constructible derived category and other concepts in this subsection.

Given a variety Y , we denote by $D_c^b(Y)$ the constructible bounded derived category of sheaves of rational vector spaces on Y , and by $DF(Y)$ its filtered variant [II 1971, Be-Be-De 1982]. We endow $D_c^b(Y)$ with the middle perversity t -structure. A functor that is exact with respect to this t -structure is said to be t -exact. The full subcategory of perverse sheaves is denoted by $P(Y)$. We employ the following standard notation for the objects associated with this t -structure: the full subcategories ${}^pD^{\leq j}(Y)$ and ${}^pD^{\geq j}(Y)$, $\forall j \in \mathbb{Z}$, and ${}^pD^{[j,k]}(Y) := {}^pD^{\geq j}(Y) \cap {}^pD^{\leq k}(Y)$, $\forall j \leq k \in \mathbb{Z}$, of $D_c^b(Y)$; the truncation functors ${}^p\tau_{\leq j} : D_c^b(Y) \rightarrow {}^pD^{\leq j}(Y)$ and ${}^p\tau_{\geq j} : D_c^b(Y) \rightarrow {}^pD^{\geq j}(Y)$; the perverse cohomology functors ${}^pH^j : D_c^b(Y) \rightarrow P(Y)$. We denote derived functors using the un-derived notation, e.g. if $f : X \rightarrow Y$ is a morphism of varieties, then the derived direct image (push-forward) functor $Rf_* : D_c^b(X) \rightarrow D_c^b(Y)$ is denoted by f_* , etc. Distinguished triangles in $D_c^b(Y)$ are denoted $G' \rightarrow G \rightarrow G'' \rightsquigarrow$. At times, we drop the space variable Y from the notation.

The following operations preserve constructibility of complexes: ordinary and extraordinary push-forward and pull-backs, hom and tensor product, Verdier duality, nearby and vanishing cycles.

When we write a Cartesian diagram of morphisms of varieties:

$$\begin{array}{ccc}
 X' & \xrightarrow{g} & X \\
 \downarrow f & & \downarrow f \\
 Y' & \xrightarrow{g} & Y,
 \end{array} \tag{1}$$

the ambiguities arising by having denoted different arrows by the same symbol, are automatically resolved by the context in which they are used; e.g. when we write the base change morphism of functors, the expression $g^* f_* \rightarrow f_* g^*$ is unambiguous.

The k th (hyper)cohomology groups of Y with coefficients in $G \in D_c^b(Y)$ are denoted by $H^k(Y, G)$. The complex computing this cohomology is denoted by $R\Gamma(Y, G)$ and it lives in the bounded derived category $D_c^b(pt)$ whose objects are complexes of vector spaces with cohomology given by finite dimensional rational vector spaces.

The filtrations we consider are finite and increasing. A sequence of morphism $G_\bullet := (\dots \rightarrow G_n \rightarrow G_{n+1} \rightarrow \dots)$ in $D_c^b(Y)$ subject to $G_i = 0 \forall i \ll 0$, and to $G_i \xrightarrow{\sim} G_{i+1} \forall i \gg 0$, gives rise to objects $(G, \mathcal{F}) \in DF(Y)$ and $(R\Gamma(Y, G), \mathcal{F}) \in D_c^b(pt)$. We call such a sequence of morphisms a system in $D_c^b(Y)$. There is the evident notion of morphism of systems $G_\bullet \rightarrow G'_\bullet$ and, for each such, the resulting morphisms $(G, \mathcal{F}) \rightarrow (G', \mathcal{F})$ in $DF(Y)$, and $(R\Gamma(Y, G), \mathcal{F}) \rightarrow (R\Gamma(Y, G'), \mathcal{F})$ in $DF(pt)$. Given $G \in D_c^b(Y)$ we have the system of perverse truncation morphisms: $\dots \rightarrow {}^p\tau_{\leq n} G \rightarrow {}^p\tau_{\leq n+1} G \rightarrow \dots \rightarrow G$. A morphism $G \rightarrow G'$ in $D_c^b(Y)$ gives rise to a morphism of systems ${}^p\tau_{\leq \bullet} G \rightarrow {}^p\tau_{\leq \bullet} G'$, which gives rise to a morphisms of filtered objects:

$$(R\Gamma(Y, G), P) \longrightarrow (R\Gamma(Y, G'), P) \quad \text{in } DF(pt). \tag{2}$$

The filtration P is called the perverse filtration. There is also the evident notion of systems of functors; e.g. the truncation functors.

Let $f : X \rightarrow Y$ be a morphism of varieties, let $F \in D_c^b(X)$ and let $G = f_*F$. Then we have the filtered objects $(R\Gamma(X, F), P) \neq (R\Gamma(X, F), P^f) := (R\Gamma(Y, f_*F), P)$. The filtration P^f is called the perverse Leray filtration associated with f . Given a morphism $F \rightarrow F'$ in $D_c^b(X)$, the analogue of (2) holds. We have the corresponding objects $P_j^f H^k(X, F)$, $\text{Gr}_j^{P^f} H^k(X, F)$.

If a statement is valid for every value of an index, e.g. the degree of a cohomology group, or the step of a filtration, then we denote such an index by a bullet-point symbol, or by a star symbol, e.g. $H^\bullet(X, F)$, P_\bullet , ${}^p\tau_{\leq \bullet}$, $P_\star H^\bullet(X, F)$, $\text{Gr}_\star^{P^f} H^\bullet(X, F)$.

We employ the following convention for shifts of filtered increasing filtrations:

$$\mathcal{F}(j)_\bullet := \mathcal{F}_{\bullet-j} \tag{3}$$

We have the evident and equivalent relations between truncations and shifts:

$$[\star] \circ {}^p\tau_{\leq \bullet} = {}^p\tau_{\leq \bullet-\star} \circ [\star], \quad {}^p\tau_{\leq \bullet} \circ [\star] = [\star] \circ {}^p\tau_{\leq \bullet+\star}, \tag{4}$$

which are valid for ${}^p\tau_{\geq \bullet}$ and ${}^pH^\bullet$ as well.

The category $P(Y)$ of perverse sheaves on Y is Abelian and Artinian, so that the Jordan–Holder theorem holds in it. The constituents of a non-zero perverse sheaf $G \in P(Y)$ are the isomorphism classes of the perverse sheaves appearing in the unique and finite collection of non-zero simple perverse sheaves appearing as the quotients in a Jordan–Holder filtration of G . The constituents of a non-zero complex $G \in D_c^b(Y)$ are defined to be the constituents of all of its non-zero perverse cohomology sheaves.

In general, we drop decorations (indices, parentheses, space variables, etc.) if it seems harmless in the context.

Given a morphism of varieties $X \rightarrow Y$ and a point $y \in Y$, we denote by X_y the fiber over y in X .

We are going to use the nearby/vanishing cycle functors. See Section 1.4 for the basic facts.

For what follows, namely the Decomposition Theorem, see: [Be-Be-De 1982], [Sa 1988], [de-Mi 2005], [Mo 2007, Sa 2005]. For a discussion, further references, and a generalization of [Mo 2007] to the proper case with rational coefficients see [de 2017].

THEOREM 1.3.1 (Decomposition and Relative Hard Lefschetz theorems). *Let $f : X \rightarrow Y$ be a proper morphism of varieties and let $F \in P(X)$ be semisimple. Then f_*F is semisimple.*

*If f is projective, then the Relative Hard Lefschetz holds: the choice of an f -ample line bundle induces isomorphisms ${}^pH^{-\bullet}(f_*F) \xrightarrow{\sim} {}^pH^\bullet(f_*(F))$.*

We define the intersection complex IC_T of a variety T as the direct sum of the intersection complexes of its irreducible components T_j : $IC_T := \bigoplus_j IC_{T_j}$. Then, as it is shown in [de 2012]: $IC_T \in P(T)$ is a semisimple perverse sheaf; it is the intermediate extension from the smooth locus of T of the direct sum of the constant sheaves on each component shifted by its dimension (see [Wu 2019] for a topological characterisation); the decomposition theorem, relative hard lefschetz theorem, and allied Hodge–theoretic facts hold for it and its

cohomology. The complex IC_T underlies a polarizable Hodge module. The simple objects in $P(Y)$ are the intersection complexes $IC_T(L)$, where $T \subseteq Y$ is closed and irreducible and L is an irreducible local systems on some Zariski dense open subset $T^o \subseteq T^{reg}$.

We define the topologist’s intersection complex as follows: $\mathcal{IC}_T := \bigoplus_j IC_{T_j}[\dim T_j]$.

We define the intersection cohomology groups by setting $IH^\bullet(T) := H^\bullet(T, IC_T)$; they start in degree $-\dim T$. We define the topologist’s intersection cohomology groups by setting $\mathcal{IH}^\bullet(T) := H^\bullet(T, \mathcal{IC}_T)$; they start in degree zero. E.g. for $T = o \amalg \Lambda$ the disjoint union of a point and a line, we have: $\mathbb{Q}_T = \mathbb{Q}_o \oplus \mathbb{Q}_\Lambda$; $IC_T = \mathbb{Q}_o \oplus \mathbb{Q}_\Lambda[1]$; $\mathcal{IC}_T = \mathbb{Q}_o \oplus \mathbb{Q}_\Lambda$; $\mathcal{IH}^\bullet(T) = H^\bullet(\mathbb{Q}_T) = H^\bullet(o) \oplus H^\bullet(\Lambda)$; $IH^\bullet(T) = H^\bullet(o) \oplus H^{\bullet+1}(T)$.

In order to avoid repeating naturality-type statements, we employ systematically the following notation. The symbol “=” denotes either equality, or a canonical isomorphism. The symbol $\xrightarrow{\sim}$ and denotes the fact that a canonical arrow is, in the context where it appears, an isomorphism. The symbol \cong denotes an isomorphism; if the direction of the arrow is relevant, we write $\xrightarrow{\sim}$. E.g.: the base change canonical isomorphism reads: $g^! f_* = f_* g^!$; the proper base change isomorphism for f proper reads: $g^* f_* \xrightarrow{\sim} f_* g^*$.

1.4. The vanishing and nearby cycles formalism

This section is an expanded version of [de-Ma 2018, Section 2.2]. Standard references for this section are [De 1972, XIII, XIV] and [Ka-Sc 1990, 8,10]. See also [II 1994] (cf. Remark 1.2.1).

Let $v = v_Y : Y \rightarrow S$ be a morphism of varieties, where S is a nonsingular curve. Let $s \in S$ be a point, usually called the special point.

We have the exact functors of triangulated categories:

$$i^*, i^!, \psi = \psi_v, \phi = \phi_v : D_c^b(Y) \longrightarrow D(Y_s), \tag{5}$$

where, $i : Y_s \rightarrow Y$ is the closed embedding of the (special) fiber of v over s , ϕ is the vanishing cycle functor and ψ is the nearby cycle functor. The functors ϕ and ψ depend on v ; e.g. [EL-Lê-Mi 2010, example 2.0.5]; this is not an issue in this paper.

We have the two, Verdier dual, canonical distinguished triangle of functors: (we often write σ instead of σ^* and $\sigma^!$)

$$i^*[-1] \xrightarrow{\sigma^*} \psi[-1] \xrightarrow{\text{can}} \phi \rightsquigarrow, \quad \phi \xrightarrow{\text{var}} \psi[-1] \xrightarrow{\sigma^!} i^![-1] \rightsquigarrow. \tag{6}$$

Fact 1.4.1 (t-exactness for $\psi[-1]$ and ϕ). The functors $\psi[-1]$ and ϕ are t-exact and commute with Verdier duality. We thus have the following canonical identifications:

$${}^p\tau_{\leq \bullet} \phi = \phi {}^p\tau_{\leq \bullet}; \quad {}^p\tau_{\leq \bullet} \psi[-1] = \psi[-1] {}^p\tau_{\leq \bullet}; \quad \text{ditto for } {}^p\tau_{\geq \bullet} \text{ and } {}^pH^\bullet. \tag{7}$$

The following is a key property of the vanishing cycle functor.

Fact 1.4.2 (Smooth morphisms and vanishing of ϕ). If $v : Y \rightarrow S$ is smooth over a neighborhood of s , and $G \in D_c^b(Y)$ has locally constant cohomology sheaves near s , then $\phi G = 0 \in D(Y_s)$. See [De 1972, XIII, 2.1.5]. In the special case where $v = id_S$, this implies that a complex $G \in D(S)$ has constant cohomology sheaves near s if and only if $\phi G = 0$.

We recall, for later use, the following simple

LEMMA 1.4.3 (**Vanishing of ϕ implies no constituents**). *Let $G \in D_c^b(Y)$. If $\phi G = 0$, then no constituent of G is supported on Y_s .*

Proof. See [de-Ma 2018, lemma 3.1.5]

Remark 1.4.4. The converse to Lemma 1.4.3 is false; see [Ka-Sc 1990, VIII.14] (Lefschetz degeneration to an ordinary double point).

Fact 1.4.5. The composition $i^*[-1] \rightarrow \psi[-1] \rightarrow i^! [1]$ yields a natural morphism of functors $D_c^b(Y) \rightarrow D(Y_s)$:

$$\nu : i^*[-1] \longrightarrow i^! [1]. \tag{8}$$

The morphism ν coincides with the morphism obtained via Verdier’s specialisation functor, (cf. [Sc 2003], for example), so that it depends only on the closed embedding $Y_s \rightarrow Y$, i.e. it is independent of v . If $\phi G = 0$, then $\sigma^*(G)$, $\sigma^!(G)$ and $\nu(G)$ are isomorphisms.

Fact 1.4.6. (**Base change diagrams for ψ and ϕ**). Let $f : X \rightarrow Y$ be an S -morphism and let $v_X = v_Y \circ f : X \rightarrow S$ be the structural morphism. The various base change natural transformation associated with i and f give rise to functorial morphisms of distinguished triangles as follows. See [De 1972, XIII, 2.1.7].

- (i) The one in $D(X_s)$ stemming from the morphism of functors $i^* f^! \rightarrow f^! i^*$, and from the isomorphism of functors $f^* i^* \xrightarrow{\cong} i^* f^*$, respectively: (we write ϕ_Y instead of ϕ_{v_Y} , etc.)

$$\begin{array}{ccc} i^*[-1]f^! \xrightarrow{\sigma \circ f^!} \psi_X[-1]f^! \longrightarrow \phi_X f^! \rightsquigarrow & & i^*[-1]f^* \xrightarrow{\sigma \circ f^*} \psi_X[-1]f^* \longrightarrow \phi_X f^* \rightsquigarrow \\ \downarrow & & \uparrow = \\ f^! i^*[-1] \xrightarrow{\sigma \circ f^!} f^! \psi_Y[-1] \longrightarrow f^! \phi_Y \rightsquigarrow, & & f^* i^*[-1] \xrightarrow{\sigma \circ f^!} f^* \psi_Y[-1] \longrightarrow f^* \phi_Y \rightsquigarrow. \end{array} \tag{9}$$

When f is étale, so that $f^* = f^!$, the two resulting morphisms of triangles are isomorphisms, inverse to each other.

- (ii) The one in $D(Y_s)$ stemming from the base change morphism of functors $i^* f_* \rightarrow f_* i^*$, and from the base change isomorphism $f_! i^* \xrightarrow{\cong} i^* f_!$:

$$\begin{array}{ccc} i^*[-1]f_* \xrightarrow{\sigma \circ f_*} \psi_Y[-1]f_* \longrightarrow \phi_Y f_* \rightsquigarrow & & i^*[-1]f_! \xrightarrow{\sigma \circ f_!} \psi_Y[-1]f_! \longrightarrow \phi_Y f_! \rightsquigarrow \\ \downarrow & & \uparrow = \\ f_* i^*[-1] \xrightarrow{f_* \circ \sigma} f_* \psi_X[-1] \longrightarrow f_* \phi_X \rightsquigarrow, & & f_! i^*[-1] \xrightarrow{f_! \circ \sigma} f_! \psi_X[-1] \longrightarrow f_! \phi_X \rightsquigarrow. \end{array} \tag{10}$$

When f is proper, so that $f_! \xrightarrow{\cong} f_*$, the two resulting morphisms of triangles are isomorphisms, inverse to each other.

(iii) We dualise the four diagrams associated in (9) and (10), and obtain the four analogous diagrams:

$$\begin{array}{ccc}
 \phi_X f^* \longrightarrow \psi_X[-1]f^* \xrightarrow{\sigma \circ f^*} i^! [1]f^* \rightsquigarrow \\
 \uparrow \qquad \qquad \qquad \uparrow \qquad \qquad \qquad \uparrow \\
 f^* \phi_Y \longrightarrow f^* \psi_Y[-1] \xrightarrow{f^* \circ \sigma} f^* i^! [1] \rightsquigarrow,
 \end{array}
 \qquad
 \begin{array}{ccc}
 \phi_X f^! \longrightarrow \psi_X[-1]f^! \xrightarrow{\sigma \circ f^!} i^! [1]f^! \rightsquigarrow \\
 \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow = \\
 f^! \phi_Y \longrightarrow f^! \psi_Y[-1] \xrightarrow{f^! \circ \sigma} f^! i^! [1] \rightsquigarrow,
 \end{array}
 \tag{11}$$

which are isomorphisms inverse to each other when f is étale, and

$$\begin{array}{ccc}
 \phi_Y f_! \longrightarrow \psi_Y[-1]f_! \xrightarrow{\sigma \circ f_!} i^! [1]f_! \rightsquigarrow \\
 \uparrow \qquad \qquad \qquad \uparrow \qquad \qquad \qquad \uparrow \\
 f_! \phi_X \longrightarrow f_! \psi_X[-1] \xrightarrow{f_! \circ \sigma} f_! i^! [1] \rightsquigarrow,
 \end{array}
 \qquad
 \begin{array}{ccc}
 \phi_Y f_* \longrightarrow \psi_Y[-1]f_* \xrightarrow{\sigma \circ f_*} i^! [1]f_* \rightsquigarrow \\
 \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \downarrow = \\
 f_* \phi_X \longrightarrow f_* \psi_X[-1] \xrightarrow{f_* \circ \sigma} f_* i^! [1] \rightsquigarrow,
 \end{array}
 \tag{12}$$

which are isomorphisms inverse to each other when f is proper.

(iv) by combining the l.h.s. square commutative diagram in (10) with the r.h.s. square commutative diagram in (12), we obtain the following commutative diagram of morphisms of functors:

$$\begin{array}{ccccc}
 i^*[-1]f_* & \longrightarrow & \psi[-1]f_* & \longrightarrow & i^! [1]f_* \\
 \downarrow & & \downarrow & & \downarrow = \\
 f_* i^*[-1] & \longrightarrow & f_* \psi[-1] & \longrightarrow & f_* i^! [1],
 \end{array}
 \tag{13}$$

where the compositions of the horizontal arrows are given by the corresponding morphisms (8), suitably pre/post-composed with f_* .

1.5. Nearby cycle functor and nearby points

Even if logically not necessary, it may be helpful to the intuition to clarify the use of the word “nearby.”

Let $t \in S$ be any point. By abuse of notation, denote by $t : t \rightarrow S$, $t : X_t \rightarrow X$ and $t : Y_t \rightarrow Y$ the resulting closed embeddings. There is the morphism (8) of functors $D(-) \rightarrow D(-_t)$:

$$t^*[-1] \longrightarrow t^! [1].
 \tag{14}$$

Let us start with the following:

Fact 1.5.1. For what follows, see [de-Ma 2018, fact 2.2.7]. Given a finite collection $G \in D_c^b(Y)$ of complexes, there is a Zariski-dense open subset $S^o(G) \subseteq S$ such that: the direct images $v_* {}^p\tau_{\leq \bullet} G$ have locally constant cohomology sheaves, and their formation commutes with arbitrary base change; the ${}^p\tau_{\leq \bullet} G$ have no constituent supported on any of the fibers of the morphisms v over $S^o(G)$; the strata on Y , of a stratification with respect to which G -hence the ${}^p\tau_{\leq \bullet} G$ - are constructible, are smooth over $S^o(G)$.

Definition 1.5.2. We say that $t \in S$ is general for G if $t \in S^o(G)$ as in fact 1.5.1.

Fact 1.5.3. Let $G \in D_c^b(Y)$. Let $t \in S^o(G)$ be a general point for G . Then:

- (i) The natural morphism $t^*[-1]G \rightarrow t^! [1]G$ is an isomorphism in $D(Y_t)$. See [de-Ma 2018, Fact 2.2.5].
- (ii) We have the identifications of [de-Ma 2018, fact 2.2.6]:

$$t^*[-1] {}^p\tau_{\leq \bullet} G = {}^p\tau_{\leq \bullet} t^*[-1]G, \quad t^! [1] {}^p\tau_{\leq \bullet} G = {}^p\tau_{\leq \bullet} t^! [1]G, \quad \text{ditto for } {}^p\tau_{\geq \bullet} \text{ and } {}^pH^\bullet. \tag{15}$$

Remark 1.5.4. In the special case where $v_Y : Y \rightarrow S$ is the identity on S , we have that $i^*, i^!, \psi, \phi : D(S) \rightarrow D(s)$ and that $t^*, t^! : D(S) \rightarrow D(t)$. In general, we have canonical identifications $D(s) = D(pt) = D(t)$, where pt is just a point, so that all three categories are naturally equivalent to the bounded derived category of finite dimensional rational vector spaces. Similarly, in the filtered case: $DF(s) = DF(pt) = DF(t)$. We use the catch-all notation $D_c^b(pt)$ and $DF(pt)$.

Given $G \in D_c^b(Y)$ and $s \in S$ our special point, we want to think: of the points $t \in S^o(G)$ as the points nearby s ; of the nearby vanishing cycle as capturing the complex G at points t nearby s . The following important fact makes this precise.

Fact 1.5.5. (Fundamental isomorphism). For what follows, see the fundamental identity [De 1972, XIV, 1.3.3.1]. Let $v : Y \rightarrow S$ be the identity, let $G \in D_c^b(Y)$, let $t \in S^o(G)$ be a general point for G . Choose a disk $\Delta \subseteq S$ centered at s and passing through t , such that $\Delta^* := \Delta \setminus \{s\} \subseteq S^o(G)$. Choose an universal covering of Δ^* and a point \tilde{t} on it over t . This choice of data gives rise to isomorphisms:

$$t^*[-1]G \xrightarrow{\sim} \psi[-1]G \xrightarrow{\sim} t^! [1]G, \quad \text{in } D(pt), \tag{16}$$

where the composition is (14), hence it is independent of the choices made above, but the indicated morphisms depend on the choice of data made above. Note that (16) is Verdier self-dual. Changing the choice of data changes the two individual arrows by the monodromy automorphism induced by an appropriate element in $\pi_1(S^o(G), t)$.

Let $v : Y \rightarrow S$ be any morphism and let $G \in D_c^b(Y)$. Then:

- (i) by taking $v_*G \in D(S)$ in (16), we get collections of isomorphisms:

$$\psi v_*G \xrightarrow{\sim} t^* v_*G = v_* t^*G = R\Gamma(Y_t, t^*G), \quad \text{in } D(pt), \tag{17}$$

where any two such identifications $\xrightarrow{\sim}$ differ by the action induced by an element of $\pi_1(S^o(G), t)$ on the r.h.s. The second $=$ sign is clear. The first can be seen by using the base change properties relative to the general point t expounded in Section 1.4 as follows. The choice of t general for G , made in Definition 1.5.2, allows us to: replace ψ with t^* (cf. (16)); use the identification $t^* v_* = v_* t^*$. One cannot select a point $t \in S$ that is general for every $G \in D_c^b(Y)$. Because of this it is preferable to work with $\psi : D(S) \rightarrow D(s)$ instead of with a general point $t \in S$. When working with finitely many complexes in $D_c^b(Y)$, or with a family of complexes constructible with respect to an arbitrary fixed stratification, we can always choose a point $t \in S$ that is general for all of them,

- (ii) by taking $v_* {}^p\tau_{\leq \bullet} G \in D(S)$ and $v_* {}^p\tau_{\leq \bullet} f_* F \in D(S)$ in (16), we reach conclusions analogous to part (1), namely, we obtain isomorphisms:

$$(\psi v_* G, P) \xrightarrow{\sim} (R\Gamma(Y_t, t^*G), P), \quad \text{in } DF(pt); \tag{18}$$

$$(\psi v_* f_* F, P) \xrightarrow{\sim} (R\Gamma(X_t, t^*F), P), \quad \text{in } DF(pt). \tag{19}$$

2. The morphisms of type δ

This section is a necessary technical interlude on the way to Section 3, where we study the behavior of the perverse filtration relative to the specialisation morphism.

The distinguished triangles (6) and the t-exactness of the functors ϕ and $\psi[-1]$ suggest the need to quantify the failure of t-exactness for the functors $i^*[-1]$ and $i^![-1]$.

In this section we quantify this failure by introducing the morphisms of functors of type δ (27) which play an important role starting with Section 3, diagram (49). When the morphisms of type δ are evaluated against a complex $G \in D_c^b(Y)$, they are isomorphisms if and only if perverse truncations and perverse cohomology commute with the appropriately shifted pull-backs.

The special fiber Y_s in Y is an effective Cartier divisor. We introduce the morphisms of type δ in the more general context of embedded effective Cartier divisors in (22). For easier bookkeeping, we use the shifted version (24).

Proposition 2.1.5 is a criterion, in the general context of Cartier divisors (i.e. when we are not necessarily working over a curve S), for the morphisms of type δ to be isomorphisms, hence it is a criterion for the aforementioned commutativity of perverse truncations/cohomology with shifted pullbacks. It states that if a complex has no constituents supported on an effective Cartier divisor, then the resulting morphisms of type δ are isomorphisms.

Lemma 2.2.1 yields, in the context of the vanishing/nearby cycle functors (here we are working over a curve S , and the Cartier divisor in question is the fiber over the special point s) the important diagram (28) of morphisms of distinguished triangles of functors involving the morphisms of type δ needed in this paper. In particular, Lemma 2.2.1.(3) combines Lemma 1.4.3 (if $\phi G = 0$, then G has no constituents on Y_s) with Proposition 2.1.5 (“if no constituents, then the δ ’s are isomorphisms”).

Proposition 2.3.2 is another general (i.e not necessarily related to a situation over a curve) criterion of a different flavor for the δ ’s to be isomorphisms in the context of a complex $G = f_* F$ being a direct image under a projective morphism f . It replaces the assumptions on constituents and/or on the vanishing of the vanishing cycle functor, with the assumptions that both F and $i^* F[-1]$ are semisimple; the proof uses the Relative Hard Lefschetz Theorem.

2.1. Pull-back to Cartier divisors and truncations: the morphisms δ

In the context of the vanishing cycle functor, i.e. $v : Y \rightarrow S$ and $s \in S$, the special fiber Y_s is a Cartier divisor in the total space Y . The vanishing cycle functor ϕ and the shifted nearby cycle functor $\psi[-1]$ are t-exact. It is important to study the defect of t-exactness of the shifted restriction to the special fiber functors $i^*[-1]$ and $i^![-1]$.

In this section, which amplifies [de-Ma 2018, section 3.1] to meet the needs of this paper, we introduce, in the general context of the closed embedding of an effective Cartier divisor, morphisms of functors that measure the aforementioned defect. We define the morphisms

of type δ in the context of embeddings of Cartier divisors in Lemma 2.1.2, and we prove a criterion for when these morphisms δ are isomorphisms in Proposition 2.1.5.

Let us emphasise that in this section we work with varieties and not with S -varieties.

We start by listing the needed general t-exactness properties related to embeddings of effective Cartier divisors on varieties.

LEMMA 2.1.1 (Inequalities for embeddings of Cartier divisors). *Let $\iota : T' \rightarrow T$ be a closed embedding of varieties such that the open embedding $T \setminus T'$ is an affine morphism (e.g. T' is an effective Weil divisor supporting an effective Cartier divisor). Then: (we omit the space variables)*

(i) *the functor ι^* is right t-exact and the functor $\iota^!$ is left t-exact:*

$$\iota^* : {}^pD^{\leq \bullet} \longrightarrow {}^pD^{\leq \bullet}, \quad \iota^! : {}^pD^{\geq \bullet} \longrightarrow {}^pD^{\geq \bullet}. \tag{20}$$

(ii) *the functor $\iota^*[-1]$ is left t-exact and the functor $\iota^![1]$ is right t-exact:*

$$\iota^* : {}^pD^{\geq \bullet} \longrightarrow {}^pD^{\geq \bullet-1}, \quad \iota^! : {}^pD^{\leq \bullet} \longrightarrow {}^pD^{\leq \bullet+1}. \tag{21}$$

In particular, we have that: $\iota^*P(T) \subseteq {}^pD^{[-1,0]}(T')$ and $\iota^!P(T) \subseteq {}^pD^{[0,1]}(T')$.

Proof. See [de-Ma 2018, lemma 3.1.1].

The following lemma extends [de-Ma 2018, Lemma 3.1.2] in the direction needed in this paper.

We denote by $\gamma_{\leq \bullet} : {}^p\tau_{\leq \bullet-1} \rightarrow {}^p\tau_{\leq \bullet}$ and $\gamma_{\geq \bullet} : {}^p\tau_{\leq \bullet} \rightarrow {}^p\tau_{\leq \bullet+1}$ the natural morphisms. They are Verdier dual to each other. We set:

$$\gamma_{\leq \bullet}^* := \gamma_{\leq \bullet} \iota^*, \quad \gamma_{\leq \bullet}^! := \gamma_{\leq \bullet} \iota^!, \quad \gamma_{\geq \bullet}^* := \gamma_{\geq \bullet} \iota^*, \quad \gamma_{\geq \bullet}^! := \gamma_{\geq \bullet} \iota^!.$$

The entries of each of the two pairs $(\gamma_{\leq \bullet}^*, \gamma_{\geq \bullet}^!)$ and $(\gamma_{\geq \bullet}^*, \gamma_{\leq \bullet}^!)$ are exchanged by Verdier duality.

LEMMA 2.1.2 (The morphisms δ). *There are natural morphisms of distinguished triangles of functors:*

which are exchanged by Verdier duality.

By iteration, the morphisms in (22) induce natural morphisms:

$$\delta_{\bullet}^* : {}^pH^{\bullet-1}[-\bullet+1] \iota^* \longrightarrow \iota^* {}^pH^{\bullet}[-\bullet], \quad \delta_{\bullet}^! : \iota^! {}^pH^{\bullet}[-\bullet] \longrightarrow {}^pH^{\bullet+1}[-\bullet-1] \iota^!, \tag{23}$$

which are exchanged by Verdier duality.

Proof. We prove the lemma for the l.h.s. of (22). The r.h.s. follows by Verdier duality.

Consider the l.h.s. diagram of (22), but with the arrows of type δ and ϵ removed. By applying [Be-Be-De 1982, Propagation 1.1.9, p.23], whose hypotheses are met by virtue of the inequalities (20) and (21), we see that we can fill in the diagram and make it commutative with unique arrows. The compositions $\epsilon\delta$ of the vertical arrows give the desired arrows of type γ because in any t -structure the structural morphism ${}^p\tau_{\leq -1} \rightarrow \text{Id}$ factors uniquely through the structural morphism ${}^p\tau_{\leq \bullet} \rightarrow \text{Id}$ via $\gamma_{\leq \bullet} : {}^p\tau_{\leq -1} \rightarrow {}^p\tau_{\leq \bullet}$.

The morphism on the l.h.s. of (23) arises by composing truncation applied to δ^* , with δ^* applied to the truncation:

$${}^p\tau_{\geq -1} {}^p\tau_{\leq -1} l^* \xrightarrow{{}^p\tau_{\geq -1} \delta_{\leq \bullet}^*} {}^p\tau_{\geq -1} l^* {}^p\tau_{\leq \bullet} \xrightarrow{\delta_{\geq -1}^* {}^p\tau_{\leq \bullet}} {}^p\tau_{\geq -1} l^* {}^p\tau_{\leq \bullet}.$$

The morphism on the r.h.s. is obtained in a similar fashion and the verification that the two arrows in (23) are Verdier dual is left to the reader.

Remark 2.1.3.

- (i) By virtue of the identity (4) concerning truncations and shifts, the morphisms of type δ in (22) and (23) may be re-written as follows:

$$\delta_{\leq \bullet}^* : {}^p\tau_{\leq \bullet} l^*[-1] \longrightarrow l^*[-1] {}^p\tau_{\leq \bullet}, \quad \delta_{\leq \bullet}^! : l^! [1] {}^p\tau_{\leq \bullet} \longrightarrow {}^p\tau_{\leq \bullet} l^! [1], \quad (24)$$

$$\delta_{\geq \bullet}^* : {}^p\tau_{\geq \bullet} l^*[-1] \longrightarrow l^*[-1] {}^p\tau_{\geq \bullet}, \quad \delta_{\geq \bullet}^! : l^! [1] {}^p\tau_{\geq \bullet} \longrightarrow {}^p\tau_{\geq \bullet} l^! [1],$$

$$\delta_{\bullet}^* : {}^pH^{\bullet-1} l^* \longrightarrow l^*[-1] {}^pH^{\bullet}, \quad \delta_{\bullet}^! : l^! [1] {}^pH^{\bullet} \longrightarrow {}^pH^{\bullet+1} l^!.$$

where Verdier duality exchanges the morphisms along the two diagonals of the first two rows, and exchanges the two terms in the third row.

- (ii) By looking at (22), and by virtue of Verdier duality, we deduce that for a given $G \in D(T)$ we have that: $\delta_{\leq \bullet}^*(G)$ is an isomorphism if and only if $\delta_{\leq \bullet}^*(G)$ is an isomorphism if and only if $\delta_{\leq \bullet}^!(G^\vee)$ is an isomorphism if and only if $\delta_{\leq \bullet}^!(G^\vee)$ is an isomorphism.
- (iii) The same example in Remark 2.2.2.(2) shows that for a given $G \in D(T)$, having $\delta_{\leq \bullet}^*(G)$ an isomorphism does not imply that $\delta_{\leq \bullet}^!(G)$ is an isomorphism.
- (iv) Proposition 2.1.5 gives a criterion for all six morphisms of type δ to be isomorphisms. When the morphisms (24) are isomorphisms, we may say that in that case “perverse truncation/cohomology commutes up to a suitable shift with restriction to the Cartier divisor”. While standard truncation commutes with such an un-shifted restriction -in fact with any pull-back-, in general perverse truncation does not.

Remark 2.1.4. In view of the l.h.s. inequality in (21), the r.h.s. vertex of the distinguished triangle in the middle row of the l.h.s. of (22) satisfies the inequality $l^* {}^p\tau_{\geq \bullet+1} : D \rightarrow {}^pD^{\geq \bullet}$. By taking the long exact sequence of perverse cohomology sheaves of said distinguished triangle, the aforementioned inequality yields the natural isomorphism of functors:

$${}^p\tau_{\leq -1} l^* {}^p\tau_{\leq \bullet} \xrightarrow{\simeq} {}^p\tau_{\leq -1} l^*. \quad (25)$$

PROPOSITION 2.1.5 (If no constituents, then the δ 's are isomorphisms). *Let $\iota : T' \rightarrow T$ be as in Lemma 2.1.1. If $G \in D(T)$ has no constituent supported on T' , then the morphisms of type δ in (24) are isomorphisms.*

Proof. This is [de-Ma 2018, proposition 3.1.4]. For the reader's convenience, we include it here.

We prove the conclusion for $\delta_{\leq \bullet}^*(G)$. We have that ${}^p\tau_{\leq \bullet} G \in {}^pD^{\leq \bullet}$, so that, by (20), we have that $\iota^* {}^p\tau_{\leq \bullet} G \in {}^pD^{\leq \bullet}$, and then, clearly, we have that:

$$\iota^* {}^p\tau_{\leq \bullet} G = {}^p\tau_{\leq \bullet} \iota^* {}^p\tau_{\leq \bullet} G. \tag{26}$$

In view of (26) and of (25), and by considering the truncation triangle ${}^p\tau_{\leq \bullet-1} \rightarrow {}^p\tau_{\leq \bullet} \rightarrow {}^pH^\bullet[-\bullet] \rightsquigarrow$ applied to $\iota^* {}^p\tau_{\leq \bullet} G$, in order to prove the desired conclusion, it is necessary and sufficient to show that ${}^pH^\bullet(\iota^* {}^p\tau_{\leq \bullet} G) = 0$.

This can be argued as follows. By taking the long exact sequence of perverse cohomology of the distinguished triangle $\iota^* {}^p\tau_{\leq \bullet-1} G \rightarrow \iota^* {}^p\tau_{\leq \bullet} G \rightarrow \iota^* {}^pH^\bullet G[-\bullet] \rightsquigarrow$, we see that it is necessary and sufficient to show that $\iota^* {}^pH^\bullet G[-\bullet] \in {}^pD^{\leq \bullet-1}$, or, equivalently, that $\iota^* {}^pH^\bullet G \in {}^pD^{\leq -1}$.

By [Be-Be-De 1982, 4.1.10.ii], we have the distinguished triangle ${}^pH^{-1}(\iota^* {}^pH^\bullet G)[1] \rightarrow \iota^* {}^pH^\bullet G \rightarrow {}^pH^0(\iota^* {}^pH^\bullet G) \rightsquigarrow$ and an epimorphism ${}^pH^\bullet G \rightarrow {}^pH^0(\iota^* {}^pH^\bullet G)$. Since G is assumed to not have constituents supported at Y_s , we must have ${}^pH^0(\iota^* {}^pH^\bullet G) = 0$, so that $\iota^* {}^pH^\bullet G[-\bullet] \in {}^pD^{\leq \bullet-1}$, as requested.

We have proved that if G has no constituents supported on T' , then $\delta_{\leq \bullet}^*(G)$ is an isomorphism. By Remark 2.1.3, we have that $\delta_{\geq \bullet}^*(G)$ is an isomorphism as well.

Note that G has no constituents supported on T' if and only if the same is true for ${}^p\tau_{\leq \bullet} G$. We thus have that $\delta_{\leq \bullet}^*({}^p\tau_{\leq \bullet} G)$ and $\delta_{\geq \bullet}^*({}^p\tau_{\leq \bullet} G)$ are isomorphisms.

Now, $\delta_{\geq \bullet}^*$ is the composition of two isomorphisms, namely ${}^p\tau_{\geq \bullet}(\delta_{\leq \bullet}^*(G))$, followed by $\delta_{\geq \bullet}^*({}^p\tau_{\leq \bullet} G)$.

We have proved that the morphisms of type δ^* are isomorphisms.

Note that G has no constituents supported on T' if and only if the same is true for G^\vee . By Remark 2.1.3.(2), it follows that the morphisms of type δ^* for G^\vee are isomorphisms, so that so are the morphisms of type $\delta^!$ for G .

The proposition is proved.

Alternatively, one could also retrace the proof given in detail above for the morphisms of type δ^* and give a proof for the morphisms of type $\delta^!$.


Proposition 2.1.5 can be informally summarised as follows: under special circumstances, the perverse truncation and restriction to a “non-special” Cartier divisor commute with each other. One can also say that we can slide one functor across the other.

2.2. The morphisms δ and vanishing/nearby cycles

Lemma 2.1.2 is about a Cartier divisor on a variety. Let us specialize to the case when the Cartier divisor is the special fiber in the context of the vanishing and nearby cycle functors as in Section 1.4. In particular, we work over a nonsingular connect curve S and we fix a point $s \in S$.

LEMMA 2.2.1. (Morphisms of type δ for ψ and ϕ).

(i) We have natural morphisms of distinguished triangles:



where each diagram is self-dual.

(ii) If we denote by $i^*[-1]$ and $i^![-1]$ the two diagrams (22) (after having replaced i^* with $i^*[-1]$ and $i^!$ with $i^![-1]$; see Remark 2.1.3.(1)), and if we denote by $\psi[-1]$ and by ϕ the two diagrams in (27), then there are morphism of diagrams $i^*[-1] \rightarrow \psi[-1] \rightarrow i^![-1]$, in the sense that all the corresponding square diagrams are commutative. In particular, we have the commutative diagram:

$$\begin{array}{ccccc}
 i^*[-1]^{p\tau_{\leq \bullet}} & \longrightarrow & \psi[-1]^{p\tau_{\leq \bullet}} & \longrightarrow & i^![-1]^{p\tau_{\leq \bullet}} & (28) \\
 \delta_{\leq \bullet}^* \uparrow & & \simeq \delta_{\leq \bullet}^{\psi} & & \delta_{\leq \bullet}^! \downarrow & \\
 p\tau_{\leq \bullet} i^*[-1] & \longrightarrow & p\tau_{\leq \bullet} \psi[-1] & \longrightarrow & p\tau_{\leq \bullet} i^![-1] &
 \end{array}$$

Similarly, for $p\tau_{\geq \bullet}$ and ${}^pH^\bullet$. The resulting diagrams are exchanged -in the case of truncations- or preserved -in the case of perverse cohomology- by Verdier Duality.

- (iii) Let $G \in D_c^b(Y)$ be such that $\phi G = 0$. Then all the vertical arrows in (28), and in its variants involving $p\tau_{\geq \bullet}$ and ${}^pH^\bullet$, evaluated at G , are isomorphisms. In particular, all the arrows of type δ evaluated at G in (24) are isomorphisms.
- (iv) Let $G \in D_c^b(Y)$ be such that G has no constituents supported on Y_s . Then all the vertical arrows in (28), and in its variants involving $p\tau_{\geq \bullet}$ and ${}^pH^\bullet$, evaluated at G are isomorphisms. In particular, all the arrows of type δ evaluated at G in (24) are isomorphisms.

Proof. (i) The proof that there are the natural diagrams (27) is the same as the one of Lemma 2.1.2. The fact that the arrows of type δ^ψ and δ^ϕ in (27) are isomorphisms simply translates the t-exactness of $\psi[-1]$ and ϕ .

(ii) Once we note that the dual to i^* is $i^!$, and that $\psi[-1]$ and ϕ are self-dual, hence the self-duality of (27), there is nothing left to prove.

(iii) That the horizontal arrows in (28) are isomorphisms is an immediate consequence of the distinguished triangles (6) and the hypothesis $\phi G = 0$. Once the horizontal arrows are isomorphisms, since one of the vertical arrows is an isomorphism, so are the remaining vertical ones.

(iv) Follows from Proposition 2.1.5.

Remark 2.2.2.

- (i) We have the following two implications: (a) if $\phi(G) = 0$, then G has no constituents on the special fiber (Lemma 1.4.3); (b) if G has no constituents on the special fiber,

then the morphisms of type δ in (28) are isomorphisms. These two implications are combined in Lemma 2.2.1.(3), which is an amplified version of [de-Ma 2018, proposition 3.1.5].

- (ii) Both the implication (a) and (b) directly above, as well the implication in Lemma 2.2.1.(3), cannot be reversed. For a), see the counterexample in Remark 1.4.4. For 2), as well as for Lemma 2.2.1.(3), consider the following counterexample: take ι to be the embedding of the origin s into a disk Δ , and G to be $j_! \mathbb{Q}[1] \in P(\Delta)$. Then G admits \mathbb{Q}_s as a constituent and $\phi G = 0$. On the other hand $i^* j_! = 0$, so that the morphisms of type δ^* are trivially isomorphisms of zero objects.
- (iii) The fact that G has no constituents supported on Y_s does not prevent the following phenomenon: the number of constituents of $\iota^* G$ may be strictly smaller than the number of constituents of $i^* G$. Consider a family $X \rightarrow Y \rightarrow S$, where X/S is a family of K3 surfaces fibered onto curves, where the general member has irreducible fibers, while a special member has some reducible fibers. Then $G := f_* \mathbb{Q}$ will exhibit this kind of behaviour. I thank Giulia Saccà, Antonio Rapagnetta and Junliang Shen for helping me sort this out.

2.3. Relative hard Lefschetz and morphisms of type δ

The aim of this section is to prove Proposition 2.3.2 which is a slight generalization of [Mi-Sh 2018, corollary 3.8], and yields another criterion, this time involving the Relative Hard Lefschetz Theorem (RHL), for the morphisms of type δ to be isomorphisms. Proposition 2.3.2 is used in the proofs of Theorems 3.3.7 and 3.5.4.

Proposition 2.3.2.(ii) is the special case of Proposition 2.3.2.(i) when the Cartier divisor is the special fiber in the context of the vanishing/nearby cycle functors. We thus start with a variety X and a complex $F \in P(X)$ perverse and semisimple such that it stays perverse and semisimple after restriction and shift by [1] to the pre-image X' on X of a Cartier divisor Y' on Y . This condition on the restriction should be thought as some kind of weakened transversality condition of the divisor Y' on Y , with respect to the morphism, and to the semisimple perverse sheaf F .

Remark 2.3.1. The hypotheses of the upcoming Proposition 2.3.2 are met, for example, when the morphism f is projective, the varieties X and X' are irreducible orbifolds and $F = \mathbb{Q}_X[\dim X]$. Note also that in this section, we do not assume that X, Y , etc., are varieties over S .

PROPOSITION 2.3.2. (RHL criterion for the morphisms δ being isomorphisms).

- (i) Let $f : X \rightarrow Y$ be a projective morphism of varieties. Let $\iota : Y' \rightarrow Y$ be a closed embedding of pure codimension one such that the resulting open embedding $Y \setminus Y' \rightarrow Y$ is affine and let:

$$\begin{array}{ccc}
 X' & \xrightarrow{\iota} & X \\
 \downarrow f & & \downarrow f \\
 Y' & \xrightarrow{\iota} & Y
 \end{array} \tag{29}$$

be the resulting Cartesian diagram. Let $F \in D_c^b(X)$ be perverse semisimple and such that $\iota^! F[1] \in D(X')$ (resp. $\iota^* F[-1]$) is perverse semisimple.

Then we have the following identities:

$${}^p\tau_{\leq \bullet} (f_* \iota^! F[1]) = (\iota^! {}^p\tau_{\leq \bullet} f_* F) [1], \quad (\text{resp. } {}^p\tau_{\leq \bullet} (f_* \iota^* F[-1]) = (\iota^* {}^p\tau_{\leq \bullet} f_* F) [-1]), \tag{30}$$

and similarly if ${}^p\tau_{\leq \bullet}$ is replaced by ${}^p\tau_{\geq \bullet}$, or by ${}^pH^\bullet$.

The complex $f_* F$ has no constituent supported on Y' .

The morphisms of type δ in (24) associated with $f_* F$ are isomorphisms.

(ii) Assume, in addition, that f is an S -morphism of S -varieties, where S is a nonsingular curve, that $s \in S$ is a point and that $Y' = Y_s$.

Then the morphisms of type δ in (24), (27) and (28) associated with $f_* F$ are isomorphisms.

The complex $f_* F$ has no constituent supported on the special fiber Y_s .

Proof. Since part (ii) is a special case of part (i), it is enough to prove part (i). We do so in the case of $\iota^! F[1]$; the case of $\iota^* F[-1]$ can be proved in the same way.

We prove the identities (30) for $\iota^!$. By the Decomposition and Relative Hard Lefschetz Theorems 1.3.1, we have isomorphisms:

$$f_* F \xrightarrow{\cong} \bigoplus_{\bullet} {}^pH^\bullet(f_* F)[- \bullet], \quad f_* \iota^! F[1] \xrightarrow{\cong} \bigoplus_{\bullet} {}^pH^\bullet(f_* \iota^! F[1])[- \bullet], \tag{31}$$

$${}^pH^{-\bullet} f_* F \cong {}^pH^\bullet f_* F, \quad {}^pH^{-\bullet} (f_* \iota^! F[1]) \cong {}^pH^\bullet (f_* \iota^! F[1]). \tag{32}$$

By the base change identity $f_* \iota^! F[1] = \iota^! f_* F[1]$, and by the splitting assumptions (31), we get an isomorphism:

$$\bigoplus_k \iota^! ({}^pH^k (f_* F_X) [1]) [-k] \cong \bigoplus_k ({}^pH^k (f_* \iota^! F[1])) [-k]. \tag{33}$$

At this juncture, it does not seem a priori clear that the desired conclusion follows by taking cohomology sheaves on both sides, the reason being that the unshifted summands on the lhs are not immediately seen to be perverse. What it is a priori clear is that, due to the shift [1], they live in ${}^pD^{[-1,0]}(Y')$ (cf. [BBD, 4.1.10.ii, p.106]).

Claim. The complex $\iota^! {}^pH^k (f_* F_X)[1]$ is a perverse sheaf, $\forall k$.

Proof of the claim. We need to show that ${}^pH^{-1}(\iota^! {}^pH^k (f_* F_X)[1]) = 0, \forall k$. To simplify the notation, we set:

$$P_{X'}^k := {}^pH^k (f_* \iota^! F[1]), \quad Q_X^k := \iota^! {}^pH^k (f_* F_X)[1] \in {}^pD^{[-1,0]}(W), \quad Q_X^{k,l} = {}^pH^l (Q_X^k), \quad l = -1, 0. \tag{34}$$

We need to show that $Q_X^{k,-1} = 0, \forall k$.

We know that ${}^pH^{-k} \cong {}^pH^k, \forall k$, both for $f_* F$ and for $f_* \iota^! F[1]$, so that we have that:

$$P_{X'}^{-k} = P_{X'}^k, \quad Q_X^{-k,l} = Q_X^{k,l}, \quad \forall k, \quad \forall l = -1, 0. \tag{35}$$

We argue by contradiction and assume that the desired conclusion fails for some k , i.e. that there is k such that $Q_X^{k,-1} \neq 0$.

By taking perverse cohomology in (33), we get an isomorphism

$$P_{X'}^k \cong Q_X^{k,0} \oplus Q_X^{k+1,-1}. \tag{36}$$

Let $-k_0$ be the smallest index $-k$ such that $Q^{-k,-1} \neq 0$. By symmetry, $k_0 \geq 0$. We thus have $P_{X'}^{-k_0-1} = Q_X^{-k_0-1,0} \oplus Q_X^{-k_0,-1} \neq Q_X^{-k_0-1,0}$. By symmetry, $P_{X'}^{k_0+1} \neq Q_X^{k_0+1,0}$.

On the other hand, we have that $P_{X'}^{k_0+1} = Q_X^{k_0+1,0} \oplus Q_X^{k_0+2,-1} = Q_X^{k_0+1,0}$, because $Q_X^{k_0+2,-1} = Q_X^{-k_0-2,-1} = 0$, by the minimality property of $-k_0$.

This contradiction establishes the claim.

At this point, (33), joint with the claim, implies, by taking perverse cohomology sheaves, that (30) holds with “=” replaced by an “ \cong ”. In order to establish (30) with = (canonical isomorphism), we simply observe that now that we know that the lhs of (33) is a direct sum of shifted perverse sheaves, we reach the desired conclusion by taking perverse cohomology in the isomorphism $\iota^! f_* F[1] = f_* \iota^! F[1]$.

We have thus proved identities (30). At this point, the two final assertions of (i) can be seen to be equivalent; we show that they hold as follows.

The assertion on the lack of constituents supported on the Cartier divisor Y' follows from the fact that any such constituent would have to show up as a direct summand supported on Y' and, as such, its $\iota^! = \iota^*$ would show up as a non-zero $Q^{k,-1}$ in the proof above, a contradiction.

The last assertion of part (i) follows from Proposition 2.1.5.

Remark 2.3.3. As the proof shows, one may replace the assumption that $\iota^! F[1]$ is perverse semisimple, with the assumption that $f_* \iota^! F[1]$ satisfies the conclusion of the RHL. Similarly, if we replace $\iota^! F[1]$ with $\iota^* F[-1]$. This assumption is met if the boundary is a simple normal crossing divisor and we take F to be the intersection complex, which, by the very definition of snc divisor, is necessarily the shifted constant sheaf near the divisor; see Corollary 2.4.2. See also the proof of part II of Theorem 3.5.4, and Remark 4.4.3.

2.4. RHL and boundary with normal crossing divisors

Let us place ourselves in the situation of Proposition 2.3.2, except that we do not assume that $\iota^* F[-1]$ is perverse semisimple on the Cartier divisor on X' . The goal of this section is to prove Lemma 2.4.1, to the effect that if we are in a simple normal crossing divisors situation on a nonsingular variety X of dimension n , so that $\iota^* \mathbb{Q}_X[n-1]$ is not perverse semisimple on X' , then we still have the useful RHL symmetry (cf. Remark 2.3.3). For an application, with details left to the reader, see Remark 4.4.3.

Let X be an irreducible variety, let n be its dimension, and let $Z \subseteq X$ be a simple normal crossing divisor; in particular, X is nonsingular near Z . Let \mathcal{I} be a finite set indexing the irreducible components Z_i of Z , let $I = \{i_1, \dots, i_k\} \subseteq \mathcal{I}$ be any subset, let $Z_I := \bigcap_{i \in I} Z_i$. We have the standard long exact sequence of sheaves on X :

$$0 \longrightarrow \mathbb{Q}_Z \longrightarrow \bigoplus_{|I|=1} \mathbb{Q}_{Z_I} \longrightarrow \dots \longrightarrow \bigoplus_{|I|=n} \mathbb{Q}_{Z_I} \longrightarrow 0. \tag{37}$$

By splicing (37), and by shifting in an appropriate manner, we get the system of short exact sequences of perverse sheaves in $P(X)$:

$$0 \longrightarrow 0 \longrightarrow K^n[0] \xrightarrow{\cong} \bigoplus_{|I|=n+1} \mathbb{Q}_{Z_I}[0] \longrightarrow 0, \tag{38}$$

$$0 \longrightarrow K^n[0] \longrightarrow K^{n-1}[1] \longrightarrow \bigoplus_{|I|=n} \mathbb{Q}_{Z_I}[1] \longrightarrow 0,$$

...

$$0 \longrightarrow K^2[d_Z - 2] \longrightarrow K^1[d_Z - 1] \longrightarrow \bigoplus_{|I|=2} \mathbb{Q}_{Z_I}[d_Z - 1] \longrightarrow 0,$$

$$0 \longrightarrow K^1[d_Z - 1] \longrightarrow \mathbb{Q}_Z[d_Z] \xrightarrow{\cong} \bigoplus_{|I|=1} \mathbb{Q}_{Z_I}[d_Z] \longrightarrow 0.$$

Note that $\mathbb{Q}_Z[d_Z]$ is not semisimple in general, even when X is nonsingular. Nevertheless, we have:

LEMMA 2.4.1. *Let $f : X \rightarrow Y$ be a proper morphism with X nonsingular and irreducible. Then $f_*\mathbb{Q}_Z[d_Z]$ has the RHL symmetry, i.e. cupping with the first Chern class L of an f -ample line bundle induces isomorphisms*

$$L^\bullet : {}^pH^{-\bullet} f_*\mathbb{Q}_Z[d_Z] \xrightarrow{\cong} {}^pH^\bullet f_*\mathbb{Q}_Z[d_Z], \tag{39}$$

so that the complex $f_*\mathbb{Q}_Z[d_Z]$ is t-split (splits as the direct sum of its shifted perverse cohomology sheaves). Moreover, we have that:

$${}^pH^\bullet (i^* {}^pH^\bullet (f_*\mathbb{Q}_Z[d_Z])) = 0, \quad \forall \bullet \neq -1, \quad \forall \star. \tag{40}$$

Proof. For the RHL symmetry: We use descending induction on n : start from the bottom of (38); go up one step at the time; at each step use the five lemma applied to:

$$\begin{array}{ccccccc} 0 & \longrightarrow & {}^pH^{-\bullet} A & \longrightarrow & {}^pH^{-\bullet} B & \longrightarrow & {}^pH^{-\bullet} C \longrightarrow 0 \\ & & \downarrow L^\bullet & & \downarrow L^\bullet & & \downarrow L^\bullet \\ 0 & \longrightarrow & {}^pH^\bullet A & \longrightarrow & {}^pH^\bullet B & \longrightarrow & {}^pH^\bullet C \longrightarrow 0. \end{array} \tag{41}$$

For the splitting: use the Deligne–Lefschetz Criterion.

For the proof of the vanishing (40), use the same kind of descending induction.

COROLLARY 2.4.2. *Let $f : X \rightarrow Y$ be a projective morphism of varieties. Consider Cartesian diagram like (29) and assume that X' is supported on a simple normal crossing divisor. Let $F = IC_{X'}$.*

Then the conclusions of Proposition 2.3.2 hold.

Proof. We simply observe that X is assumed to be nonsingular near X' , so that, near X' , $IC_{X'}$ is the constant sheaf, up to shifts. The shifts are locally constant integers, and this does not affect the arguments. Now, F is semisimple on X . While $i^*F[-1]$ is not semisimple, Lemma 2.4.1 applies, so that the RHL holds, and we can repeat verbatim the proof of Proposition 2.3.2.

3. The specialisation morphism as a filtered morphism

In this section, we initiate a systematic discussion of when the specialisation morphism is defined and when it is a filtered isomorphism for the perverse Leray filtrations associated with a morphism. The case of the perverse filtration is the special case when said morphism is the identity.

Section 3.1 revisits the classical specialisation morphism $sp^*(-)$ and frames it in a formalism suited to handle it together with perverse truncation in the later sections. The specialisation morphism and its filtered counterparts are not defined for an arbitrary complex. The main aim of this paper is to identify conditions that ensure that these morphisms are defined, and that when they are defined they are isomorphisms and, in the filtered context, filtered isomorphisms.

Section 3.2 introduces the notion of P -filtered specialisation morphism $sp_p^*(-)$, where P stands for the perverse filtration. This can be viewed as a special case of the perverse Leray specialisation morphism seen later in Section 3.3. However, it seems useful to discuss it separately, without the complication arising from the additional data of a morphism; this is similar to what happens with the Grothendieck and Leray spectral sequences/filtrations.

Section 3.3 introduces the central-to-this-paper notion of (perverse Leray) P^f -filtered specialisation morphism $sp_{P^f}^*(-)$. In order to streamline the discussion, we introduce the commutative diagram (55). We then prove Theorem 3.3.7 which lists criteria for the specialisation morphism to be a filtered isomorphism for the perverse Leray filtration; the earlier Propositions 3.1.9 and 3.2.7 are the analogous criteria for the specialisation and (perverse) P -filtered specialisation morphisms $sp_p^*(-)$ and $sp_{P^f}^*(-)$.

Section 3.4 is devoted to give a criterion for the specialisation morphism to exist and to be a filtered isomorphism when a suitable compactification is available: see Set-up 3.4.1 and Proposition 3.4.2.

In Section 3.5, we continue the theme of Section 3.4 -the existence of a suitable compactification- and consider the problem of the resulting long exact sequence of cohomology associated with the compactification -boundary, space, open part-. We prove Theorem 3.5.4 which gives criteria for the three specialisation morphisms involved to exist, to be filtered isomorphisms, and to fit into an isomorphism of the resulting long exact sequences for the triples, at the special point s and at the general point t .

We discuss the relation between the various criteria in Remark 3.3.10.

We apply these criteria to the Hitchin morphism in Section 4.

3.1. The specialisation morphism revisited

Let things be as in Section 1.4. In particular, we have: a morphism $v : Y \rightarrow S$, a point $s \in S$, a complex $G \in D_c^b(Y)$.

By combining the base change properties in (10) and (12), and by considering the morphisms of type v in (8), we obtain the commutative diagram of morphisms of functors $D_c^b(Y) \rightarrow D_c^b(pt)$ (cf. Remark 1.5.4):

$$\begin{array}{ccccc}
 i^*[-1]v_* & \xrightarrow{\sigma^*} & \psi[-1]v_* & \xrightarrow{\sigma^!} & i^![-1]v_* & (\phi v_*) & (42) \\
 \downarrow bc^{i^*v_*} & \dashrightarrow^{?sp^*?} & \downarrow bc^{\psi v_*} & \searrow sp^! & \downarrow bc^{i^!v_*} & & \\
 v_*i^*[-1] & \xrightarrow{\sigma^*} & v_*\psi[-1] & \xrightarrow{\sigma^!} & v_*i^![-1] & (v_*\phi), & \\
 & \searrow^{v=\nu} & & & & &
 \end{array}$$

where: there is a ν for each of the two rows, i.e. ${}^1\nu$ and ${}^2\nu$, but we have indicated only the one for the second row; the terms ϕv_* and $v_*\phi$ in parentheses are, up to shift, the cones of the morphisms of type σ (cf. Remark 3.1.1).

We have the functors $\nu, sp^1 : D_c^b(Y) \rightarrow D_c^b(pt)$. Due to the directions of the arrows on the l.h.s. of (42), the broken arrow “ ${}^?sp^*$ ” is not defined as a morphism of functors, but, as it is discussed below, it can happen to be defined for some $G \in D_c^b(Y)$.

Remark 3.1.1. (Cones in (42)). By (6), the cones of the morphisms σ^* and σ^1 on the top row of (42) are ϕv_* and $\phi v_*[1]$, respectively. Then, for $G \in D_c^b(Y)$, the morphism $\sigma^*(v_*G)$ is an isomorphism, if and only if the morphism $\sigma^1(v_*G)$ is an isomorphism, if and only if $\phi v_*G = 0$. What above remains true for the bottom row, with $v_*\phi$ replacing ϕv_* , etc.

Remark 3.1.2. (Interrelations between vanishings). In general, the two conditions $v_*\phi G = 0$ and $\phi v_*G = 0$ are independent. We have the following implications:

$$(\phi G = 0) \implies (v_*\phi G = 0); \nu \text{ proper} \implies ((v_*\phi G = 0) \iff (\phi v_*G = 0)).$$

Since in general the base change morphism $bc^{i^*v_*}$ is not an isomorphism, the sought-after morphisms ${}^?sp^*$ of functors $D_c^b(Y) \rightarrow D_c^b(pt)$ is not defined.

Definition 3.1.3. (The specialisation morphism $sp^*(G)$). Let $G \in D_c^b(Y)$. We say that the specialisation morphism $sp^*(G)$ is defined for G if the base change morphism $bc^{i^*v_*}(G)$ is an isomorphism. In this case, we define the morphism in $D_c^b(pt)$:

$$sp^*(G) := \sigma^*(bc^{i^*v_*})^{-1} : R\Gamma(Y_s, i^*G) = v_*i^*G \longrightarrow \psi v_*G = R\Gamma(s, \psi v_*G). \tag{43}$$

When $sp^*(G)$ is defined we have that:

$$\nu(G) = sp^1(G) \circ sp^*(G). \tag{44}$$

Remark 3.1.4. The part of diagram (42) that is needed to define specialisation morphisms is:

$$v_*i^*[-1] \xleftarrow{bc^{i^*v_*}} i^*[-1]v_* \xrightarrow{\sigma^*} \psi[-1]v_*. \tag{45}$$

The harmless shift by $[-1]$ is convenient later on, when dealing with the morphisms of type δ ; see diagram (49). The reason for having embedded (45) in (42) is explained in Remark 3.1.10.

Remark 3.1.5.

- (i) By using (17), we may re-write (43) as follows:

$$sp^*(G) : R\Gamma(Y_s, i^*G) \longrightarrow \psi v_*G \xrightarrow{\sim} R\Gamma(Y_t, t^*G). \tag{46}$$

The ambiguity involved in this re-writing (see Fact 1.5.5, especially (17)) plays no role in this paper: firstly, the image lands in the monodromy invariants; secondly, we have limited ourselves to criteria for when the specialisation morphism is: defined; an isomorphism; filtered for the perverse filtrations; a filtered isomorphism for the perverse filtrations. All of these issues can be settled at a point t general for G , independently of the monodromy action at t .

(ii) Even when $sp^*(G)$ is defined, I do not see a reason for the base change morphism

$$bc^{\psi v_*} : (R\Gamma(Y_t, t^*G) \simeq) \psi v_* G \longrightarrow v_* \psi G = R\Gamma(Y_s, \psi G) \tag{47}$$

to be necessarily an isomorphism.

(iii) In view of the isomorphism $\psi v_* G \simeq R\Gamma(Y_t, t^*G)$, typically, and this paper is no exception, we are interested in $\psi v_* G$, and not as much in $v_* \psi G$. This is important to keep in mind when, later we consider the filtered version of the specialisation morphisms see (49), (50) and (53) for the perverse version, and see (55), (56) and (59) for the perverse Leray version. Of course, the base change morphism $bc^{\psi v_*} : \psi v_* \rightarrow v_* \psi$ is an isomorphism when v is proper.

Remark 3.1.6. (v proper). If $v : Y \rightarrow S$ is proper, then, by the proper base change theorem, we get the morphism of functors sp^* .

Remark 3.1.7. (The morphism of functors $sp^1 : \psi[-1]v_ \rightarrow v_*i^1[1]$).* Regardless of whether v is proper or not, the functor $sp^1 : D_c^b(Y) \rightarrow D_c^b(pt)$ is defined. In analogy with (46), and taking into account the definition of cohomology with supports on the closed set Y_s , for a given $G \in D_c^b(Y)$, it takes the form of a morphism:

$$sp^1(G) : R\Gamma(Y_t, t^*G) \xrightarrow{\simeq} \psi v_* G \longrightarrow R\Gamma(Y_s, i^1[2]G) = R\Gamma_{Y_s}(Y, G[2]). \tag{48}$$

Most of the results proved in this paper for the morphism $sp^*(G)$ (when defined), hold for the morphism sp^1 . In fact, the proofs are simpler in this case, since we do not need to establish the existence of sp^1 along the way, the way we must do for a $sp^*(G)$. On the other hand, in general, the r.h.s. of (48) is not easily relatable to, say, $R\Gamma(Y_s, i^*G)$. For this reason, we have not included in this paper the sp^1 -versions of the results. We simply observe that when, for some necessarily special G , the natural morphism $v : i^*[-1](G) \rightarrow i^1[1](G)$ is an isomorphism, then, we do have that $sp^1(G) : R\Gamma(Y_t, t^*G) \rightarrow R\Gamma(Y_s, i^*G)$ and this can be a very useful tool, especially, when $sp^*(G)$ fails to be defined. This proved to be crucial in the paper [de-Ma-Sh 2020] (where due to the particular set-up there, we opted for a more direct construction of the morphism sp^1).

Remark 3.1.8. In what follows, “($\Leftarrow (\phi G = 0)$)” is short for “(which is implied by $(\phi G = 0)$)”.

PROPOSITION 3.1.9. (Criteria for the existence of $sp^*(G)$ and for it being an isomorphism). *Let $G \in D_c^b(Y)$. Recall Remark 3.1.8.*

- (i) *If v is proper, then $sp^* : D_c^b(Y) \rightarrow D(pt)$ is defined as a functor. If in addition $\phi v_* G = 0$ ($\Leftarrow (\phi G = 0)$), then $sp^*(G)$ is an isomorphism.*
- (ii) *If $\phi v_* G = 0$ and $\phi G = 0$, then $sp^*(G)$ is defined and an isomorphism.*

Proof. (i). For the first assertion, see Remark 3.1.6. For the second one, we are left with showing that the morphism $\sigma^*(v_*G)$ on the top row of (42) is an isomorphism. This follows from the fact that its cone $\phi v_* G = 0$.

(ii). The hypotheses imply that the cones of the four horizontal arrows in (42) are zero. These four arrows are then all isomorphisms. Since one of the three vertical arrows in (42) is an isomorphism, so are the remaining two, and we are done.

Remark 3.1.10 (Vanishing of ϕ and base change). In order to define the specialisation morphism, we only need the l.h.s. of diagram (42). The proof of Proposition 3.1.9.(2) shows that the two conditions $\phi G = 0$ and $\phi v_* G = 0$ combined, imply that the base change morphism $bc^{i^*v_*}(G)$ for G is an isomorphism. This is the reason why we have introduced diagram (42): if, after having plugged some $G \in D_c^b(Y)$, the four horizontal arrows in (42) are isomorphisms, then, because of the presence of base change isomorphism $bc^{i^*v_*}$, the three vertical arrows evaluated at G are also isomorphism, i.e. the base change morphism $bc^{i^*v_*}$ is also an isomorphism, and the specialisation morphism $sp^*(G)$ is defined.

3.2. The specialisation morphism and the perverse filtration

Let things be as in Section 1.4. In particular, we have: a morphisms $v : Y \rightarrow S$, a point $s \in S$, a complex $G \in D_c^b(Y)$.

By analogy to (42), and by using the naturality properties of the morphisms of type δ in Section 2, we obtain the commutative diagram of morphism of systems of functors:

$$\begin{array}{ccccc}
 i^*[-1]v_* \mathcal{P}\tau_{\leq \bullet} & \xrightarrow{1\sigma_{\leq \bullet}^*} & \psi[-1]v_* \mathcal{P}\tau_{\leq \bullet} & \xrightarrow{1\sigma'_{\leq \bullet}} & i^![1]v_* \mathcal{P}\tau_{\leq \bullet} & (\phi v_* \mathcal{P}\tau_{\leq \bullet}) \\
 \downarrow bc_{\leq \bullet}^{i^*v_*} & \nearrow ?sp_{\leq \bullet}^*? & \downarrow bc^{\psi v_*} & \searrow sp'_{\leq \bullet} & \uparrow = bc_{\leq \bullet}^{i^!v_*} & \\
 v_* i^*[-1] \mathcal{P}\tau_{\leq \bullet} & \xrightarrow{2\sigma_{\leq \bullet}^*} & v_* \psi[-1] \mathcal{P}\tau_{\leq \bullet} & \xrightarrow{2\sigma'_{\leq \bullet}} & v_* i^![1] \mathcal{P}\tau_{\leq \bullet} & (v_* \phi \mathcal{P}\tau_{\leq \bullet}) \\
 \uparrow \delta_{\leq \bullet}^* & \nearrow & \downarrow = \delta_{\leq \bullet}^{\psi} & \searrow & \downarrow \delta'_{\leq \bullet} & \\
 v_* \mathcal{P}\tau_{\leq \bullet} i^*[-1] & \xrightarrow{3\sigma_{\leq \bullet}^*} & v_* \mathcal{P}\tau_{\leq \bullet} \psi[-1] & \xrightarrow{3\sigma'_{\leq \bullet}} & v_* \mathcal{P}\tau_{\leq \bullet} i^![1], & (\phi). \\
 & \searrow v_{\leq \bullet} = \mathfrak{v}_{\leq \bullet} & & & &
 \end{array}$$

(49)

Remark 3.2.1 (Cones/not cones in (49)). In the top two rows of diagram (49), we have indicated in parentheses the cones -up to shift- of the morphisms of type σ : apply (42) and Remark 3.1.1 to $\mathcal{P}\tau_{\leq \bullet}$. Since truncation is not an exact functor, the cones of the morphisms of type σ in the third row are not shifts of $v_* \mathcal{P}\tau_{\leq \bullet} \phi$; the term in parentheses on the third bottom row is a term whose vanishing when evaluated at G ensures that the arrows of type $\sigma(G)$ on the third row are isomorphisms.

The reason why (49) has three rows, instead of the two in (cf. 42), is that $\mathcal{P}\tau_{\leq \bullet}$ does not commute with $i^*[-1]$ -nor with any other shift-; the morphisms of type δ measure the failure of this commutativity.

Remark 3.2.2. (Interrelations between vanishings) . The conditions $\phi v_* \mathcal{P}\tau_{\leq \bullet} G = 0$ and $v_* \phi \mathcal{P}\tau_{\leq \bullet} G = 0$ are independent; if v is proper, then they are equivalent. By the t-exactness of ϕ and the additivity of v_* , we have that: $v_* \phi \mathcal{P}\tau_{\leq \bullet} = v_* \mathcal{P}\tau_{\leq \bullet} \phi$, and that: $(\phi G = 0) \iff (\phi \mathcal{P}\tau_{\leq \bullet} G = 0) \implies v_* \phi \mathcal{P}\tau_{\leq \bullet} G = 0$.

Since in general the base change morphism $bc_{\leq \bullet}^{i^*v_*}$ are not isomorphisms, the sought-after morphism of systems of functors $?sp_{\leq \bullet}^*?$ is not defined. On the other hand, there can be a corresponding arrow for special $G \in D_c^b(Y)$. Recall the notion of “systems” (cf. Section 1.3).

Definition 3.2.3. (The P-filtered specialisation morphism $sp_p^(G)$)* . Let $G \in D_c^b(Y)$. We say that the P -filtered specialisation morphism $sp_p^*(G)$ is defined for G if the base change morphisms $bc_{\leq \bullet}^{i^*v_*}(G)$ are isomorphisms, in which case, since the inverse isomorphisms form an isomorphism of systems, we can define the arrow in $DF(pt)$: ($bc_P^{i^*v_*}$ denotes the evident resulting filtered isomorphism, and δ_P the evident resulting filtered morphism)

$$sp_p^*(G) := \sigma_p^*(bc_P^{i^*v_*})^{-1} \delta_P^* : v_* i^* G = (R\Gamma(Y_s, i^*G), P) \longrightarrow (\psi v_* G, P), \tag{50}$$

where the last term is the one associated with the system $\psi v_* {}^P\tau_{\leq \bullet} G$.

We have functors $v_P, sp_p^1 : D_c^b(Y) \rightarrow DF(pt)$. When $sp_p^*(G)$ is defined, we have that:

$$v_P(G) = sp_p^1(G) \circ sp_p^*(G). \tag{51}$$

Remark 3.2.4. Remark 3.1.4 holds essentially verbatim in the context of diagram (49). In particular, we have:

$$v_* {}^P\tau_{\leq \bullet} i^*[-1] \xrightarrow{\delta_{\leq \bullet}^*} v_* i^*[-1] {}^P\tau_{\leq \bullet} \xleftarrow{bc^{i^*v_*}} i^*[-1] v_* {}^P\tau_{\leq \bullet} \xrightarrow{\sigma^*} \psi[-1] v_* {}^P\tau_{\leq \bullet}. \tag{52}$$

Remark 3.2.5.

- (i) By using (18), the filtered analogue of (43) takes the following form:

$$sp_p^*(G) : (R\Gamma(Y_s, i^*G), P) \longrightarrow (\psi v_* G, P) \xrightarrow{\sim} (R\Gamma(Y_t, i^*G), P). \tag{53}$$

The analogue of Remark 3.1.5 on the ambiguities due to monodromy holds in this context.

- (ii) Even when $sp_p^*(G)$ is defined, the base change morphism:

$$bc^{\psi v_*} : (R\Gamma(Y_t, i^*G) \simeq) \psi v_* G \longrightarrow v_* \psi G = R\Gamma(Y_s, \psi G) \tag{54}$$

is not necessarily an isomorphism. In particular, the filtration P in $(\psi v_* G, P)$ is not the perverse filtration for a complex on Y_s .

Remark 3.2.6. (v proper) . The evident analogue of Remark 3.1.6 holds in this perverse filtered context: if $v : Y \rightarrow S$ is proper, then, by the proper base change theorem, we get the functor: $sp_p^* : D_c^b(Y) \rightarrow DF(pt)$.

Proposition 3.2.7 below is the P -filtered counterpart to the un-filtered Proposition 3.1.9.

PROPOSITION 3.2.7. (Criteria for the existence of $sp_p^*(G)$ and for it being a filtered isomorphism) .

- (i) If v is proper, then $sp_p^* : D_c^b(Y) \rightarrow DF(pt)$ is defined as a functor.
If in addition $\phi G = 0$, then $sp_p^*(G)$ is a filtered isomorphism.
- (ii) If $\phi v_* {}^P\tau_{\leq \bullet} G = 0$ and $v_* \phi {}^P\tau_{\leq \bullet} G = 0$, then $sp_p^*(G)$ is defined.
If in addition $\phi G = 0$, then $sp_p^*(G)$ is a filtered isomorphism.
- (iii) If G is semisimple, $\phi v_* G = 0$ and $v_* \phi G = 0$, then $sp_p^*(G)$ is defined.
If in addition $\phi G = 0$, then $sp_p^*(G)$ is a filtered isomorphism.

Proof. (i) For the first assertion, see Remark 3.2.6. If we assume that $\phi G = 0$, then, in view of Remarks 3.2.1 and 3.2.2, all horizontal arrows in the commutative diagram (49) are isomorphisms. Since each row of vertical arrows contains an isomorphisms, all arrows in (49) are isomorphisms, so that sp_p^* is a filtered isomorphism.

(ii) The two vanishing assumptions imply that all the four horizontal arrows in the first two rows of (49) are isomorphisms. As seen above, this implies that the vertical base change arrows are isomorphisms as well, so that sp_p^* is defined. If we we assume that $\phi G = 0$, then we conclude as above that sp_p^* is a filtered isomorphism.

(iii) Since G is semisimple, we have that: $\phi v_* G = 0$ is equivalent to $\phi v_* {}^p\tau_{\leq \bullet} G = 0$; $v_* \phi G = 0$ is equivalent to $v_* \phi {}^p\tau_{\leq \bullet} G = 0$, Now (iii) follows from (ii).

Remark 3.2.8. (Vanishing of ϕ and base change) . The analogue of Remark 3.1.10, on the vanishing of ϕ assumptions implying base change isomorphisms, holds in the context of the proof of Proposition 3.2.7.

Question 3.2.9. I do not know of an example where: v is proper -so that sp^* and sp_p^* are defined-, where $\phi v_* G = 0$ -so that $sp(G)$ is an isomorphism-, but where $sp_p(G)$ is not an isomorphism (i.e. in the filtered sense). The issue is that while $\delta_{\leq \bullet}^*(G)$ in (49) is an isomorphism for $\bullet \gg 0$, it is not clear to me what happens for other values of \bullet . We know that if in addition $\phi G = 0$, then sp_p^* is an isomorphism.

3.3. The specialisation morphism and the perverse Leray filtration

Let things be as in Section 1.4. In particular, we have: morphisms $f : X \rightarrow Y$ and $v : Y \rightarrow S$, a point $s \in S$, a complex $G := f_* F \in D_c^b(Y)$ for $F \in D_c^b(X)$.

By analogy to (42) and (49), and by using the naturality properties of the morphisms of type δ in Section 2, we obtain the commutative diagram of morphisms of systems of functors:

$$\begin{array}{ccccc}
 i^*[-1]v_* {}^p\tau_{\leq \bullet} f_* & \xrightarrow{1\sigma_{\leq \bullet}^*} & \psi[-1]v_* {}^p\tau_{\leq \bullet} f_* & \xrightarrow{1\sigma_{\leq \bullet}^!} & i^![-1]v_* {}^p\tau_{\leq \bullet} f_* & ((\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0) \Leftrightarrow 1\sigma \text{ isos}) \\
 \downarrow bc_{\leq \bullet}^{i^* v_*} & & \downarrow 4 & & \downarrow bc_{\leq \bullet}^{i^! v_*} & \\
 v_* i^*[-1] {}^p\tau_{\leq \bullet} f_* & \xrightarrow{2\sigma_{\leq \bullet}^*} & v_* \psi[-1] {}^p\tau_{\leq \bullet} f_* & \xrightarrow{2\sigma_{\leq \bullet}^!} & v_* i^![-1] {}^p\tau_{\leq \bullet} f_* & ((v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0) \Leftrightarrow 2\sigma \text{ isos}) \\
 \uparrow \delta_{\leq \bullet}^* & \nearrow ?sp_{p^*}^* & \uparrow = & \searrow sp_{p^*}^! & \downarrow \delta_{\leq \bullet}^! & \\
 v_* {}^p\tau_{\leq \bullet} i^*[-1] f_* & \xrightarrow{3\sigma_{\leq \bullet}^*} & v_* {}^p\tau_{\leq \bullet} \psi[-1] f_* & \xrightarrow{3\sigma_{\leq \bullet}^!} & v_* {}^p\tau_{\leq \bullet} i^![-1] f_* & ((\phi f_* F = 0) \Rightarrow 3\sigma \text{ isos}) \\
 \downarrow bc_{\leq \bullet}^{i^* f_*} & & \downarrow 4' & & \downarrow bc_{\leq \bullet}^{i^! f_*} & \\
 v_* {}^p\tau_{\leq \bullet} f_* i^*[-1] & \xrightarrow{4\sigma_{\leq \bullet}^*} & v_* {}^p\tau_{\leq \bullet} f_* \psi[-1] & \xrightarrow{4\sigma_{\leq \bullet}^!} & v_* {}^p\tau_{\leq \bullet} f_* i^![-1] & ((f_* \phi F = 0) \Rightarrow 4\sigma \text{ isos}). \\
 & \searrow v_{\leq \bullet} = 4v_{\leq \bullet} & & & &
 \end{array}$$

Remark 3.3.1. (Cones/not cones in (55)) . This remark is analogous to Remark 3.2.1. The cones of the morphisms of type σ appearing on the first two rows are indicated in parentheses on the r.h.s. The terms in parentheses on the third and fourth row are not the cones of the morphisms of type σ (truncation is not an exact functor), but their vanishing implies that the morphisms of type σ are isomorphisms.

Remark 3.3.2. (Interrelations between vanishings). The conditions $\phi F = 0$ and $\phi f_* F = 0$ are independent; if f is proper, then $\phi F = 0$ implies $\phi f_* F = f_* \phi F = 0$. Note that $\phi f_* F = 0$ is equivalent to $\phi {}^p\tau_{\leq \bullet} f_* F = 0$, and it implies $v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$.

The conditions $v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$ and $\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0$ are independent; if v is proper, then they are equivalent.

Definition 3.3.3. (The P^f -filtered specialisation morphism $sp_{P^f}^(F)$)*. We say that the P^f -filtered specialisation morphism $sp_{P^f}^*(F)$ is defined for $F \in D_c^b(X)$ if the base change morphisms $bc_{\leq \bullet}^{i^* v_*}(F)$ and $bc_{\leq \bullet}^{i^* f_*}(F)$ are isomorphisms. In this case, since the inverse isomorphisms form a morphism of systems, we can define the arrow in $DF(pt)$:

$$sp_{P^f}^*(F) := {}^1\sigma_P^*(bc_P^{i^* v_*})^{-1} \delta_{P^f}^*(bc_P^{i^* f_*})^{-1} : v_* i^* G = (R\Gamma(X_s, i^* F), P^f) \longrightarrow (\psi v_* f_* F, P), \tag{56}$$

where the last term is the one associated with the system $\psi v_* {}^p\tau_{\leq \bullet} f_* F$.

We have functors $v_{P^f}, sp_{P^f}^! : D_c^b(X) \rightarrow DF(pt)$. When $sp_{P^f}^*(F)$ is defined, we have that:

$$v_{P^f}(F) = sp_{P^f}^!(F) \circ sp_{P^f}^*(F). \tag{57}$$

Remark 3.3.4. Remark 3.1.4 holds essentially verbatim in the context of diagram (55). In particular, we have:

$$v {}^p\tau_{\leq \bullet} f_* i^*[-1] \xleftarrow{bc_{\leq \bullet}^{i^* f_*}} v_* {}^p\tau_{\leq \bullet} i^*[-1] f_* \xrightarrow{\delta_{\leq \bullet}^*} v_* i^*[-1] {}^p\tau_{\leq \bullet} f_* \xleftarrow{bc_{\leq \bullet}^{i^* v_*}} i^*[-1] v_* {}^p\tau_{\leq \bullet} f_* \downarrow \sigma^* \psi[-1] v_* {}^p\tau_{\leq \bullet} f_* \tag{58}$$

Remark 3.3.5. By using (19), we may re-write (56) as follows:

$$sp_{P^f}^*(F) : (R\Gamma(X_s, i^* F), P^{f_s}) \longrightarrow (\psi v_* f_* F, P) \xrightarrow{\sim} (R\Gamma(X_t, i^* F), P^{f_t}). \tag{59}$$

The analogue of Remark 3.2.5, on the ambiguities due to monodromy and on the meaning of P in $(\psi v_* f_* F, P)$, holds in this context.

Remark 3.3.6. (v proper). The evident analogue of Remarks 3.1.6 and 3.2.6 holds in this perverse Leray filtered context: if f and v are proper, then we get the functor: $sp_{P^f}^* : D_c^b(X) \rightarrow DF(pt)$.

The following theorem is a perverse Leray analogue of Proposition 3.2.7. Recall the notational Remark 3.1.8.

THEOREM 3.3.7. (Criteria for the existence of $sp_{P^f}^*(F)$ and for it being a filtered isomorphism).

- (i) If v and f are proper, then $sp_{P^f}^* : D_c^b(X) \rightarrow DF(pt)$ is a functor. If in addition $\phi f_* F = 0$ ($\Leftarrow (\phi F = 0)$) then $sp_{P^f}^*(F)$ is defined and a filtered isomorphism.
- (ii) If v proper, $\phi f_* F = 0$, and $f_* \phi F = 0$ ($\Leftarrow (\phi F = 0)$), then $sp_{P^f}^*(F)$ is defined and a filtered isomorphism.

- (iii) If f is proper, $\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0$, and $v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$ ($\Leftarrow (\phi f_* F = 0) \Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is defined.
 If in addition $\phi f_* F = 0$ ($\Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is defined and a filtered isomorphism.
- (iv) If $\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0$, $\phi f_* F = 0$, and $f_* \phi F = 0$ ($\Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is defined and a filtered isomorphism.
- (v) If f is proper and F is semisimple, $\phi v_* f_* F = 0$ and $v_* \phi f_* F = 0$ ($\Leftarrow (\phi f_* F = 0) \Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is defined.
 If in addition, $\phi f_* F = 0$ ($\Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is a filtered isomorphism.
- (vi) If v is proper, f is projective and F and $i^*F[-1]$ are semisimple, and $\phi v_* f_* F = 0$ ($\Leftarrow (\phi f_* F = 0) \Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is defined and a filtered isomorphism.
- (vii) If f is projective and F and $i^*F[-1]$ are semisimple, $\phi v_* f_* F = 0$ and $v_* \phi f_* F = 0$ ($\Leftarrow (\phi f_* F = 0) \Leftarrow (\phi F = 0)$), then $sp_{p_f}^*(F)$ is defined and a filtered isomorphism.

Proof. (i) For the first assertion in (i), see Remark 3.3.6. We now prove the second assertion in (i). Keep in mind that the base change morphisms are all isomorphisms. Since f is proper, $\phi f_* F = 0$ implies that -it is equivalent to- $f_* \phi F = 0$, as well as $v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$. It follows that the four horizontal arrows on row two and three in (55) are isomorphisms. This implies that so are the vertical arrows of type δ . It remains to show that the morphisms $\iota_{\leq \bullet}^*$ are isomorphisms, which follows from $\phi v_* {}^p\tau_{\leq \bullet} f_* F = v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$.

(ii) Since v is proper, rows one and two are identified. The vanishing assumptions imply that all seven arrows on rows three and four are isomorphisms. In particular, the six base change morphisms in (55) are isomorphisms, so that $sp_{p_f}^*$ is defined. The assumption $\phi f_* F = 0$ implies that the cones of the four horizontal arrows in rows one and two are isomorphisms, so that all seven arrows on these two rows are isomorphisms. The same argument shows that the seven arrows in rows two and three are isomorphisms. In particular, the morphisms of type δ and the morphisms $\iota_{\leq \bullet}^*$ are isomorphisms, and the conclusion follows.

(iii) We prove the first assertion in (iii). The base change morphisms for f are isomorphisms. The vanishing assumptions imply that we can identify the first two rows, so that $sp_{p_f}^*$ is defined. We prove the second assertion in (iii). The hypothesis $\phi f_* F = 0$ implies that the morphisms of type δ are isomorphisms, and we are done.

(iv) The vanishing assumptions imply that all horizontal arrows in (55) are isomorphisms. It follows that all arrows in (55) are isomorphisms and the conclusion follows.

(v) Since f is proper and F is semisimple, $f_* F$ is semisimple by the Decomposition Theorem 1.3.1, so that ${}^p\tau_{\leq \bullet} f_* F$ is a direct summand of $f_* F$. It follows that: $\phi v_* f_* F = 0$ is equivalent to $\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0$; $v_* \phi f_* F = 0$ is equivalent to $v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$. This shows that (v) follows from (iii).

(vi) Since v and f are proper, $sp_{p_f}^*$ is defined. The semisimplicity assumptions on F and on $i^*F[-1]$ and the projectivity of f imply, via Proposition 2.3.2, that the morphisms of type δ are isomorphisms. It remains to observe that the assumption $\phi v_* f_* F = 0$, coupled with the semisimplicity of $f_* F$, implies, as seen in the proof of (v), that $\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0$. It follows that σ^* is an isomorphisms and we are done.

(vii) Since f is proper, the corresponding base change morphisms are isomorphisms. As seen in the proof of (vi), the morphisms of type δ are isomorphisms. As seen in the proof of (v), the vanishing assumptions imply that $\phi v_* {}^p\tau_{\leq \bullet} f_* F = 0$ and $v_* \phi {}^p\tau_{\leq \bullet} f_* F = 0$. As in the proof of (iii), the base change morphisms for v are isomorphisms and, as in the proof of (i), the morphisms $\iota_{\leq \bullet}^*$ are isomorphisms. The proof of (vii) is complete.

Remark 3.3.8. (Vanishing of ϕ and base change) . The analogue of Remark 3.1.10 holds in the context of the proof of Theorem 3.3.7.

Remark 3.3.9. Recall that a condition of type $\phi v_*(-) = 0$ means that $v_*(-)$ has locally constant cohomology sheaves $R^\bullet v_*(-)$ on S near s .

Remark 3.3.10. (i) [de-Ma 2018, theorem 3.2.1 parts (i,ii,ii)] are implied by the slightly more precise Theorem 3.3.7 parts (i,iii,v), respectively.
 (ii) Proposition 3.2.7 parts (i,ii,iii) are also the special cases of Theorem 3.3.7 parts (i,iii,v) when one takes $f : X \rightarrow Y$ to be the identity.

3.4. *The specialisation morphism via a compactification*

Consider Theorem 3.3.7: parts (iii,iv,v,vii) do not assume that $v = v_X : X \rightarrow S$ is proper, but make some local constancy assumptions for certain direct images to S , namely $\phi v_*(-) = 0$.

In this subsection we provide, in the context of a proper S -morphism $f : X \rightarrow Y$ and of a non proper morphisms $v_X : X \rightarrow S$, sufficient conditions ensuring that we obtain well-defined $sp_p^*(G)$ or $sp_{p,f}^*(F)$, but that do not require local constancy assumptions. On the other hand, they require the existence of good compactification. As we shall see, here “good” is made precise by conditions relating vanishing cycles at the boundary. In general, it may be difficult to achieve such a vanishing. See Remark 3.4.5 for one situation in which this is possible. They are also achieved in the compactification of Dolbeault moduli spaces constructed in [de 2018]; see Section 4.

We start with a proper S -morphism $f : X^o \rightarrow Y^o$, and we do not assume that $v = v_{Y^o} : Y^o \rightarrow S$ is proper.

In order to circumvent base change issues, it is natural to first compactify the picture. We start with $f : X^o \rightarrow Y^o$ and we get, by Nagata’s completion theorems: Zariski open and dense embeddings $X^o \subseteq X$ and $Y^o \subseteq Y$; a proper morphism $f : X \rightarrow Y$ extending $f : X^o \rightarrow Y^o$; a proper morphism $v_Y : Y \rightarrow S$ extending $v_{Y^o} : Y^o \rightarrow S$. By blowing-up Y , if necessary, we may assume that $W := Y \setminus Y^o$ supports an effective Cartier divisor.

It follows that we can place ourselves in the following:

SET-UP 3.4.1. *Let S be a variety. Consider a Cartesian diagram of S -morphisms, with $(a : W \rightarrow Y \leftarrow Y^o : b)$ closed/open complementary immersions:*

$$\begin{array}{ccccc}
 Z & \xrightarrow{a} & X & \xleftarrow{b} & X^o \\
 \downarrow f & & \downarrow f & & \downarrow f \\
 W & \xrightarrow{a} & Y & \xleftarrow{b} & Y^o
 \end{array} \tag{60}$$

such that: all morphisms f are proper; the varieties Y and W proper over S ; the boundary W on Y supports an effective Cartier divisor. In particular, the functors $b_!, b_ : Y^o \rightarrow Y$ are t-exact.*

Many of us are used to denote the pair of closed/open embeddings (a, b) by (i, j) . However, in this paper, and in large part of the vanishing cycle literature, the closed embedding of the special fiber is systematically denoted by i .

We denote by ad the distinguished triangle of endofunctors of $D_c^b(Y)$:

$$\text{ad} := a_* a^! \xrightarrow{a} \text{Id} \xrightarrow{b} b_* b^* \rightsquigarrow ; \tag{61}$$

by plugging $G \in D_c^b(Y)$ in (61), we obtain the distinguished triangle $\text{ad}(G)$ in $D_c^b(Y)$, functorial in G .

Proposition 3.4.2 below is an “ f -proper, but v_X -non-proper” analogue to Theorem 3.3.7.(i), where it was assumed that f and v are proper and that $\phi F = 0$. Recall Remark 3.3.5, which allows us to use a general point t to express the target of specialisation morphisms.

Note that the assumptions (B) in Proposition 3.4.2 imply the assumptions of Theorem 3.3.7.(v) which lead to $sp_{P^f}^*$ being an isomorphism. We have included (B) for completeness in the context of this subsection. Recall the notational Remark 3.1.8.

PROPOSITION 3.4.2. *Assume we are in the Set-up 3.4.1. Assume that S is a nonsingular and connected curve and let $s \in S$ be a point. Let $F \in D_c^b(X)$.*

Consider the following two sets of conditions:

- (A) $\phi b_* f_* b^* F = 0$ ($\Leftarrow (\phi b_* b^* F = 0)$), and $\phi f_* b^* F = 0$ ($\Leftarrow (\phi b^* F = 0) \Leftarrow (\phi b^* F = 0) \Leftarrow (\phi F = 0)$);
- (B) F semisimple, $\phi v_* b_* f_* b^* F = 0$ ($\Leftarrow (\phi f_* b_* b^* F = 0) \Leftarrow (\phi b_* b^* F = 0)$), and $\phi f_* b^* F = 0$ ($\Leftarrow (\phi b^* F = 0) \Leftarrow (\phi F = 0)$).

If either conditions (A) or (B) are met, then the P^f -filtered specialisation morphism:

$$sp_{P^f}^*(F) : (R\Gamma(X_s^o, i^* b^* F), P^{f_s}) \longrightarrow (\psi v_* f_* i^* F, P) \simeq (R\Gamma(X_t^o, t^* b^* F), P^{f_t}), \tag{62}$$

where $t \in S$ is general for F with respect to X/S and for $f_* F$ with respect to Y/S , is defined and it is a filtered isomorphism for the perverse Leray filtrations.

If $f : X \rightarrow Y$ is the identity, then the same conclusion holds with $sp_{P^f}^(F) = sp_P^*(F)$.*

Proof. Since the proof is analogous to the proof of Theorem 3.3.7.(i), we only indicate the main line of the arguments.

In complete analogy with the formation of the commutative diagram (55), we form the commutative diagram: (with many decorations omitted)

$$\begin{array}{ccccc}
 i^*[-1]v_* b_* P_{\tau_{\leq \bullet}} f_* b^* & \xrightarrow{\sigma^*} & \psi[-1]v_* b_* P_{\tau_{\leq \bullet}} f_* b^* & \xrightarrow{\sigma^i} & i^![-1]v_* b_* P_{\tau_{\leq \bullet}} f_* b^* & (\phi v_* b_* P_{\tau_{\leq \bullet}} f_* b^* F), \\
 \downarrow bc & & \swarrow / & \downarrow bc & \downarrow = bc & \\
 v_* i^*[-1]b_* P_{\tau_{\leq \bullet}} f_* b^* & \xrightarrow{\sigma^*} & v_* \psi[-1]b_* P_{\tau_{\leq \bullet}} f_* b^* & \xrightarrow{\sigma^i} & v_* i^![-1]b_* P_{\tau_{\leq \bullet}} f_* b^* & (v_* \phi b_* P_{\tau_{\leq \bullet}} f_* b^* F), \\
 \downarrow bc^{i^* b_*} & & \swarrow / & \downarrow bc & \downarrow = bc & \\
 v_* b_* i^*[-1]P_{\tau_{\leq \bullet}} f_* b^* & \xrightarrow{\sigma^*} & v_* b_* \psi[-1]P_{\tau_{\leq \bullet}} f_* b^* & \xrightarrow{\sigma^i} & v_* b_* i^![-1]P_{\tau_{\leq \bullet}} f_* b^* & (\phi f_* b^* F), \\
 \uparrow \delta_{\leq \bullet}^* & & \swarrow / & \downarrow = & \downarrow \delta_{\leq \bullet}^i & \\
 v_* b_* P_{\tau_{\leq \bullet}} i^*[-1]f_* b^* & \xrightarrow{\sigma^*} & v_* b_* P_{\tau_{\leq \bullet}} \psi[-1]f_* b^* & \xrightarrow{\sigma^i} & v_* b_* P_{\tau_{\leq \bullet}} i^![-1]f_* b^* & (\phi f_* b^* F), \\
 \downarrow bc / & & \downarrow bc & & \downarrow = bc & \\
 v_* b_* P_{\tau_{\leq \bullet}} f_* i^*[-1]b^* & \xrightarrow{\sigma^*} & v_* b_* P_{\tau_{\leq \bullet}} f_* \psi[-1]b^* & \xrightarrow{\sigma^i} & v_* b_* P_{\tau_{\leq \bullet}} f_* i^![-1]b^* & (\phi b^* F),
 \end{array} \tag{63}$$

the properties of which are similar to the ones of (55). The second row from the top in (63) does not have a counterpart in (55), and this is because here we need to consider the base change morphisms for i^*b_* . The terms in parentheses for the first three rows are, up to shift, the cones of the arrows of type σ . After having plugged an $F \in D_c^b(X)$ in (63), the vanishing of the term in parentheses implies that the corresponding morphisms of type σ are isomorphisms.

Since $v : X \rightarrow S$ is proper, the first two rows get identified by proper base change. Since f is proper, the same is true for the last two.

Since every row contains a vertical arrow which is an isomorphism, we only need to show that all horizontal arrows in (63) are isomorphisms. It is enough to show that the corresponding terms in parentheses vanish. By the identifications above, we need to do so only for rows two, three and four.

The assumption $\phi b^* F = 0$, which is common to (A) and (B), together with the properness of f , implies the vanishing of the terms in parentheses for the bottom three rows (for the third row, one uses that ϕ commutes with perverse truncation).

We are left with proving that the morphisms of type σ on row two are isomorphisms.

We prove that (A) implies the desired conclusion. The assumption $\phi b_* b^* F = 0$ implies the vanishing of the terms in parentheses for the second row because of what follows:

$$0 = f_* \phi b_* b^* F = \phi f_* b_* b^* F = \phi b_* f_* b^* F = \phi b_* b^* f_* F,$$

so that:

$$0 = {}^p\tau_{\leq \bullet} \phi b_* b^* f_* F = \phi {}^p\tau_{\leq \bullet} b_* b^* f_* F = \phi b_* {}^p\tau_{\leq \bullet} b^* f_* F = \phi b_* {}^p\tau_{\leq \bullet} f_* b^* F,$$

where in the identities above we have used proper base change for f , the fact that $fb = bf$ in the r.h.s. of (60), the t-exactness of b_* (b is affine and quasi-finite) and of ϕ .

(B) has already been proved in Theorem 3-3-7.(v).

Remark 3-4-3. The proof of Proposition 3-4-2 shows that under its hypotheses, (A) or (B), the base change morphism $bc^{i^*b_*} : v_* i^* b_* {}^p\tau_{\leq \bullet} f_* b^* F \rightarrow v_* b_* i^* {}^p\tau_{\leq \bullet} f_* b^* F$ in (65) is an isomorphism.

Remark 3-4-4. Remark 3-1-4, on the part of (55) that is needed to define $sp_{p_f}^*(F)$, holds essentially verbatim in the context of diagram (63).

Remark 3-4-5. The conditions $\phi b_* b^* F = 0 = \phi b^* F$ in Proposition 3-4-2 are met if, for example, X/S is smooth, $X \setminus X^o$ is a simple normal crossing divisor over S and F has locally constant cohomology sheaves. See [De 1972, XIII, Lemme 2-1-11 (dualised)]. We do not know of a weaker, but similar set of conditions that leads to $sp_{p_f}^*$ being defined, but not necessarily an isomorphisms.

COROLLARY 3-4-6. *Let things be as in Proposition 3-4-2. Then we have the following commutative diagram, with horizontal arrows given by restriction, and with vertical arrows given by the well-defined P^f -filtered specialisation morphisms:*

$$\begin{array}{ccc}
 (R\Gamma(X_t, t^*F), P^{f_i}) & \longrightarrow & (R\Gamma(X_t^o, t^*b^*F), P^{f_i}) \\
 \uparrow sp_{P_f}^* & & \uparrow sp_{P_f}^* \cong \\
 (R\Gamma(X_s, i^*F), P^{f_s}) & \longrightarrow & (R\Gamma(X_s^o, i^*b^*F), P^{f_s}).
 \end{array} \tag{64}$$

Proof. We artificially add to digram (55) one row identical to the second row from the top and place it between the second and third row; there are now five rows; we connect the second row to the new third row by the identity; we connect the new third row to the new fourth row the same way as rows two and three in (55) are connected. The reader can verify that: if we apply the adjunction $Id \rightarrow b_*b^*$ to this new diagram, call it (55'), then we obtain what can be called the adjunction morphism between (55') and (63). The conclusion follows.

3.5. specialisation morphisms and the long exact sequence of a triple

In the context of the compactification Set-up 3.4.1, with S a nonsingular curve and $s \in S$ a point, it is natural to ask about the relation between specialisation morphisms and the long exact sequence in cohomology associated with the triple (Z, X, X^o) . Because of base change issues, and because of the possible failure of the morphisms of type δ for the closed embedding a to be isomorphisms, the relation is not always the expected one, i.e. the filtered cones of the horizontal morphisms in (64) are not necessarily the corresponding objects for $i^*a^!F$ and $t^*a^!F$.

Recall Remarks 3.1.4, 3.2.4, 3.3.4 and 3.4.4 on the parts of the diagrams involved in the definitions of the specialisation morphisms sp^* , sp_P^* , $sp_{P_f}^*$.

We discuss the case of $sp_{P_f}^*$. The case sp_P^* is then the special case when $f : X \rightarrow Y$ is the identity. The case sp^* is the special case, when we remove ${}^p\tau_{\leq \bullet}$ from the picture.

LEMMA 3.5.1. *Let things be as in Set-up 3.4.1, except that we do not assume that v and f are proper; nor that W is Cartier. Assume that S is a nonsingular curve and $s \in S$ is a point. The following diagram is commutative:*

$$\begin{array}{ccccc}
 \psi[-1]v_*a_!a^!{}^p\tau_{\leq \bullet}f_* & \longrightarrow & \psi[-1]v_* & {}^p\tau_{\leq \bullet}f_* & \longrightarrow & \psi[-1]v_*b_*b^*{}^p\tau_{\leq \bullet}f_* & \rightsquigarrow \\
 \uparrow \sigma^* & & \uparrow \sigma^* & & & \uparrow \sigma^* & \\
 i^*[-1]v_*a_!a^!{}^p\tau_{\leq \bullet}f_* & \longrightarrow & i^*[-1]v_* & {}^p\tau_{\leq \bullet}f_* & \longrightarrow & i^*[-1]v_*b_*b^*{}^p\tau_{\leq \bullet}f_* & \rightsquigarrow \\
 \downarrow bc^{i^*}v_* & & \downarrow bc^{i^*}v_* & & & \downarrow bc^{i^*}v_* & \\
 v_*i^*[-1]a_!a^!{}^p\tau_{\leq \bullet}f_* & \longrightarrow & v_*i^*[-1] & {}^p\tau_{\leq \bullet}f_* & \longrightarrow & v_*i^*[-1]b_*b^*{}^p\tau_{\leq \bullet}f_* & \rightsquigarrow \\
 \downarrow i^*a_!a^! \rightarrow a_!a^!i^* & & \downarrow = & & & \downarrow bc^{i^*}b_* \rightarrow i^*b_*b^* \rightarrow b_*b^*i^* & \\
 v_*a_!a^!i^*[-1]{}^p\tau_{\leq \bullet}f_* & \longrightarrow & v_*i^*[-1] & {}^p\tau_{\leq \bullet}f_* & \longrightarrow & v_*b_*b^*i^*[-1]{}^p\tau_{\leq \bullet}f_* & \rightsquigarrow \\
 \uparrow \delta^{i^*} & & \uparrow \delta^{i^*} & & & \uparrow \delta^{i^*} & \\
 v_*a_!a^!{}^p\tau_{\leq \bullet}i^*[-1]f_* & \longrightarrow & v_*{}^p\tau_{\leq \bullet} & i^*[-1]f_* & \longrightarrow & v_*b_*b^*{}^p\tau_{\leq \bullet}i^*[-1]f_* & \rightsquigarrow \\
 \downarrow bc^{i^*}f_* & & \downarrow bc^{i^*}f_* & & & \downarrow bc^{i^*}f_* & \\
 v_*a_!a^!{}^p\tau_{\leq \bullet}f_*i^*[-1] & \longrightarrow & v_*{}^p\tau_{\leq \bullet} & f_*i^*[-1] & \longrightarrow & v_*b_*b^*{}^p\tau_{\leq \bullet}f_*i^*[-1] & \rightsquigarrow .
 \end{array} \tag{65}$$

Proof. This follows at once from the following list of formal properties.

Let things be as in Set-up 3.4.1. Recall (61). Let $i : T \rightarrow S$ be a morphism of varieties. The verification of the following facts is formal and is left to the reader.

- (i) σ^* is a morphism of functors.
- (ii) The attaching triangles are compatible with base change. More precisely, the base change morphisms $i^*v_* \rightarrow v_*i^*$ applied to the distinguished triangle of functors (61) (recall that it contains the morphisms a, b) yield morphisms of the corresponding distinguished triangles.
- (iii) Let $\mu : i^*a_!a^! \rightarrow a_!a^!i^*$ be the composition of the base change isomorphism $i^*a_!a^! \rightarrow a_!i^*a^!$ with the natural morphism $a_!i^*a^! \rightarrow a_!a^!i^*$ [Ka-Sc 1990, Proposition 3.1.9-iii]. Then $i^*a = a_i^* \circ \mu$.
- (iii)' Let $\mu' : i^*b_*b^* \rightarrow b_*b^*i^*$ be the composition of the base change isomorphism $i^*b_*b^* \rightarrow b_*i^*b^*$ with the natural isomorphism $b_*i^*b^* \rightarrow b_*b^*i^*$. Then $\mu' \circ i^*b = b_i^*i^*$.
- (iv) The attaching triangles are compatible with the morphisms of type δ . More precisely, the morphisms of type δ applied to the distinguished triangle of functors (61) yield morphisms of the corresponding distinguished triangles.
- (v) Part (ii) holds for the base change morphisms $i^*f_* \rightarrow f_*i^*$.

We need the following remark and lemma in the proof of Proposition 3.5.4.

Remark 3.5.2. The base change morphism $bc^{i^*b^*}$ in (65) and (63) coincide.

LEMMA 3.5.3. *Let things be as in the compactification Set-up 3.4.1 and assume that S is a nonsingular connected curve and $s \in S$ is a point. Let $G \in D_c^b(Y)$ be such that $\phi G = 0$ and $\phi a^!G = 0$. Then the natural morphisms $i^*a^! \tau_{\leq \bullet} G \rightarrow a^!i^* \tau_{\leq \bullet} G$ are isomorphisms.*

Proof. The assumptions on vanishing are equivalent to assuming that $\phi \tau_{\leq \bullet} G = 0$ and $\phi \tau_{\leq \bullet} a^!G = 0$, so that it is enough to prove the conclusion without truncations. We have $i^*[-1]G \cong i^![1]G$ and $i^*[-1]a^!G \cong i^![1]a^*G$. The conclusion follows from the natural identification $i^!a^! = a^!i^!$ as follows: $i^*[-1]a^!G = i^![1]a^!G = a^!i^![1]G = a^!i^*[-1]G$.

Recall Remark 3.3.5, which allows us to use a general point t to express the target of specialisation morphisms. Recall the notational Remark 3.1.8, and convention (iii) on shifts of filtrations.

THEOREM 3.5.4 (specialisation and long exact sequence of a triple). *Let things be as in the compactification Set-up 3.4.1 and assume that S is a nonsingular connected curve and $s \in S$ is a point. Let $F \in D_c^b(X)$.*

Consider the following three sets of conditions:

- (I)
 - a. f_*F has no constituents supported on W ;
 - b. f_*i^*F has no constituents supported on W_s ;
 - c. $\phi f_*F = 0$ ($\Leftarrow (\phi F = 0)$);
 - d. $\phi f_*a^!F = 0$ ($\Leftarrow (\phi a^!F = 0)$);
- (II)
 - a. f is projective;
 - b. $F, a^!F[1], i^*F[-1], a^!i^*F$ are perverse semisimple;
 - c. $\phi f_*F = 0$ ($\Leftarrow (\phi F = 0)$);
 - d. $\phi f_*a^!F = 0$ ($\Leftarrow (\phi a^!F = 0)$);

- (III) a. f is projective;
- b. $F, a^1F[1], i^*F[-1]$ and a^1i^*F are perverse semisimple;
- c. $\phi v_* b_* f_* b^* F = 0$ (cf. Remark 3.3.9) ($\Leftarrow (\phi f_* b_* b^* F = 0) \Leftarrow (\phi b_* b^* F = 0)$);
- d. $\phi f_* b^* F = 0$;
- e. $\phi f_* a^1 F = 0$.

Assume that either (I), (II), or (III) holds. Then, for $t \in S$ general, we have the isomorphism of distinguished triangles in $DF(pt)$:

$$\begin{array}{ccccc}
 (R\Gamma(Z_t, t^* a^1 F), P^{f_t}(-1)) & \longrightarrow & (R\Gamma(X_t, t^* F), P^{f_t}) & \longrightarrow & (R\Gamma(X_t^a, t^* b^* F), P^{f_t}) \rightsquigarrow \\
 \uparrow \cong & & \uparrow \cong & & \uparrow \cong \\
 sp_{p_f}^*(a^1 F) & & sp_{p_f}^*(F) & & sp_{p_f}^*(b^* F) \\
 (R\Gamma(Z_s, i^* a^1 F), P^{f_s}(-1)) & \longrightarrow & (R\Gamma(X_s, i^* F), P^{f_s}) & \longrightarrow & (R\Gamma(X_s^a, i^* b^* F), P^{f_s}) \rightsquigarrow . \\
 & & & & (66)
 \end{array}$$

Proof. We plug $F \in D_c^b(X)$ in diagram (65). We still denote the resulting diagram by (65) in what follows. Then (65) consists of three columns, C_1, C_2 and C_3 , and six rows R_1, \dots, R_6 . Each row is a distinguished triangle. The vertical arrows yield morphisms of distinguished triangles.

The aim is to prove that, under the assumptions of the theorem: all the vertical arrows in (65) are isomorphisms. In fact, then, by considering the compositum of the arrows from R_6 to R_1 , and in view of (19), we obtain the following system of isomorphisms of distinguished triangles:

$$\begin{array}{ccccc}
 v_* a_! {}^{p\tau}_{\leq +1} f_* t^*[-1] a^1 F & \longrightarrow & v_* {}^{p\tau}_{\leq \bullet} f_* t^*[-1] F & \longrightarrow & v_* b_* {}^{p\tau}_{\leq \bullet} f_* t^*[-1] b^* F \rightsquigarrow \\
 \uparrow \cong & & \uparrow \cong & & \uparrow \cong \\
 \psi[-1] v_* a_! {}^{p\tau}_{\leq +1} f_* a^1 F & \longrightarrow & \psi[-1] v_* {}^{p\tau}_{\leq \bullet} f_* F & \longrightarrow & \psi[-1] v_* b_* {}^{p\tau}_{\leq \bullet} f_* b^* F \rightsquigarrow \\
 \uparrow \cong & & \uparrow \cong & & \uparrow \cong \\
 sp^*({}^{p\tau}_{\leq +1} a^1 F) & & sp^*({}^{p\tau}_{\leq \bullet} F) & & sp^*({}^{p\tau}_{\leq \bullet} b^* F) \\
 v_* a_! {}^{p\tau}_{\leq +1} f_* i^*[-1] a^1 F & \longrightarrow & v_* {}^{p\tau}_{\leq \bullet} f_* i^*[-1] F & \longrightarrow & v_* b_* {}^{p\tau}_{\leq \bullet} f_* i^*[-1] b^* F \rightsquigarrow , \\
 & & & & (67)
 \end{array}$$

which yields the desired conclusion (66).

We first prove that (I) implies that desired goal that all the vertical arrows are isomorphisms.

The plan of the proof is as follows. Prove that columns C_1, C_2 and C_3 coincide with diagrams (58) involved in the definitions of $sp_{p_f}^*(a^1 F), sp_{p_f}^*(F)$ and $sp_{p_f}^*(b^* F)$, respectively. Prove that all arrows in said columns are isomorphisms, so that all three morphisms $sp_{p_f}^*$ are defined and isomorphisms, and the composition of the arrows from R_6 to R_1 yields (66).

Step 1. We carry out the plan for C_1 . In this case, $P^f(-1)$ will enter the picture naturally. Let us identify C_1 with (58) for $a^1 F$. In order to accomplish this, we need to show that:

- (i) the top of C_1 , i.e. $\psi[-1] v_* a_! {}^{p\tau}_{\leq \bullet} f_* F$, coincides with $\psi[-1] v_* a_! {}^{p\tau}_{\leq +1} f_* a^1 F$;
- (ii) the morphism $v_* i^*[-1] a_! a^1 {}^{p\tau}_{\leq \bullet} f_* F \rightarrow v_* a_! a^1 i^*[-1] {}^{p\tau}_{\leq \bullet} f_* F$ is an isomorphism;
- (iii) the bottom of C_1 , i.e. $v_* a_! a^1 {}^{p\tau}_{\leq \bullet} f_* i^*[-1] F$, coincides with $v_* a_! {}^{p\tau}_{\leq +1} f_* i^*[-1] a^1 F$.

To prove (i) it is enough to prove that $a^1 {}^{p\tau}_{\leq \bullet} f_* F$ coincides with ${}^{p\tau}_{\leq \bullet} f_* a^1 F$. This follows from the fact that the natural morphism (22) of type δ^{a^1} for the closed embedding a is an

isomorphism in view of our assumptions that f_*F has no constituents supported on W , so that Proposition 2.1.5 applies.

For (ii), we argue as follows. Since $i^*a_1 = a_1i^*$, it is enough to prove that $i^*a^1 p\tau_{\leq \bullet} f_*F$ coincides with $a^1 i^* p\tau_{\leq \bullet} f_*F$. This follows from our vanishing hypotheses and from Lemma 3.5.3, in view of the assumptions $\phi f_*F = 0$ and $\phi f_*a^1 F = 0$.

For (iii), we argue as we have done for (i), by using the assumption that f_*i^*F has no supports on W_s .

We have shown that C_1 agrees with (58) for a^1F . Since f and v are proper, we have that $sp_{p_f}^*(a^1F)$ is defined.

Let us now prove that all arrows in C_1 are isomorphisms.

The base change maps are isomorphisms. The assumption that $\phi F = 0$ implies $\phi f_*F = 0$, so that δ^{i^*} is an isomorphism by Lemma 2.2.1.(iii). Finally, we claim that σ^* on the top of C_1 is an isomorphism. This follows from the fact that its cone $\phi v_*a_1a^1 p\tau_{\leq \bullet} f_*F = 0$: in fact, by using the same kind of argument employed in (i) above, this is implied by the assumption $\phi f_*a^1 F = 0$.

This completes the proof of Step 1: all the vertical arrows in C_1 are isomorphisms and give rise to $sp_{p_f}^*(a^1F)$, which is an isomorphism for the shifted $P^f(-1)$'s.

Step 2. Clearly, C_2 is on the nose the collection of morphisms in (58) involved in the definition of $sp_{p_f}^*(F)$. Moreover, in view of our hypotheses, Theorem 3.3.7.(i) applies and $sp_{p_f}^*(F)$ is defined and an isomorphism. In particular, all the arrows in C_2 are isomorphisms.

Step 3: We carry out the plan for C_3 by following the template of Step 1.

Since b^* is étale, b^* is t-exact and it commutes with f_* . Clearly, b^* commutes with i^* . We thus have the identities:

$$\psi[-1]v_*b_*b^* p\tau_{\leq \bullet} f_* = \psi[-1]v_*b_* p\tau_{\leq \bullet} f_*b^*, \quad v_*b_*b^* p\tau_{\leq \bullet} f_*i^*[-1] = v_*b_* p\tau_{\leq \bullet} f_*i^*[-1]b^*. \tag{68}$$

This implies that the top and bottom of C_3 give rise to the domain and target of the putative $sp_{p_f}^*(b^*F)$ as in Step 1.

Since the morphisms of rows are morphisms of distinguished triangles and the vertical arrows in C_1 and in C_2 are isomorphisms, then, by the Five Lemma, so are the ones in C_3 .

This concludes Step 3 and we have shown that (I) implies the desired goal that all the vertical arrows are isomorphisms.

We prove that (II) implies the same desired goal.

To do so, it is enough to prove that (II) implies (I). Since (c) and (d) are common to (I) and (II), we need to show that (II) implies (Ia) and (Ib). By Proposition 2.3.2, the assumptions f projective, F and $a^1F[1]$ perverse semisimple imply (Ia). For the same reason, the assumptions f projective, $i^*F[-1]$ and a^1i^*F perverse semisimple imply (Ib). The desired goal is thus met.

Note that instead of assuming that $a^1F[1]$ (resp. a^1i^*F) is perverse semisimple, it is enough to assume that $f_*a^1F[1]$ (resp. $f_*a^1i^*F$) satisfies the conclusion of the relative Hard Lefschetz Theorem; see Remark 2.3.3.

We prove that (III) implies the desired goal that all the vertical arrows are isomorphisms.

We go back to (65). Again, the goal is to prove that the bottom and top rows are as in the proof that (I) implies the desired conclusion, and that all vertical arrows are isomorphisms.

The top and bottom of C_2 are already in the desired form.

The top and bottom of C_3 are already in the desired form; see (68), which uses only that b^* is t-exact and that it commutes with f_* .

By Remarks 3.5.2 and 3.4.3, the base change morphism $bc^{i^*b^*}$ in (65) is an isomorphism. As in the case of (II), all the morphisms of type δ , connecting R_5 to R_4 are isomorphisms. The base change morphisms for i^*f_* and i^*v_* are also isomorphisms in all columns, including C_3 .

All the arrows in C_3 are thus isomorphisms as soon as σ^* is. This follows from the hypothesis that $\phi v_* b_* f_* b^* F = 0$ as follows. Since F is semisimple, so is b^*F . The Decomposition Theorem implies that $\phi v_* b_* {}^p\tau_{\leq \bullet} f_* b^* F$ is a direct summand of $\phi v_* b_* f_* b^* F = 0$, so that $\phi v_* b_* b^* {}^p\tau_{\leq \bullet} f_* F = \phi v_* b_* {}^p\tau_{\leq \bullet} f_* b^* F = 0$, where we have used the t-exactness of b^* and proper base change. The cone of σ^* in C_3 is zero and σ^* in C_3 is an isomorphism.

We have proved that C_3 has the desired form and that all of the arrows in C_3 are isomorphisms, so that $sp_{p_f}(b^*G)$ is an isomorphism.

Since the base change arrow $bc^{i^*v_*}$ is an isomorphism in C_3 it follows that all arrows connecting R_3 to R_4 are isomorphisms. In particular, the arrow denoted by $i^*a_1 a^1 \rightarrow a_1 a^1 i^*$ is an isomorphism.

The arrow σ^* in C_1 is an isomorphism if and only if $\phi v_* a_1 a^1 {}^p\tau_{\leq \bullet} f_* F = 0$, which we now prove. As in the case of (II), we have that $f_* F$ has no constituents supported on W , so that $\phi v_* a_1 a^1 {}^p\tau_{\leq \bullet} f_* F = \phi v_* a_1 {}^p\tau_{\leq \bullet+1} f_* a^1 F = v_* a_1 {}^p\tau_{\leq \bullet+1} f_* \phi a^1 F = 0$, by the assumption $0 = \phi f_* a^1 F = f_* \phi a^1 F$ (f is proper).

What above also implies that the top of C_1 is $\psi[-1]v_* a_1 {}^p\tau_{\leq \bullet+1} f_* a^1 F$, i.e. the system giving rise to the target of the specialisation map. As to the bottom of C_1 , since all arrows are isomorphisms, it is identified with $v_* i^*[-1]a_1 a^1 {}^p\tau_{\leq \bullet} f_* F$, which, by what above, is identified with $v_* a_1 i^*[-1] {}^p\tau_{\leq \bullet+1} f_* a^1 F$, which –using the fact that $f_* a^1 F$ has no constituents supported on W_s by virtue of the hypotheses that f is projective, $a^1 F[1]$ and $i^* a^1 F$ are perverse semisimple, coupled with Proposition 2.3.2– equals $v_* a_1 {}^p\tau_{\leq \bullet+1} f_* i^*[-1] a^1 F$, i.e. the system giving rise to the source of $sp_{p_f}^*(a^1 F)$.

We have proved that C_1 has the desired form and that all arrows in C_1 are isomorphisms. We have already proved that the same holds for C_3 . As seen earlier by using the Five Lemma, it follows that the same holds for C_2 . The conclusion follows.

Remark 3.5.5. Condition (I) in Proposition 3.5.4 is met if: f is projective, F is perverse semisimple on X and $a^*[-1]F$ is perverse semisimple on Z . Condition (II) in Proposition 3.5.4 is met if: f is projective, $i^*F[-1]$ is perverse semisimple on X_s and $a^* i^*[-2]F$ is perverse semisimple on Z_s . Both cases follow from Proposition 2.3.2.

Remark 3.5.6. Proposition 3.5.4 yields a different proof, under different hypotheses, of the conclusions of both Proposition 3.4.2 and Corollary 3.4.6.

Remark 3.5.7. The toy-model for Theorem 3.5.4 is when we assume that v_X, v_Y, v_Z, v_W and f are proper and smooth. We do not assume that v_{Y° is proper. In this case, we leave to the reader to verify that if we set $F = \mathbb{Q}_X$, and we consider the usual Leray filtration on the cohomology of X_s and X_t (Z_s, Z_t , resp.), with respect to $f_s : X_s \rightarrow Y_s$ and $f_t : X_t \rightarrow Y_t$, ($f_s : Z_s \rightarrow W_s$ and $f_t : Z_t \rightarrow W_t$, resp.), then we end up with: the specialisation morphisms are defined, they are filtered isomorphisms, and they fit in the long exact Gysin sequences: (in what follows \star is arbitrary, but fixed)

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & L_{\star-2}H^{\bullet-2}(Z_t, \mathbb{Q}) & \xrightarrow{\text{Gysin}} & L_{\star}H^{\bullet}(X_t, \mathbb{Q}) & \longrightarrow & L_{\star}H^{\bullet}(X_t^o, \mathbb{Q}) \xrightarrow{+1} \dots \\
 & & \uparrow \text{sp}_Z & & \uparrow \text{sp}_X & & \uparrow \text{sp}_{X^o} \\
 \dots & \longrightarrow & L_{\star-2}H^{\bullet-2}(Z_s, \mathbb{Q}) & \xrightarrow{\text{Gysin}} & L_{\star}H^{\bullet}(X_s, \mathbb{Q}) & \longrightarrow & L_{\star}H^{\bullet}(X_s^o, \mathbb{Q}) \xrightarrow{+1} \dots
 \end{array} \tag{69}$$

where L stands for the increasing classical Leray filtration (which classically starts at zero).

4. Applications to the Hitchin morphism

In this section we apply the results of Section 3, especially equation (66) in Theorem 3.5.4 to the triples arising from the compactification of Dolbeault moduli spaces in families [de 2018]. The main result is Theorem 4.4.2, to the effect that the triple given by the Dolbeault moduli spaces over a curve, its compactification and the boundary gives rise to a specialisation morphism of long exact sequences for the (intersection) cohomology of the special and general points that is a filtered isomorphisms of triples for the perverse Leray filtrations. This is what one would obtain if we were in the oversimplified and ideal situation of a fiber bundle with boundaries, and we were to take the Leray filtrations.

The compactification [de 2018] is obtained by taking a \mathbb{G}_m -quotient; this is quickly reviewed in Section 4.2. In order to work with such quotients, we need some results on descending objects and properties along these quotients, which may be of some independent interest; see Section 4.1. Section 4.3 contains some special properties of the constant sheaf and of the intersection cohomology complex for the quotients we obtain. Finally, we put everything together in Section 4.4, where we prove Theorem 4.4.2.

4.1. Descending along \mathbb{G}_m -quotients

In this section, we assume we have two Cartesian diagrams of morphisms over a variety S as in the compactification Set-up 3.4.1:

$$\begin{array}{ccccc}
 \mathcal{X} & \xrightarrow{a} & \mathcal{X} & \xleftarrow{b} & \mathcal{X}^o \\
 \downarrow f & & \downarrow f & & \downarrow f \\
 \mathcal{W} & \xrightarrow{a} & \mathcal{Y} & \xleftarrow{b} & \mathcal{Y}^o
 \end{array}
 \qquad
 \begin{array}{ccccc}
 Z & \xrightarrow{a} & X & \xleftarrow{b} & X^o \\
 \downarrow f & & \downarrow f & & \downarrow f \\
 W & \xrightarrow{a} & Y & \xleftarrow{b} & Y^o,
 \end{array} \tag{70}$$

such that: \mathbb{G}_m -acts equivariantly on the l.h.s. with finite stabilisers; the actions cover the trivial action on S ; the r.h.s. is obtained by taking the quotient under the \mathbb{G}_m -action; the quotient morphisms π are geometric quotients.

LEMMA 4.1.1. *Each quotient morphism π above factors as $\pi = qp$, where p is a quotient S -morphism by a finite subgroup C of \mathbb{G}_m , and q is a smooth quotient S -morphism by a \mathbb{G}_m -action, so that, for example, we have:*

$$\pi : \mathcal{X} \xrightarrow{p} \mathcal{X}' := \mathcal{X} / C \xrightarrow{q} X = \mathcal{X} / \mathbb{G}_m = \mathcal{X}' / \mathbb{G}'_m (:= \mathbb{G}_m / C). \tag{71}$$

Proof. Let $C \subseteq \mathbb{G}_m$ be any finite subgroup containing all the stabilizers (recall that we are in a finite type situation). The variety $\mathcal{X}' := \mathcal{X} / C$ over S is endowed with the induced $\mathbb{G}'_m := \mathbb{G}_m / C \simeq \mathbb{G}_m$ -action, and we have diagram (71). The quotients morphism p is finite. By Luna Étale Slice Theorem, the quotient morphism q is smooth: the \mathbb{G}'_m -action has trivial stabilisers, so that q is étale locally a projection map with scheme-theoretic fiber \mathbb{G}_m .

We record the following remark for use in the proof of Lemma 4.3.3.

Remark 4.1.2. The following commutative diagrams are Cartesian in the category of topological spaces:

$$\begin{array}{ccccc}
 \mathcal{Z} & \xrightarrow{a} & \mathcal{X} & \xleftarrow{b} & \mathcal{X}^o \\
 p \downarrow & & p \downarrow & & p \downarrow \\
 \mathcal{Z}' & \xrightarrow{a} & \mathcal{X}' & \xleftarrow{b} & \mathcal{X}'^o \\
 q \downarrow & & q \downarrow & & q \downarrow \\
 Z & \xrightarrow{a} & X & \xleftarrow{b} & X^o.
 \end{array} \tag{72}$$

The same is true if we replace (Z, X, X^o) etc. with (W, Y, Y^o) etc.

LEMMA 4.1.3. (**Descending the lack of constituents on the boundary to a quotient by \mathbb{G}_m**). *Let $\mathcal{G} \in D(\mathcal{Y})$ and $G \in D_c^b(Y)$ be such that q^*G is a direct summand of $p_*\mathcal{G}$. If \mathcal{G} has no constituents supported on the boundary \mathcal{W} , then G has no constituents supported on the boundary W .*

Proof. Claim: $p_\mathcal{G}$ has no constituents supported on \mathcal{W}' .* Since p is finite, p_* is t-exact, so that p_* and ${}^pH^*$ commute. It follows that we may assume that \mathcal{G} is perverse. Let $J_\bullet\mathcal{G}$ be a Jordan–Holder filtration for \mathcal{G} , so that the non trivial graded objects $Gr_\bullet^J\mathcal{G}$ are simple, i.e. they are intersection complexes supported at irreducible closed subvarieties of \mathcal{Y} and with simple coefficients. By the t-exactness of p_* , we have that $p_*J_\bullet\mathcal{G}$ is a filtration and $Gr_\bullet^{p_*J}p_*\mathcal{G} = p_*Gr_\bullet^J\mathcal{G}$. Since p is finite, hence small, we have that p_* sends the intersection complex of a closed subvariety \mathcal{T} of \mathcal{Y} with twisted coefficients, to the intersection complex of the image $p(\mathcal{T})$ with appropriately twisted coefficients. By construction, we have that $\mathcal{W} = p^{-1}(\mathcal{W}')$. The subvarieties \mathcal{T} appearing as the supports of the simple $Gr_\bullet^J\mathcal{G}$ above are not contained in \mathcal{W} by assumption, so that their $p(\mathcal{T})$'s are not contained in \mathcal{W}' . The claim follows.

By construction, we have that $\mathcal{W}' = q^{-1}(W)$. Since q^*G is assumed to be a direct summand of $p_*\mathcal{G}$, to prove the lemma we need to show that the constituents of q^*G are $q^*[1]$ of the constituents of G . But this follows from the fact that, since q is smooth of relative dimension one with connected fibers, $q^*[1]$ is t-exact and it preserves simple objects (cf. [Be-Be-De 1982, p.106, bottom]).

We need the following general result in the proof of Lemma 4.1.5.

LEMMA 4.1.4. *Let C be a finite group acting on a variety X , let $p : X \rightarrow X'$ be the quotient morphism. Then: (i) $p_*\mathbb{Q}_X$ admits $\mathbb{Q}_{X'}$ as a direct summand; (ii) p_*IC_X admits $IC_{X'}$ as a direct summand.*

Proof. We prove (i). By [Gr 1956, (5.1.1), p.108], we have that $\mathbb{Q}_{X'} = (p_*(\mathbb{Q}_X))^C$. We split the natural adjunction morphism $\mathbb{Q}_{X'} \rightarrow p_*\mathbb{Q}_X$ by sending a section $s \in \mathbb{Q}_X(p^{-1}(U))$ to $(1/|C|) \sum_{c \in C} c^* \cdot s$.

We prove (ii). One can argue as above; we leave this to the reader. We prove a stronger statement, which may be of independent interest, namely that the conclusion remains valid

if we assume that p is a small morphism that has the property that every irreducible component of X maps onto an irreducible component of X' . Since IC_X is the direct sum of the intersection complexes of the irreducible components of X and similarly for $IC_{X'}$, we may assume that X and X' are irreducible, and that p is small and surjective. By the conditions of support/co-support characterising intersection complexes, the fact that p is small and surjective implies that p_*IC_X is an intersection complex with some twisted coefficients L . Since p is small and surjective, there is a Zariski dense open subset $j : U' \subseteq X'$ that is nonsingular, and over which p is finite and étale. Then L can be taken to be the local system on U associated with the topological covering, i.e. the direct image of the constant sheaf. Note that this splits off \mathbb{Q}_U by a standard trace argument. We now show that this splitting extends uniquely to a splitting of the corresponding intersection complexes, which proves our contention. Let $I', J' \in P(X')$ be two intersection complexes with support precisely X' . We have the standard identity: $Hom(I', J') = Hom(j^*I', j^*J')$: we have $j_*j^*J' \in {}^pD^{\geq 0}(X)$; so $Hom(I', j_*j^*J') = Hom(I', {}^pH^0(j_*j^*J'))$; we have the short exact sequence $0 \rightarrow J' \rightarrow {}^pH^0(j_*j^*J') \rightarrow Q \rightarrow 0$, where $Q \in P(X')$ has support strictly contained in X' ; we thus have $Hom(I', J') = Hom(I', {}^pH^0(j_*j^*J'))$ because $Hom(I', Q) = 0$ by reasons of support, and $Hom(I', Q[-1]) = 0$ by the axioms of t-structure. We apply the standard identity above to $I' = p_*IC_X$ and to $J' = IC_{X'}$, and we lift the splitting off of \mathbb{Q}_U from L to the splitting off of IC_X from p_*IC_X .

LEMMA 4.1.5 (**Descending $\phi = 0$ to a quotient by \mathbb{G}_m**). *Let \mathcal{X}/S be as in (70). Assume S is a nonsingular curve, and let $s \in S$ be a point. Then we have the following implications:*

- (i) *if $\phi\mathbb{Q}_{\mathcal{X}} = 0$, then $\phi\mathbb{Q}_X = 0$;*
- (ii) *if $\phi IC_{\mathcal{X}} = 0$, then $\phi IC_X = 0$;*
- (iii) *let $\mathcal{F} \in D(\mathcal{X})$, $F \in D_c^b(X)$ be such that q^*F is a direct summand of $p_*\mathcal{F}$, then: if $\phi\mathcal{F} = 0$, then $\phi F = 0$.*

Proof. We prove (iii). Since p is proper, we have that $\phi p_*\mathcal{F} = p_*\phi\mathcal{F} = 0$. It follows that, since ϕ is additive, we have that $\phi q^*F = 0$. Since q is smooth, we have $q^*\phi F = \phi q^*F = 0$. Since q is surjective, we have that $q^*\phi F = 0$ implies the desired conclusion (iii). Now, (i) and (ii) follow from (iii) coupled with Lemma 4.1.4, and the facts $q^*\mathbb{Q}_X = \mathbb{Q}_{\mathcal{X}'}$ and $q^*IC_X = IC_{\mathcal{X}'}[-1]$ (q is smooth of relative dimension one; the shifts are innocuous).

LEMMA 4.1.6. (**Descending other identities**).

- (i) *If $a^1\mathbb{Q}_{\mathcal{X}} = \mathbb{Q}_{\mathcal{X}}[-2]$, then $a^1\mathbb{Q}_X = \mathbb{Q}_Z[-2]$.*
- (ii) *If $a^1IC_{\mathcal{X}} = IC_{\mathcal{X}}[-1]$, then $a^1IC_X = IC_Z[-1]$.*

Proof. We prove (i).

We apply p_* and take invariants: $(p_*a^1\mathbb{Q}_{\mathcal{X}})^C = (a^1p_*\mathbb{Q}_{\mathcal{X}})^C = a^1(p_*\mathbb{Q}_{\mathcal{X}})^C = a^1\mathbb{Q}_{\mathcal{X}'}$; $(p_*\mathbb{Q}_{\mathcal{X}})^C = \mathbb{Q}_{\mathcal{X}'}$. We thus have that: $a^1\mathbb{Q}_{\mathcal{X}'} = \mathbb{Q}_{\mathcal{X}'}[-2]$.

Since q is smooth of relative dimension 1, we have that $q^*[2] = q^!$, so that q^* and a^1 commute. It follows that we also have that: $q^*a^1\mathbb{Q}_X = a^1\mathbb{Q}_{\mathcal{X}'} = \mathbb{Q}_{\mathcal{X}'}[-2] = q^*\mathbb{Q}_Z[-2]$, or $q^*a^1\mathbb{Q}_X[2] = q^*\mathbb{Q}_Z$. Since q is surjective, we have that $a^1\mathbb{Q}_X[2]$ is a sheaf. Since q is smooth and surjective, we have that $a^1\mathbb{Q}_X[2]$ is locally constant of rank one. Since the fibers of q are connected, we deduce that $a^1\mathbb{Q}_X[2]$ is constant and this proves (i).

We prove (ii).

By repeating the first part of the proof of part (i), we see that: $a^1IC_{\mathcal{X}'} = IC_{\mathcal{X}'}[-1]$.

By repeating the second part of the proof of part (i), we see that: $q^*a^1IC_X = q^*IC_Z[-1]$, which we re-write as $q^*[1]a^1IC_X[1] = q^*[1]IC_Z$.

By [Be-Be-De 1982, corollaire 4.1.12], $a^1IC_X[1]$ is perverse. By [Be-Be-De 1982, Proposition 4.2.5], $q^*[1]: P(Z) \rightarrow P(\mathcal{X}')$ is fully faithful, hence conservative. This implies the desired conclusion (ii).

4.2. Compactification of Dolbeault moduli spaces via \mathbb{G}_m -actions

Let us summarise the main features of the compactification in [de 2018] that we need in this section. The notation we use here is adapted to the present needs, and differs from the one in [de 2018].

Let X^o/S be the Dolbeault moduli space associated with a reductive group G and a smooth projective family over S . Let $f: X^o \rightarrow Y^o$ be the corresponding Hitchin S -morphism; it is projective.

The main construction in [de 2018] yields two diagrams as in (70), so that, in particular, X/S is a projective completion over S of X^o/S , with boundary the S -relative Cartier divisor Z , which is proper over S . Similarly for (W, Y, Y^o) . Let us give some more details. Let $o_S \rightarrow Y^o$ be the canonical section: each Y^o_s has a canonical distinguished point; e.g. if $G = GL_n$, then this point corresponds to the characteristic polynomial t^n , which in turn corresponds to nilpotent Higgs fields. We have the closed S -subvariety $f^{-1}(o_S) \subseteq X^o$. We have \mathbb{G}_m -equivariant open inclusions and equalities as follows:

$$X^o \times (\mathbb{A}^1 \setminus \{0\}) = \mathcal{X}^o \subsetneq \mathcal{X}^o \cup (X^o \setminus f^{-1}(o_S)) \times \{0\} = \mathcal{X} \subsetneq X^o \times \mathbb{A}^1, \tag{73}$$

$$Y^o \times (\mathbb{A}^1 \setminus \{0\}) = \mathcal{Y}^o \subsetneq \mathcal{Y}^o \cup (Y^o \setminus o_S) \times \{0\} = \mathcal{Y} \subsetneq Y^o \times \mathbb{A}^1, \tag{74}$$

$$\mathcal{Z} = (X^o \setminus f^{-1}(o_S)) \times \{0\} \subsetneq X^o \times \{0\}, \quad \mathcal{W} = (Y^o \setminus o_S) \times \{0\} \subsetneq Y^o \times \{0\}. \tag{75}$$

Remark 4.2.1. Note that we are not ruling out that $\mathcal{Z} = \emptyset$. In this case, our goal, i.e. Theorem 4.4.2, still holds and it is not trivial, for it states that the P^f -filtered specialisation morphism is an isomorphism, whether we take singular or intersection cohomology groups with rational coefficients. If \mathcal{Z} is not empty, it could still happen that there are points $s \in S$ such that \mathcal{Z}_s is not dense in every irreducible component of $X^o_s \times \{0\}$; this is not an issue in what follows.

4.3. Special properties of \mathbb{Q}_X and IC_X for the compactification

LEMMA 4.3.1.

$$a^1\mathbb{Q}_X = \mathbb{Q}_Z[-2], \quad a^1IC_X = IC_Z[-1]. \tag{76}$$

Proof. By virtue of Lemma 4.1.6, it is enough to show that: (i) $a^1\mathbb{Q}_{\mathcal{X}} = \mathbb{Q}_{\mathcal{X}}[-2]$, and (ii) $a^1IC_{\mathcal{X}} = IC_{\mathcal{X}}[-1]$.

In view of the open immersions $\mathcal{Z} = (X^o \setminus f^{-1}(o_S)) \times \{0\} \subseteq X^o \times \{0\}$, and $\mathcal{X} \subseteq X^o \times \mathbb{A}^1$, it is enough to prove (i,ii) with $a: \mathcal{Z} \xrightarrow{\cong} \mathcal{X}$ replaced by $a: X^o \times \{0\} \xrightarrow{\cong} X^o \times \mathbb{A}^1$. One is easily reduced to the case $a: \{0\} \xrightarrow{\cong} \mathbb{A}^1$, where the desired conclusions are trivial.

The following lemma makes precise the relation between the irreducible components of the varieties in the triple (Z, X, X^o) . Let J (resp. K , resp. J^o) be the set of irreducible components of X (resp. Z , resp. X^o).

- LEMMA 4.3.2. (i) *The function $J \rightarrow J^o, X_j \mapsto X_j \cap X^o =: X_j^o$ is well-defined and a bijection. We have that $X_j \cap Z =: Z_j$ is either empty, or an irreducible component of Z ; if $j, j' \in J$ are such that $Z_j = Z_{j'} \neq \emptyset$, then $j = j'$; in particular, this defines an injection $K \rightarrow J$, where $Z_k = X_{j(k)} \cap Z$.*
- (ii) *The natural adjunction distinguished triangle (61) $a_1 a^1 IC_X \rightarrow IC_X \rightarrow b_* b^* IC_X \rightsquigarrow$ reads as follows:*

$$\bigoplus_K a_1 IC_{Z_k}[-1] \longrightarrow \bigoplus_J IC_{X_j} \longrightarrow \bigoplus_J b_* IC_{X_j^o} \rightsquigarrow, \tag{77}$$

where the arrows are direct sum arrows (i.e. the components of each arrow with pairs of distinct indices are zero).

Proof. Since \mathcal{Z} is open in $X^o = X^o \times \{0\}$, we have that the set of irreducible component of \mathcal{Z} injects in the evident fashion into the one of X^o . Since \mathcal{X} is open and dense in $X^o \times \mathbb{A}^1$, we have that the set of irreducible component of \mathcal{X} is naturally identified with the one of $X^o \times \mathbb{A}^1$, which in turn, since \mathbb{A}^1 is irreducible, is naturally identified with the one of X^o . It follows that the desired statement holds if we replace (Z, X, X^o) with $(\mathcal{Z}, \mathcal{X}, \mathcal{X}^o)$. The first triple is obtained from the second by dividing by the action of the connected group \mathbb{G}_m ; the quotient morphism has the orbits as fibers (in fact, it is a geometric quotient). It follows that the sets of irreducible components and their mutual relations are preserved by passing to the quotient, and (i) is proved.

We have the Cartesian diagram:

$$\begin{CD} \widehat{Z} := \coprod_K Z_k @>a>> \widehat{X} := \coprod_J X_j @<b<< \widehat{X}^o := \coprod_J X_j^o \\ @VVvV @VVvV @VVvV \\ Z = \cup_K Z_k @>a>> X = \cup_J X_j @<b<< X^o = \cup_J X_j^o \end{CD} \tag{78}$$

where: v are the evident morphisms induced by the closed embeddings of the irreducible components $Z_j \rightarrow Z$ and $X_j \rightarrow X$; a (resp. b) are the evident closed (resp. open) embeddings. We have $IC_X = v_* IC_{\widehat{X}}$ and $IC_Z = v_* IC_{\widehat{Z}}$. By base change, by the additivity of adjunction morphisms, and by the identity $a^1 IC_{X_k} = IC_{Z_k}[-1]$ –which is due to Lemma 4.3.1, which remains valid for the irreducible components, by virtue of the just-proved part (i)–, we have that the distinguished triangle (61) applied to IC_X reads as (77), and (ii) is proved.

LEMMA 4.3.3. *Let S be a nonsingular curve and let $s \in S$ be a point. Let $F \in D_c^b(X)$ be either \mathbb{Q}_X , or IC_X . Then:*

$$\phi a^1 F = 0, \quad \phi F = 0, \quad \phi b_* b^* F = 0. \tag{79}$$

In particular, we have:

$$\phi v_* a_1 f_* a^1 F = 0, \quad \phi v_* f_* F = 0, \quad \phi v_* b_* f_* b^* F = 0. \tag{80}$$

Proof. We refer to [de-Ma 2018] and to [de 2018] for references.

We have the evident morphisms $\mathcal{X} \rightarrow X^o \times \mathbb{A}^1 \rightarrow X^o$. Let $F^o := b^*F = F|_{X^o}$. Let $\mathcal{F} \in D(\mathcal{X})$ be the pull back of F^o to \mathcal{X} ; if $F^o = \mathbb{Q}_{X^o}$, then $\mathcal{F} = \mathbb{Q}_{\mathcal{X}}$; if $F^o = IC_{X^o}$, then $\mathcal{F} = IC_{\mathcal{X}}[-1]$. By the Non Abelian Hodge Theorem, X^o is topologically locally trivial over S , so that $\phi F^o = 0$: this is clear when $F = \mathbb{Q}_X$, so that $F^o = \mathbb{Q}_{X^o}$; when $F = IC_X$, so that $F^o = IC_{X^o}$, we invoke the topological invariance of the intersection complex in the not necessarily irreducible context [de-Ma 2018]. Since $pr_{X^o} : X^o \times \mathbb{A}^1 \rightarrow X^o$ is smooth, we have that $\phi pr_{X^o}^* F^o = pr_{X^o}^* \phi F^o = 0$. Since \mathcal{X} is open in $X^o \times \mathbb{A}^1$, we have that $\phi \mathcal{F} = 0$. By Lemma 4.1.5.(i),(ii), applied to $\pi = qp : \mathcal{X} \rightarrow \mathcal{X}' \rightarrow X$ and \mathcal{F} and F , we see that $\phi F = 0$. Since v and f are proper, this implies that $\phi v_* f_* F = v_* f_* \phi F = 0$.

We now wish to apply Lemma 4.1.5 to $\pi = qp : \mathcal{Z} \rightarrow \mathcal{Z}' \rightarrow Z$, $a^! \mathcal{F}$ and $a^! F$. By the base change identities associated with (72), we have that $p_* a^! \mathcal{F} = a^! p_* \mathcal{F}$ (always valid) and that $q^* a^! F = a^! q^* F$ (because q is smooth). From what above, we know that $p_* \mathcal{F}$ admits $q^* F$ as a direct summand, so that $p_* a^! \mathcal{F}$ admits $q^* a^! F$ as a direct summand. We claim that $\phi a^! \mathcal{F} = 0$. Since \mathcal{Z} is open in $X^o \times \{0\}$, it is enough to show that $\phi \tilde{a}^! \mathcal{F} = 0$, where $\tilde{a} : X^o \times \{0\} \rightarrow X^o \times \mathbb{A}^1$. Since pr_{X^o} is smooth of relative dimension one, so that $pr_{X^o}^! = pr_{X^o}^*[-2]$, we see that $\tilde{a}^! \mathcal{F} = F^o[-2]$. Since we know that $\phi F^o = 0$, we see that $\phi \tilde{a}^! \mathcal{F} = 0$, as desired. By Lemma 4.1.5, we have that $\phi a^! F = 0$. Since v, a and f are proper, this implies that $\phi v_* a_* f_* a^! F = v_* a_* f_* \phi a^! F = 0$.

The remaining statements follow from what we have proved and the distinguished triangle (61): we get $\phi b_* b^* F = 0$, and then we apply again $v_* f_*$ to $\phi b_* b^* F$ to conclude.

LEMMA 4.3.4. *Let $F \in D_c^b(X)$. Then: (i) $f_* F$ has no constituents supported on W ; (ii) $f_* i^* F$ has no constituents supported on W_s .*

Proof. We have Cartesian diagrams:

$$\begin{array}{ccccc}
 \mathcal{X} & \longrightarrow & X^o \times \mathbb{A}^1 & \longrightarrow & X^o \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathcal{Y} & \longrightarrow & Y^o \times \mathbb{A}^1 & \longrightarrow & Y^o \\
 \end{array}
 \qquad
 \begin{array}{ccccc}
 \mathcal{X}_s & \longrightarrow & X_s^o \times \mathbb{A}^1 & \longrightarrow & X_s^o \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathcal{Y}_s & \longrightarrow & Y_s^o \times \mathbb{A}^1 & \longrightarrow & Y_s^o.
 \end{array}
 \tag{81}$$

Let $F^o := b^*F = F_{X^o}$. Denote by \mathcal{F} the pul-back of F^o to \mathcal{X} . By proper base change, $f_* pr_{X^o}^* F^o = pr_{X^o}^* f_* F$ has no constituent on $Y^o \times \{0\}$. In view of (75), the same is true for $f_* \mathcal{F}$. We apply Lemma 4.1.3, so that $f_* F$ has no constituents supported on W . The proof of the second assertion is identical in view of the fact that: if $s \in S$ is a point, then we have the $-_s$ -version of (70) and it enjoys similar properties; in particular, Lemma 4.1.3 applies to it.

4.4. The long exact sequence of the triple (Z, X, X^o)

In this section, we start with a smooth family of projective manifolds over a nonsingular connected curve S , so that, according to Section 4.2, we have the situation in (70), where X^o/S is the Dolbeault moduli space for the family. We fix a point $s \in S$ (the special point) and we let $t \in S$ be a general point.

[de-Ma 2018, theorem 1.1.2] proves that the specialisation morphism $sp_{P_f}^*$ in intersection cohomology for the Dolbeault moduli spaces of a smooth family of projective manifolds is an isomorphism:

$$sp_{P_f}^* : (IH^\bullet(X_s^o, \mathbb{Q}), P^{f_s}) \xrightarrow{\sim} (IH^\bullet(X_t^o, \mathbb{Q}), P^{f_t}). \quad (82)$$

This is accomplished by applying [de-Ma 2018, theorem 3.2.1, part (ii), or part (iii)]. In fact, part (ii) implies (iii); this latter seems like a more natural statement, at least when dealing with semisimple coefficients $F \in D_c^b(X)$. Since the semisimplicity is essential in what above, the methods of [de-Ma 2018] do not seem to afford the same kind of results for the non-semisimple $F = \mathbb{Q}_{X^o}$, i.e. for singular cohomology.

As it is noted in Remark 3.3.10, Theorem 3.3.7 is an amplification of [de-Ma 2018, Theorem 3.2.1] and it can thus be used to prove (82): in fact we can use any of the parts (iii,iv,v,vii) of Theorem 3.3.7 to deduce (82). However, none of these approaches yields a genuine new proof of (82) since, as it turns out, in verifying the assumptions, we end up verifying the assumptions of part (5), and thus end up using [de-Ma 2018, Theorem 3.2.1.(iii)].

In this section, we show that:

- (i) by using a good compactification of Dolbeault moduli spaces, Proposition 3.4.2 yields a new proof of (82); see Proposition 4.4.1. In fact, (82) is made more precise in (83). The proof in Proposition 4.4.1 uses Proposition 3.4.2.(A); the reader can verify that, in the case of (topologist's) intersection cohomology, one can use Proposition 3.4.2.(B) as well.
- (ii) Proposition 3.4.2 also leads to the proof of a new fact -not seemingly affordable by the methods of [de-Ma 2018], nor of Theorem 3.3.7-, namely that (83) holds for ordinary singular cohomology as well.
- (iii) The good compactification of Dolbeault moduli spaces gives rise to the main new fact proved in this section that the specialisation morphisms $sp_{P_f}^*$ for the triple (Z, X, X^o) , in (the topologist's) intersection cohomology as well as in rational singular cohomology, give rise to the isomorphism (84) of long exact sequences of Theorem 4.4.2

Recall that P^f on $H^\bullet(X, F)$ is defined by considering the system ${}^p\tau_{\leq \bullet} f_* F$.

We shall make implicit use of (iv) and (iii).

In the remainder of this section: when dealing with intersection cohomology groups, all morphisms are direct sum morphisms with respect to the canonical decomposition according to irreducible components; in Proposition 4.4.1 and Theorem 4.4.2, the horizontal arrows are well-defined without ambiguities, the vertical ones are well-defined modulo the ambiguities introduced by the action of monodromy exchanging the irreducible components. These ambiguities are harmless for the purposes of this paper; see Remark 3.1.5. Moreover, they are removed if instead of using a general point t , we use the nearby cycle functor ψ .

We state the following proposition using (hyper)cohomology; the reader should have no difficulty re-writing the stronger version of (83) in $DF(pt)$ that uses $(R\Gamma(-, -), P^f)$. Even ignoring the l.h.s. of (83), as pointed out above, this result, i.e. that the perverse filtered specialisation morphisms exists and is an isomorphism for the singular cohomology of Dolbeault moduli spaces, is new.

PROPOSITION 4.4.1. *Let $\mathcal{H}(-)$ denote either the singular cohomology, the intersection cohomology, or the topologist's intersection cohomology groups. We have the following commutative diagram of filtered finite dimensional rational vector spaces, where the horizontal arrows are the natural restriction morphisms, the vertical arrows are the well-defined perverse-filtered specialisation morphisms and they are both isomorphisms:*

$$\begin{array}{ccc}
 (\mathcal{H}^\bullet(X_t), P^{f_t}) & \longrightarrow & (\mathcal{H}^\bullet(X_t^o), P^{f_t}) \\
 \uparrow \simeq \scriptstyle sp_{P^f}^* & & \uparrow \simeq \scriptstyle sp_{P^f}^* \\
 (\mathcal{H}^\bullet(X_s), P^{f_s}) & \longrightarrow & (\mathcal{H}^\bullet(X_s^o), P^{f_s})
 \end{array} \tag{83}$$

In the case of (the topologist's) intersection cohomology, the arrows are direct sum arrows for the canonical decompositions into irreducible components (77).

Proof. By Lemma 4.3.3, coupled with Theorem 3.3.7.(i), we have that the perverse filtered specialisation morphism on the l.h.s. of (83) is defined and it is an isomorphism. We wish to apply Proposition 3.4.2.(A) and its Corollary 3.4.6. We need to verify that $\phi b_* b^* F = 0$ and that $\phi f_* b^* F = 0$. Both follow, again, from Lemma 4.3.3. This proves diagram (83). That the arrows are direct sum arrows follows from Lemma 4.3.2.

The following result identifies the filtered cone of the restriction morphism in (83). We state the following proposition using morphisms of long exact sequences in cohomology. The reader should have no difficulty re-writing the stronger version of (84) that uses $(R\Gamma(-, -), P^f)$ and distinguished triangles in $DF(pt)$.

THEOREM 4.4.2. *Let $\mathcal{H}(-)$ denote either the singular cohomology, the intersection cohomology, or the topologist's intersection cohomology groups. In what follows, for intersection cohomology, when dealing with Z_s, Z_t , replace $P^f(1)$ with P^f and $\bullet - 2$ with $\bullet - 1$. We have as isomorphism of long exact sequences of filtered morphisms:*

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & (\mathcal{H}^{\bullet-2}(Z_t), P^{f_t}(1)) & \longrightarrow & (\mathcal{H}^\bullet(X_t), P^{f_t}) & \longrightarrow & (\mathcal{H}^\bullet(X_t^o), P^{f_t}) & \longrightarrow & \dots \\
 & & \uparrow \simeq \scriptstyle sp_{P^f}^* & & \uparrow \simeq \scriptstyle sp_{P^f}^* & & \uparrow \simeq \scriptstyle sp_{P^f}^* & & \\
 \dots & \longrightarrow & (\mathcal{H}^{\bullet-2}(Z_s), P^{f_s}(1)) & \longrightarrow & (\mathcal{H}^\bullet(X_s), P^{f_s}) & \longrightarrow & (\mathcal{H}^\bullet(X_s^o), P^{f_s}) & \longrightarrow & \dots
 \end{array} \tag{84}$$

For (the topologist's) intersection cohomology, the morphisms are direct sum morphisms for the decomposition according to irreducible components (77).

Proof. The plan is to use (66) in Theorem 3.5.4, which stems from (67), and plug (76) into the result.

In order to use (66), we need to verify that the conditions (I) in Theorem 3.5.4 are met. Even if it is not necessary, let us note that conditions (II) and (III) are also met in the case where we deal with intersection cohomology via the use of IC_X ; however, they are not necessarily met if we deal with singular cohomology via the use of the non semisimple \mathbb{Q}_X .

Conditions (Ia,Ib) are met by virtue of Lemma 4.3.4. Conditions (Ic,Id) are met by virtue of Lemma 4.3.3.

Plug $F = \mathbb{Q}_X[1]$ in (67), use $i^*\mathbb{Q}_X = \mathbb{Q}_{X_s}$ and $i^*a^1\mathbb{Q}_{X_s} = \mathbb{Q}_{Z_s}[-2]$ (cf. 76), and similarly for t^* , and get the system of isomorphisms of distinguished triangles:

$$\begin{array}{ccccccc}
 p\tau_{\leq \bullet+1} f_* \mathbb{Q}_{Z_t}[-2] & \longrightarrow & p\tau_{\leq \bullet} f_* \mathbb{Q}_{X_t} & \longrightarrow & b_* p\tau_{\leq \bullet} f_* \mathbb{Q}_{X_t^o} & \rightsquigarrow & \\
 \uparrow \simeq & & \uparrow \simeq & & \uparrow \simeq & & \\
 p\tau_{\leq \bullet+1} f_* \mathbb{Q}_{Z_s}[-2] & \longrightarrow & p\tau_{\leq \bullet} f_* \mathbb{Q}_{X_s} & \longrightarrow & b_* p\tau_{\leq \bullet} f_* \mathbb{Q}_{X_s^o} & \rightsquigarrow & .
 \end{array} \tag{85}$$

Then the reader can use (iv) and (iii), to observe that $P^f(-1)$ in (66) then becomes $P^f(1)$ as in (84), which is thus proved when \mathcal{H} is singular cohomology.

Plug $F = IC_X$ in (67), use $i^*[-1]IC_X = IC_{X_s}$ and $i^*a^1IC_X = IC_{Z_s}[-1]$ (cf. 76), and similarly for t^* , and get the system of isomorphisms of distinguished triangles:

$$\begin{array}{ccccccc}
 p\tau_{\leq \bullet+1} f_* IC_{Z_t}[-1] & \longrightarrow & p\tau_{\leq \bullet} f_* IC_{X_t} & \longrightarrow & b_* p\tau_{\leq \bullet} f_* IC_{X_t^o} & \rightsquigarrow & \\
 \uparrow \simeq & & \uparrow \simeq & & \uparrow \simeq & & \\
 p\tau_{\leq \bullet+1} f_* IC_{Z_s}[-1] & \longrightarrow & p\tau_{\leq \bullet} f_* IC_{X_s} & \longrightarrow & b_* p\tau_{\leq \bullet} f_* IC_{X_s^o} & \rightsquigarrow & .
 \end{array} \tag{86}$$

Then the reader can use (iv) and (iii) to observe that $P^f(-1)$ in (66) becomes P^f as in (84) -and that $\bullet - 2$ should be replaced by $\bullet - 1$ -, which is thus proved when \mathcal{H} is intersection cohomology. That the morphisms are direct sum morphisms for the decompositions into irreducible components follows from (77).

Similarly, if we plug $\mathcal{I}C_X = \bigoplus_j IC_{X_j}[-\dim X_j]$, we end up with the desired conclusion.

Remark 4.4.3.

- (i) The results of this section hold if we replace the Dolbeault moduli space of Higgs bundles for $G = GL_n, SL_n, PGL_n$ for families of curves with their twisted counterparts of [de 2018, remark 2.1.2], where the degree and rank of the Higgs bundles are coprime.
- (ii) If $G = GL_n, SL_n$, these twisted Dolbeault moduli spaces are nonsingular, their compactifications in Section 4.2 are orbifolds, and they can be resolved with boundary a simple normal crossing divisor over the base of the family S ; see [de 2018, theorem 3.1.1.(6)]. What follows is certainly redundant in our set-up of the Hitchin morphism, but may be useful in other set-ups: the proof of Theorem 4.4.2 uses conditions (I) in Theorem 3.5.4; since we are in a simple normal crossing divisor situation, in view of Corollary 2.4.2, we could use the variant of conditions (II,III) in Theorem 3.5.4 where one only assumes that $f_*i^*F[-1]$ satisfies the conclusion of the Hard Lefschetz Theorem. We leave the details to the reader.

Acknowledgements. The author thanks: Michel Brion, Victor Ginzburg, Jochen Heinloth, Luca Migliorini, Jörg Schürmann and Geordie Williamson for useful suggestions. Special thanks to Davesh Maulik, with whom the author has discussed various preliminary versions of Theorem 4.4.2. The author, who is partially supported by N.S.F. D.M.S. Grants n. 1600515 and 1901975, would like to thank the Freiburg Research Institute for Advanced Studies for the perfect working conditions; the research leading to these results has received funding from the People Programme (Marie Curie Actions) of the European Union’s Seventh

Framework Programme (FP7/2007-2013) under REA grant agreement n. [609305]. Finally, we thank very warmly the referee for pointing out some inaccuracies in the first submission.

REFERENCES

- [Be-Be-De 1982] *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), 5-171, Astérisque, 100, (Soc. Math. France, Paris, 1982).
- [de 2012] M.A. DE CATALDO, “The perverse filtration and the Lefschetz Hyperplane Theorem, II.,” *J. Algebraic Geometry* 21 (2012), 305–345.
- [de 2017] M.A. DE CATALDO, “Decomposition theorem for semi-simples.,” *J. Singul.* 14 (2016), 194–197.
- [de 2018] M.A. DE CATALDO, “Compactification of Dolbeault moduli spaces,” to appear in *Int. Mat. Res. Nat.* [arXiv:1812.07086](https://arxiv.org/abs/1812.07086).
- [de-Ma 2018] M.A. DE CATALDO, D. MAULIK, “The perverse filtration for the Hitchin fibration is locally constant,” to appear in *Pure and Applied Mathematics Quarterly*, [arXiv:1808.02235](https://arxiv.org/abs/1808.02235).
- [de-Ma-Sh 2020] M.A. DE CATALDO, D. MAULIK, J. SHEN “Hitchin fibrations, abelian surfaces, and the P=W conjecture.,” [arXiv:1909.11885](https://arxiv.org/abs/1909.11885).
- [de-Mi 2005] M.A. DE CATALDO, L. MIGLIORINI, “The Hodge theory of algebraic maps.,” *Ann. Sci. École Norm. Sup. (4)* 38 (2005), no. 5, 693–750.
- [de-Mi 2009] M.A. DE CATALDO, L. MIGLIORINI, “The decomposition theorem, perverse sheaves and the topology of algebraic maps.,” *Bull. Amer. Math. Soc. (N.S.)* 46 (2009), no. 4, 535–633.
- [De 1972] A. GROTHENDIECK et. al, SGA 7.I., *Groupes de Monodromie en Géométrie Algébrique.*, Lecture Notes in Math. 288, (Springer-Verlag, Heidelberg 1972).
- [Ed-Gr 1998] D. EDIDIN, W. GRAHAM, “Algebraic cuts,” *Proc. Amer. Math. Soc.* 126 (1998), no. 3, 677–685.
- [El-Lê-Mi 2010] F. EL ZEIN, D.T. LÊ, L. MIGLIORINI, “A topological construction of the weight filtration,” *Manuscripta Math.* 133 (2010), no. 1-2, 173–182.
- [Gr 1956] A. GROTHENDIECK, “Sur quelques points d’algèbre homologique.” *Tôhoku Math. J. (2)* 9 (1957), 119–221.
- [Il 1971] L. ILLUSIE, *Complexe cotangent et déformations, I.*, (French) Lecture Notes in Mathematics, Vol. 239. (Springer-Verlag, Berlin-New York, 1971). xv+355 pp.
- [Il 1994] L. ILLUSIE, “Autour du théorème de monodromie locale.,” in *Périodes p-adiques* (Bures-sur-Yvette, 1988)., *Astérisque* 223 (1994), 9–57.
- [Ka-Sc 1990] M. KASHIWARA, P. SCHAPIRA, *Sheaves on manifolds. With a chapter in French by Christian Houzel.*, *Grundlehren Math. Wiss.*, 292. (Springer-Verlag, Berlin, 1990). x+512 pp.
- [Mi-Sh 2018] L. MIGLIORINI, V. SHENDE, “Higher discriminants and the topology of algebraic maps,” *Algebr. Geom.* 5 (2018), no. 1, 114–130.
- [Mo 2007] T. MOCHIZUKI, *Asymptotic behaviour of tame harmonic bundles and an application to pure twistor D-modules, I*, *Mem. Amer. Math. Soc.* 185 (2007), no. 869, xii+324 pp.
- [Sa 2005] C. SABBAH, Polarizable twistor D-modules, *Asterisque No.* 300 (2005), vi+208 pp.
- [Sa 1988] M. SAITO, “Modules de Hodge polarizables,” *Publ. Res. Inst. Math. Sci.* 24 (1988), no. 6, 849–995 (1989).
- [Sc 2003] J. SCHÜRMAN, *Topology of singular spaces and constructible sheaves*, Instytut Matematyczny Polskiej Akademii Nauk. Monografie Matematyczne (New Series), 63, (Birkhäuser Verlag, Basel, 2003). x+452 pp.
- [Wu 2019] B. WU, “A topological characterization of the middle perversity intersection complex for arbitrary complex algebraic varieties.,” *Journal of Singularities*, 20 (2020), 54–78.