

## EDGE WEIGHTING FUNCTIONS ON THE SEMITOTAL DOMINATING SET OF CLAW-FREE GRAPHS

JIE CHEN , HONGZHANG CHEN  and SHOU-JUN XU  

(Received 1 November 2023; accepted 13 November 2023; first published online 12 February 2024)

### Abstract

In an isolate-free graph  $G$ , a subset  $S$  of vertices is a *semitotal dominating set* of  $G$  if it is a dominating set of  $G$  and every vertex in  $S$  is within distance 2 of another vertex of  $S$ . The *semitotal domination number* of  $G$ , denoted by  $\gamma_{t2}(G)$ , is the minimum cardinality of a semitotal dominating set in  $G$ . Using edge weighting functions on semitotal dominating sets, we prove that if  $G \neq N_2$  is a connected claw-free graph of order  $n \geq 6$  with minimum degree  $\delta(G) \geq 3$ , then  $\gamma_{t2}(G) \leq \frac{4}{11}n$  and this bound is sharp, disproving the conjecture proposed by Zhu *et al.* [*Semitotal domination in claw-free cubic graphs*, *Graphs Combin.* **33**(5) (2017), 1119–1130].

2020 *Mathematics subject classification*: primary 05C69.

*Keywords and phrases*: claw-free graph, edge weighting function, semitotal domination.

### 1. Introduction

Domination and its variations have been extensively studied (see, for example, [1, 2, 4, 6, 11]). A subset  $D$  of vertices in a graph  $G$  is a *dominating set* of  $G$  if every vertex of  $V(G) \setminus D$  is adjacent to a vertex in  $D$ . The minimum cardinality  $\gamma(G)$  of a dominating set is called the *dominating number* of  $G$ . A subset  $D$  of vertices in a graph  $G$  is a *total dominating set* of  $G$  if every vertex of  $V(G)$  is adjacent to a vertex in  $D$ . The minimum cardinality  $\gamma_t(G)$  of a total dominating set is called the *total dominating number* of  $G$ . It is worth noting that the study of total dominating sets is meaningful only on an isolate-free graph.

Semitotal domination, introduced by Goddard *et al.* [3] in 2014, is a relaxed form of total domination. A subset  $D$  of vertices in an isolate-free graph  $G$  is a *semitotal dominating set*, abbreviated semi-TD-set, of  $G$  if it is a dominating set of  $G$  and every vertex in  $D$  is within distance 2 of another vertex of  $D$ . The *semitotal domination number* of  $G$ , denoted by  $\gamma_{t2}(G)$ , is the minimum cardinality of a semi-TD-set in  $G$ .

---

This work was funded in part by the National Natural Science Foundation of China (Grant No. 12071194) and the Chongqing Natural Science Foundation Innovation and Development Joint Fund (Municipal Education Commission) (Grant No. CSTB2022NSCQ-LZX0003).

© The Author(s), 2024. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc.




FIGURE 1. Two graphs:  $N_2$  and  $N'_2$ , where the black vertices form a minimum semi-TD-set of their respective graphs.

We refer to a minimum semi-TD-set of  $G$  as a  $\gamma_{t2}(G)$ -set. Since every total dominating set is a semi-TD-set and every semi-TD-set is a dominating set,  $\gamma(G) \leq \gamma_{t2}(G) \leq \gamma_t(G)$ . However, the semitotal domination number is very different from the domination and total domination number. For example, the total domination number cannot be compared with the matching number, while the semitotal domination number is comparable with the matching number and cannot be greater than the matching number plus one (see [7, 8]). That makes the study of semitotal domination interesting.

There is much interest in bounds for the semitotal domination number of graphs. For example, Goddard *et al.* [3] proved that if  $G$  is a connected graph of order  $n \geq 4$ , then  $\gamma_{t2}(G) \leq \frac{1}{2}n$  and proposed Conjecture 1.1 below. Henning and Marcon [9] proved that if  $G$  is a connected claw-free cubic graph of order  $n \geq 10$ , then  $\gamma_{t2}(G) \leq \frac{4}{11}n$ , and conjectured that this bound can be improved to  $\frac{1}{3}n$  if  $G \notin \{K_4, N_2\}$ , where  $N_2$  is a graph shown in Figure 1(a). This conjecture was solved by Zhu *et al.* [13] and they proposed Conjecture 1.2 below. Zhu and Liu [12] proved that Conjectures 1.1 and 1.2 hold for line graphs with minimum degree 3 and 4, respectively. In [5], Henning established the tight upper bounds on the upper semitotal domination number of a regular graph using edge weighting functions. For algorithmic aspects of semitotal domination in graphs, Henning and Pandey [10] showed the semitotal domination problem is NP-complete for planar graphs, chordal bipartite graphs and split graphs.

**CONJECTURE 1.1** [3]. If  $G \neq K_4$  is a graph of order  $n$  with minimum degree  $\delta(G) \geq 3$ , then  $\gamma_{t2}(G) \leq \frac{2}{5}n$ .

**CONJECTURE 1.2** [13]. If  $G \neq N_2$  is a connected claw-free graph of order  $n \geq 6$  with minimum degree  $\delta(G) \geq 3$ , then  $\gamma_{t2}(G) \leq \frac{1}{3}n$ .

Inspired by [5], using edge weighting functions, we establish the tight upper bound on the semitotal domination number of a connected claw-free graph with minimum degree at least 3. In Section 2, we give some basic definitions and a lemma as preliminaries. In Section 3, we prove that if  $G \neq N_2$  is a connected claw-free graph of order  $n \geq 6$  with minimum degree  $\delta(G) \geq 3$ , then  $\gamma_{t2}(G) \leq \frac{4}{11}n$ . Also, we construct a graph attaining this bound and thus disprove Conjecture 1.2.

## 2. Preliminaries

In this section, we introduce some basic definitions and a useful lemma.

Let  $G = (V(G), E(G))$  be a connected finite simple undirected graph with vertex set  $V(G)$  and edge set  $E(G)$  of order  $n = |V(G)|$ . For a vertex  $v \in V(G)$ , we denote by  $N_G(v) = \{u \in V(G) \mid uv \in E(G)\}$  the *neighbourhood* of  $v$  and by  $N_G[v] = N_G(v) \cup \{v\}$  the *closed neighbourhood* of  $v$ . The *degree* of  $v$  is  $d_G(v) = |N_G(v)|$  and the number  $\delta(G) = \min\{d_G(v) \mid v \in V(G)\}$  is the *minimum degree* of  $G$ . We call a path connecting vertices  $u$  and  $v$  a  $(u, v)$ -*path*. The *distance*  $d_G(u, v)$  between  $u$  and  $v$  is the length of a shortest  $(u, v)$ -path in  $G$ . For a subset  $S$  of  $V(G)$ , we denote by  $N_S(v)$  the neighbourhood of  $v$  restricted on  $S$  and by  $G[S]$  the subgraph of  $G$  induced by  $S$ , while the graph  $G - S$  is the graph obtained from  $G$  by deleting the vertices in  $S$  and all edges incident with  $S$ . A graph is *claw-free* if it does not contain the complete bipartite graph  $K_{1,3}$  as an induced subgraph. If there is no confusion, then the subscript  $G$  is omitted in the notation, such as  $N(v)$ ,  $d(v)$ ,  $d(u, v)$  and so on.

Now consider  $S_1$  and  $S_2$  which are two disjoint subsets of  $V(G)$ . Let  $E[S_1, S_2] = \{u_1u_2 \mid u_1 \in S_1 \text{ and } u_2 \in S_2\}$ . For a vertex  $v$  of  $S$ , the  $S$ -*external private neighbourhood* of  $v$ , denoted by  $e_{pn}(v, S)$ , is the set of all vertices in  $V(G) \setminus S$  that are adjacent to  $v$  but to no other vertex of  $S$ . In other words, if  $u \in e_{pn}(v, S)$ , then  $u \in V(G) \setminus S$  and  $N_G(u) \cap S = \{v\}$ . The  $S$ -*internal private 2-neighbourhood* of  $v$ , denoted by  $ipn_2(v, S)$ , is the set of all vertices in  $S \setminus \{v\}$  that are within distance 2 of  $v$  in  $G$  but at a distance greater than 2 from every other vertex of  $S$ . In other words, if  $u \in ipn_2(v, S)$ , then  $u \in S \setminus \{v\}$ ,  $d(v, u) \leq 2$  and  $d(u, w) > 2$  for any vertex  $w \in S \setminus \{u, v\}$ .

A semi-TD-set in a graph  $G$  is a *minimal semi-TD-set* if it contains no semi-TD-set of  $G$  as a proper subset. The following result in [8] provides a characterisation of minimal semi-TD-sets.

**LEMMA 2.1** [8]. *Let  $S$  be a semi-TD-set in a graph  $G$ . Then,  $S$  is a minimal semi-TD-set of  $G$  if and only if every vertex  $v \in S$  satisfies at least one of the following three properties:*

- (a) *the vertex  $v$  is isolated in  $G[S]$ ;*
- (b)  *$ipn_2(v, S) \neq \emptyset$ ;*
- (c)  *$e_{pn}(v, S) \neq \emptyset$ .*

## 3. Main result

In this section, we establish the tight upper bound on the semitotal domination number of a connected claw-free graph with minimum degree at least 3 using edge weighting functions. Before that, we define two graphs  $N_2$  and  $N'_2$  as in Figure 1. Note that  $N'_2$  is a graph attaining the bound of Theorem 3.1. This shows that Conjecture 1.2 is not true.

**THEOREM 3.1.** *If  $G \neq N_2$  is a connected claw-free graph of order  $n \geq 6$  with minimum degree  $\delta(G) \geq 3$ , then  $\gamma_{12}(G) \leq \frac{4}{11}n$ , and this bound is sharp.*

**PROOF.** Suppose that the theorem is false. Let  $G$  be a counterexample such that  $|V(G)|$  is as small as possible. By the choice of  $G$ ,  $G \neq N_2$  is a connected claw-free graph of order  $n \geq 6$  with  $\delta(G) \geq 3$  such that  $\gamma_{12}(G) > \frac{4}{11}n$ , and any connected claw-free graph  $G' \neq N_2$  of order  $n' < n$  with  $\delta(G') \geq 3$  has  $\gamma_{12}(G') \leq \frac{4}{11}n'$ , where  $n' \geq 6$ .

For a  $\gamma_{12}(G)$ -set  $S$ , set  $\bar{S} = V(G) \setminus S$ . Define sets  $A^S = \{v \in S \mid ipn_2(v, S) \neq \emptyset\}$ ,  $A_1^S = \{v \in A^S \mid v \in ipn_2(v', S) \text{ for some vertex } v' \in S\}$  and  $A_2^S = A^S \setminus A_1^S$ . Let  $v \in A_1^S$  and  $v \in ipn_2(v', S)$ . Then  $v'$  is the only vertex of  $S$  within distance 2 from  $v$  in  $G$ . Since  $ipn_2(v, S) \neq \emptyset$ ,  $v' \in ipn_2(v, S)$  and  $v' \in A_1^S$ . Further,  $ipn_2(v, S) = \{v'\}$  and  $ipn_2(v', S) = \{v\}$ . This implies that the vertices in  $A_1^S$  are paired off. For each vertex  $u \in A_2^S$ , let  $S_u = ipn_2(u, S) \cup \{u\}$ . If  $u' \in ipn_2(u, S)$ , then  $u' \notin A^S$ . Otherwise,  $u' \in A^S$  and  $u \in ipn_2(u', S)$ , for  $u$  is the only vertex of  $S$  within distance 2 from  $u'$  in  $G$ , which contradicts the fact that  $u \in A_2^S$ . We note that if  $u_1$  and  $u_2$  are two distinct vertices in  $A_2^S$ , then  $ipn_2(u_1, S) \cap ipn_2(u_2, S) = \emptyset$ . Hence,  $S_{u_1} \cap S_{u_2} = \emptyset$  for each pair of different vertices  $u_1, u_2 \in A_2^S$ . Let  $B^S = \bigcup_{u \in A_2^S} S_u$  and  $C^S = S \setminus (A_1^S \cup B^S)$ . Further, we partition  $C^S$  into three subsets:  $C_0^S = \{z \mid z \in C^S \text{ and } |epn(z, S)| = 0\}$ ,  $C_1^S = \{z \mid z \in C^S \text{ and } |epn(z, S)| = 1\}$ , and  $C_2^S = \{z \mid z \in C^S \text{ and } |epn(z, S)| \geq 2\}$ . Then  $S = A_1^S \cup B^S \cup C_0^S \cup C_1^S \cup C_2^S$ .

In particular, a vertex  $u$  of  $A_2^S$  is special if  $|S_u| = 2$ ,  $d(u') = 3$  and  $|N_{\bar{S}}(u) \setminus N_{\bar{S}}(u')| = 1$ , where  $\{u'\} = S_v \setminus \{u\}$ . Further, we define sets  $A_2^S = \{u \mid u \in A_2^S \text{ and } u \text{ is special}\}$  and  $C_i^S = \{z \mid z \in C_i^S \text{ and } d(z) = 3\}$  for  $i \in \{0, 1\}$ .

A diamond in  $G$  is an induced graph of  $G$  isomorphic to  $K_4 - e$ . We call a diamond of  $G$  a special diamond if each of its vertices has degree 3 in  $G$ . Let  $\mathcal{D}$  be the set of vertices in a special diamond. Among all  $\gamma_{12}(G)$ -set, we choose a  $\gamma_{12}(G)$ -set  $S$  satisfying the following conditions:

- (1) the number of edges in  $G[S]$ , denoted by  $\lambda(S)$ , is minimised;
- (2) subject to condition (1),  $|\mathcal{D} \cap S|$  is minimised;
- (3) subject to condition (2),  $|C_0^S|$  is minimised;
- (4) subject to condition (3),  $|C_0^S|$  is minimised;
- (5) subject to condition (4),  $|C_1^S|$  is minimised.

We prove the following claim about the set  $S$ . □

*Claim 1.*  $S$  is an independent set of  $G$ .

Suppose to the contrary that there exist two adjacent vertices  $v_1$  and  $v_2$  in  $S$ . If  $epn(v_1, S) \neq \emptyset$  or  $epn(v_2, S) \neq \emptyset$ , then without loss of generality, consider  $epn(v_1, S) \neq \emptyset$ . Let  $x_1$  be a vertex in  $epn(v_1, S)$ . Since  $G$  is claw-free, each vertex of  $N(v_1) \setminus \{x_1, v_2\}$  is adjacent to either  $x_1$  or  $v_2$ . Thus,  $S_1 = (S \setminus \{v_1\}) \cup \{x_1\}$  is a  $\gamma_{12}(G)$ -set. However,  $\lambda(S_1) < \lambda(S)$ , for  $x_1$  is adjacent to no vertex of  $S \setminus \{v_1\}$ , which contradicts the choice of  $S$ . Hence,  $epn(v_1, S) = \emptyset$  and  $epn(v_2, S) = \emptyset$ . By Lemma 2.1,  $ipn_2(v_1, S) \neq \emptyset$  and  $ipn_2(v_2, S) \neq \emptyset$ .

If  $ipn_2(v_1, S) \neq \{v_2\}$ , then there exists a vertex  $v_3 \in ipn_2(v_1, S) \setminus \{v_2\}$ . Combined with  $v_1v_2 \in E(G)$ ,  $d(v_1, v_3) = 2$ . As  $v_3 \in ipn_2(v_1, S)$ , any vertex of  $N(v_3)$  belongs to  $\bar{S}$  and is

adjacent to no vertex of  $S \setminus \{v_1, v_3\}$ . Let  $x_2$  be a vertex connecting  $v_1$  and  $v_3$ . Then  $x_2v_2 \notin E(G)$ . Since  $G$  is claw-free, each vertex of  $N(v_1) \setminus \{x_2, v_2\}$  is adjacent to either  $x_2$  or  $v_2$ . When all vertices of  $N(v_3) \setminus \{x_2\}$  are adjacent to  $x_2$ ,  $(S \setminus \{v_1, v_3\}) \cup \{x_2\}$  is a semi-TD-set of  $G$ , which contradicts the minimality of  $S$ . However, when there exists a vertex  $x_3 \in N(v_3) \setminus \{x_2\}$  such that  $x_2x_3 \notin E(G)$ , each vertex of  $N(v_3) \setminus \{x_2, x_3\}$  is adjacent to either  $x_2$  or  $x_3$  as  $G$  is claw-free. Then  $S_1 = (S \setminus \{v_1, v_3\}) \cup \{x_2, x_3\}$  is a  $\gamma_{t2}(G)$ -set. But,  $\lambda(S_1) < \lambda(S)$ , which is a contradiction.

Hence,  $ipn_2(v_1, S) = \{v_2\}$ . Similarly,  $ipn_2(v_2, S) = \{v_1\}$ . Further, all vertices in  $N_{\bar{S}}(v_1) \cup N_{\bar{S}}(v_2)$  are adjacent to no vertex of  $S \setminus \{v_1, v_2\}$ . Recall that  $epn(v_1, S) = \emptyset$  and  $epn(v_2, S) = \emptyset$ . Thus,  $N_{\bar{S}}(v_1) = N_{\bar{S}}(v_2)$ . Since  $n \geq 6$ ,  $\gamma_{t2}(G) \geq \frac{4}{11}n > 2$ . This implies that  $\{v_1, v_2\}$  is not a  $\gamma_{t2}(G)$ -set. As  $G$  is connected, there exists a vertex  $x_4$  in  $\bar{S} \setminus N_{\bar{S}}(v_1)$  such that  $x_4$  is adjacent to a vertex  $x_5$  in  $N_{\bar{S}}(v_1)$ . We note that  $x_4$  has a neighbour in  $S \setminus \{v_1, v_2\}$ . When all vertices of  $N_{\bar{S}}(v_1)$  are adjacent to  $x_5$ ,  $S_1 = (S \setminus \{v_1, v_2\}) \cup \{x_5\}$  is a semi-TD-set of  $G$  with  $|S_1| < |S|$ , which is a contradiction. When there exists a vertex  $x_6 \in N_{\bar{S}}(v_1)$  such that  $x_5x_6 \notin E(G)$ , each vertex of  $N_{\bar{S}}(v_1) \setminus \{x_5, x_6\}$  is adjacent to either  $x_5$  or  $x_6$  as  $G$  is claw-free. Then  $S_1 = (S \setminus \{v_1, v_2\}) \cup \{x_5, x_6\}$  is a  $\gamma_{t2}(G)$ -set with  $\lambda(S_1) < \lambda(S)$ , which is a contradiction. This completes the proof of Claim 1.

Combining Claim 1 and the claw-freeness of  $G$ , we see that  $x$  has at most two neighbours in  $S$  for any vertex  $x$  of  $\bar{S}$ . We define an edge weighting function  $w$  on  $G: [\bar{S}, S] \rightarrow [0, 1]$ . For each vertex  $x \in \bar{S}$ , the function  $w$  assigns weight for each edge  $e \in [\{x\}, S]$  as follows.

- If  $x$  is an  $S$ -external private neighbour, then for the unique edge  $e \in [\{x\}, S]$ ,  $w(e) = 1$ .
- If  $x$  is not an  $S$ -external private neighbour and  $N_{C_0^S}(x) = \emptyset$ , then  $w(e) = \frac{1}{2}$  for each edge  $e \in [\{x\}, S]$ .
- Assume that  $x$  is not an  $S$ -external private neighbour and  $N_{C_0^S}(x) \neq \emptyset$ . Let  $N_{\bar{S}}(x) = \{y_1, y_2\}$ , where  $y_1 \in N_{C_0^S}(x)$ . It follows from the partition of  $S$  that  $y_2 \in A_2^S \cup C_0^S \cup C_1^S \cup C_2^S$ .
  - If  $y_2 \in A_2^S \cup C_0^S$ , then  $w(xy_1) = w(xy_2) = \frac{1}{2}$ .
  - If either  $y_2 \in (A_2^S \setminus A_2^S) \cup (C_1^S \setminus C_1^S)$ , or  $y_2 \in C_1^S$  and  $|\{u \mid u \in N_{\bar{S}}(y_2) \text{ and } N_{C_0^S}(u) \neq \emptyset\}| = 1$ , then  $w(xy_1) = \frac{3}{4}$  and  $w(xy_2) = \frac{1}{4}$ .
  - If either  $y_2 \in C_0^S \setminus C_0^S$  and  $|\{u \mid u \in N_{\bar{S}}(y_2) \text{ and } N_{C_0^S}(u) \neq \emptyset\}| \leq 2$ , or  $y_2 \in C_1^S$  and  $|\{u \mid u \in N_{\bar{S}}(y_2) \text{ and } N_{C_0^S}(u) \neq \emptyset\}| = 2$ , then  $w(xy_1) = \frac{5}{8}$  and  $w(xy_2) = \frac{3}{8}$ .
  - If  $y_2 \in C_0^S \setminus C_0^S$  and  $|\{u \mid u \in N_{\bar{S}}(y_2) \text{ and } N_{C_0^S}(u) \neq \emptyset\}| \geq 3$ , then  $w(xy_1) = \frac{9}{16}$  and  $w(xy_2) = \frac{7}{16}$ .
  - If  $y_2 \in C_2^S$ , then  $w(xy_1) = 1$  and  $w(xy_2) = 0$ .

From the definition of the edge weighting functions, the sum of the weights assigned to the edges joining  $x$  to  $S$  is 1. For any subset  $S_1$  of  $S$ , we define a weighting function  $f$  on  $S_1$  with  $f(S_1) = \sum_{e \in [\bar{S}, S_1]} w(e)$ . We prove the following claims.

**Claim 2.**  $f(A_1^S) > \frac{7}{4}|A_1^S|$ .

Recall that the vertices in  $A_1^S$  are paired off. Let  $v_1$  and  $v_2$  be a pair of vertices in  $A_1^S$ . Then  $ipn_2(v_1, S) = \{v_2\}$  and  $ipn_2(v_2, S) = \{v_1\}$ . This implies that all vertices in  $N_{\bar{S}}(v_1) \cup N_{\bar{S}}(v_2)$  are adjacent to no vertex of  $S \setminus \{v_1, v_2\}$ . Further, we have  $f(\{v_1, v_2\}) = |N_{\bar{S}}(v_1) \cup N_{\bar{S}}(v_2)|$ . Combining Claim 1 and  $\delta(G) \geq 3$ , we have  $|N_{\bar{S}}(v_1) \cup N_{\bar{S}}(v_2)| \geq 3$ . If  $|N_{\bar{S}}(v_1) \cup N_{\bar{S}}(v_2)| = 3$ , then  $N_{\bar{S}}(v_1) = N_{\bar{S}}(v_2)$  and  $|N_{\bar{S}}(v_1)| = 3$ . In this case,  $n = 5$  as  $G$  is claw-free and  $G$  is connected, which is a contradiction. Thus,  $|N_{\bar{S}}(v_1) \cup N_{\bar{S}}(v_2)| \geq 4$ . Hence  $f(\{v_1, v_2\}) \geq 4 > \frac{7}{2}$  and then  $f(A_1^S) > \frac{7}{4}|A_1^S|$ .

**Claim 3.**  $f(B^S) \geq \frac{7}{4}|B^S|$ .

Note that  $B^S = \bigcup_{u \in A_2^S} S_u$  and  $S_u \cap S_{u'} = \emptyset$  for any two different vertices  $u, u' \in A_2^S$ . We show that for any vertex  $u_1$  of  $A_2^S$ ,  $f(S_{u_1}) \geq \frac{7}{4}|S_{u_1}|$ . Let  $S_{u_1} = \{u_1, \dots, u_r\}$ , where  $r = |S_{u_1}| \geq 2$ . Since  $\{u_2, \dots, u_r\} \subseteq ipn_2(u_1)$ , all neighbours of  $u_i$  in  $\bar{S}$  are adjacent to no vertex of  $S \setminus \{u_1, u_i\}$ , where  $i \in \{2, \dots, r\}$ . Combined with Claim 1,  $f(S_{u_1}) \geq \sum_{i \in \{2, \dots, r\}} d(u_i)$ . If  $u_1 \in A_2^S$ , then  $f(S_{u_1}) = f(\{u_1, u_2\}) = w(x_1 u_1) + d(u_2) = w(x_1 u_1) + 3$ , where  $\{x_1\} = N_{\bar{S}}(u_1) \setminus N_{\bar{S}}(u_2)$ . Since  $u_1 \notin ipn_2(u_2, S)$ ,  $x_1$  has a neighbour in  $S$  other than  $u_1$ . Thus,  $w(x_1 u_1) = \frac{1}{2}$ . Further,  $f(\{u_1, u_2\}) = \frac{7}{2}$  and  $f(S_{u_1}) = \frac{7}{4}|S_{u_1}|$ , as desired. Thus, we may assume that  $u_1 \in A_2^S \setminus A_2^S$ . Then either  $r \geq 3$ , or  $r = 2$  and  $d(u_2) \geq 4$ , or  $r = 2$  and  $d(u_2) = 3$  and  $|N_{\bar{S}}(u_1) \setminus N_{\bar{S}}(u_2)| \geq 2$ .

If  $r \geq 3$ , then  $3r - 3 > \frac{7}{4}r$ . Since  $\delta(G) \geq 3$ ,  $f(S_{u_1}) \geq \sum_{i \in \{2, \dots, r\}} d(u_i) \geq 3(r - 1) = 3r - 3$ . Further,  $f(S_{u_1}) > \frac{7}{4}r$ . When  $r = 2$  and  $d(u_2) \geq 4$ ,  $f(S_{u_1}) = f(\{u_1, u_2\}) \geq d(u_2) \geq 4 > \frac{7}{4}r$ . When  $r = 2$ ,  $d(u_2) = 3$  and  $|N_{\bar{S}}(u_1) \setminus N_{\bar{S}}(u_2)| \geq 2$ , let  $x_1$  be a vertex in  $N_{\bar{S}}(u_1) \setminus N_{\bar{S}}(u_2)$ . From the definition of the edge weighting functions, we have  $w(x_1 u_1) \geq \frac{1}{4}$ . Thus,  $f(S_{u_1}) = f(\{u_1, u_2\}) \geq 2w(x_1 u_1) + d(u_2) \geq \frac{1}{2} + 3 = \frac{7}{2} \geq \frac{7}{4}r$ . This completes the proof of Claim 3.

**Claim 4.**  $f(C_0^S \setminus C_0^S) \geq \frac{7}{4}|C_0^S \setminus C_0^S|$ .

Let  $z_1$  be a vertex in  $C_0^S \setminus C_0^S$  and let  $N_{\bar{S}}(z_1) = \{x_1, \dots, x_r\}$ , where  $r \geq 4$ . If we have  $|\{x \mid x \in N_{\bar{S}}(z_1) \text{ and } N_{C_0^S}(x) \neq \emptyset\}| \leq 2$ , then  $|\{x \mid x \in N_{\bar{S}}(z_1) \text{ and } N_{C_0^S}(x) = \emptyset\}| \geq r - 2 \geq 2$ . Without loss of generality, consider  $N_{C_0^S}(x_1) = \emptyset$  and  $N_{C_0^S}(x_2) = \emptyset$ . By the definition of the edge weighting functions,  $w(x_1 z_1) = \frac{1}{2}$ ,  $w(x_2 z_1) = \frac{1}{2}$  and  $w(x_i z_1) \geq \frac{3}{8}$  for any  $i \in \{3, \dots, r\}$ . Hence,  $f(\{z_1\}) \geq w(x_1 z_1) + w(x_2 z_1) + \sum_{i \in \{3, \dots, r\}} w(x_i z_1) \geq 1 + \frac{3}{8}(r - 2) \geq \frac{7}{4}$ . When  $|\{x \mid x \in N_{\bar{S}}(z_1) \text{ and } N_{C_0^S}(x) \neq \emptyset\}| \geq 3$ ,  $w(x_i z_1) \geq \frac{7}{16}$  for any  $i \in \{1, \dots, r\}$ . Then  $f(\{z_1\}) \geq \frac{7}{16}r \geq \frac{7}{4}$ . In all cases, we have  $f(\{z_1\}) \geq \frac{7}{4}$ . Therefore,  $f(C_0^S \setminus C_0^S) \geq \frac{7}{4}|C_0^S \setminus C_0^S|$ .

**Claim 5.**  $f(C_1^S) \geq \frac{7}{4}|C_1^S|$ .

Let  $z_1$  be a vertex in  $C_1^S$  and  $N_{\bar{S}}(z_1) = \{x_1, x_2, \dots, x_r\}$ , where  $\{x_1\} = epn(z_1, S)$  and  $r \geq 3$ . According to the definition of the edge weighting functions,  $w(x_1 z_1) = 1$ . When

$z_1 \in C_1^S \setminus C_1^S$ , we have  $r \geq 4$  and  $w(x_i z_1) \geq \frac{1}{4}$  for any  $i \in \{2, \dots, r\}$ . Thus,  $f(\{z_1\}) = w(x_1 z_1) + \sum_{i \in \{2, \dots, r\}} w(x_i z_1) \geq 1 + \frac{1}{4}(r - 1) \geq \frac{7}{4}$ . When  $z_1 \in C_1^S$  and either  $N_{C_0^S}(x_2) = \emptyset$  or  $N_{C_0^S}(x_3) = \emptyset$ , without loss of generality, we can take  $N_{C_0^S}(x_2) = \emptyset$ . Then  $w(x_2 z_1) = \frac{1}{2}$  and  $w(x_3 z_1) \geq \frac{1}{4}$ . Further,  $f(z_1) = w(x_1 z_1) + w(x_2 z_1) + w(x_3 z_1) \geq 1 + \frac{1}{2} + \frac{1}{4} = \frac{7}{4}$ . When both  $z_1 \in C_1^S$  and  $N_{C_0^S}(x_2) \neq \emptyset$  and  $N_{C_0^S}(x_3) \neq \emptyset$ , we have  $w(x_2 z_1) = w(x_3 z_1) = \frac{3}{8}$ . Further,  $f(\{z_1\}) = w(x_1 z_1) + w(x_2 z_1) + w(x_3 z_1) = \frac{7}{4}$ . In both cases,  $f(\{z_1\}) \geq \frac{7}{4}$ . Therefore,  $f(C_1^S) \geq \frac{7}{4}|C_1^S|$ .

**Claim 6.**  $f(C_2^S) > \frac{7}{4}|C_2^S|$ .

Let  $z_1$  be a vertex in  $C_2^S$  and  $x_1, x_2$  be two vertices in  $epn(z_1, S)$ . Then  $w(x_1 z_1) = w(x_2 z_1) = 1$  and further  $f(\{z_1\}) \geq 2$ . Hence,  $f(C_2^S) \geq 2|C_2^S| > \frac{7}{4}|C_2^S|$ .

If  $f(C_0^S) \geq \frac{7}{4}|C_0^S|$ , then  $f(S) \geq \frac{7}{4}|S|$  by Claims 2–6. From the definition of the edge weighting functions,  $f(S) = n - |S|$ . It follows that  $|S| \leq \frac{4}{11}n$ , which is a contradiction. Thus,  $f(C_0^S) < \frac{7}{4}|C_0^S|$  and there exists a vertex  $y_1 \in C_0^S$  such that  $f(\{y_1\}) < \frac{7}{4}$ . Suppose that  $N_S(y_1) = \{x_1, x_2, x_3\}$  and  $N_S(x_i) = \{y_1, y_{i+1}\}$  for any  $i \in \{1, 2, 3\}$ . If  $\{y_2, y_3, y_4\} \cap ((A_2^S \setminus A_2^S) \cup (C_1^S \setminus C_1^S) \cup C_2^S) \neq \emptyset$ , then at least one edge of  $\{x_1 y_1, x_2 y_1, x_3 y_1\}$  has a weight of at least  $\frac{3}{4}$ . Further,  $f(\{y_1\}) = w(x_1 y_1) + w(x_2 y_1) + w(x_3 y_1) \geq \frac{3}{4} + \frac{1}{2} + \frac{1}{2} \geq \frac{7}{4}$ , which is a contradiction. Thus,  $\{y_2, y_3, y_4\} \cap ((A_2^S \setminus A_2^S) \cup (C_1^S \setminus C_1^S) \cup C_2^S) = \emptyset$ . Next, we prove two claims about the set  $\{y_2, y_3, y_4\}$ .

**Claim 7.**  $\{y_2, y_3, y_4\} \cap A_2^S = \emptyset$ .

In contrast, we may assume that  $y_2 \in A_2^S$ . Let  $S_{y_2} = \{y_2, y_5\}$  and  $N(y_5) = \{x_4, x_5, x_6\}$ , where  $x_4$  is a vertex connecting  $y_2$  and  $y_5$ . According to the definition of  $A_2^S$ ,  $N(y_2) \subseteq \{x_1, x_4, x_5, x_6\}$ . Note that  $ipn_2(y_1, S) = \emptyset$  and  $epn(y_1, S) = \emptyset$ . If  $d(x_1, y_5) \leq 2$ , then  $(S \setminus \{y_1, y_2\}) \cup \{x_1\}$  is a semi-TD-set of  $G$ , which contradicts the minimality of  $S$ . Thus,  $d(x_1, y_5) \geq 3$  which implies that  $x_1 x_i \notin E(G)$  for any  $i \in \{4, 5, 6\}$ . Combining  $\delta(G) \geq 3$  and the claw-freeness of  $G$ , we have  $x_1 x_2 \in E(G)$  or  $x_1 x_3 \in E(G)$  and  $x_4 x_5 \in E(G)$  or  $x_4 x_6 \in E(G)$ . Without loss of generality, consider  $x_1 x_2 \in E(G)$  and  $x_4 x_5 \in E(G)$ .

If  $x_4 x_6 \in E(G)$ , then  $S_1 = (S \setminus \{y_1, y_2, y_5\}) \cup \{x_1, x_4\}$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $x_4 x_6 \notin E(G)$ . Since  $G$  is claw-free,  $y_2 x_6 \notin E(G)$ . Then  $y_2 x_5 \in E(G)$  as  $N(y_2) \subseteq \{x_1, x_4, x_5, x_6\}$  and  $d(y_2) \geq 3$ . Further,  $d(y_2) = 3$ . If  $x_5 x_6 \in E(G)$ , then  $S_1 = (S \setminus \{y_1, y_2, y_5\}) \cup \{x_1, x_5\}$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $x_5 x_6 \notin E(G)$ . Since  $G$  is claw-free,  $N(x_4) = \{y_2, y_5, x_5\}$  and  $N(x_5) = \{y_2, y_5, x_4\}$ . We note that  $d(x_4) = d(x_5) = 3$  and then  $G[\{y_2, y_5, x_4, x_5\}]$  is a special diamond.

Since  $y_5 \in ipn_2(y_2, S)$  and  $x_6 y_2 \notin E(G)$ ,  $x_6$  is adjacent to no vertex of  $S \setminus \{y_5\}$ . If  $x_6$  is not in a special diamond, then  $S_1 = (S \setminus \{y_2, y_5\}) \cup \{x_4, x_6\}$  is a  $\gamma_{12}(G)$ -set with  $\lambda(S_1) = \lambda(S)$  but with  $|\mathcal{D} \cap S_1| < |\mathcal{D} \cap S|$ , which contradicts our choice of  $S$ . Thus,  $x_6$  is in a special diamond  $D$ . Observe that  $x_6 x_2 \notin E(G)$  and  $x_6 x_3 \notin E(G)$ . Let  $V(D) = \{x_6, x_7, x_8, y_6\}$ , where  $x_6 y_6$  is the missing edge in the special diamond  $D$ .

Then  $\{x_7, x_8\} \subseteq \bar{S}$ . To dominate  $x_7$  and  $x_8$ , we must have  $y_6 \in S$ . If  $y_6x_2 \in E(G)$ , then  $y_6 \in ipn_2(y_1, S)$  which contradicts  $y_1 \in C_0^S$ . Thus,  $y_6x_2 \notin E(G)$ . Similarly,  $y_6x_3 \notin E(G)$ .

Let  $N(y_6) = \{x_7, x_8, x_9\}$ . We note that  $y_6 \in ipn_2(v, S)$  for some vertex  $v \in S$  so call  $v = y_7$ . Then  $y_7x_9 \in E(G)$ . Recall that  $\{y_3, y_4\} \cap (A_2^S \setminus A_2^S) = \emptyset$ . If  $y_7x_2 \in E(G)$  or  $y_7x_3 \in E(G)$ , then  $y_7 = y_3$  or  $y_4$  and  $y_7 \in A_2^S \setminus A_2^S$ , which is a contradiction. Thus,  $y_7x_2 \notin E(G)$  and  $y_7x_3 \notin E(G)$ . If  $y_7 \notin ipn_2(y_6, S)$ , then  $S_1 = (S \setminus \{y_1, y_2, y_6\}) \cup \{x_1, x_7\}$  is a semi-TD-set of  $G$ , which is a contradiction. Hence,  $y_7 \in ipn_2(y_6, S)$ . Let  $S_1 = (S \setminus \{y_5\}) \cup \{x_6\}$ . Clearly,  $S_1$  is a  $\gamma_{t_2}(G)$ -set,  $\lambda(S_1) = \lambda(S)$  and  $|\mathcal{D} \cap S_1| = |\mathcal{D} \cap S|$ . We note that  $y_2 \in ipn_2(y_1, S_1)$  and  $\{x_6, y_7\} \subset ipn_2(y_6, S_1)$ . Further,  $\{y_1, y_2, x_6, y_6, y_7\} \subseteq B^{S_1}$  and  $|C_0^{S_1}| < |C_0^S|$ , which contradicts the choice of  $S$ .

By Claim 7, each vertex of  $\{y_2, y_3, y_4\}$  belongs to  $C_0^S \cup C_1^S$ .

*Claim 8.*  $\{y_2, y_3, y_4\} \cap C_1^S \neq \emptyset$ .

Suppose to the contrary that  $\{y_2, y_3, y_4\} \cap C_1^S = \emptyset$ . Then  $\{y_2, y_3, y_4\} \subseteq C_0^S$ . Since  $G$  is claw-free, there exists an edge in  $G[\{x_1, x_2, x_3\}]$ , say  $x_1x_2$ . If  $y_2 \neq y_3$ , then  $x_3y_2 \notin E(G)$  or  $x_3y_3 \notin E(G)$ , as  $x_3$  has only one neighbour in  $S \setminus \{y_1\}$ . By symmetry, consider  $y_2x_3 \notin E(G)$  (that is,  $y_2 \neq y_4$ ). Then  $S_1 = (S \setminus \{y_1, y_2\}) \cup \{x_1\}$  is a dominating set of  $G$  as  $y_2 \in C_0^S$  and  $y_1 \in C_0^S$ . Since  $|S_1| < |S|$ ,  $S_1$  cannot be a semi-TD-set of  $G$ . Combined with  $\{y_1, y_2\} \subseteq C_0^S$ ,  $y_1$  and  $y_2$  are the only two vertices of  $S$  within distance 2 from  $y_4$  in  $G$ , and  $y_4 \neq y_3$ . Thus, all vertices of  $N_{\bar{S}}(y_4) \setminus \{x_3\}$  are adjacent to  $y_2$ . Further,  $S_1 = (S \setminus \{y_1, y_3\}) \cup \{x_2\}$  is a semi-TD-set of  $G$  as  $y_3 \in C_0^S$  and  $y_1 \in C_0^S$ , which contradicts the minimality of  $S$ . Hence,  $y_2 = y_3$ .

Since  $y_1 \notin ipn_2(y_2, S)$ ,  $y_2x_3 \notin E(G)$  (that is,  $y_2 \neq y_4$ ). Let  $S_2 = (S \setminus \{y_1, y_4\}) \cup \{x_3\}$ . As  $y_4 \in C_0^S$ ,  $S_2$  is a dominating set of  $G$ . If  $d(y_2, x_3) \leq 2$ , then  $S_2$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $d(y_2, x_3) > 2$ . Further,  $x_1x_3 \notin E(G)$  and  $x_2x_3 \notin E(G)$ . Combining  $d(x_3) \geq 3$  and the claw-freeness of  $G$ , there exists a vertex  $x_4$  such that  $x_4x_3 \in E(G)$  and  $x_4y_4 \in E(G)$ . By Claim 1,  $x_4 \in \bar{S}$ . We note that  $y_2x_4 \notin E(G)$  as  $d(y_2, x_3) > 2$ . Since  $y_4 \in C_0^S$ ,  $x_4 \notin epn(y_4, S)$  and  $x_4$  has a neighbour  $y_5$  in  $S$  other than  $y_4$ . As  $S_2$  cannot be a semi-TD-set of  $G$ ,  $y_1$  and  $y_4$  are the only two vertices of  $S$  within distance 2 from  $y_2$  in  $G$ . Thus, all vertices of  $N_{\bar{S}}(y_2) \setminus \{x_1, x_2\}$  are adjacent to  $y_4$ .

Let  $x_5$  be a vertex in  $N_{\bar{S}}(y_2) \setminus \{x_1, x_2\}$ . Then  $x_5y_4 \in E(G)$ . If  $x_1x_5 \in E(G)$ , then  $(S \setminus \{y_1, y_2\}) \cup \{x_1\}$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $x_1x_5 \notin E(G)$ . Recall that  $x_1x_3 \notin E(G)$ . This implies that  $d(x_1) = 3$ . Similarly,  $d(x_2) = 3$ . Note that  $x_5x_3 \notin E(G)$  as  $d(y_2, x_3) > 2$ . Since  $G$  is claw-free,  $x_5x_4 \notin E(G)$  and each vertex of  $N(y_4) \setminus \{x_3, x_5\}$  is adjacent to either  $x_3$  or  $x_5$ . Let  $S_3 = (S \setminus \{y_1, y_2, y_4\}) \cup \{x_1, x_3, x_5\}$ . Then  $S_3$  is a  $\gamma_{t_2}(G)$ -set and  $\lambda(S_3) = \lambda(S)$ . If  $d(y_2) = 3$ , then  $G[\{y_1, y_2, x_1, x_3\}]$  is a special diamond. Further,  $|\mathcal{D} \cap S_2| < |\mathcal{D} \cap S|$ , which contradicts the choice of  $S$ . Thus,  $d(y_2) \geq 4$ . Let  $x_6$  be a vertex in  $N_{\bar{S}}(y_2) \setminus \{x_1, x_2, x_5\}$ . Then  $x_6y_4 \in E(G)$ . We note that  $\{y_2, y_4\} \in C_0^S \setminus C_0^S$  and  $|\{x \mid x \in N_{\bar{S}}(y_2) \text{ and } N_{C_0^S}(x) \neq \emptyset\}| \leq 2$ . From the definition



of the edge weighting functions, we have  $w(x_1y_1) = w(x_2y_1) = \frac{5}{8}$  and  $w(x_3y_1) \geq \frac{9}{16}$ . Further,  $f(\{y_1\}) = w(x_1y_1) + w(x_2y_1) + w(x_3y_1) > \frac{7}{4}$ , which contradicts the choice of  $y_1$ .

By Claim 8, we may assume that  $y_2 \in C_1^S$ . Then  $w(x_1y_1) \geq \frac{5}{8}$ . If  $y_3 \in C_1^S$ , then  $w(x_2y_1) \geq \frac{5}{8}$  and further  $f(\{y_1\}) = w(x_1y_2) + w(x_2y_2) + w(x_3y_2) \geq \frac{5}{8} + \frac{5}{8} + \frac{1}{2} \geq \frac{7}{4}$ , which is a contradiction. Thus,  $y_3 \in C_0^S$ . Similarly,  $y_4 \in C_0^S$ . This implies that  $y_2 \neq y_3$  and  $y_2 \neq y_4$ . Let  $N(y_2) = \{x_1, x_4, x_5\}$ , where  $\{x_4\} = epn(y_2, S)$ . If  $N_{C_0^S}(x_5) = \emptyset$ , then by the definition of the edge weighting functions, we have  $w(x_1y_1) = \frac{3}{4}$ . Further,  $f(\{y_1\}) = w(x_1y_2) + w(x_2y_2) + w(x_3y_2) \geq \frac{3}{4} + \frac{1}{2} + \frac{1}{2} \geq \frac{7}{4}$ , which is a contradiction. Hence,  $N_{C_0^S}(x_5) \neq \emptyset$ . We proceed with a series of claims that culminate in a contradiction.

**Claim 9.**  $|E(G[\{x_1, x_2, x_3\}])| = 1$ .

Since  $G$  is claw-free,  $|E(G[\{x_1, x_2, x_3\}])| \geq 1$ . If  $x_2x_1 \in E(G)$  and  $x_2x_3 \in E(G)$ , then  $(S \setminus \{y_1, y_3\}) \cup \{x_2\}$  is a semi-TD-set of  $G$  as  $y_1 \in C_0^S$  and  $y_3 \in C_0^S$ , which contradicts the minimality of  $S$ . Thus,  $x_2x_1 \notin E(G)$  or  $x_2x_3 \notin E(G)$ . Similarly,  $x_3x_1 \notin E(G)$  or  $x_3x_2 \notin E(G)$ . Suppose that  $|E(G[\{x_1, x_2, x_3\}])| \geq 2$ . Then  $x_1x_2 \in E(G)$ ,  $x_1x_3 \in E(G)$  and  $x_2x_3 \notin E(G)$ . This means  $G$  has a claw, which contradicts the claw-freeness of  $G$ . Hence,  $|E(G[\{x_1, x_2, x_3\}])| = 1$ .

**Claim 10.**  $y_3 = y_4$ .

Assume, to the contrary, that  $y_3 \neq y_4$ . By Claim 9, without loss of generality, we consider  $E(G[\{x_1, x_2, x_3\}]) = \{x_1x_2\}$  or  $\{x_2x_3\}$ . Let  $S_1 = (S \setminus \{y_1, y_3\}) \cup \{x_2\}$ . Since  $y_3 \in C_0^S$ ,  $S_1$  is a dominating set of  $G$ . Note that  $S_1$  cannot be a semi-TD-set of  $G$ . Thus, there exists a vertex  $y$  such that  $y_1$  and  $y_3$  are the only two vertices of  $S$  within distance 2 from  $y$  in  $G$  and  $d(y, x_2) > 2$ . If  $E(G[\{x_1, x_2, x_3\}]) = \{x_2x_3\}$ , then  $d(x_2, y_4) \leq 2$  and  $y = y_2$ . This implies that  $x_5y_3 \in E(G)$ . However, then  $(S \setminus \{y_1, y_4\}) \cup \{x_3\}$  is a semi-TD-set of  $G$  as  $y_4 \in C_0^S$ , which contradicts the minimality of  $S$ . Hence,  $E(G[\{x_1, x_2, x_3\}]) = \{x_1x_2\}$ . In this case,  $d(y_2, x_2) \leq 2$  and  $y = y_4$ . Thus, all vertices of  $N_{\bar{S}}(y_4) \setminus \{x_3\}$  are adjacent to  $y_3$ . Let  $x$  be a vertex in  $N_{\bar{S}}(y_4) \setminus \{x_3\}$ . Then  $xy_3 \in E(G)$ . Recall that  $d(y, x_2) > 2$ . Thus,  $d(y_4, x_2) > 2$  and  $x_2x \notin E(G)$ . Since  $G$  is claw-free,  $N_{\bar{S}}(y_4) \setminus \{x_3\}$  is a clique of  $G$ . Combining  $d(x_3) \geq 3$  and the claw-freeness of  $G$ , there exists a vertex  $x'$  in  $N_{\bar{S}}(y_4) \setminus \{x_3\}$  such that  $x_3x' \in E(G)$ . Then  $(S \setminus \{y_3, y_4\}) \cup \{x'\}$  is a semi-TD-set of  $G$  as  $y_3 \in C_0^S$ , which is a contradiction.

If  $y_3 (= y_4) \in C_0^S \setminus C_0^S$ , then  $w(x_2y_1) \geq \frac{9}{16}$  and  $w(x_3y_1) \geq \frac{9}{16}$ . Further,  $f(\{y_1\}) = w(x_1y_2) + w(x_2y_1) + w(x_3y_1) \geq \frac{5}{8} + \frac{9}{16} + \frac{9}{16} \geq \frac{7}{4}$ , which contradicts the choice of  $y_1$ . Thus,  $y_3 \in C_0^S$ .

**Claim 11.**  $x_2x_3 \notin E(G)$ .

For the sake of contradiction, suppose that  $x_2x_3 \in E(G)$ . If  $d(y_2, x_2) \leq 2$ , then  $(S \setminus \{y_1, y_3\}) \cup \{x_2\}$  is a semi-TD-set of  $G$  as  $y_3 \in C_0^S$ , which contradicts the minimality of  $S$ . Thus,  $d(y_2, x_2) \geq 3$ . Similarly,  $d(y_2, x_3) \geq 3$ . This implies that  $E[\{x_1, x_4, x_5\}, \{x_2, x_3\}] = \emptyset$ . If  $d(x_2) > 3$ , then there exists a vertex  $x$  in  $\bar{S}$  adjacent to  $x_2$ . Since  $G$  is claw-free,  $xy_3 \in E(G)$ . Thus,  $x$  has a neighbour in  $S$  different from  $y_3$  as  $y_3 \in C_0^S$ . Combined with  $N_{C_0^S}(x_5) \neq \emptyset$ ,  $(S \setminus \{y_1, y_3\}) \cup \{x_2\}$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $d(x_2) = 3$ . Similarly,  $d(x_3) = 3$ . Observe that  $y_1$  and  $y_3$  are in the same special diamond of  $G$ .

If  $x_1x_4 \in E(G)$ , then  $S_1 = (S \setminus \{y_1, y_2, y_3\}) \cup \{x_1, x_2\}$  is a dominating set of  $G$ . Otherwise,  $x_5$  is not dominated by  $S_1$ , and further  $x_5x_1 \notin E(G)$  and  $x_5y_3 \in E(G)$  as  $y_2 \in C_1^S$  and  $y_3 \in C_0^S$ . Since  $d(x_5) \geq 3$  and  $G$  is claw-free,  $N(x_5) = \{y_2, y_3, x_4\}$ . In this case,  $G = N_2$ , which is a contradiction. As  $N_S(x_5) \setminus \{y_2\} \subseteq C_0^S$  and  $y_3 \in C_0^S$ , there does not exist a vertex in  $S \setminus \{y_1\}$  such that  $y_2$  and  $y_3$  are the only two vertices of  $S$  within distance 2 from it in  $G$ . Thus,  $S_1$  is a semi-TD-set of  $G$  which contradicts the minimality of  $S$ . Hence,  $x_1x_4 \notin E(G)$ . Since  $d(x_1) \geq 3$  and  $G$  is claw-free,  $x_1x_5 \in E(G)$ . Note that  $d(x_1) = 3$ . If  $x_4x_5 \in E(G)$ , then  $G$  has a claw, for  $x_5$  has a neighbour in  $S$  other than  $y_2$ , which is a contradiction. Thus,  $x_4x_5 \notin E(G)$ . This implies that  $x_1$  is not in a special diamond.

Let  $S_2 = (S \setminus \{y_1, y_2, y_3\}) \cup \{x_1, x_2, x_4\}$ . Observe that  $S_2$  is a  $\gamma_{I2}(G)$ -set and  $\lambda(S_2) = \lambda(S)$ . If  $x_4$  is not in a special diamond, then  $|\mathcal{D} \cap S_2| < |\mathcal{D} \cap S|$ , which contradicts the choice of  $S$ . Thus,  $x_4$  is in a special diamond  $D$  of  $G$ . Let  $V(D) = \{x_4, x_6, x_7, y_5\}$ , where  $x_4y_5$  is the missing edge in the special diamond  $D$ . Clearly,  $\{x_6, x_7\} \subseteq \bar{S}$  and  $y_5 \in S$ . We note that  $y_5$  is an  $S$ -internal private neighbour. Let  $y_5 \in ipn_2(y_6, S)$  and  $x_8$  be the vertex of  $\bar{S}$  connecting  $y_5$  and  $y_6$ .

If  $epn(y_6, S) = \emptyset$ , then  $S_3 = (S \setminus \{y_1, y_2, y_5, y_6\}) \cup \{x_1, x_4, x_8\}$  is a dominating set of  $G$ . Since  $d(x_8) \geq 3$ , there exists a vertex  $x$  adjacent to  $xx_8 \in E(G)$ . Further,  $xy_6 \in E(G)$  as  $G$  is claw-free. Combined with  $epn(y_6, S) = \emptyset$ ,  $x$  has a neighbour in  $S \setminus \{y_1, y_5, y_6\}$ . Thus, there exists a vertex in  $S_3$  within distance 2 from  $x_8$ . It follows from the minimality of  $S$  that  $S_3$  cannot be a semi-TD-set of  $G$ . Thus,  $y_3$  is at a distance greater than 2 from every other vertex of  $S_3$ . This implies that  $y_3x_5 \notin E(G)$ ,  $x'y_6 \in E(G)$  and  $x'x_8 \notin E(G)$ , where  $\{x'\} = N(y_3) \setminus \{x_2, x_3\}$ . We note that neither  $x_8$  nor  $x'$  are in a special diamond. Otherwise,  $epn(y_6, S) \neq \emptyset$  or  $G$  has a claw, which is a contradiction. Further,  $(S \setminus \{y_1, y_2, y_3, y_5, y_6\}) \cup \{x_1, x_2, x_6, x_8, x'\}$  is a semi-TD-set of  $G$  with  $\lambda(S_3) = \lambda(S)$  but with  $|\mathcal{D} \cap S_3| < |\mathcal{D} \cap S|$ , which contradicts the choice of  $S$ . Hence,  $epn(y_6, S) \neq \emptyset$ .

Let  $x_9$  be a vertex in  $epn(y_6, S)$ . Then  $|\mathcal{D} \cap S_2| = |\mathcal{D} \cap S|$ ,  $\{x_1, x_2\} \cap C_0^{S_2} = \emptyset$  and  $y_6 \notin C_0^{S_2}$ . Further,  $|C_0^{S_2}| \leq |C_0^S|$  and  $|C_0^{S_2}| \leq |C_0^S|$ . If  $y_6 \notin C_1^{S_2}$ , then  $|C_1^{S_2}| < |C_1^S|$ , which contradicts the choice of  $S$ . Thus,  $y_6 \in C_1^{S_2}$ . Let  $N(y_6) = \{x_8, x_9, x_{10}\}$  (possibly,  $x_{10} = x_5$ ). Then  $x_{10}$  has a neighbour in  $S_2 \setminus \{y_6, y_5, x_4, x_2\}$ . Hence,  $x_{10}$  has a neighbour in  $S \setminus \{y_5, y_6\}$ . Let  $S_4 = (S \setminus \{y_5\}) \cup \{x_6\}$ . Then  $S_4$  is a  $\gamma_{I2}(G)$ -set. Now,  $\lambda(S_4) = \lambda(S)$ ,  $|\mathcal{D} \cap S_4| = |\mathcal{D} \cap S|$ ,  $x_6 \in epn_2(y_2, S_4)$  and  $y_6 \in ipn_2(y, S_4)$  for some vertex  $y$  of  $S_4$ . Thus,  $|C_0^{S_4}| \leq |C_0^S|$ ,  $|C_0^{S_4}| \leq |C_0^S|$  and  $|C_1^{S_4}| < |C_1^S|$ , which contradicts the choice of  $S$ .

By Claims 9–11, we may assume that  $E(G[\{x_1, x_2, x_3\}]) = \{x_1x_2\}$ . If  $x_1x_4 \in E(G)$ , then  $(S \setminus \{y_1, y_2\}) \cup \{x_1\}$  is a semi-TD-set of  $G$  as  $y_2 \in C_1^S$ , which contradicts the minimality of  $S$ . Thus,  $x_1x_4 \notin E(G)$ .

Suppose first that  $x_5y_3 \in E(G)$ . Since  $d(x_3) \geq 3$  and  $G$  is claw-free,  $x_5x_3 \in E(G)$ . If  $x_5x_4 \in E(G)$ , then  $(S \setminus \{y_1, y_2, y_3\}) \cup \{x_1, x_5\}$  is a semi-TD-set of  $G$ , which contradicts the choice of  $S$ . Thus,  $x_5x_4 \notin E(G)$ . Combining the claw-freeness of  $G$  and  $x_1x_4 \notin E(G)$ , we have  $x_5x_1 \in E(G)$ . Since  $G$  is claw-free,  $N(x_1) = \{y_1, y_2, x_2, x_5\}$ ,  $N(x_2) = \{y_1, y_3, x_1\}$ ,  $N(x_3) = \{y_1, y_3, x_5\}$ ,  $N(x_5) = \{y_2, y_3, x_1, x_3\}$  and  $X_1 := N(x_4) \setminus \{y_2\}$  is a clique of  $G$ . We construct  $G'$  from  $G$  by removing all vertices of  $\{y_1, y_3, x_1, x_2, x_3, x_5\}$  and adding the edges between  $\{y_2\}$  and  $X_1$  such that  $\{y_2\} \cup X_1$  is a clique of  $G'$ . Since  $d(x_4) \geq 3$ , we have  $|X_1| \geq 2$ . Thus,  $G' \neq N_2$  is a connected claw-free graph of order  $n' = n - 6$  with  $\delta(G') \geq 3$ . Note that  $X_1 \subseteq \bar{S}$  as  $x_4 \in epn(y_2, S)$ . Since  $S$  is a semi-TD-set of  $G$ , there exist a vertex  $y$  of  $S \setminus \{y_1, y_2, y_3\}$  adjacent to some vertices of  $X_1$  and a vertex  $y'$  of  $S \setminus \{y_1, y_2, y_3, y\}$  within distance 2 from  $y$  in  $S$ . Hence,  $n' \geq 6$ . By the minimality of  $G$ ,  $\gamma_{t_2}(G') \leq \frac{4}{11}n'$ . Let  $S'$  be a  $\gamma_{t_2}(G')$ -set. When  $y_2 \notin S'$ ,  $S' \cup \{y_1, x_5\}$  is a semi-TD-set of  $G$ . When  $y_2 \in S'$ ,  $(S' \setminus \{y_2\}) \cup \{y_1, x_5, x_4\}$  is a semi-TD-set of  $G$ . In both cases,  $\gamma_{t_2}(G) \leq \frac{4}{11}n' + 2 = \frac{4}{11}(n - 6) + 2 < \frac{4}{11}n$ , which contradicts the choice of  $G$ .

Suppose next that  $x_5y_3 \notin E(G)$ . Let  $N_S(x_5) \setminus \{y_2\} = \{y_5\}$  and  $N(y_3) \setminus \{x_2, x_3\} = \{x_6\}$ . Recall that  $N_{C_0^S}(x_5) \neq \emptyset$ . Thus,  $y_5 \in C_0^S$ . Since  $d(x_3) \geq 3$  and  $G$  is claw-free,  $x_3x_6 \in E(G)$ . If  $y_5x_6 \in E(G)$ , then  $(S \setminus \{y_3, y_5\}) \cup \{x_6\}$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $y_5x_6 \notin E(G)$ . Let  $N(y_5) = \{x_5, x_7, x_8\}$  and  $N_S(x_6) \setminus \{y_3\} = \{y_6\}$ . If  $x_5x_1 \notin E(G)$ , then  $x_4x_5 \in E(G)$  as  $G$  is claw-free and  $x_1x_4 \notin E(G)$ . In this case,  $(S \setminus \{y_1, y_2, y_5\}) \cup \{x_1, x_5\}$  is a semi-TD-set of  $G$ , which contradicts the minimality of  $S$ . Thus,  $x_5x_1 \in E(G)$ . If  $x_5x_4 \in E(G)$ , then  $(S \setminus \{y_2, y_5\}) \cup \{x_5\}$  is a semi-TD-set of  $G$ , which is a contradiction. Thus,  $x_5x_4 \notin E(G)$ . Since  $G$  is claw-free,  $d(x_1) = 4$ ,  $d(x_2) = 3$ ,  $d(x_3) = 3$ ,  $N(x_5) \subseteq \{y_2, y_5, x_1, x_7, x_8\}$  and  $X_2 := N(x_6) \setminus \{y_3, x_3\}$  is a clique of  $G$ . Let  $G'$  be the graph obtained from  $G$  by removing all vertices of  $\{x_1, x_2, x_3, x_6, y_1, y_3\}$  and adding the edges between  $\{y_2, x_5\}$  and  $X_2$  such that  $\{y_2, x_5\} \cup X_2$  is a clique of  $G'$ . We observe that  $G' \neq N_2$  is a connected claw-free graph of order  $n' = n - 6 \geq 6$  with  $\delta(G') \geq 3$ . Then  $G'$  has a  $\gamma_{t_2}(G')$ -set  $S'$  with at most  $\frac{4}{11}n'$  vertices by the minimality of  $G$ . When  $X_2 \cap S' \neq \emptyset$ ,  $S' \cup \{x_1, y_3\}$  is a semi-TD-set of  $G$ . When  $X_2 \cap S' = \emptyset$ ,  $S' \cup \{y_1, x_6\}$  is a semi-TD-set of  $G$ . In either case,  $\gamma_{t_2}(G) \leq \frac{4}{11}n' + 2 = \frac{4}{11}(n - 6) + 2 < \frac{4}{11}n$ , which is a contradiction.

### Acknowledgement

The authors thank the referees for their careful review and helpful suggestions.

### References

- [1] J. Chen and S.-J. Xu, 'A characterization of 3- $\gamma$ -critical graphs which are not bicritical', *Inform. Process. Lett.* **166** (2021), Article no. 106062.
- [2] Y. X. Dong, E. F. Shan, L. Y. Kang and S. Li, 'Domination in intersecting hypergraphs', *Discrete Appl. Math.* **251** (2018), 155–159.

- [3] W. Goddard, M. A. Henning and C. A. McPillan, 'Semitotal domination in graphs', *Util. Math.* **94** (2014), 67–81.
- [4] T. W. Haynes, S. T. Hedetniemi and P. J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker Inc., New York, 1998).
- [5] M. A. Henning, 'Edge weighting functions on semitotal dominating sets', *Graphs Combin.* **33** (2017), 403–417.
- [6] M. A. Henning, 'Bounds on domination parameters in graphs: a brief survey', *Discuss. Math. Graph Theory* **42** (2022), 665–708.
- [7] M. A. Henning, L. Kang, E. Shan and A. Yeo, 'On matching and total domination in graphs', *Discrete Math.* **308** (2008), 2313–2318.
- [8] M. A. Henning and A. J. Marcon, 'On matching and semitotal domination in graphs', *Discrete Math.* **324** (2014), 13–18.
- [9] M. A. Henning and A. J. Marcon, 'Semitotal domination in claw-free cubic graphs', *Ann. Comb.* **20**(4) (2016), 1–15.
- [10] M. A. Henning and A. Pandey, 'Algorithmic aspects of semitotal domination in graphs', *Theoret. Comput. Sci.* **766** (2019), 46–57.
- [11] M. A. Henning and A. Yeo, *Total Domination in Graphs* (Springer, New York, 2013).
- [12] E. Zhu and C. Liu, 'On the semitotal domination number of line graphs', *Discrete Appl. Math.* **254** (2019), 295–298.
- [13] E. Zhu, Z. Shao and J. Xu, 'Semitotal domination in claw-free cubic graphs', *Graphs Combin.* **33**(5) (2017), 1119–1130.

JIE CHEN, School of Mathematics and Statistics,  
Gansu Center for Applied Mathematics, Lanzhou University,  
Lanzhou, Gansu 730000, PR China  
e-mail: [ChenJieJie2023@hotmail.com](mailto:ChenJieJie2023@hotmail.com)

HONGZHANG CHEN, School of Mathematics and Statistics,  
Gansu Center for Applied Mathematics, Lanzhou University,  
Lanzhou, Gansu 730000, PR China  
e-mail: [mnhzchern@gmail.com](mailto:mnhzchern@gmail.com)

SHOU-JUN XU, School of Mathematics and Statistics,  
Gansu Center for Applied Mathematics, Lanzhou University,  
Lanzhou, Gansu 730000, PR China  
e-mail: [shjxu@lzu.edu.cn](mailto:shjxu@lzu.edu.cn)