

## 0-HECKE ALGEBRAS

P. N. NORTON

(Received 8 March 1978)

Communicated by D. E. Taylor

### Abstract

The structure of a 0-Hecke algebra  $H$  of type  $(W, R)$  over a field is examined.  $H$  has  $2^n$  distinct irreducible representations, where  $n = |R|$ , all of which are one-dimensional, and correspond in a natural way with subsets of  $R$ .  $H$  can be written as a direct sum of  $2^n$  indecomposable left ideals, in a similar way to Solomon's (1968) decomposition of the underlying Coxeter group  $W$ .

*Subject classification (Amer. Math. Soc. (MOS) 1970):* 16 A 48.

### 1. Introduction

NOTATION.  $\{i_1, \dots, i_s, \dots, i_n\}$  denotes the set  $\{i_1, \dots, i_n\} - \{i_s\}$ ,  $\cup$  denotes set union and  $\cap$  denotes set intersection.  $(xyx\dots)_n$  denotes the product of the first  $n$  terms of the sequence  $x, y, x, y, x, \dots$ . ACC denotes the ascending chain condition and DCC denotes the descending chain condition. Let  $S$  be a set and  $A$  a subset of  $S$ . Then  $|A|$  denotes the number of elements in  $A$ , and  $\hat{A}$  denotes the complement of  $A$  in  $S$ .

Let  $K$  be any field, and let  $(W, R)$  be a finite Coxeter system, with root system  $\Phi$ , positive system  $\Phi^+$  and simple system  $\Pi$ . For each  $J \subseteq R$ , let  $\Phi_J$ ,  $\Phi_J^+$  and  $\Pi_J$  be the corresponding root system, positive system and simple system.  $w_i \in R$  is the reflection in the hyperplane perpendicular to  $r_i \in \Pi$ . For each  $J \subseteq R$ , let

$$X_J = \{w \in W: w(\Pi_J) \subseteq \Phi^+\} \quad \text{and} \quad Y_J = \{w \in W: w(\Pi_J) \subseteq \Phi^+, w(\Pi_{\hat{J}}) \subseteq \Phi^-\},$$

where  $\hat{J} = R - J$ . We shall assume all the standard results on finite Coxeter systems, as found in Bourbaki (1968), Carter (1972) and Steinberg (1967).

1.1 DEFINITION. The 0-Hecke algebra  $H$  over  $K$  of type  $(W, R)$  is the associative algebra over  $K$  with identity 1 generated by  $\{a_i: w_i \in R\}$  subject to the relations:

- (i)  $a_i^2 = -a_i$  for all  $w_i \in R$ ,
- (ii)  $(a_i a_j a_i \dots)_{n_{ij}} = (a_j a_i a_j \dots)_{n_{ij}}$  for all  $w_i, w_j \in R, w_i \neq w_j$ , where  $n_{ij}$  = the order of  $w_i w_j$  in  $W$ .

For all  $w \in W$ , define  $a_w = a_{i_1} \dots a_{i_r}$ , where  $w = w_{i_1} \dots w_{i_r}$  is a reduced expression for  $w \in W$  in terms of the elements of  $R$ . Note that  $a_{1_W} = 1$ , where  $1_W$  denotes the identity element of  $W$ . It is easy to show that  $a_w$  is independent of the reduced expression for  $w$ , and that every element of  $H$  is a  $K$ -linear combination of elements  $a_w$ , for  $w \in W$ .

By Bourbaki (1968) (Exercise 23, p. 55),  $\{a_w: w \in W\}$  are linearly independent over  $K$  and so form a  $K$ -basis of  $H$ .

1.2 SOME EXAMPLES. (i) Let  $G = G(q)$  be a Chevalley group over the finite field  $F = GF(q)$  of  $q$  elements, where  $q = p^m$  for some prime  $p$  and positive integer  $m$ . Then  $G$  has a  $(B, N)$  pair  $(G, B, N, R)$  and Weyl group  $W$  such that for each  $w_i \in R$  there is a positive integer  $c_i$  such that  $|B: B \cap B^{w_i}| = q^{c_i}$ . If  $K$  is a field of characteristic  $p$ , then the Hecke algebra  $H_K(G, B)$  is a 0-Hecke algebra.

(ii) Let  $G$  be a finite group with a split  $(B, N)$  pair  $(G, B, N, R, U)$  of rank  $n$  and characteristic  $p$  with Weyl group  $W$ , and let  $K$  be a field of characteristic  $p$ . Then the Hecke algebra  $H_K(G, B)$  is a 0-Hecke algebra of type  $(W, R)$  over  $K$ .

1.3 LEMMA. For all  $w_i \in R$  and all  $w \in W$ ,

$$a_i a_w = \begin{cases} a_{w_i w} & \text{if } l(w_i w) = l(w) + 1, \\ -a_w & \text{if } l(w_i w) = l(w) - 1; \end{cases}$$

$$a_w a_i = \begin{cases} a_{w w_i} & \text{if } l(w w_i) = l(w) + 1, \\ -a_w & \text{if } l(w w_i) = l(w) - 1. \end{cases}$$

PROOF. If  $l(w_i w) = l(w) + 1$ , then  $a_{w_i w} = a_i a_w$  by the definition of  $a_{w_i w}$ . Suppose  $l(w_i w) = l(w) - 1$ ; then there is a reduced expression for  $w$  beginning with  $w_i$ : say  $w = w_i w'$  where  $l(w) = l(w') + 1$ . Then  $a_w = a_i a_{w'}$ , and so

$$a_i a_w = a_i a_i a_{w'} = -a_i a_{w'} = -a_w.$$

Similarly for  $a_w a_i$ .

1.4 COROLLARY. (1) For all  $w, w' \in W$ ,

- (a)  $a_w a_{w'} = \pm a_{w''}$  for some  $w'' \in W$ , with  $l(w'') \geq \max(l(w), l(w'))$ ;
- (b)  $a_w a_{w'} = a_{ww'}$  if and only if  $l(ww') = l(w) + l(w')$ ;

- (c)  $a_w a_{w'} = (-1)^{l(w')} a_w$  if and only if  $w(r_i) \in \Phi^-$  for each  $r_i \in \Pi_J$ , where  $J = \{w_i \in R: w_i \text{ occurs in some reduced expression for } w'\}$ .
  - (d)  $a_w a_{w'} = (-1)^{l(w)} a_{w'}$  if and only if  $(w')^{-1}(r_i) \in \Phi^-$  for each  $r_i \in \Pi_J$ , where  $J = \{w_i \in R: w_i \text{ occurs in some reduced expression for } w\}$ ;
  - (e)  $a_w a_{w'} = \pm a_{w'}$  with  $l(w'') > l(w)$ , where  $l(w) \geq l(w')$ , if and only if there exists  $r_i \in \Pi_J$  such that  $w(r_i) \in \Phi^+$ , where  $J = \{w_j \in R: w_j \text{ occurs in some reduced expression for } w'\}$ .
- (2) Let  $w_0$  be the unique element of maximal length in  $W$ . Then for all  $w \in W$ ,

$$a_w a_{w_0} = (-1)^{l(w)} a_{w_0} \text{ and } a_{w_0} a_w = (-1)^{l(w)} a_{w_0}.$$

### 2. The nilpotent radical of $H$

Let  $N$  be the nilpotent radical of  $H$ . Since  $H$  is a finite-dimensional algebra over  $K$ ,  $H$  has the DCC and ACC and so  $N$  is also the Jacobson radical of  $H$ , and is the unique maximal nilpotent ideal of  $H$ .

There is a natural composition series for  $H$ , consisting of (two-sided) ideals of  $H$  such that every factor is a one-dimensional  $H$ -module. This series arises as follows: list the basis elements  $\{a_w: w \in W\}$  in order of increasing length of  $w$ , and if  $w, w' \in W$  have the same length it does not matter in which order  $a_w$  and  $a_{w'}$  occur on the list. Rename these elements  $h_1, h_2, \dots, h_{|W|}$  respectively. Note that  $h_1 = 1$  and  $h_{|W|} = a_{w_0}$ . Let  $H_j$  be the ideal of  $H$  generated by  $\{h_m: m \geq j\}$ .  $H_j$  has  $K$ -basis  $\{h_m: m \geq j\}$  and dimension  $|W| - j + 1$ . Then

2.1 
$$H = H_1 > H_2 > \dots > H_{|W|} = a_{w_0} H > 0$$

is the natural composition series of  $H$  described above.  $H_i/H_{i+1}$  is a one-dimensional  $H$ -module,  $1 \leq i \leq |W|$ , where  $H_{|W|+1} = 0$ , with basis  $h_i + H_{i+1}$ , where  $h_i = a_w$  for some  $w \in W$ . Either  $a_w^2 = (-1)^{l(w)} a_w$  or  $a_w^2 \in H_{i+1}$ ; in the first case, the factor ring  $H_i/H_{i+1}$  is generated by an idempotent, and in the second case it is nilpotent.

2.2 LEMMA. *The number of factors which are generated by an idempotent is equal to  $2^n$ , where  $n = |R|$ .*

PROOF. The factors generated by idempotents correspond to elements  $w \in W$  such that  $a_w^2 = (-1)^{l(w)} a_w$ . Let  $w \in W$  be such an element. Write  $w = w_{i_1} \dots w_{i_s}$ , where  $l(w) = s$ , and let  $J = \{w_{i_j}: 1 \leq j \leq s\}$ . Then  $w \in W_J$ , and by 1.4(1c),  $w(\Pi_J) \subseteq \Phi^-$ . Hence  $w = w_{0,J}$ , the unique element of maximal length in  $W_J$ . Conversely, for each subset  $J$  of  $R$ ,  $a_{w_{0,J}}^2 = (-1)^{l(w_{0,J})} a_{w_{0,J}}$ . Hence the number of factors which are generated by an idempotent is equal to the number of subsets of  $R$ , that is,  $2^n$ , where  $n = |R|$ .

By Schreier’s theorem, any series of ideals of  $H$  can be refined to a composition series, and all so obtained have the same number of terms in them as the natural series, and with the factors in one–one correspondence with those of the natural series. In particular, consider  $H > N \geq 0$ . This can be refined to a composition series  $H = H'_1 > \dots > H'_{|W|} > H'_{|W|+1} = 0$ , where  $N = H'_r$ ,  $2 < r \leq |W| + 1$ . Now each factor  $H'_i/H'_{i+1}$ ,  $i \geq r$ , is nilpotent as  $H'_i \leq N$ , and each factor  $H'_i/H'_{i+1}$ ,  $i + 1 \leq r$ , must be generated by an idempotent as  $H'_i/N \leq H/N$ , a semi-simple ring. Hence the number of factors which are nilpotent is equal to the dimension of  $N$ . Thus,  $\dim N = |W| - 2^n$ , where  $n = |R|$ .

We can, however, give a precise basis of  $N$ .

2.3 THEOREM. Let  $w \in W$ , and suppose  $w \neq w_{0J}$  for any  $J \subseteq R$ . Write  $w = w_{i_1} \dots w_{i_s}$ ,  $l(w) = s$ , and let  $J(w) = \{w_{i_j} : 1 \leq j \leq s\}$ . Then  $E(w) = a_w + (-1)^{l(w_{0J(w)})+l(w)+1} a_{w_{0J(w)}}$  is nilpotent, and  $\{E(w) : w \in W, w \neq w_{0J} \text{ for any } J \subseteq R\}$  is a basis of  $N$ .

PROOF. Show  $E(w)$  is nilpotent by induction on  $l(w_{0J(w)}) - l(w)$ . Note that if  $w = w_{0J}$  for some  $J \subseteq R$  then  $E(w) = 0$ . Suppose  $l(w_{0J(w)}) - l(w) = 1$ . Then since a reduced expression for  $w$  involves all  $w_i \in J(w)$ ,  $w \neq w_{0J(w)}$ , there exists  $r_j \in \Pi_{J(w)}$  such that  $w(r_j) \in \Phi^+$ . So  $a_w^2 = (-1)^{l(w)-1} a_{w_{0J(w)}}$ . Thus

$$\begin{aligned} E(w)^2 &= a_w^2 + a_w a_{w_{0J(w)}} + a_{w_{0J(w)}} a_w + a_{w_{0J(w)}}^2 \\ &= a_{w_{0J(w)}}^b \quad \text{where } b = (-1)^{l(w)-1} + 2(-1)^{l(w)} + (-1)^{l(w_{0J(w)})} \\ &= 0 \text{ as } l(w_{0J(w)}) = l(w) + 1. \end{aligned}$$

Now suppose  $l(w_{0J(w)}) - l(w) > 1$ . Consider the product  $a_w a_w$ . Since  $w \neq w_{0J(w)}$ , there exists  $r_j \in \Pi_{J(w)}$  such that  $w(r_j) \in \Phi^+$ . As any reduced expression for  $w$  involves all  $w_i \in J(w)$ , we have  $a_w a_w = (-1)^{2l(w)-l(w')} a_{w'}$ , with  $w' \in W_{J(w)}$  and  $l(w') > l(w)$ . Further,  $J(w') = J(w)$ . Then

$$\begin{aligned} E(w)^2 &= a_w^2 + 2(-1)^{l(w_{0J(w)})+1} a_{w_{0J(w)}} + (-1)^{l(w_{0J(w)})} a_{w_{0J(w)}} \\ &= (-1)^{l(w')} a_{w'} + (-1)^{l(w_{0J(w)})+1} a_{w_{0J(w)}} \\ &= (-1)^{l(w')} (a_{w'} + (-1)^{l(w_{0J(w'))+l(w')+1} a_{w_{0J(w')}}) \\ &= (-1)^{l(w')} E(w'). \end{aligned}$$

As  $l(w') > l(w)$ , either  $w' = w_{0J(w)}$  and thus  $E(w)^2 = 0$  or  $w' \neq w_{0J(w)}$  and then by induction  $E(w')$  is nilpotent. Thus  $E(w)$  is nilpotent.

Finally, note that we get a nilpotent element for each  $w \in W$ ,  $w \neq w_{0J}$  for any  $J \subseteq R$ . The set of all  $E(w)$ ,  $w \neq w_{0J}$  for any  $J \subseteq R$ , is obviously linearly independent, and there are  $|W| - 2^n$  elements in all, where  $n = |R|$ . Hence they are a  $K$ -basis for  $N$ .

2.4 COROLLARY.  $H/N$  is commutative.

PROOF. We show that  $a_i a_j - a_j a_i \in N$  for all  $w_i, w_j \in R$ . If  $a_i a_j = a_j a_i$ , the result is obvious. So suppose  $a_i a_j \neq a_j a_i$ . Then we can form  $E(w_i w_j)$  and  $E(w_j w_i)$  and  $E(w_i w_j) - E(w_j w_i) = a_i a_j - a_j a_i \in N$  as each of  $E(w_i w_j)$  and  $E(w_j w_i)$  is in  $N$ .

### 3. The irreducible representations of $H$

Consider the one-dimensional  $H$ -modules which arise from the natural composition series of  $H$ . Let the factor  $H_i/H_{i+1}$  be generated as left  $H$ -module by  $a_w + H_{i+1}$ . The action of  $H$  on this element is determined as follows: for each  $w_i \in R$ ,

$$a_i(a_w + H_{i+1}) = \begin{cases} -(a_w + H_{i+1}) & \text{if } w^{-1}(r_i) \in \Phi^-, \\ 0 & \text{if } w^{-1}(r_i) \in \Phi^+. \end{cases}$$

For any  $w \in W$ , let  $J(w) = \{w_j : 1 \leq j \leq s\}$  where  $w = w_{i_1} \dots w_{i_s}$  is a reduced expression for  $w$ . Then for  $w' \in W$ ,

$$a_{w'}(a_w + H_{i+1}) = \begin{cases} (-1)^{l(w')} (a_w + H_{i+1}) & \text{if } w^{-1}(\prod_{J(w')} r_i) \subseteq \Phi^-, \\ 0 & \text{if there exists } r_i \in \prod_{J(w')} \text{ such} \\ & \text{that } w^{-1}(r_i) \in \Phi^+. \end{cases}$$

Hence the action of  $H$  on  $a_w + H_{i+1}$  depends on  $w^{-1}$ .

3.1 DEFINITION. For each  $J \subseteq R$ , let  $\lambda_J$  be the one-dimensional representation of  $H$  defined by

$$\lambda_J(a_i) = \begin{cases} 0 & \text{if } w_i \in J, \\ -1 & \text{if } w_i \in \hat{J}. \end{cases}$$

For all  $w \in W$ , let  $w = w_{i_1} \dots w_{i_s}$  with  $l(w) = s$ . Then  $\lambda_J(a_w) = \lambda_J(a_{i_1}) \dots \lambda_J(a_{i_s})$ . Extend  $\lambda_J$  to  $H$  by linearity.

For each  $J \subseteq R$ , let  $H_{i(J)}/H_{i(J)+1}$  be the factor of the natural series which is generated by  $a_{w_J} + H_{i(J)+1}$ . Then the left  $H$ -module  $H_{i(J)}/H_{i(J)+1}$  affords the representation  $\lambda_J$  of  $H$ .

Since each composition factor of  $H$  is one-dimensional, it follows that all irreducible representations of  $H$  are one-dimensional. Let  $\mu$  be an irreducible representation of  $H$ . Then  $\mu$  is completely determined by the values  $\mu(a_i)$  for all  $w_i \in R$ . Since  $\mu$  is an algebra homomorphism,  $\mu(a_i)^2 = -\mu(a_i)$  for all  $w_i \in R$ . Let  $\mu(a_i) = u_i \in K$  for all  $w_i \in R$ . Then  $u_i^2 = -u_i$  in  $K$  implies that  $u_i = 0$  or  $u_i = -1$ .

Thus each irreducible representation of  $H$  can be described by an  $n$ -tuple  $(u_1, \dots, u_n)$ , where  $n = |R|$ , with  $u_i = 0$  or  $-1$  for all  $i$ . In particular,  $\lambda_J$  corresponds to the  $n$ -tuple  $(u_1, \dots, u_n)$  where  $u_i = 0$  if  $w_i \in J$  and  $u_i = -1$  if  $w_i \in \bar{J}$ . There are  $2^n$  such irreducible representations, and they all occur in the natural series of  $H$ .

$2^n$  maximal ideals of  $H$  are determined as follows: for each  $J \subseteq R$ , form the  $n$ -tuple  $(u_1, \dots, u_n)$ , where  $u_i = 0$  if  $w_i \in J$  and  $u_i = -1$  otherwise. Let  $M_J$  be the left ideal of  $H$  generated by  $\{a_i - u_i 1 : w_i \in R\}$ . Then  $M_J = \ker \lambda_J$ , and as each  $\lambda_J$  is irreducible,  $M_J$  is a maximal left ideal of  $H$ .

Now  $H/N$  is semi-simple Artinian. So by extending  $K$  to its algebraic closure  $\bar{K}$  and considering  $H$  as an algebra over  $\bar{K}$ , we deduce that

$$H/N \cong \bar{K} \oplus \bar{K} \oplus \dots \oplus \bar{K}, \text{ a direct sum of } 2^n \text{ fields.}$$

(Actually, we will show that

$$H/N \cong K \oplus K \oplus \dots \oplus K, \text{ } 2^n \text{ copies of } K,$$

regardless of which field  $K$  is.)

#### 4. Some decompositions of $H$

For each  $J \subseteq R$ , let  $H_J$  be the subalgebra of  $H$  generated by  $\{a_i : w_i \in J\}$ .

4.1 DEFINITION. For each  $J \subseteq R$ , let

$$e_J = \sum_{w \in W_J} a_w, \quad o_J = (-1)^{l(w_{0J})} a_{w_{0J}}.$$

4.2 LEMMA. For all  $w_i \in J$ ,

$$a_i e_J = 0 = e_J a_i \quad \text{and} \quad a_i o_J = -o_J = o_J a_i.$$

PROOF. Use 1.3.

4.3 LEMMA. Let  $w_{0J} = w_{i_1} \dots w_{i_r}$ ,  $l(w_{0J}) = s$ . Then

$$e_J = (1 + a_{i_1}) \dots (1 + a_{i_r})$$

and is independent of the reduced expression for  $w_{0J}$ .

NOTATION. For all  $w \in W$ , if  $w = w_{i_1} \dots w_{i_t}$  with  $l(w) = t$ , write

$$[1 + a_w] = (1 + a_{i_1}) \dots (1 + a_{i_t}).$$

By the following proof it follows that  $[1 + a_w]$  is independent of the reduced expression for  $w$ .

PROOF. Firstly, we show that  $[1 + a_{w_0j}]$  is independent of the reduced expression for  $w_{0j}$ . Since we can pass from one reduced expression for  $w_{0j}$  to another by substitutions of the form  $(w_i w_j w_i \dots)_{n_{ij}} = (w_j w_i w_j \dots)_{n_{ij}}$ ,  $i \neq j$ , where  $n_{ij}$  is the order of  $w_i w_j$  in  $W$ , we need to show that

$$[1 + a_{(w_i w_j w_i \dots)_{n_{ij}}}] = [1 + a_{(w_j w_i w_j \dots)_{n_{ij}}}]$$

To do this, we use induction on  $n$ ,  $n \leq n_{ij}$ , to show that

$$[1 + a_{(w_i w_j w_i \dots)_n}] = 1 + \sum_{m=1}^n a_{(w_i w_j w_i \dots)_m} + \sum_{m=1}^{n-1} a_{(w_j w_i w_j \dots)_m}$$

This is clearly true for  $n = 1$ . Suppose it is true for all integers  $\leq k$ , and suppose that  $k$  is odd. Then

$$\begin{aligned} [1 + a_{(w_i w_j w_i \dots)_{k+1}}] &= [1 + a_{(w_i w_j w_i \dots)_k}] (1 + a_j) \\ &= \left( 1 + \sum_{m=1}^k a_{(w_i w_j w_i \dots)_m} + \sum_{m=1}^{k-1} a_{(w_j w_i w_j \dots)_m} \right) (1 + a_j) \\ &= \left( 1 + \sum_{m=1}^k a_{(w_i w_j w_i \dots)_m} + \sum_{m=1}^{k-1} a_{(w_j w_i w_j \dots)_m} \right) + a_j \\ &\quad + \sum_{m=0}^{\frac{1}{2}(k-1)} a_{(w_i w_j w_i \dots)_{2m+1}} a_j + \sum_{m=1}^{\frac{1}{2}(k-1)} a_{(w_i w_j w_i \dots)_{2m}} a_j \\ &\quad + \sum_{m=1}^{\frac{1}{2}(k-1)} a_{(w_j w_i w_j \dots)_{2m-1}} a_j + \sum_{m=1}^{\frac{1}{2}(k-1)} a_{(w_j w_i w_j \dots)_{2m}} a_j. \end{aligned}$$

Now,

$$a_{(w_i w_j w_i \dots)_{2m-1}} a_j = -a_{(w_i w_j w_i \dots)_{2m}} a_j, \quad 1 \leq m \leq \frac{1}{2}(k-1),$$

and

$$a_{(w_j w_i w_j \dots)_{2m-1}} a_j = -a_{(w_j w_i w_j \dots)_{2m-2}} a_j, \quad 1 \leq m \leq \frac{1}{2}(k-1),$$

where  $a_{(w_i w_j w_i \dots)_0} = 1$ . Then

$$\begin{aligned} [1 + a_{(w_i w_j w_i \dots)_{k+1}}] &= 1 + \sum_{m=1}^k a_{(w_i w_j w_i \dots)_m} + \sum_{m=1}^{k-1} a_{(w_j w_i w_j \dots)_m} \\ &\quad + a_{(w_i w_j w_i \dots)_k} a_j + a_{(w_j w_i w_j \dots)_{k-1}} a_j \\ &= 1 + \sum_{m=1}^{k+1} a_{(w_i w_j w_i \dots)_m} + \sum_{m=1}^k a_{(w_j w_i w_j \dots)_m}. \end{aligned}$$

Similarly, we get the above result if we assume  $k$  is even.

Similarly, for all  $n \leq n_{ij}$ ,

$$[1 + a_{(w_j w_i w_j \dots)_n}] = 1 + \sum_{m=1}^n a_{(w_j w_i w_j \dots)_m} + \sum_{m=1}^{n-1} a_{(w_i w_j w_i \dots)_m}$$

Then, for all  $n \leq n_{ij}$ ,

$$[1 + a_{(w_i w_j w_i \dots)_n}] - [1 + a_{(w_j w_i w_j \dots)_n}] = a_{(w_i w_j w_i \dots)_n} - a_{(w_j w_i w_j \dots)_n}$$

When  $n = n_{ij}$ , this difference is zero, and so

$$[1 + a_{(w_i w_j w_i \dots)_{n_{ij}}}] = [1 + a_{(w_j w_i w_j \dots)_{n_{ij}}}]$$

and thus  $[1 + a_{w_{0j}}]$  is independent of the reduced expression for  $w_{0j}$  chosen.

Finally,  $[1 + a_{w_{0j}}]$  is a linear combination of certain  $a_w$  with  $w \in W_J$ . We show by induction on  $l(w)$  for all  $w \in W_J$  that  $a_w$  occurs in the expansion of  $[1 + a_{w_{0j}}]$  with coefficient 1. If  $l(w) = 0$ , then  $w = 1$  and obviously 1 occurs with coefficient 1. Suppose  $l(w) > 0$ . Let  $w = w'w_j$ ,  $w' \in W_J$ ,  $w_j \in J$ , where  $l(w) = l(w') + 1$ . By induction  $a_{w'}$  occurs in  $[1 + a_{w_{0j}}]$  with coefficient 1. Choose an expression for  $w_{0j}$  ending in  $w_j$ , and then  $[1 + a_{w_{0j}}] = [1 + a_{w_{0j}w_j}](1 + a_j)$ . Since  $l(w'w_j) > l(w')$ , the only contribution to  $a_{w'}$  from the last bracket is from the 1. If instead we take  $a_j$  from the last bracket, we get  $a_w$ , with coefficient 1. Now suppose  $a_w$  occurs in  $[1 + a_{w_{0j}w_j}]$  with coefficient  $m$ . Then

$$ma_w(1 + a_j) = ma_w + ma_w a_j = ma_w - ma_w = 0 \text{ as } w(r_j) \in \Phi^-.$$

Thus  $a_w$  occurs in the expansion of  $[1 + a_{w_{0j}}]$  with coefficient 1, and hence  $e_J = [1 + a_{w_{0j}}]$ .

4.4 COROLLARY. (1) If  $J, L \subseteq R$ ,  $J \cap L \neq \emptyset$ , then  $o_J e_L = 0$  and  $e_J o_L = 0$ .

(2) If  $L \subseteq J \subseteq R$ , then  $e_L e_J = e_J = e_J e_L$  and  $o_L o_J = o_J = o_J o_L$ .

PROOF. Use 4.2 and 4.3.

4.5 LEMMA. Let  $y \in Y_J$  for some  $J \subseteq R$ . Then  $a_y o_J = a_y$  and  $a_y o_J e_J = \sum_{w \in W_J} a_y w$ , with  $l(yw) = l(y) + l(w)$  for all  $w \in W_J$ , that is,  $a_y o_J e_J$  is equal to  $a_y$  plus a sum of certain  $a_w$  with  $l(w) > l(y)$ .

PROOF. If  $y \in Y_J$ , then  $y = ww_{0j}$  for some  $w \in W$  with  $l(y) = l(w) + l(w_{0j})$ . Hence  $a_y o_J = (-1)^{l(w_{0j})} a_w a_{w_{0j}} a_{w_{0j}}$ , and so  $a_y o_J = a_y$ . Now for all  $w \in W_J$ , as  $y \in Y_J \subseteq X_J$ , we have  $l(yw) = l(y) + l(w)$ . So for all  $w \in W_J$ ,  $a_y a_w = a_{yw}$ . Thus

$$a_y o_J e_J = a_y e_J = \sum_{w \in W_J} a_y a_w = \sum_{w \in W_J} a_y w = a_y + \sum_{w \in W_J, w \neq 1} a_y w$$

and  $l(yw) > l(y)$  if  $w \neq 1$ ,  $w \in W_J$ .



4.6 LEMMA. For  $y \in Y_J$ ,  $a_y$  occurs in the expansion of  $a_y e_J o_{\mathfrak{J}}$  with coefficient 1, and if, for any  $w \in W$ ,  $a_w$  occurs in the expansion of  $a_y e_J o_{\mathfrak{J}}$  with non-zero coefficient, then  $w = y$  or  $l(w) > l(y)$ .

PROOF. By 4.5,  $a_y e_J = \sum_{w \in W_J} a_{yw} a_w$ , with  $l(yw) = l(y) + l(w)$  for all  $w \in W_J$ . So

$$a_y e_J o_{\mathfrak{J}} = \sum_{w \in W_J} a_{yw} o_{\mathfrak{J}} = a_y o_{\mathfrak{J}} + \sum_{w \in W_J, w \neq 1} a_{yw} o_{\mathfrak{J}}.$$

From the proof of 4.5,  $a_y o_{\mathfrak{J}} = a_y$ , and for all  $w \in W_J$ ,  $w \neq 1$ ,

$$a_{yw} o_{\mathfrak{J}} = a_{yw} (-1)^{l(w_{o_{\mathfrak{J}}})} a_{w_{o_{\mathfrak{J}}}} = \pm a_{w'}$$

for some  $w' \in W$  with  $l(w') \geq l(yw) > l(y)$ .

4.7 THEOREM. (i) The elements  $\{a_y o_{\mathfrak{J}} e_J = a_y e_J : y \in Y_J, J \subseteq R\}$  are linearly independent and form a basis of  $H$ .

(ii) The elements  $\{a_y e_J o_{\mathfrak{J}} : y \in Y_J, J \subseteq R\}$  are linearly independent and form a basis of  $H$ .

PROOF. (i) Suppose that for each  $y \in Y_J$  and each  $J \subseteq R$  there is an element  $k_y \in K$  such that  $\sum_{J \subseteq R} \sum_{y \in Y_J} k_y a_y e_J = 0$ . Let

$$S_n = \sum_{J \subseteq R} \sum_{y \in Y_J, l(y) \geq n} k_y a_y e_J.$$

We show that if  $S_n = 0$ , then  $k_y = 0$  whenever  $l(y) = n$  and hence  $S_{n+1} = 0$ .

Let  $y_1, \dots, y_t$  be those elements of  $W$  for which  $l(y_i) = n$ . Then by 4.5, if  $y_i \in Y_{J(i)}$  for some  $J(i) \subseteq R$ ,

$$a_{y_i} e_{J(i)} = a_{y_i} + (\text{a linear combination of certain } a_w \text{ where } l(w) > l(y_i)).$$

Hence,

$$S_n = \sum_{i=1}^t k_{y_i} a_{y_i} + (\text{a linear combination of certain } a_w \text{ with } l(w) > n).$$

If  $S_n = 0$ , then as  $\{a_w : w \in W\}$  are a basis of  $H$ , we must have  $k_{y_i} = 0$  for all  $i$ ,  $1 \leq i \leq t$ . Then  $S_{n+1} = 0$ .

Since  $S_0 = 0$ ,  $k_y = 0$  for all  $y$  whenever  $l(y) = 0$ , and then  $S_1 = 0$ . By induction, all  $k_y$  are zero, and so  $\{a_y e_J : y \in Y_J, J \subseteq R\}$  is a set of linearly independent elements. As there are  $|W|$  of them, they must form a basis of  $H$ .

(ii) This is proved using similar arguments.

4.8 COROLLARY. (i) For any  $L \subseteq R$ , the elements of the set

$$\{a_y o_j e_J o_{\hat{L}} = a_y e_J o_{\hat{L}} : y \in Y_J, J \subseteq L\}$$

are linearly independent.

(ii) For any  $L \subseteq R$ , the elements of the set  $\{a_y e_J o_j e_L : y \in Y_J, J \supseteq L\}$  are linearly independent.

PROOF. (i)  $a_y e_J o_{\hat{L}} = \sum_{w \in W_J} a_{yw} o_{\hat{L}}$ . As  $J \subseteq L$ ,  $\hat{L} \subseteq \hat{J}$  and so  $a_{w_0} o_{\hat{L}} = a_{w_0}$ . Then

$$\begin{aligned} a_y e_J o_{\hat{L}} &= a_y o_{\hat{L}} + \sum_{w \in W_J, w \neq 1} a_{yw} o_{\hat{L}} \\ &= a_y + \sum_{w \in W_J, w \neq 1} a_{yw} o_{\hat{L}} \text{ as } y \in Y_J \\ &= a_y + (\text{a linear combination of certain } a_w \text{ with } l(w) > l(y)). \end{aligned}$$

The result now follows by using an argument similar to that used in the proof of 4.7.

(ii) For any  $y \in Y_J$ ,  $a_y e_J o_j = a_y + (\sum_{w \in W} k_w a_w)$ , where  $k_w \in K$  and  $k_w = 0$  if  $l(w) \leq l(y)$ . Then

$$\begin{aligned} a_y e_J o_j e_L &= a_y e_L + (\sum_{w \in W} k_w a_w) e_L, \quad k_w \in K \text{ given as above,} \\ &= a_y + (\sum_{w \in W} k'_w a_w) \text{ for certain } k'_w \in K, \text{ with } k'_w = 0 \text{ if } l(w) \leq l(y). \end{aligned}$$

Once again the result is given using an argument similar to that given in the proof of 4.7.

4.9 THEOREM. (i) For each  $a \in H$  and for any  $J \subseteq R$ , there exist elements  $k_y \in K$  such that

$$a o_j e_J = \sum_{y \in Y_J} k_y a_y e_J = (\sum_{y \in Y_J} k_y a_y o_j e_J).$$

(ii) For each  $a \in H$  and for any  $J \subseteq R$ , there exist elements  $k_y \in K$  such that

$$a e_J o_j = \sum_{y \in Y_J} k_y a_y e_J o_j.$$

PROOF. (i) As  $\{a_w : w \in W\}$  is a basis of  $H$ , we may write  $a = \sum_{w \in W} u_w a_w$  with  $u_w \in K$  for all  $w \in W$ . It is thus sufficient to express  $a_w o_j e_J$  as a linear combination of the elements  $\{a_y e_J : y \in Y_J\}$  for all  $w \in W$ . Use induction on  $l(w)$  to prove this.

If  $l(w) = 0$ , then  $w = 1$  and  $1 o_j e_J = (-1)^{j(w_0 j)} a_{w_0 j} e_J$ . The result is true for  $w = 1$  as  $w_0 j \in Y_J$ .

Suppose  $l(w) > 0$ . Let  $w = w_i w'$  for some  $w_i \in R$ ,  $w' \in W$ ,  $l(w) = l(w') + 1$ . By induction,

$$a_{w'} o_{\hat{j}} e_J = \sum_{y \in Y_J} u_y a_y e_J \text{ for some } u_y \in K.$$

Then

$$a_w o_{\hat{j}} e_J = a_i a_{w'} o_{\hat{j}} e_J = \sum_{y \in Y_J} u_y a_i a_y e_J.$$

Hence for each  $y \in Y_J$  we have to express  $a_i a_y e_J$  as a combination of  $\{a_v e_J : v \in Y_J\}$ . Now for any  $y \in Y_J$ ,

$$(4.10) \quad a_i a_y e_J = \begin{cases} -a_y e_J, & \text{if } y^{-1}(r_i) \in \Phi^-, \\ 0, & \text{if } y^{-1}(r_i) = r_j \text{ for some } r_j \in \Pi_J, \\ & \text{as then } a_i a_y = a_y a_i, \\ a_{w_i y} e_J, & \text{where } w_i y \in Y_J \text{ if } y^{-1}(r_i) \in \Phi^+, \\ & y^{-1}(r_i) \neq r_j \text{ for any } r_j \in \Pi_J. \end{cases}$$

The result follows.

(ii) Since  $\{a_y e_L o_{\hat{L}} : y \in Y_L, L \subseteq R\}$  is a basis of  $H$ , there exist elements  $u_y \in K$  such that

$$a e_J o_{\hat{j}} = \sum_{L \subseteq R} \sum_{y \in Y_L} u_y a_y e_L o_{\hat{L}}.$$

Choose any  $M \subseteq R$  with  $M \cap \hat{J} \neq \emptyset$ . Then  $a e_J o_{\hat{j}} e_M = 0$ ; so

$$\sum_{L \subseteq R} \sum_{y \in Y_L} u_y a_y e_L o_{\hat{L}} e_M = 0.$$

But  $o_{\hat{L}} e_M = 0$  if  $\hat{L} \cap M \neq \emptyset$ . So the only non-zero terms in the above equation involve those  $L \subseteq R$  for which  $\hat{L} \cap M = \emptyset$ . Thus

$$\sum_{L, M \subseteq L \subseteq R} \sum_{y \in Y_L} u_y a_y e_L o_{\hat{L}} e_M = 0.$$

By 4.8(ii),  $u_y = 0$  for all  $y \in Y_L$ ,  $M \subseteq L \subseteq R$ . Hence we have that  $u_y = 0$  for all  $y \in Y_L$ , with  $L \cap \hat{J} \neq \emptyset$ . Thus

$$a e_J o_{\hat{j}} = \sum_{L \subseteq J} \sum_{y \in Y_L} u_y a_y e_L o_{\hat{L}}.$$

Let  $S_J = \{w \in W : u_w \neq 0, w \in Y_L \text{ for some } L \subset J\}$ . Suppose  $S_J \neq \emptyset$ . Choose an element  $y_0 \in S_J$  of minimal length, and suppose  $y_0 \in Y_{J_0}$  for some  $J_0 \subset J$ . Consider

$$a e_J o_{\hat{j}} o_{\hat{j}_0} = \sum_{L \subseteq J} \sum_{y \in Y_L} u_y a_y e_L o_{\hat{L}} o_{\hat{j}_0}.$$

As  $J_0 \subset J$ ,  $e_J o_{\hat{j}} o_{\hat{j}_0} = e_J o_{\hat{j}_0} = 0$ . Then

$$(*) \quad \sum_{L \subset J} \sum_{y \in Y_L} u_y a_y e_L o_{\hat{L}} o_{\hat{j}_0} = 0.$$

Now if  $L \subset J$  and  $y \in Y_L$ ,

$$a_y e_L o_{\hat{L}} o_{\hat{J}_0} = a_y o_{\hat{J}_0} + \sum_{w \in W, l(w) > l(y)} k_w a_w$$

where  $k_w \in K$ , and  $a_y o_{\hat{J}_0} = \pm a_w$ , for some  $w \in W$  with  $l(w) \geq l(y)$ .

Since  $y_0$  is of minimal length in  $S_J$ , the coefficient of  $a_{y_0}$  on the left side of (\*) is  $u_{y_0}$ . As  $\{a_w : w \in W\}$  is a basis of  $H$ , so  $u_{y_0} = 0$ , which is a contradiction. Hence  $S_J = \emptyset$  and  $ae_J o_{\hat{J}} = \sum_{y \in Y_J} u_y a_y e_J o_{\hat{J}}$ .

REMARK. Let  $z \in Z$ . Then  $z$  can be regarded as an element of  $K$  in a natural way—it is the element  $z1_K = 1_K + \dots + 1_K$  ( $z$  times), where  $1_K$  is the identity of  $K$ .

4.11 COROLLARY. (1) For each  $w \in W$ , there exist rational integers  $u_y = u_y(w)$  such that  $a_w o_{\hat{J}} e_J = \sum_{y \in Y_J} u_y a_y o_{\hat{J}} e_J$ .

(2) For each  $w \in W$ , there exist rational integers  $u_y = u_y(w)$  such that

$$a_w e_J o_{\hat{J}} = \sum_{y \in Y_J} u_y a_y e_J o_{\hat{J}}$$

PROOF. (1) Follows from the proof of 4.9(i).

(2) List the elements  $y_1, \dots, y_m$  of  $Y_J$  in order of increasing length; if  $i < j$  then  $l(y_i) \leq l(y_j)$ . Let  $c_{ij}$  be the coefficient of  $a_{y_i}$  in  $a_{y_j} e_J o_{\hat{J}}$ . Clearly  $c_{ij}$  is an integer as  $a_{y_j} e_J o_{\hat{J}}$  is an integral combination of certain elements  $a_w$ ,  $w' \in W$ . Also,  $c_{ii} = 1$  for all  $i$ ,  $1 \leq i \leq m$ , and  $c_{ij} = 0$  if  $i < j$  by 4.6. Let  $h_i$  be the coefficient of  $a_{y_i}$  in  $a_w e_J o_{\hat{J}}$ . Clearly  $h_i$  is an integer, and

$$h_i = \sum_{j=1}^m k_j c_{ij} \quad \text{where} \quad a_w e_J o_{\hat{J}} = \sum_{i=1}^m k_i a_{y_i} e_J o_{\hat{J}}$$

for some  $k_i \in K$ . Hence,  $h_i = \sum_{j=1}^{i-1} k_j c_{ij} + k_i$ . Let  $i = 1$ . Then  $h_1 = k_1$ , an integer. Now use increasing induction on  $i$  to show  $k_i$  is an integer for all  $i$ ,  $1 \leq i \leq m$ .

4.12 THEOREM. (1)  $Ho_{\hat{J}} e_J$  is a left ideal of  $H$  with  $K$ -basis  $\{a_y o_{\hat{J}} e_J = a_y e_J : y \in Y_J\}$ . Hence  $\dim Ho_{\hat{J}} e_J = |Y_J|$ . Let  $Y_J = \{y_1, \dots, y_s\}$ , with  $l(y_i) \leq l(y_j)$  if  $i < j$ , and let  $H_{J,i} = \{\sum_{j=i}^s k_j a_{y_j} o_{\hat{J}} e_J : k_j \in K\}$ ; then

$$Ho_{\hat{J}} e_J = H_{J,1} > H_{J,2} > \dots > H_{J,s} > 0$$

is a composition series of  $Ho_{\hat{J}} e_J$  of left  $H$ -modules, and  $H_{J,i}/H_{J,i+1}$  affords the representation  $\lambda_{M_i}$  of  $H$ , where  $y_i^{-1} \in Y_M$ , and  $H_{J,s+1} = 0$ . Finally,  $H = \sum_{J \in R}^{\oplus} Ho_{\hat{J}} e_J$ , a direct sum of  $2^n$  left ideals, where  $n = |R|$ .

(2)  $He_J o_{\hat{J}}$  is a left ideal of  $H$  with  $K$ -basis  $\{a_y e_J o_{\hat{J}} : y \in Y_J\}$ . Hence  $\dim He_J o_{\hat{J}} = |Y_J|$ . Let  $Y_J = \{y_1, \dots, y_s\}$ , with  $l(y_i) \leq l(y_j)$  if  $i < j$ , and let

$$H_{J,i} = \left\{ \sum_{j=i}^s k_j a_{y_j} e_J o_{\hat{J}} : k_j \in K \right\};$$

then

$$He_J o_{\hat{j}} = H_{J,1} > H_{J,2} > \dots > H_{J,s} > 0$$

is a composition series of  $He_J o_{\hat{j}}$  of left  $H$ -modules, and  $H_{J,i}/H_{J,i+1}$  affords the representation  $\lambda_M$  of  $H$ , where  $y_i^{-1} \in Y_M$ , and  $H_{J,s+1} = 0$ . Finally,  $H = \sum_{J \subseteq R}^{\oplus} He_J o_{\hat{j}}$ , a direct sum of  $2^n$  left ideals, where  $n = |R|$ .

PROOF. The results follow by 4.7, 4.8, 4.10 and the fact that

$$\dim H = |W| = \sum_{J \subseteq R} |Y_J|.$$

4.13 COROLLARY.  $Ho_{\hat{j}} e_J$  and  $He_J o_{\hat{j}}$  are indecomposable left ideals of  $H$ , for all  $J \subseteq R$ , and they are isomorphic as left ideals of  $H$ .

PROOF. From the theory of Artinian rings and the fact that  $H/N$  is a direct sum of  $2^n$  irreducible components (see remarks at the end of Section 3), it follows that  $H$  can be expressed as the direct sum of  $2^n$  indecomposable left ideals. Hence  $Ho_{\hat{j}} e_J$  and  $He_J o_{\hat{j}}$  must be indecomposable left ideals of  $H$  for all  $J \subseteq R$ .

To show they are isomorphic, first note that  $He_J o_{\hat{j}} = Ho_{\hat{j}} e_J o_{\hat{j}}$ . Then define the homomorphism  $f_J: Ho_{\hat{j}} e_J \rightarrow He_J o_{\hat{j}}$  by  $f_J(ao_{\hat{j}} e_J) = ao_{\hat{j}} e_J o_{\hat{j}}$ , for all  $ao_{\hat{j}} e_J \in Ho_{\hat{j}} e_J$ . As  $f_J$  is given by right multiplication by  $o_{\hat{j}}$ , it is well defined and is a homomorphism of left ideals of  $H$ .  $f_J$  is onto, since  $He_J o_{\hat{j}} = Ho_{\hat{j}} e_J o_{\hat{j}}$  and an element  $ao_{\hat{j}} e_J o_{\hat{j}} \in He_J o_{\hat{j}}$  is the image under  $f_J$  of  $ao_{\hat{j}} e_J$ .  $f_J$  is one-one as  $\dim Ho_{\hat{j}} e_J = \dim He_J o_{\hat{j}}$ . Hence  $f_J$  is an isomorphism of left ideals of  $H$ .

4.14 COROLLARY. (1) For any  $L \subseteq R$ ,

$$Ho_{\hat{L}} = \sum_{J \subseteq L}^{\oplus} Ho_{\hat{j}} e_J o_{\hat{L}}, \quad \text{and} \quad \dim Ho_{\hat{L}} = \sum_{J \subseteq L} |Y_J| = |X_{\hat{L}}|.$$

(2) For any  $L \subseteq R$ ,

$$He_L = \sum_{J \supseteq L}^{\oplus} He_J o_{\hat{j}} e_L, \quad \text{and} \quad \dim He_L = \sum_{J \supseteq L} |Y_J| = |X_L|.$$

PROOF. Use 4.12 and 4.8.

4.15 THEOREM. For any  $J \subseteq R$ ,

$$\begin{aligned} He_J &= \{a \in H: aa_i = 0 \text{ for all } w_i \in J\} \\ &= \{a \in H: a(1 + a_i) = a \text{ for all } w_i \in J\}. \end{aligned}$$

Further,  $He_J = \sum_{J \subseteq L}^{\oplus} Ho_{\hat{L}} e_L$ , and  $He_J$  has basis  $\{a_w e_J : w \in X_J\}$  and dimension  $|X_J|$ . Finally,

$$\begin{aligned} Ho_{\hat{J}} e_J &= \{a \in H : aa_i = 0 \text{ for all } w_i \in J, ae_L = 0 \text{ for all } L \supset J\} \\ &= He_J \cap \left( \bigcap_{J \supset L} \ker e_L \right), \end{aligned}$$

where  $\ker e_L = \{a \in H : ae_L = 0\}$ .

PROOF. Clearly,  $He_J \leq \{a \in H : aa_i = 0 \text{ for all } w_i \in J\}$ . Conversely, take  $a \in H$  and suppose  $aa_i = 0$  for all  $w_i \in J$ . Then  $a(1 + a_i) = a$  for all  $w_i \in J$ , and so  $ae_J = a$ , and so  $a \in He_J$ . Thus the first part is proved.

Now  $Ho_{\hat{L}} e_L \leq He_J$  for all  $L \supseteq J$ , and so  $\sum_{L \supseteq J}^{\oplus} Ho_{\hat{L}} e_L \leq He_J$ . By 4.14,  $\dim He_J = |X_J|$ , and as  $\dim Ho_{\hat{L}} e_L = |Y_L|$ , we have  $He_J = \sum_{L \supseteq J}^{\oplus} Ho_{\hat{L}} e_L$ .

Let  $a = \sum_{w \in W} u_w a_w \in He_J$ , where  $u_w \in K$ . Let  $w_i \in J$ . Then  $aa_i = 0$ , and so  $\sum_{w \in W} u_w a_w a_i = 0$ . Now

$$\sum_{w \in W} u_w a_w a_i = \sum_{w \in W, w(r_i) \in \Phi^+} u_w a_{ww_i} - \sum_{w \in W, w(r_i) \in \Phi^-} u_w a_w = 0.$$

That is,

$$\sum_{w \in W, w(r_i) \in \Phi^-} u_w a_w - \sum_{w \in W, w(r_i) \in \Phi^+} u_w a_w = 0.$$

Since  $\{a_w : w \in W\}$  form a basis of  $H$ , we have  $u_{ww_i} = u_w$  for all  $w \in W$  with  $w(r_i) \in \Phi^-$ . Hence  $u_w = u_{ww_i}$  for all  $w \in W$ , with  $w(r_i) \in \Phi^+$ . Now if  $w \in W$ ,  $w$  can be expressed uniquely in the form  $w = yw_J$ , where  $y \in X_J$ ,  $w_J \in W_J$  and  $l(w) = l(y) + l(w_J)$ . Write  $w_J = w_{i_1} \dots w_{i_t}$ ,  $w_{i_j} \in J$ ,  $l(w_J) = t$ . By the above, we have

$$u_y = u_{yw_{i_1}} = \dots = u_{yw_J} = u_w.$$

Hence  $a = \sum_{y \in X_J} u_y a_y e_J$ . Conversely, for each  $y \in X_J$ ,  $a_y e_J \in He_J$ , and as  $\{a_y e_J : y \in X_J\}$  is linearly independent and  $\dim He_J = |X_J|$ ,  $\{a_y e_J : y \in X_J\}$  is a basis of  $He_J$ .

Finally,  $Ho_{\hat{J}} e_J \leq \{a \in H : aa_i = 0 \text{ for all } w_i \in J, ae_L = 0 \text{ for all } L \supset J\}$ . Let  $a = \sum_L \sum_{y \in Y_L} u_y a_y o_{\hat{L}} e_L$ ,  $u_y \in K$ , satisfy  $aa_i = 0$  for all  $w_i \in J$  and  $ae_L = 0$  for all  $L \supset J$ . Since  $a \in He_J$ ,  $u_y = 0$  for all  $y \in Y_L$  if  $J \not\subseteq L$ . So  $a = \sum_{L \supseteq J} \sum_{y \in Y_L} u_y a_y o_{\hat{L}} e_L$ . Set  $S_J = \{w \in W : u_w \neq 0, w \in Y_L, L \supset J\}$ . Suppose  $S_J \neq \emptyset$ . Then there exists an element  $y_0$  of minimal length in  $S_J$ ; suppose  $y_0 \in Y_M$ ,  $M \supset J$ . Then  $ae_M = 0$ . Also  $o_{\hat{J}} e_J e_M = 0$  as  $M \supset J$ . For other  $L \supset J$ , if  $y \in Y_L$ ,

$$\begin{aligned} a_y o_{\hat{L}} e_L e_M &= a_y e_L e_M = a_y + (\text{a combination of certain } a_w, \\ &w \in W, \text{ with } l(w) > l(y)). \end{aligned}$$

Then  $ae_M = 0$  gives  $\sum_{L \supset J} \sum_{y \in Y_L} u_y a_y o_{\hat{L}} e_L e_M = 0$ . As  $y_0$  is of minimal length in  $S_J$ , the coefficient of  $a_{y_0}$  in the left-hand side of the last equation is  $u_{y_0}$ . By the linear independence of  $\{a_w : w \in W\}$ , we have  $u_{y_0} = 0$ , which is a contradiction. Hence  $S_J = \emptyset$  and  $a = \sum_{y \in Y_J} u_y a_y o_{\hat{J}} e_J \in Ho_{\hat{J}} e_J$ . Thus

$$Ho_{\hat{J}} e_J = \{a \in He_J : ae_L = 0 \text{ for all } L \supset J\}.$$

4.16 THEOREM. For any  $J \subseteq R$ ,

$$Ho_J = \{a \in H : a(1 + a_i) = 0 \text{ for all } w_i \in J\}.$$

$Ho_J$  has basis  $\{a_w : w \in Y_{\hat{L}}, \hat{L} \subseteq \hat{J}\}$ , dimension  $|X_J|$  and  $Ho_J = \sum_{\hat{L} \supseteq J} He_{\hat{L}} o_{\hat{L}}$ . Finally,  $He_{\hat{J}} o_{\hat{J}} = \{a \in Ho_J : ao_L = 0 \text{ for all } L \supset J\}$ .

PROOF. Similar to the proof of 4.15.

4.17 LEMMA. Let  $\psi_J$  be the character of the representation of  $H$  on  $Ho_{\hat{J}} e_J$ . Then  $\psi_J$  takes values as follows: for each  $w \in W$ , let  $w = w_{i_1} \dots w_{i_t}$  be a reduced expression for  $w$ , and set  $J(w) = \{w_{i_j} : 1 \leq j \leq t\}$ . Then  $\psi_J(a_w) = (-1)^{l(w)} N_J(w)$ , where  $N_J(w) =$  the number of elements  $y \in Y_J$  such that  $y^{-1}(\prod_{J(w)}) \subseteq \Phi^-$ .

PROOF. Use 4.10.

4.18 LEMMA. Let  $\phi_J$  be the character of the representation of  $H$  on  $He_J$ . Then  $\phi_J$  takes values as follows: for  $w \in W$  let  $w = w_{i_1} \dots w_{i_t}$  be a reduced expression for  $w$ . Set  $J(w) = \{w_{i_j} : 1 \leq j \leq t\}$ . Then  $\phi_J(a_w) = (-1)^{l(w)} M_J(w)$ , where  $M_J(w) =$  the number of elements  $x \in X_J$  such that  $x^{-1}(\prod_{J(w)}) \subseteq \Phi^-$ . Also,  $M_J(w) = \sum_{L \supseteq J} N_L(w)$ .

PROOF.  $He_J$  has basis  $\{a_w e_J : w \in X_J\}$ . For any  $w_i \in R$ ,

$$a_i a_w e_J = \begin{cases} -a_w e_J & \text{if } w^{-1}(r_i) < 0, \\ a_{w_i w} e_J, & \text{where } w_i w \in X_J \text{ if } w^{-1}(r_i) > 0, \text{ and} \\ & w^{-1}(r_i) \neq r_j \text{ for any } r_j \in \Pi, \\ 0 & \text{if } w^{-1}(r_i) = r_j \text{ for some } r_j \in \Pi_J, \text{ for then} \\ & a_i a_w = a_w a_j \text{ and } a_j e_J = 0. \end{cases}$$

The result now follows.

4.19 LEMMA. Let  $\mu_J$  be the character of the representation of  $H$  on  $Ho_J$ . Then  $\mu_J$  takes values as follows: for each  $w \in W$ , let  $w = w_{i_1} \dots w_{i_t}$  be a reduced expression for  $w$ , and set  $J(w) = \{w_{i_j} : 1 \leq j \leq t\}$ . Then  $\mu_J(a_w) = (-1)^{l(w)} L_J(w)$ , where  $L_J(w) =$  the number of elements  $z \in Z_J$  such that  $z^{-1}(\prod_{J(w)}) \subseteq \Phi^-$ , and  $Z_J = \{w \in W : w(\Pi_J) \subseteq \Phi^-\}$ . Note that  $Z_J = \sum_{L \subseteq \hat{J}} Y_L$ .

PROOF.  $Ho_J$  has basis  $\{a_w : w \in Z_J\}$ . For all  $w_i \in R$ ,

$$a_i a_w = \begin{cases} -a_w & \text{if } w^{-1}(r_i) < 0, \\ a_{w_i w} & \text{if } w^{-1}(r_i) > 0. \end{cases}$$

If  $w \in Z_J$ ,  $w_i \in R$  and  $w^{-1}(r_i) > 0$ , then  $w_i w \in Z_J$ , for if  $r_j \in \Pi_J$ ,  $w(r_j) = -s$  for some  $s \in \Phi^+$ , and  $w_i(s) < 0$  if and only if  $s = r_i$ . But if  $s = r_i$ ,  $w^{-1}(r_i) = -r_j$ —impossible. The result now follows.

- 4.20 COROLLARY. (1)  $\phi_J = \sum_{J \supseteq L} \psi_L$  for all  $J \subseteq R$ .  
 (2)  $\mu_J = \sum_{J \supseteq L} \psi_L$  for all  $J \subseteq R$ .

A direct sum decomposition of  $H$  into indecomposable left ideals is equivalent to expressing the identity of  $H$  as a sum of mutually orthogonal primitive idempotents. Let  $1 = \sum_{J \subseteq R} q_J$  and  $1 = \sum_{J \subseteq R} p_J$  be the decompositions of 1 corresponding to the decompositions  $H = \{\sum_{J \subseteq R} Ho_J e_J$  and  $H = \sum_{J \subseteq R} He_J o_J$  respectively, where  $Hq_J = Ho_J e_J$  and  $Hp_J = He_J o_J$ . (There does not appear to be a specific expression for the  $q_J$  or the  $p_J$  in terms of  $\{a_y o_J e_J : y \in Y_J\}$  or  $\{a_y e_J o_J : y \in Y_J\}$  respectively).

4.21 THEOREM. Let  $\{q_J : J \subseteq R\}$  be a set of mutually orthogonal primitive idempotents with  $q_J \in Ho_J e_J$  for all  $J \subseteq R$  such that  $1 = \sum_{J \subseteq R} q_J$ . Then  $Ho_J e_J = Hq_J$ , and if  $N$  is the nilpotent radical of  $H$ ,  $No_J e_J = Nq_J$  is the unique maximal left ideal of  $Hq_J$ , and  $Hq_J/Nq_J \cong K$ .  $Hq_J/Nq_J$  affords the representation  $\lambda_J$  of  $H$  defined in 3.1. Finally,

$$H/N \cong \sum_{J \subseteq R}^{\oplus} Hq_J/Nq_J \cong K \oplus K \oplus \dots \oplus K, \quad 2^n \text{ summands, where } n = R.$$

PROOF. By the theory of Artinian rings,  $Nq_J$  is the unique maximal left ideal of  $Hq_J$ , and  $H/N \cong \sum_{J \subseteq R}^{\oplus} Hq_J/Nq_J$ . Since  $q_J \in Ho_J e_J$ ,  $Hq_J \subseteq Ho_J e_J$ . As

$$H = \sum_{J \subseteq R}^{\oplus} Hq_J = \sum_{J \subseteq R}^{\oplus} Ho_J e_J,$$

we must have  $Hq_J = Ho_J e_J$  for all  $J \subseteq R$ . Then  $Nq_J = NHq_J = NHo_J e_J = No_J e_J$  is the unique maximal left ideal of  $Hq_J$ . But

$$\left\{ \sum_{y \in Y_J, y \neq w_0 J} u_y a_y o_J e_J : u_y \in K \right\}$$

is a maximal left ideal of  $Ho_J e_J$  (see 4.10), and so

$$Nq_J = \left\{ \sum_{y \in Y_J, y \neq w_0 J} u_y a_y o_J e_J : u_y \in K \right\}.$$



Then  $Hq_J/Nq_J$  is a one-dimensional  $H$ -module generated by  $a_{w_0} o_{\bar{J}} e_J + Nq_J$  which affords the representation  $\lambda_J$  of  $H$ , and since every element of  $Hq_J/Nq_J$  is of the form  $ka_{w_0} o_{\bar{J}} e_J + Nq_J$  for some  $k \in K$ ,  $Hq_J/Nq_J \cong K$  for all  $J \subseteq R$ . Hence the result.

**4.22 THEOREM.** *Let  $\{p_J : J \subseteq R\}$  be a set of mutually orthogonal primitive idempotents with  $p_J \in He_J o_{\bar{J}}$  for all  $J \subseteq R$  such that  $1 = \sum_{J \subseteq R} p_J$ . Then  $He_J o_{\bar{J}} = Hp_J$ , and if  $N$  is the nilpotent radical of  $H$ ,  $Ne_J o_{\bar{J}} = Np_J$  is the unique maximal left ideal of  $Hp_J$ , and  $Hp_J/Np_J \cong K$ .  $Hp_J/Np_J$  affords the representation  $\lambda_J$  of  $H$  defined in 3.1. Finally,  $H/N \cong \sum_{J \subseteq R}^{\oplus} Hp_J/Np_J \cong K \oplus K \oplus \dots \oplus K$ ,  $2^n$  summands, where  $n = |R|$ .*

**4.23 LEMMA.**  *$\{ka_{w_0 w_0} o_{\bar{J}} e_J : k \in K\}$  and  $\{ka_{w_0 w_0} e_J o_{\bar{J}} : k \in K\}$  are minimal submodules of  $Ho_{\bar{J}} e_J$  and  $He_J o_{\bar{J}}$  respectively, where  $w_0 w_0$  is the unique element of maximal length in  $Y_J$ . These minimal left ideals both afford the representation  $\lambda_{\bar{J}}$  of  $H$ , where  $J = \{w_i \in R : \text{there exists } w_j \in J \text{ with } w_0 w_j = w_i w_0\}$ , or, alternatively,  $\Pi_{\bar{J}}$  is defined by  $w_0(\Pi_J) = -\Pi_{\bar{J}}$ .*

**4.24 NOTE.** By the same methods,  $H = \sum_{J \subseteq R}^{\oplus} e_J o_{\bar{J}} H$  and  $H = \sum_{J \subseteq R}^{\oplus} o_{\bar{J}} e_J H$ , both being direct sum decompositions of  $H$  into  $2^n$  right ideals, where  $n = |R|$ . Further,  $e_J o_{\bar{J}} H$  has  $K$ -basis  $\{e_J o_{\bar{J}} a_y : y^{-1} \in Y_J\}$ , and  $o_{\bar{J}} e_J H$  has  $K$ -basis  $\{o_{\bar{J}} e_J a_y : y^{-1} \in Y_J\}$ . All the results for the left ideals  $He_J, Ho_J, He_J o_{\bar{J}}$  and  $Ho_{\bar{J}} e_J$  have analogues for the right ideals  $e_J H, o_J H, o_{\bar{J}} e_J H$  and  $e_J o_{\bar{J}} H$  respectively.

Let  $G$  be a finite group with a split  $(B, N)$  pair of rank  $n$  and characteristic  $p$  with Weyl group  $W$ , and let  $K$  be a field of characteristic  $p$ . Then the above decomposition of  $H = H_K(G, B)$  gives a decomposition of  $1_B^G$ , where  $1_B$  is the principal character of the subgroup  $B$  of  $G$ , which will be discussed in a later paper.

### 5. The Cartan matrix of $H$

We have that  $H = \sum_{J \subseteq R}^{\oplus} U_J$ , where  $U_J = Ho_{\bar{J}} e_J$  is an indecomposable left  $H$ -module. Thus  $\{U_J : J \subseteq R\}$  are the principal indecomposable  $H$ -modules.  $\{U_J/\text{rad } U_J : J \subseteq R\}$ , where  $\text{rad } U_J$  is the unique maximal submodule of  $U_J$ , are irreducible  $H$ -modules, such that  $M_J = U_J/\text{rad } U_J$  affords the representation  $\lambda_J$  of  $H$ .

**DEFINITION.** The *Cartan matrix*  $C$  of  $H$ , where  $H$  is of type  $(W, R)$ , with  $|R| = n$ , is a  $2^n \times 2^n$  matrix with rows and columns indexed by the subsets of  $R$ , and if we write  $C = (c_{JL})$ , then

$$c_{JL} = \text{the number of times } M_L \text{ is a composition factor of } U_J.$$

5.1 THEOREM. For all  $J, L \subseteq R$ ,

$$c_{JL} = |Y_J \cap (Y_L)^{-1}| = |Y_L \cap (Y_J)^{-1}| = c_{LJ}.$$

Hence  $C$  is a symmetric matrix.

PROOF.  $U_J$  has  $K$ -basis  $\{a_y o_j e_J = a_y e_J : y \in Y_J\}$ . Let  $y_1, \dots, y_s$  be all the elements of  $Y_J$  written in order of increasing length; if  $i > j$  then  $l(y_i) \geq l(y_j)$ . Then set  $U_J(i) = \{\sum_{j \geq i} k_{y_j} a_{y_j} e_J : k_{y_j} \in K\}$ .  $U_J(i)$  is a left ideal of  $H$  for all  $i$ , and  $U_J(i) > U_J(i+1)$  for all  $i, 1 \leq i \leq s-1$ . Then  $U_J = U_J(1) > U_J(2) > \dots > U_J(s) > 0$  is a composition series of  $U_J$ , with  $U_J(i)/U_J(i+1)$  being an irreducible  $H$ -module with basis  $a_{y_i} e_J + U_J(i+1)$  and affording the irreducible representation  $\lambda_L$ , defined in 3.1, where  $L$  is determined as follows: recall 4.10; let  $w_j \in R$  and  $y_i \in Y_J$ . Then

$$a_j a_{y_i} e_J = \begin{cases} -a_{y_i} e_J & \text{if } y_i^{-1}(r_j) < 0, \\ 0 & \text{if } y_i^{-1}(r_j) = r_k \text{ for some } r_k \in \Pi, \\ a_{w_j y_i} e_J & \text{where } w_j y_i = y_l \text{ for some } y_l \in Y_J \text{ with } i < l, \text{ if} \\ & y_i^{-1}(r_j) > 0 \text{ but } y_i^{-1}(r_j) \neq r_k \text{ for any } r_k \in \Pi. \end{cases}$$

Hence

$$\lambda_L: a_j \rightarrow \begin{cases} -1 & \text{if } y_i^{-1}(r_j) < 0, \\ 0 & \text{if } y_i^{-1}(r_j) > 0. \end{cases}$$

That is,  $y_i^{-1} \in Y_L$ .

Hence  $c_{JL}$  = the number of elements  $y \in Y_J$  such that  $y^{-1} \in Y_L$

$$= |Y_J \cap (Y_L)^{-1}| = |Y_L \cap (Y_J)^{-1}|$$

since if  $y \in Y_J \cap (Y_L)^{-1}$ , then  $y^{-1} \in Y_L \cap (Y_J)^{-1}$ .

5.2 THEOREM. Let  $H$  be the 0-Hecke algebra over the field  $K$  of type  $(W, R)$ , where  $W$  is indecomposable. Then if  $|R| > 1$ ,  $H$  has three blocks. If  $|R| = 1$ , then  $H$  has two blocks.

PROOF. If  $|R| = 1$ , then  $W = W(A_1)$  and  $H = H(1 + a_1) \oplus H(-a_1)$ , where  $R = \{w_1\}$ . Both  $(1 + a_1)$  and  $(-a_1)$  are primitive idempotents as well as being central. Hence  $H$  has only two blocks.

Now suppose that  $|R| > 1$ .  $e_R = [1 + a_{w_0}]$  and  $(-1)^{l(w_0)} a_{w_0}$  are primitive and centrally primitive idempotents in  $H$  and so correspond to two distinct blocks.

The other primitive idempotents in  $H$ , that is,  $\{q_J: J \neq \emptyset, R\}$  as in 4.21, determine at least one other block. We will show that provided  $W$  is indecomposable the Cartan matrix  $C'$  corresponding to the indecomposables  $U_J$  for  $J \neq \emptyset, R$  and the irreducibles  $M_L$  for  $L \neq \emptyset, R$  cannot be expressed in the form  $C' = \begin{bmatrix} C_1 & 0 \\ 0 & C_2 \end{bmatrix}$  (see Dornhoff (1972), Theorem 46.3).

Suppose that  $C'$  can be put in the form above. Let

$$S_1 = \{J \subset R: U_J \text{ and } M_J \text{ index the rows and columns of } C_1\},$$

$$S_2 = \{J \subset R: U_J \text{ and } M_J \text{ index the rows and columns of } C_2\}.$$

Suppose for some  $J \subset R$ ,  $|J| = n - 1$  (where  $n = |R|$ ), that  $J \in S_1$ . Then we show

(1) for all  $L \subset R$  with  $|L| = n - 1$ ,  $L \in S_1$ ,

(2) by decreasing induction on  $|J|$  for all  $J \neq \emptyset, R$  that  $J \in S_1$ .

(a) Suppose  $J = \{w_1, \dots, \hat{w}_j, \dots, w_n\}$  and  $L = \{w_1, \dots, \hat{w}_{j+1}, \dots, w_n\}$ , where the nodes corresponding to  $w_j$  and  $w_{j+1}$  in the graph of  $W$  are joined. Then the order of  $w_j w_{j+1}$  is greater than 2. Now  $w_{0J} = w_j \in Y_J$  and  $w_{0L} = w_{j+1} \in Y_L$ . Since the order of  $w_j w_{j+1}$  is greater than 2,  $w_{j+1} w_j \in Y_J$  and  $w_j w_{j+1} \in Y_L$ ; that is,  $w_{j+1} w_j \in Y_J \cap (Y_L)^{-1}$ . Hence  $J \in S_1$  if and only if  $L \in S_1$ .

Hence if there is some  $J \in S_1$ , with  $|J| = n - 1$ , then all  $L \subset R$  with  $|L| = n - 1$  are in  $S_1$  by the above.

(b) Suppose that for all  $J \subset R$  with  $|J| > m$  that  $J \in S_1$ . Choose  $L \subset R$  with  $|L| = m$ . We show  $L \in S_1$ . Suppose  $L = \{w_{i_1}, \dots, w_{i_m}\}$  with  $1 \leq i_1 < \dots < i_m \leq n$ . Since  $W$  is indecomposable and  $L \neq \emptyset, R$ , then  $|Y_L| > 1$ . Choose some  $w_{i_j} \in L$  and  $w_k \in \hat{L}$  such that  $w_{i_j} w_k$  has order  $r$ , where  $r \geq 3$ . Then  $w_{i_j} w_{0\hat{L}} \in Y_L$  (as  $w_{0\hat{L}}(r_{i_j}) \neq r_{i_j}$  for any  $r_{i_j} \in \Pi_L$ , for  $w_{0\hat{L}}(r_{i_j}) = r_{i_j}$  for some  $r_{i_j} \in \Pi_L$  implies that  $r_{i_j} = r_{i_j}$  and  $w_{0\hat{L}}$  is a product of reflections corresponding to roots orthogonal to  $r_{i_j}$ , and so for all  $w_k \in \hat{L}$ ,  $w_{i_j} w_k = w_k w_{i_j}$ , which is a contradiction). Now consider  $(w_{i_j} w_{0\hat{L}})^{-1} = w_{0\hat{L}} w_{i_j}$ . Then suppose  $w_{i_j} \in L$ ,  $w_{i_j} \neq w_{i_j}$ . Then  $w_{0\hat{L}} w_{i_j}(r_{i_j}) \in \Phi^+$ . Also  $w_{0\hat{L}} w_{i_j}(r_{i_j}) \in \Phi^-$ . Suppose  $w_k \in \hat{L}$ . Then

$$\begin{aligned} w_{0\hat{L}} w_{i_j}(r_k) &= w_{0\hat{L}}(r_k + u r_{i_j}) \quad \text{with } u \geq 0 \\ &= w_{0\hat{L}}(r_k) + u w_{0\hat{L}}(r_{i_j}). \end{aligned}$$

If  $u = 0$ , that is, if  $w_{i_j} w_k = w_k w_{i_j}$ , then  $w_{0\hat{L}} w_{i_j}(r_k) \in \Phi^-$ . If  $u > 0$ , as  $w_{0\hat{L}}(r_k) = -r_k$  for some  $r_k \in \Pi_{\hat{L}}$ , and  $w_{0\hat{L}}(r_{i_j}) \in \Phi^+$ ,  $w_{0\hat{L}}(r_{i_j}) \neq r_{i_j}$  for any  $r_{i_j} \in \Pi_L$ , we have  $w_{0\hat{L}} w_{i_j}(r_k) \in \Phi^+$ . Hence  $w_{0\hat{L}} w_{i_j} \in Y_M$ , where

$$\begin{aligned} M &= \{L - \{w_{i_j}\}\} \cup \{w_k \in \hat{L}: w_{i_j} w_k \text{ has order } > 2\} \\ &= \{L - \{w_{i_j}\}\} \cup \{w_k \in \hat{L}: \text{the node corresponding to } w_k \text{ in the graph of} \\ &\quad W \text{ is joined to that corresponding to } w_{i_j}\}. \end{aligned}$$

Now  $|M| > |L|$  if the node corresponding to  $w_{i_j}$  is joined to at least two nodes corresponding to elements of  $\hat{L}$ , and then  $L \in S_1$  by induction.

Let  $P_i$  be the node of the graph of  $W$  which corresponds to  $w_i \in R$ ,  $1 \leq i \leq n$ . Then suppose  $P_{i_j}$  is joined to only one  $P_k$  for all  $w_k \in \hat{L}$ . Then the above argument shows that  $L = \{w_{i_1}, \dots, w_{i_m}\}$  and  $M = \{w_{i_1}, \dots, \hat{w}_{i_j}, \dots, w_{i_m}, w_k\}$  belong to the same  $S_i$ , where  $i = 1$  or  $i = 2$ . Since  $|L| \leq n - 2$ ,  $|\hat{L}| \geq 2$ . Let  $w_{k_1}$  and  $w_{k_2}$  be any two elements of  $\hat{L}$ , such that there exists a sequence  $P_{k_1} = P_{j_0}, P_{j_1}, \dots, P_{j_r} = P_{k_2}$  of nodes such that  $P_{j_i}$  and  $P_{j_{i+1}}$  are joined for all  $i$ ,  $0 \leq i \leq r - 1$ , and  $P_{j_i}$  corresponds to an element of  $L$  for all  $i$ ,  $1 \leq i \leq r - 1$ . If  $r = 1$ , then  $P_{k_1}$  and  $P_{k_2}$  are joined. Without loss of generality, we may suppose there exists  $w_{i_s} \in L$  such that  $P_{i_s}$  is joined to  $P_{k_1}$ . Then let  $M = \{L - \{w_{i_s}\} \cup \{w_{k_1}\}\}$ .  $M$  and  $L$  belong to the same  $S_i$ , and by the above, as  $M$  has an element  $w_{k_1}$  such that  $w_{k_1} w_{i_s}$  and  $w_{k_1} w_{k_2}$  both have order  $> 2$ , where  $w_{i_s}, w_{k_2} \in \hat{M}$ ,  $w_{i_s} \neq w_{k_2}$ , then  $M \in S_1$ . If  $r = 2$ , then  $L$  and  $M$  are in the same  $S_i$ , where  $M = \{L - \{w_{j_1}\} \cup \{w_{k_1}, w_{k_2}\}\}$ , and by induction  $M \in S_1$ . If  $r > 2$ , define

$$L_0 = L,$$

$$L_1 = \{L - \{w_{j_1}\}\} \cup \{w_{j_0}\},$$

...

$$L_{r-2} = \{L_{r-3} - \{w_{j_{r-3}}\}\} \cup \{w_{j_{r-3}}\}.$$

Then  $L_0, L_1, \dots, L_{r-2}$  are all in the same  $S_i$ , and by the above,  $L_{r-2} \in S_1$ .

Hence  $L \in S_1$ . Then  $S_2 = \emptyset$ , and so  $H$  has precisely three blocks.

**5.3 THEOREM.** *Let  $H$  be a 0-Hecke algebra of type  $(W, R)$ . Suppose  $W$  is decomposable, and let  $W = W_1 \times W_2 \times \dots \times W_r$ , where each  $W_i$  is an indecomposable Coxeter group, and the corresponding Coxeter system is  $(W_i, R_i)$ . Let  $H_i$  be the 0-Hecke algebra of type  $(W_i, R_i)$ , and let  $m_i$  be the number of blocks of  $H_i$ . Then  $H$  has  $m_1 m_2 \dots m_r$  blocks.*

**PROOF.** Suppose that  $1 = \sum_{i=1}^t e_i$  where the  $e_i$  are mutually orthogonal central primitive idempotents in  $H$ . Then the number of blocks of  $H$  is equal to  $t$ .

Now for all  $w \in W_i, w' \in W_j$ , where  $1 \leq i, j \leq r$  and  $i \neq j$ , we have that

$$a_w a_{w'} = a_{ww'} = a_{w'w} = a_{w'} a_w,$$

and so it follows that if  $f_i$  is a centrally primitive idempotent of  $H_i$ , then  $f_1 \dots f_r$  is a centrally primitive idempotent of  $H$ . Suppose  $1_{H_i} = \sum_{j=1}^{t(i)} f_{ij}$  where for a fixed  $i$ ,  $\{f_{ij} : 1 \leq j \leq t(i)\}$  is a set of mutually orthogonal central primitive idempotents in  $H_i$ . Then  $1_H = \sum_{j_1=1}^{t(1)} \dots \sum_{j_r=1}^{t(r)} f_{1j_1} \dots f_{rj_r}$ , a sum of mutually orthogonal central primitive idempotents in  $H$ , and so  $H$  has  $t(1)t(2) \dots t(r)$  blocks, where  $t(i) = m_i$ .

### Acknowledgement

This work was carried out while the author was at the Mathematics Institute, University of Warwick, Coventry, England, supported by a scholarship from the Association of Commonwealth Universities.

### References

- N. Bourbaki (1968), *Groupes et algèbres de Lie*, Chapitres 4, 5 et 6 (Hermann, Paris).  
R. W. Carter (1972), *Simple groups of Lie type* (John Wiley and Sons, New York).  
C. W. Curtis and I. Reiner (1962), *Representation theory of finite groups and associative algebras* (Interscience Publishers, New York).  
L. Dornhoff (1972), *Group representation theory*, Part B. Marcel Decker, Inc., New York).  
L. Solomon (1968), 'A decomposition of the group algebra of a finite Coxeter group', *J. Algebra*, **9**, 220–239.  
R. Steinberg (1967), *Lectures on Chevalley groups* (Yale University).

Technical Education Division  
Education Department of W.A.  
36 Parliament Place  
West Perth, Western Australia, 6005,  
Australia