

Size-density strategy displayed by *Diadema africanum* linked with the stability of urchin-barrens in the Canary Islands

NANCY CABANILLAS-TERÁN¹, JOSÉ A. MARTÍN², RUBER RODRÍGUEZ-BARRERAS³ AND ÁNGEL LUQUE²

¹Laboratory of Marine Resources, Central Department of Research, Universidad Laica Eloy Alfaro de Manabí, Ciudadela Universitaria, Vía a San Mateo, Manta, Manabí, Ecuador, EC130802, ²Department of Biology, Marine Sciences Faculty, University of Las Palmas de Gran Canaria, 35017 Las Palmas de G.C., Canary Islands, Spain, ³Department of Biology, University of Puerto Rico, Río Piedras. PO Box 23360, San Juan 00931-3360, Puerto Rico

The sea urchin Diadema africanum is considered a key herbivore in sublittoral ecosystems of the Canary Islands. Spatial and temporal variability in population structure was carried out at Gran Canaria. We performed a morphometric and population density analysis during 2005, 2006 and 2007 at four sites in zones of Gran Canaria. The study considered a vertical gradient (5, 10 and 20 m depth) during both seasons, the cold season (February and March) and the warm season (October and November). The sea urchin D. africanum in Gran Canaria exhibited an overall density of 7.59 ± 2.92 urchin m^{-2} . A two-way ANOVA evidenced spatial differences in mean abundance of the species, while seasonality was not relevant. The vertical analysis of the abundance of D. africanum showed differences, the smaller sizes appeared at greater depths. The Aristotle's lantern width decreased in a vertical gradient, being remarkable between 10 and 20 m. Findings of uniformity in size over time, a stable range of high densities and the lack of a relationship between the size of the sea urchins and the season reveals that the density–size strategy displayed by D. africanum which explains in turns the high stability of the urchin barrens, which, once developed, remain as areas of permanent desertification in subtidal depths throughout the Canary Archipelago.

Keywords: *Diadema africanum*, urchin barren, morphometry, abundance, Canary Islands

Submitted 6 June 2014; accepted 26 July 2014; first published online 10 September 2014

INTRODUCTION

Echinoderms have been called a ‘boom–bust’ phylum because of the large fluctuations of some species (Uthicke *et al.*, 2009). There are many factors involved in the regulation of the structure of urchin population; where predation, recruitment, pollution, diseases, large-scale oceanographic events, food supplies, niche variability are some of them (Hyman, 1955; Dayton *et al.*, 1984; Hughes, 1994; Sala & Zabala, 1996; Sala & Ballesteros, 1997; 2007; Lessios, 2013; García-Sanz *et al.*, 2014). Hence, the ecological processes triggering these population explosions cannot be easily determined (Lessios *et al.* (2001); Hernández *et al.*, 2008; Lessios, 2013). Expansion and crashes are more common than was usually thought. Overpopulation of sea urchins may generate extensive areas devoid of macroalgae (Lessios, 2013) called ‘urchin barren’.

The presence of barrens has been extensively documented in the coastal zones of temperate environments and in subtropical and tropical regions (Sammarco, 1982; McClanahan & Curtis, 1991; McClanahan, 1994; Alves *et al.*, 2001; Brito *et al.*, 2004; Tuya *et al.*, 2005; Clemente *et al.*, 2010; Lessios, 2013; García-Sanz *et al.*, 2014). The development of barrens has been reported on rocky substrates of the Macaronesian islands,

without associated anthropogenic factors (Lawrence, 1975). However, higher densities of the sea urchins have been linked with overfishing and areas protected from high wave action (Bortone *et al.*, 1991; McClanahan *et al.*, 1996; Tuya *et al.*, 2004a, 2005, 2007). Barrens are considered undesirable due to the negative impact on fisheries productivity (Clemente *et al.*, 2010). Several studies have evaluated the role of *Diadema africanum* in subtidal coastal ecosystems, because it has become the most voracious consumer of the sublittoral vegetation in the Canary Archipelago with densities up to 12 urchin m^{-2} (Alves *et al.*, 2003; Brito *et al.*, 2004; Hernández *et al.*, 2008; Clemente *et al.*, 2010).

The long-spined sea urchin *Diadema africanum* (Rodríguez *et al.*, 2013) is considered one of the most well known echinoid because of its important role as a benthic grazer in the Eastern Atlantic (Randall *et al.*, 1964; Tuya *et al.*, 2005; Lessios, 2013). The spatial distribution of *Diadema* has been linked to a variety of factors, including predation (Dayton *et al.*, 1984; Sala & Ballesteros, 1997), settlement and recruitment (Young & Chia, 1982), availability of trophic resources (Menge, 1992) and competitive interactions (Hagen & Mann, 1992). Changes in density may influence changes in the dimensions of some morphological structures of sea urchins, particularly the Aristotle's lantern (Garrido, 2003). This pentagonal structure forms an efficient scraping tool to acquire food, and its analysis provides important information about food limitation or plastic resource allocation (Ebert, 1980; Levitan, 1991).

Corresponding author:

N. Cabanillas-Terán

Emails: nanchamex@gmail.com/nancy.cabanillas@uleam.edu.ec

Therefore, the main objectives of this study were to: (1) determine the current population density of *D. africanum*; and (2) compare differences in the Aristotle's lantern across different space, time and vertical conditions.

MATERIALS AND METHODS

Study area

Surveys were conducted during 2005, 2006 and 2007 in four different areas on the island of Gran Canaria in the Canary Islands (Figure 1). The sites were La Catedral (LC: 28°10'57"N 15°24'11"W), Sardina del Norte (SAR: 28°08'56"N 15°42'02"W), Risco Verde (RV: 27°51'24"N 15°23'9"W) and Puerto Rico (PR: 27°47'47"N 15°43'59"W). Gran Canaria Island is nearly a round island with different environmental characteristics at the four sampling sites. LC and SAR are located far from urban areas, and are under the influence of north-east winds. Both sites have deep waters and are characterized by extensive rocky substrates. RV is a shallow water area on the east side of the island, protected from the wind, has a sandy substrate below 15 m and is adjacent to an urban centre. The fourth site, PR, is located to the south-west and is shallow, with a mixed substrate of rocks, gravel and sand below 10 m depth.

LC and SAR are colonized by sea urchins and exhibit no erect algae, but there are some encrusting red algae; whereas RV and PR are characterized by erect algae, predominantly of the genera *Padina*, *Dictyota* and *Stypocaulon* spp. (see Cabanillas-Terán, 2009 for more details). Censuses were conducted at three different depth ranges according to the specific topography of each site. Ten metres was the common depth for all sites (Table 1). Samples were taken in two seasons: warm (October–November) and cold (March–April), according to the time when the water exhibits its maximum (23–24°C) and minimum (17–18°C) peaks of temperature (Hernández et al., 2011).

Survey methodology and population structure

Mean density was estimated using a belt-transect methodology (Vanderklift & Kendrick, 2004). At each site, eight random transects of 5 m² were placed parallel to the shoreline

Table 1. Sampling depths for the four sites in Gran Canaria.

Sites	Depth 1	Depth 2	Depth 3
Risco Verde	5 m	10 m	–
Puerto Rico	5 m	10 m	–
La Catedral	–	10 m	20 m
Sardina del Norte	–	10 m	20 m

and separated 10 m from each other. All crevices and small holes were carefully inspected to avoid missing any individuals. Censuses were performed in 2005, 2006 and 2007 at each of the sampling sites in both the cold and warm seasons at different depths (Table 1). A total of 792 *Diadema africanum* were collected randomly across all sites for morphometric analysis during cold and warm seasons of 2005, 2006 and the cold season of 2007 (25 individuals by site by depth). Individuals were collected randomly, brought to the laboratory, labelled and frozen for later analyses. Once in the laboratory, the Aristotle's lantern diameters were measured (caliper error: 0.05 mm). Depth range in Table 1 was established according to the presence of *D. africanum* at these depths. We divided individuals in four size-classes following Tuya et al. (2004a, b): class 1: 1.5–3.5 cm; class 2: 3.5–5.4 cm; class 3: 5.5–7.4 cm; and class 4: 7.5–10 cm.

Statistical analysis

The Kolmogorov–Smirnov and Levene tests were run to test normality and homogeneity of variance for all biometric parameters (Zar, 2010). We ran a two-way analysis of variance (ANOVA) to test spatial and temporal differences in mean abundance and (10 m) mean size. This model incorporates time as the fixed and site as the random factor. In addition, we ran different one-way ANOVAs to determine potential differences in depth at each site of mean density, but also differences in depth and among sites for the Aristotle's lantern width. Multiple *post-hoc* Tukey's HSD analyses were run when ANOVAs were significant to identify specific subsets. Furthermore, we ran the parametric Pearson correlation among mean density of *D. africanum* and depth. All statistical




Fig. 1. Study area and sampling sites in Gran Canaria: Sardina (SAR); PR (Puerto Rico); Risco Verde (RV); La Catedral (LC).

Table 2. A two-way analysis of variance on the abundances, and sizes of the sea urchin *Diadema africanum* at four sites of Gran Canaria. SS, sum of squares; MS, mean of squares.

	Mean density*			Mean size*		
	df	F	P	df	F	P
Time (t)	5	1.768	0.092	4	16.481	<0.001
Site (s)	7	35.18	<0.001	3	147.837	<0.001
S × T	17	3.306	<0.001	11	18.198	<0.001
		SS	MS		SS	MS
Residuals	355	3758.2	10.59	415	196,562	0.474

*, comparisons were run only for 10 m data (the only common depth among sites).

analyses were performed in the free-license statistical software R-3.0.1 with a P value of 0.05 (R Core Team, 2013).

RESULTS

The sea urchin *D. africanum* exhibited an overall density of 7.59 ± 2.92 (mean \pm standard deviation) urchin m^{-2} at Gran Canaria. The overall abundance varied from 2.09 ± 1.42 urchin m^{-2} (PR) to 10.12 ± 3.17 urchin m^{-2} (SAR). Seasonal highest density of *D. africanum* was recorded at RV during the cold season of 2005 at 5 m (12.98 ± 4.47 urchin m^{-2}), while PR displayed the lowest density during the cold season in the same depth and year (0.75 ± 0.45 urchin m^{-2}) (Figure 2). A two-way ANOVA found spatial differences in mean abundance of the species, while seasonality was not relevant (Table 2). The *post-hoc* analysis

determined that the most important differences in abundances occurred between RV with respect to LC/PR ($P < 0.001$), and between PR/SAR ($P < 0.001$). On the other hand, the abundance across a vertical gradient (depth) exhibited higher densities at 20 m in LC and in RV at 5 m (Figure 2). Differences in depth were found within PR (ANOVA, $F = 38.69$, $df = 1$, $P < 0.001$) and SAR ($F = 13.20$, $df = 1$, $P < 0.001$), while LC ($F = 2.01$, $df = 1$, $P = 0.159$) and RV ($F = 0.0008$, $df = 1$, $P < 0.997$) did not exhibit a great vertical changes in abundance. Furthermore, we did not find a strong relationship between depth and mean density of *D. africanum* ($r = 0.247$, $N = 384$, $P < 0.01$).

Overall mean size of *D. africanum* during the study was 5.06 ± 1.01 cm. The maximum average mean size was recorded in RV (5.53 ± 0.92 cm), followed by PR, LC and lastly SAR, with 4.29 ± 0.68 cm. The spatial variability showed that RV and PR were the two sites with greater mean sizes at some intervals, whereas SAR and LC remained below 6.0 cm (Figure 3). The classes two and three together represented 94% of the whole sample, and class 2 alone represented more than 65% of the individuals collected during the study.

Differences in depth of mean sizes showed that in PR and RV mean sizes increase between 5 and 10 m, whereas in LC and SAR sizes diminished between 10 and 20 m (Figure 3). The two-way ANOVA performed only at 10 m found significant spatial and temporal mean sizes (Table 2). All sites exhibited differences ($P < 0.001$) among the others in a *post-hoc* analysis, whereas temporal differences were not remarkable except between the warm seasons of 2005 and 2006 ($P = 0.095$), and between the cold seasons of 2006 and 2007 ($P = 0.997$).

The morphometric analysis of the Aristotle’s lantern structure found that the width of the structure decreased in a




Fig. 2. Spatial and temporal density of *Diadema africanum* in Gran Canaria from cold season 2005 to winter of 2007 at three depths (5, 10, and 20 m) at four sites (see Materials and Methods for sites acronyms). The X axis represents time with cold-2005 (c-05), warm-2006 (w-06), etc. Depths are represented by square (5 m), circle (10 m), and triangle (20 m). Bars represent 95% of confidence interval of the mean.




Fig. 3. Seasonal mean size distribution of *Diadema africanum* at four sites in Gran Canaria. The X axis represents time with cold-2005 (c-05), warm-2006 (w-06), etc. Depths are represented by square (5 m), circle (10 m), and triangle (20 m). Bars represent 95% of confidence interval of the mean. There are some missing data because not all sites were measured in all seasons. Lack of information (SAR c-05, PR w-06 and c-07 at 10 m) represents missing data.

vertical gradient, notably between 10 and 20 m ($F = 12.307$, $d = 2$, $P < 0.001$). A *post hoc* comparison found a lack of difference only between 5 and 10 m ($P = 0.245$) (Figure 3A). On the other hand, the width of the lantern was greater at PR and RV and lower in LC and SAR (Figure 3B). A one-way ANOVA found spatial differences ($F = 8.43$, $d = 3$, $P < 0.001$), whereas *post hoc* comparisons did not find differences between either RV and PR or LC and SAR ($P > 0.05$).

DISCUSSION

The general stability displayed at all sites in abundance confirms the high seasonal stability of *D. africanum* population on Gran Canaria. Temporal stability in abundance of this invertebrate agrees with previous studies conducted throughout the Canary Archipelago (Brito *et al.*, 1984; Bortone *et al.*, 1991; Tuya *et al.*, 2006, 2007; Hernández *et al.*, 2007; Clemente *et al.*, 2010; Hernández *et al.*, 2011). The existence of a depth gradient was notable at least for PR and SAR. This result agrees with other studies conducted in Canary Islands (Tuya *et al.*, 2004b; Hernández *et al.*, 2005, 2007). The hydrodynamic action in intertidal and subtidal environments affects the distribution patterns of marine organisms (Roberts *et al.*, 2006).

The existence of higher densities of *D. africanum* in deeper areas in PR and SAR could be supported by the negative relationship between depth and wave effect (Roberts *et al.*, 2006). The sea urchin *D. africanum* is weakly resistant to unidirectional hydrodynamic forces, because its morphology does not allow a large adhesive surface area to attach to the bottom (Tuya *et al.*, 2007). Higher hydrodynamics in shallow areas also may limit *D. africanum* recruitment, according to Tuya *et al.* (2007), which affects population

growth. Nevertheless, the vertical distribution of *D. africanum* in the Macaronesia Archipelago showed that the species was more abundant in shallow water areas (Entrambasaguas *et al.*, 2008), as occurs with *Diadema antillarum* in the Caribbean (Hendler *et al.*, 1995).

The vertical Aristotle's lantern results (Figure 4) agreed with other studies that indicate a limited growth of *D. africanum* at greater depths (Levitan, 1988; Garrido, 2003; Hernández *et al.*, 2005, 2007). However, we only found this behaviour in LC, whereas SAR displayed an inverse relationship and PR and RV did not exhibit a clear pattern (Figure 3). The reduction in width of this structure may be a result of food restrictions due to a decrease in vegetation cover and/or lower primary production limited by light intensity (Alves *et al.*, 2001; Lessios *et al.*, 2001a, b). In addition, Levitan (1991) found that *Diadema* might adjust its body size as a function of available resources, as an adaptive strategy to protect against the harmful effects of increased density.

A limiting factor shaping abundance and size of *D. africanum* are food resources, and in barrens this factor is determinant. Urchin barrens have greater development in deeper zones due to continuous grazing activity, whereas in shallower waters, grazing remains restricted during night-time (Tuya *et al.*, 2004c). It has been demonstrated for sea urchins of temperate zones that recruitment may be favoured under high density conditions, due to higher larval settlement success (Balch *et al.*, 1998; Tuya *et al.*, 2006). However, when densities reached 10 urchin m^{-2} or higher, sea urchins may collapse benthic production (Tuya *et al.*, 2004b). The existence of a correlation between size and urchin barrens development found here is consistent with other studies (Black *et al.*, 1982; Ebert, 1983; McClanahan (1991); Levitan & Sewell, 1998; Garrido, 2003). For instance, we found that barrens with less algal cover have higher densities but smaller urchins.




Fig. 4. The Aristotle's lantern width variation on a depth gradient (A) and among the four sites in Gran Canaria (B). Grey columns represent 95% of confidence interval of the mean.

Further studies on the trophic ecology would provide a better understanding of how *D. africanum* can persist over long periods with a high reproductive success and seasonal stability, even at the limit of the carrying capacity of the environment. Despite extensive studies on the biological role of this species, there are still some questions as to how this species is dominant and is the key genera on both sides of the Atlantic.

To conclude, population density of *D. africanum* exhibited high temporal stability in Gran Canaria, while changing across a vertical gradient and between sites. The existence of higher densities and smaller sizes in deeper areas explains how the density–size strategy displayed by *Diadema* sp. (Levitan, 1989, 1991) accounts for the high stability of the urchin barrens, once developed, remaining as areas of permanent desertification at subtidal depths throughout the Canary Archipelago.

ACKNOWLEDGEMENTS

We would like to thank the logistic support provided by the Department of Biology, Marine Sciences Faculty, University of Las Palmas de Gran Canaria and the University Foundation. We also want to express our gratitude to the students of scientific diving for supporting fieldworks.

FINANCIAL SUPPORT

The first author expresses gratitude to the Mexican National Council for Science and Technology (CONACyT) for a scholarship.

REFERENCES

Alves F.M.A., Chicharo L.M., Serrão E. and Abreu A.D. (2001) Algal cover and sea urchin spatial distribution at Madeira Island (NE Atlantic). *Scientia Marina* 65, 383–392.

Alves F.M.A., Chicharo L.M., Serrão E. and Abreu A.D. (2003) Grazing by *Diadema antillarum* (Philippi) upon algal communities on rocky substrates. *Scientia Marina* 67, 307–311.

Balch T., Scheibling R.E., Harris L.G., Chester C.M. & Robinson S.M.C. (1998) Variation in settlement of *Strongylocentrotus droebachiensis* in

the northwest Atlantic: effect of spatial scale and sampling method. In Mooi R. and Telford M. (eds) *Echinoderms*. Rotterdam: A.A. Balkema, pp. 555–560.

Black R., Johnson M.S. and Trendall J.T. (1982) Relative size of Aristotle's lantern in *Echinometra mathaei* occurring at different densities. *Marine Biology* 71, 101–106.

Bortone S.A., Van Tassell J., Brito A., Falcon J.M. and Bundrick C.M. (1991) A visual assessment of the inshore fishes and fishery resources of El Hierro, Canary Islands: a Baseline Survey. *Scientia Marina* 55, 529–541.

Brito A., Hernández J.C., Falcón J.M., García N., González-Lorenzo G., Gil-Rodríguez M.C., Cruz-Reyes A., Herrera G., Sancho A., Clemente S., Cubero E., Girard D. and Barquín J. (2004) El Erizo de lima (*Diadema antillarum*) una especie clave en los fondos rocosos litorales de Canarias. *Makaronesia* 6, 68–86.

Brito T., Cruz E. and Moreno-Pérez J.M. (1984) Fauna marina de las Islas Canarias. In Bacallado J.J. (ed.) *Fauna marina y terrestre del Archipiélago Canario*. Edirca: Las Palmas de Gran Canaria, pp. 42–65.

Cabanillas-Terán N. (2009) *Ecología y estatus trófico del erizo de mar Diadema antillarum* (Philippi, 1845) en los fondos rocosos de las Islas Canarias (Gran Canaria, España). PhD thesis. Universidad de Las Palmas de Gran Canaria, Spain, available at: <http://hdl.handle.net/10553/2589> (accessed 8 August 2014).

Clemente S., Hernández J.C., Rodríguez A. and Brito A. (2010) Identifying keystone predators and the importance of preserving functional diversity in sublittoral rocky bottoms. *Marine Ecology Progress Series* 413, 55–67.

Dayton P.K., Currie V., Gerrodette T., Keller B.D., Rosenthal R. and Tresca D.V. (1984) Patch dynamics and stability of some California kelp. *Ecological Monographs* 54, 254–289.

Ebert T.A. (1980) Relative growth of sea urchin jaws: an example of plastic resource allocation. *Bulletin of Marine Science* 30, 467–474.

Ebert T.A. (1983) Recruitment in echinoderms. In Jangoux M. and Lawrence J.M. (eds) *Echinoderm studies*. Rotterdam: A.A. Balkema, pp. 169–203.

Entrambasaguas L., Pérez-Ruzafa A., García-Charton J.A., Stobart B. and Bacallado J.J. (2008) Abundance, spatial distribution and habitat relationships of echinoderms in the Cabo Verde Archipelago (eastern Atlantic). *Marine and Freshwater Research* 59, 477–488.

García-Sanz S., Navarro P.G. and Tuya F. (2014) Contrasting recruitment seasonality of sea urchin species in Gran Canaria, Canary Islands (eastern Atlantic). *Mediterranean Marine Science* 15, 475–481.

- Garrido M.J.** (2003) *Contribución al conocimiento de Diadema antillarum Philippi 1845, en Canarias*. PhD thesis. Universidad de Las Palmas de Gran Canaria, Spain.
- Hagen N.T. and Mann K.H.** (1992) Functional response of the predators american lobster *Homarus americanus* (Milne-Edwards) and Atlantic wolffish *Anarhichas lupus* (L.) to increasing numbers of the green sea urchin *Strongylocentrotus droebachiensis* (Müller). *Journal of Marine Experimental Biology and Ecology* 159, 89–112.
- Hendler G., Miller J.E., Pawson D.L. and Kier P.M.** (1995) *Sea stars, sea urchins, and allies. Echinoderms of Florida and the Caribbean*. Washington, DC: Smithsonian Institution Press. Washington, DC and London, 390 pp.
- Hernández J.C., Clemente S. and Brito A.** (2011) Effects of seasonality on the reproductive cycle of *Diadema* aff. *antillarum* in two contrasting habitats: implications for the establishment of a sea urchin fishery. *Marine Biology* 158, 2603–2615.
- Hernández J.C., Clemente S., Brito A., Falcon J.M., García N. and Barquin J.** (2005) Estado de las Poblaciones de *Diadema antillarum* (Echinoidea: Diademataceae) y del recubrimiento de macroalgas en las Reservas Marinas de Canarias: Patrones de distribución espacial. *Vieraea* 33, 367–383.
- Hernández J.C., Clemente S., Sangil C. and Brito A.** (2007) Actual status of *Diadema* aff. *antillarum* populations and macroalgal cover in the Marine Protected Areas comparing to a Highly Fished Area (Canary Islands–Eastern Atlantic Ocean). *Aquatic Conservation: Marine and Freshwater Ecosystems* 18, 1091–1108.
- Hernández J.C., Clemente S., Sangil C. and Brito A.** (2008) The key role of the sea urchin *Diadema* aff. *antillarum* in controlling macroalgae assemblages throughout the Canary Islands (eastern subtropical Atlantic): an spatio-temporal approach. *Marine Environmental Research* 66, 259–270.
- Hughes T.P.** (1994) Catastrophes, phase shifts and large-scale degradation of a Caribbean coral reef. *Science* 265, 1547–1551.
- Hyman L.H.** (1955) *The invertebrates: Echinodermata* (Volume 4). New York: McGraw-Hill.
- Lawrence J.M.** (1975) On the relationships between marine plants and sea urchins. *Oceanography and Marine Biology: an Annual Review* 13, 213–286.
- Lessios H.A.** (2013) ‘Natural’ population density fluctuations of echinoids. Do they help predict the future? In Fernández-Palacios J.M., de Nascimento L., Hernández J.C., Clemente S., González A. and Díaz-González J.P. (eds) *Climate change perspectives from the Atlantic: past, present and future*. Servicios de Publicaciones, Universidad de La Laguna, pp. 341–359.
- Lessios H.A., Garrido M.J. and Kessing B.D.** (2001a) Demographic history of *Diadema antillarum*, a keystone herbivore on Caribbean reefs. *Proceedings of the Royal Society of London Series B—Biological Sciences* 268, 2347–2353.
- Lessios H.A., Kessing B.D. and Pearse J.S.** (2001b) Population structure and speciation in tropical seas: global phylogeography of the sea urchin *Diadema*. *Evolution* 55, 955–975.
- Levitan D.R.** (1988) Density-dependent size regulation and negative growth in the sea urchin *Diadema antillarum* Philippi. *Oecologia* 76, 627–629.
- Levitan D.R.** (1989) Density-dependent size regulation in *Diadema antillarum*: effects on fecundity and survivorship. *Ecology* 70, 1414–1424.
- Levitan D.R.** (1991) Skeletal changes in the test and jaws of the sea urchin *Diadema antillarum* when food-limited. *Marine Biology* 106, 431–435.
- Levitan D.R. and Sewell M.A.** (1998) Male abundance and fertilization success in free-spawning marine invertebrates: review of the evidence and fisheries implications. In Jamieson G.S. and Campbell A. (eds) *Proceedings of the North Pacific Symposium on Invertebrate Stock Assessment and Management. Volume 125*. Canadian Special Publication in Fisheries and Aquatic Science, National Research Press, Canada, pp. 159–164.
- McClanahan T.R.** (1994) Kenyan coral reef lagoon fish: associations with reef management, complexity and sea urchins. *Coral Reefs* 13, 231–241.
- McClanahan T.R. and Curtis J.D.** (1991) Population regulation of the rock-boring sea urchin *Echinometra mathaei* de Blainville. *Journal of Experimental Marine Biology and Ecology* 147, 121–146.
- McClanahan T.R., Kamukuru A.T., Muthiga N.A., Yebio M.G. and Obura D.** (1996) Effect of sea urchin reductions on algae, coral, and fish populations. *Conservation Biology* 10, 136–154.
- Menge B.A.** (1992) Community regulation: under what conditions are bottom-up factors important on rocky shores? *Ecology* 73, 755–765.
- Randall J.E., Schroeder R.E. and Starck W.A.** (1964) Notes on the biology of the echinoid *Diadema antillarum*. *Caribbean Journal of Science* 4, 421–433.
- R Core Team** (2013) *R: a language and environment for statistical computing*. Vienna: R Foundation for statistical computing. Available at: <http://www.R-project.org/> (accessed 8 August 2014).
- Roberts D.E., Cummins S.P., Davis A.R. and Chapman M.G.** (2006) Structure and dynamics of sponge-dominated assemblages on exposed and sheltered temperate reefs. *Marine Ecology Progress Series* 321, 19–30.
- Rodríguez A., Hernández J.C., Clemente S. and Coppard S.E.** (2013) A new species of *Diadema* (Echinodermata: Echinoidea: Diademataceae) from the eastern Atlantic Ocean and a neotype designation of *Diadema antillarum* (Philippi, 1845). *Zootaxa* 3636, 144–170.
- Sala E. and Ballesteros E.** (1997) Partitioning of space and food resources by three fish of the genus *Diplodus* (Sparidae) in a Mediterranean rocky intertidal ecosystem. *Marine Ecology Progress Series* 152, 273–283.
- Sala E., Boudouresque C.F. and Harmelin-Vivien M.** (1998) Fishing, trophic cascades and the structure of algal assemblages: evaluation of an old but untested paradigm. *Oikos* 82, 425–439.
- Sala E. and Zabala M.** (1996) Fish predation and the structure of sea urchin *Paracentrotus lividus* populations in the NW Mediterranean. *Marine Ecology Progress Series* 140, 71–81.
- Sammarco P.W.** (1982) Echinoid grazing as a structuring force in coral communities: whole reef manipulations. *Journal of Experimental Marine Biology and Ecology* 61, 31–55.
- Tuya F., Boyra A., Sánchez-Jerez P., Barbera C. and Haroun R.J.** (2004a) Relationships between rocky-reef fish assemblages, the sea urchin *Diadema antillarum* and macroalgae throughout the Canarian Archipelago. *Marine Ecology Progress Series* 278, 157–169.
- Tuya F., Boyra A., Sánchez-Jerez P., Barbera C. and Haroun R.J.** (2004b) Can one species determine the structure of the benthic community on a temperate rocky reef? The case of the long-spined sea-urchin *Diadema antillarum* (Echinodermata: Echinoidea) in the eastern Atlantic. *Hydrobiologia* 519, 211–214.
- Tuya F., Cisneros-Aguirre J., Ortega-Borges L. and Haroun R.J.** (2007) Bathymetric segregation of sea urchins on reefs of the Canarian Archipelago: role of flow-induced forces. *Estuarine, Coastal Shelf Science* 73, 481–488.
- Tuya F., Martín J.A. and Luque A.** (2004c) Patterns of nocturnal movement of the long-spined sea urchin *Diadema antillarum* (Phillipi) in Gran Canaria (the Canary Islands, Central East Atlantic Ocean). *Helgoland Marine Research* 58, 26–31.

- Tuya F., Ortega-Borges L., Del Rosario-Pinilla A.B. and Haroun R.J.** (2006) Spatio-temporal variability in a key herbivore, the long-spined black sea urchin (*Diadema antillarum*, Echinodermata: Echinoidea) in the Canary Islands. *Journal of the Marine Biological Association of the United Kingdom* 86, 791–797.
- Tuya F., Sánchez-Jerez P. and Haroun R.J.** (2005) Influence of fishing and functional group of algae on sea urchin control of algal communities in the eastern Atlantic. *Marine Ecology Progress Series* 287, 255–260.
- Uthicke S., Schaffelke B. and Byrne M.** (2009) A boom-bust phylum? Ecological and evolutionary consequences of density variations in echinoderms. *Ecological Monographs* 79, 3–24.
- Vanderklift M.A. and Kendrick G.A.** (2004) Variation in abundances of herbivorous invertebrates in temperate subtidal rocky reef habitats. *Marine and Freshwater Research* 55, 93–103.
- Young C.M. and Chia F.S.** (1982) Factors controlling spatial distribution of the sea cucumber *Psolus chitonoides*: settling and post-settling behavior. *Marine Biology* 69, 195–205.
- and
- Zar J.H.** (2010) *Biostatistical analysis*. 5th edition. Upper Saddle River, NJ: Prentice-Hall.
- Correspondence should be addressed to:**
N. Cabanillas-Terán
Departamento Central de Investigación, Universidad Laica Eloy Alfaro de Manabí, Ciudadela Universitaria, Vía San Mateo, Manta, Manabí, Ecuador
emails: nanchamex@gmail.com/nancy.cabanillas@uleam.edu.ec