

Otitis externa: what is the problem with getting it right? A mixed-methods study in primary and secondary care

H Mohammed¹, M W Mather¹, J Lumb¹, C C Butler² and J A Wilson³

¹Department of ENT, Freeman Hospital, Newcastle upon Tyne, ²Nuffield Department of Primary Care Health Sciences, Oxford University and ³Department of Otolaryngology, Population Health Sciences Institute, Newcastle University, UK

Main Article

Dr H Mohammed takes responsibility for the integrity of the content of the paper

Cite this article: Mohammed H, Mather MW, Lumb J, Butler CC, Wilson JA. Otitis externa: what is the problem with getting it right? A mixed-methods study in primary and secondary care. *J Laryngol Otol* 2022;**136**: 486–491. <https://doi.org/10.1017/S0022215121003649>

Accepted: 26 May 2021

First published online: 25 November 2021

Key words:

Otitis Externa; Primary Health Care; *Pseudomonas Aeruginosa*; Osteomyelitis

Author for correspondence:

Dr H Mohammed,
Freeman Hospital, Freeman Road,
High Heaton,
Newcastle upon Tyne NE7 7DN, UK
E-mail: hmohammed@nhs.net

Abstract

Objective. Otitis externa accounts for 1.1–1.3 per cent of patient presentations in primary care and 25 per cent of urgent referrals to ENT. This study aimed to explore otitis externa clinical decision-making at the primary-secondary care interface, otitis externa prevalence and recent trends in antimicrobial resistance in otitis externa related bacterial isolates and ototopical prescribing.

Method. This is a mixed-methods study drawing on data from primary and secondary care and open National Health Service sources.

Results. A total of 101 general practitioner survey respondents reported frequently prescribing oral antibiotics for otitis externa. General practitioner consultations for otitis externa increased 25 per cent over 15 years. General practitioner ototopical preparations cost the National Health Service £7 410 440 in 2006 and £11 325 241 in 2016. A total of 162 consecutive hospital otitis externa-related bacterial isolates yielded 128 *pseudomonas* species, with 18 that were resistant to gentamicin and 7 that were resistant to ciprofloxacin. Ten guidelines reviewed showed systematic inconsistencies.

Conclusion. General practitioners reported regularly prescribing oral antibiotics for otitis externa. Antimicrobial drug resistance is common in otitis externa. The available guidance is suboptimal.

Introduction

Diffuse inflammation of the external auditory meatus (otitis externa) has been estimated to have an annual UK primary care prevalence of 1 per cent, with 3 per cent of sufferers referred to secondary care.¹ Key recommended treatments are ototopical drops, analgesia and avoidance of precipitating factors.²

We were concerned with the increased promotion of fluoroquinolone drops for the treatment of ear infections because they are still a key part of systemic therapy for necrotising otitis externa, and resistance driven by prior topical use may reduce efficacy in severe infection. In this study, we therefore aimed to explore the following. In primary care, we aimed to analyse: (1) otitis externa decision-making, guideline content, awareness and adherence, and (2) consultation rates and costs. In secondary care, we aimed to analyse: (1) otitis externa and necrotising otitis externa inpatient activity, and (2) the antimicrobial resistance profile of otitis externa-related bacterial isolates and content of available guidelines.

Materials and methods

Relevant components of the study were registered with the Clinical Effectiveness and Audit department of the Newcastle hospitals.

Data sources

Primary care

For primary care, the data sources were as follows. The first data source was from an otitis externa decision-making in primary care online survey (Figure 1) of a convenience sample of general practitioners and general practitioner trainees in North East England. It comprised research contacts in 12 clinical commissioning groups in North East England and North Cumbria. The office that distributed the questionnaire was not able to provide the total number of general practitioners to whom the questionnaire was sent. The questionnaire was designed based on the contents of the National Institute for Health and Care Excellence (NICE) clinical knowledge scenario. It was then discussed with and revised by the senior author and then piloted in a group of general practitioner trainees. The final version was posted online. A link to the survey was emailed to 500 general practitioner trainees in North East England.

Table 1. Summary of surveys of general practitioner trainees and general practitioners

Parameter	General practitioner trainees	General practitioners	Total
Respondents (<i>n</i>)	64	37	101
Post-graduate experience (mean; years)	6.7	14.5	9.6
Indications for ear swab (<i>n</i> (%))			
– Recurrent episodes	47 (73)	30 (81)	77 (76)
– Failure of first choice	38 (59)	23 (62)	61 (60)
Routinely prescribe (<i>n</i> (%))			
Otomize® spray	60 (94)	37 (100)	97 (96)
EarCalm® spray	10 (16)	13 (35)	23 (23)
Oral antibiotics	27 (42)	19 (51)	46 (46)
Follow guidelines (<i>n</i> (%))			
– National Institute for Health and Care Excellence	36 (56)	2 (5)	38 (38)
– Regional	22 (34)	15 (41)	37 (37)
– None	12 (19)	21 (57)	33 (33)
Advise patients (<i>n</i> (%))			
– Avoid use of cotton buds	51 (80)	35 (95)	86 (85)
– Keep ears dry when bathing or showering	47 (73)	32 (86)	79 (78)
– Avoid swimming or use ear plugs	50 (78)	29 (78)	79 (78)

Table 2. Incidence of otitis externa by age group*

Parameter	0–18 years	19–64 years	65+ years	All
	Patients (<i>n</i> (%))	Patients (<i>n</i> (%))	Patients (<i>n</i> (%))	Patients (<i>n</i> (%))
Diagnosed in last 12 months	318 (0.21)	1147 (0.75)	433 (0.28)	1898 (1.25)
Diagnosed at any point in their lifetime	1852 (1.22)	10742 (7.06)	4467 (2.93)	17061 (11.21)

*North West Primary Care Data, *n* = 152 238

Results

Primary care

Otitis externa decision-making in primary care

Our 101 respondents (Table 1) showed that 42 per cent of general practitioner trainees and 54 per cent of general practitioners prescribe oral antibiotics, most often amoxicillin (14 of 27 trainees) and flucloxacillin (12 of 19 general practitioners). Systemic treatments were used similarly (40 per cent) among the 20 respondents who had an ENT trainee placement.

Workload of otitis externa

The primary care data are summarised in Table 2. The annual incidence of otitis externa in the North West primary care catchment area was 1.25 per cent. Extrapolation to the population of England in 2015 equates to around 700 000 patients.

Cost of topical preparations

From 2006 to 2016, there was a steady increase in total prescription costs for community otitis externa topical prescriptions, from £7 million to £11 million (Table 3). Expenditure on ciprofloxacin drops rose most rapidly, increasing 12-fold in 3 years.

Secondary care

Hospital Episode Statistics data

In-patient otitis externa admissions in England showed a 122 per cent overall increase from 2611 patients in 2002 to 5786 patients in 2020 (Table 4). In 2002 to 2003, the mean length of stay was two days. The 2019 to 2020 bed occupancy equates to approximately 3.1 million at £346 per day.⁶ Necrotising otitis externa workload increased by 950 per cent in the same interval, at a cost of approximately £6.1 million.

Antimicrobial resistance profile

Pseudomonas species were isolated in almost 80 per cent (128 of 162) of the consecutive hospital swabs. *Staphylococcus aureus* was isolated from 41 swabs (25.3 per cent). Three out of five isolated streptococcus species were resistant to gentamicin. Multiple pathogens were grown from 20 swabs. One *pseudomonas* isolate was resistant to all antibiotics tested. Table 5 summarises resistance patterns in isolated bacteria according to the European Committee on Antimicrobial Susceptibility Testing guidelines for systemic antibiotic use.⁷

Guidelines

The guidelines reviewed (Table 6) show discrepancies in first line and second line recommendations. Some⁸ lacked clarity

Table 3. National Health Service indicative price of topical ear preparations ranked by 2016 total expenditure

Topical drug	NHS IP (£)	Volume (ml)	Price per ml (£)	Total expenditure (£)	
				2006 unless stated	2016
Otomize® (dexamethasone, acetic acid, neomycin sulfate) spray	3.27	5	0.65	2 983 774	4 297 422
Sofradex® (gramicidin, framycetin sulfate, dexamethasone) drops	7.50	8	0.94	1 502 165	2 657 946
Gentamicin 0.3% / hydrocortisone acetate 1% ear drops	26.19	10	2.62	712 190 (in 2015)	1 838 203
Gentisone HC® (gentamicin 0.3% / hydrocortisone acetate 1%) drops	4.76	10	0.47	1 418 689	16 108
Flumetasone 0.02%/clioquinol/1% drops	11.34	7.5	1.51	187 589 (in 2014)	1 751 250
Locorten-Vioform® (clioquinol, flumetasone pivalate) drops	1.76	7.5	0.23	453 339	3 987
Earcalm®* (glacial acetic acid) spray	7.99 [†]	5	1.60	54 982	345 183
Ciloxan® (ciprofloxacin) drops	4.70	5	0.94	12 270 (in 2011)	145 557
Chloramphenicol 5% drops	63.69	10	6.37	4 081	90 147
Betnesol-N® (betamethasone and neomycin) drops	2.39	10	0.24	280 997	82 759
Chloramphenicol 10% drops	84.70	10	8.47	110	37 589
Aluminium acetate 8% drops	No Information	No information		16 370	31 506
Aluminium acetate 13% drops	75 [‡]	No information	N/A	18 013	10 617
Otosporin® (hydrocortisone, polymyxin B, neomycin) drops	7.45	10	0.75	555 687	500
Total				7 290 455	11 308 777

*Available over the counter; [†]over the counter price; [‡]price quoted from pharmacy. NHS = National Health Service; IP = indicative price

Table 4. Hospital Episode Statistics data for admissions with otitis externa and necrotising otitis externa

Year	Otitis externa (n)	Bed days* (n)	Necrotising otitis externa (n)	Bed days* (n)
2002–2003	2611	5312	107	1130
2019–2020	5786	9019	1127	15 346

*Finished consultant episode bed days

concerning prophylaxis (i.e. water exclusion from the ears). Some lacked specificity; for example, the local Hertfordshire guidelines⁹ advise consideration of an oral antibiotic if otitis externa is 'severe' but without defining severity. Suspected tympanic membrane perforation was associated with considerable divergence: some guidelines⁸ recommend non-ototoxic topical treatment while others^{10,11} state any topical antibiotic class can be used for up to seven days.

Discussion

This study is a multi-method review of several facets of the diagnosis and management of otitis externa. It included prospectively assessed data from general practitioners, national level data for both primary and secondary care, and detailed investigations into the microbiology and cost implications of otitis externa.

The 101 general practitioners who responded to the survey reported frequently prescribing oral antibiotics for otitis externa. General practitioner consultations for otitis externa increased 25 per cent over 15 years. General practitioner ototoxic preparations cost over £11.3 million in 2016. The 1.25 per cent otitis externa prevalence we estimated in North West

Table 5. Resistance patterns in 169 isolated bacteria

Bacteria	Antibiotic	Resistance (n (%))
Pseudomonas species*	Ciprofloxacin	7 (5.5)
	Gentamicin	18 (14)
	Meropenem	2 (1.5)
	Tazocin	2 (1.5)
Staphylococcus aureus [†]	Flucloxacillin	2 (4.7)
	Gentamicin	1 (2)
	Tetracycline	4 (10)
	Erythromycin	5 (12)

*n = 128; [†]n = 41

England is similar to a prior UK report¹ and higher than the prevalence in the USA.¹² Hospital admissions for necrotising otitis externa increased 700 per cent over 14 years, and otitis externa admissions increased in number and duration.

Our survey of general practitioners and general practitioner trainees highlighted a number of significant issues around implementing otitis externa guidelines, which have also been noted in the USA.¹³ Most of our respondent general practitioners report having accessed guidelines, yet 42 per cent of surveyed general practitioner trainees and 51 per cent of general practitioners would prescribe oral antibiotics, which is notably higher than previous surveys^{1,14,15} and contrary to most guidance. Not only is oral treatment less effective than topical treatment,¹⁶ it encourages the development of antimicrobial resistance.¹⁷ There was also some variation in terms of prescription of topical agents and the type of agent prescribed. A number of factors could influence the selection of topical agent in primary care, such as being mentioned in

Table 6. Comparison of 10 commonly used otitis externa UK guidelines

Guideline	First line recommendations	Second line recommendations	Detailed prophylaxis	Data source
NICE CKS ²⁵	Avoiding aggravation factors. Cleaning ear canal and analgesia. Then antibiotics and steroid drops. Acetic acid stated to be equivalent to topical antibiotics in week 1	Not mentioned	Detailed advice including recommendation of using ear plugs	BMJ; Cochrane American
BMJ best practice ⁸	Topical antibiotic solutions, non-ototoxic (quinolones) if TM perforation suspected and based on patient preference	Not mentioned	Prevention of water ingress	Cochrane American and other references
North American ²⁴	Topical and systemic analgesia, topical non-ototoxic antibiotics and steroids. Systemic antibiotics if infection extends beyond ear canal or if specific host factors exist	Not mentioned	Yes, including cotton wool with Vaseline®	Cochrane and other references
Public Health England ²⁶	Analgesia for pain relief. Apply localised heat (e.g. a warm flannel). Systemic antibiotics if systemic symptoms or spreading cellulitis	Topical acetic acid or neomycin sulphate with corticosteroid	Nil	American, NICE CKS and Cochrane
North Yorkshire ¹¹	Acetic acid 2% spray for 7 days and analgesia. Topical antibiotics could be used in case of TM perforation for less than 10 days if infection is present. Systemic antibiotics if systemic symptoms or spreading cellulitis	Locorten-Vioform®	Nil	NICE CKS
North East London ²⁷	Acetic acid 2%	Neomycin sulphate with corticosteroid	Nil	NICE CKS and public health
Hertfordshire ⁹	Adequate analgesia. Do not prescribe topical quinolones (unlicensed)	Acetic acid 2% spray. In more severe otitis externa, topical antibiotics with or without steroids. Oral antibiotics for severe cases	Nil	NICE CKS
North Tyneside ²⁸	Consider 2% acetic acid ear drops if mild symptoms (over the counter). Topical antibiotic and steroid drops	Refer to nurse practitioner ear care clinic	General advice (do not poke or let shampoo or soap into ears)	Nil
Northern Lincolnshire ²⁹	Acetic acid 2%	Neomycin sulphate with corticosteroid (maximum 2 weeks)	Not mentioned	NICE CKS
North Cumbria referral guidelines ¹⁰	Swab and dry-mop. Topical treatment drops or spray (topical antibiotics could be used in case of TM perforation for less than 10 days if infection is present). If pinna involved, oral antibiotics long-term or as prophylaxis	Not mentioned	Consider underlying skin condition, close diabetic control, water precautions, review shampoo, regular (acetic acid)	Nil

*This is prescription guidance. NICE CKS = National Institute for Health and Care Excellence Clinical Knowledge Scenario; BMJ = British Medical Journal; TM = tympanic membrane

available guidance, cost and availability in local formularies. In total, around a quarter will routinely prescribe EarCalm® spray (acetic acid). It is possible that its cost (£6–8 per bottle) might be a contributing factor as it is relatively expensive in comparison to other topical preparations as shown in Table 3.

- Otitis externa is the commonest cause of ear discharge in adults
- Otitis externa carries a significant financial burden
- Recommended first line treatment for otitis externa is with topical preparations
- Otitis externa incidence and rate of complications are increasing
- Oral antibiotics in primary care are used commonly for treatment
- Available guidance is inconsistent

Hospital Episode Statistics data confirm a marked increase in otitis externa secondary care referrals. Potential factors may include increased antimicrobial resistance including to neomycin (Otomize®),¹⁸ the observed steady increase of earphone or headphone users in the UK from 12 million in 2013 to 16 million in

2016,¹⁹ and the increased popularity of showering which may allow more water ingress into the external auditory canal.

The frequency of resistance to gentamicin (18 per cent of *Pseudomonas* species) is noteworthy as it is one of the more widely prescribed ototopical agents. Resistance to ciprofloxacin (7 per cent in this study) was higher than in a previous report.²⁰ As otitis externa is so common, guideline adherence is important to achieve good antimicrobial stewardship.²¹ The rapidly expanding use of ciprofloxacin drops (National Health Service Digital prescribing data) is concerning. The use of topical fluoroquinolones is being actively promoted by pharmaceutical companies on the basis of a relatively low ototoxicity compared with aminoglycoside preparations²² and the fact that non-specialists may find it difficult to exclude a tympanic membrane perforation. However, such increased use will increase the risk of evolved resistance to fluoroquinolones. This would deprive clinicians of one of very few oral treatments for necrotising otitis externa,²³ potentially leading in turn to longer in-patient admissions for intravenous therapy.

We found marked variation in guidelines. NICE has a 'clinical knowledge skills document' rather than formal guidelines. This less structured format leaves more room for interpretation. Neither NICE nor the North American guidelines recommend topical antibiotics as first line treatment, while five of six local and regional guidelines more explicitly recommended acetic acid as a first line treatment. Avoidance of topical antibiotics aligns with the UK Commissioning for Quality and Innovation initiative to tackle antimicrobial resistance.²⁴ Prophylactic measures against otitis externa are alluded to only in some of the available guidelines.^{10,12} Even where the use of earplugs or cotton wool smeared with water repellent materials, such as Vaseline® (petroleum jelly)⁸ are referred to, consistency and adequate explanation are lacking.²⁵

Limitations

We sampled selected guidelines and were not in a position to explore the reasons behind their variation, although lack of robust evidence is a likely contributor both to their deficiencies and to lack of adherence by community healthcare professionals. A further limitation is the local nature of the microbiological data, which may not be reflected nation-wide, but could form the basis for similar studies at other institutions.

Conclusion

In this study, we highlighted a number of important issues in otitis externa practice, including the increasing disease burden, complications and cost of topical treatment, the overuse of oral antibiotics in primary care and the promotion of broad-spectrum topical quinolone antibiotics in the community. The proportion of otitis externa isolated with an antibiotic resistant pathogen in this study is greater than has been reported in other studies. There is a need for improvement in evidence-based guidelines and their implementation in otitis externa, especially in terms of the avoidance of oral antibiotics.

Further research is needed to explore the perceptions of stakeholders concerning the misuse of systemic and topical antibiotic preparations, and the optimal methods of prevention.

Acknowledgements. The authors thank Jade Barker, Healthcare Science Associate, Freeman Hospital, Newcastle upon Tyne.

Competing interests. None declared

References

- Rowlands S, Devalia H, Smith C, Hubbard R, Dean A. Otitis externa in UK general practice: a survey using the UK General Practice Research Database. *Br J Gen Pract* 2001;**51**:533–8
- Kaushik V, Malik T, Saeed SR. Interventions for acute otitis externa. *Cochrane Database Syst Rev* 2010;(1):CD004740
- North West EHealth. A new direction in clinical trial design. In: <https://nweh.co.uk> [26 September 2020]
- Prescribing and Medicine Team. Prescriptions Dispensed in the Community - England, 2006 to 2016. In: <https://files.digital.nhs.uk/publication/s/q/pres-disp-com-eng-2006-16-trendtabs.xls> [15 December 2019]
- NHS Digital. Hospital Episode Statistics. In: <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/hospital-episode-statistics> [26 September 2020]
- NHS Improvement. Reference costs 2017/18: highlights, analysis and introduction to the data. In: https://improvement.nhs.uk/documents/1972/1_-_Reference_costs_201718.pdf [25 Mar 2021]
- European Committee on Antimicrobial Susceptibility Testing. Clinical breakpoints and dosing of antibiotics. In: https://www.eucast.org/clinical_breakpoints/ [15 July 2020]
- Ghossaini S. Otitis externa. BMJ Best Practice. In: <https://bestpractice.bmj.com/topics/en-gb/40> [15 December 2019]
- East and North Hertfordshire CCG. Guidelines for the management of infection in primary care within Hertfordshire. In: https://hertsvalleysccg.nhs.uk/application/files/2215/7183/1216/Infection_in_Primary_Care_Guidelines_HMMC_201906.pdf [15 December 2019]
- North Cumbria University Hospital Trust. Management and referral guidelines for ENT in Primary Care. In: <https://documents.in/document/management-and-referral-guidelines-for-ent-in-primary-and-referral-guidelines.html> [15 Dec 2019]
- Medicine Management Team. North Yorkshire antibiotic prescribing guideline for primary care. In: <http://www.valeofyorkccg.nhs.uk/rss/data/uploads/prescribing/north-yorkshire-antibiotic-prescribing-guideline-for-primary-care/ny-antibiotic-guidance-published.pdf> [15 December 2019]
- Centres for Disease Control and Prevention. Estimated burden of acute otitis externa—United States, 2003–2007. *Morb Mortal Wkly Rep* 2011;**60**:605–9
- Bhattacharyya N, Kepnes LJ. Initial impact of the acute otitis externa clinical practice guideline on clinical care. *Otolaryngol Neck Surg* 2011;**145**:414–17
- Collier SA, Hlavsa MC, Piercefield EW, Beach MJ. Antimicrobial and analgesic prescribing patterns for acute otitis externa, 2004–2010. *Otolaryngol Head Neck Surg* 2013;**148**:128–34
- Pabla L, Jindal M, Latif K. The management of otitis externa in UK general practice. *Eur Arch Otorhinolaryngology* 2012;**269**:753–6
- Yelland MJ. The efficacy of oral cotrimoxazole in the treatment of otitis externa in general practice. *Med J Aust* 1993;**158**:697–9
- Simpson KL, Markham A. Ofloxacin otic solution. *Drugs* 1999;**58**:509–31
- Cantrell HF, Lombardy EE, Duncanson FP, Katz E, Barone J. Declining susceptibility to neomycin and polymyxin b of pathogens recovered in otitis externa clinical trials. *South Med J* 2004;**97**:465–71
- Statista. Number of people using headphones/earphones in Great Britain from 2013 to 2018, by product feature (in 1,000s). In: <https://www.statista.com/statistics/308327/headphones-earphones-usage-by-feature-in-the-uk/> [15 December 2019]
- Berenholz L, Katzenell U, Harell M. Evolving resistant pseudomonas to ciprofloxacin in malignant otitis externa. *Laryngoscope* 2002;**112**:1619–22
- NICE. Antimicrobial stewardship: systems and processes for effective antimicrobial medicine use. In: <https://www.nice.org.uk/guidance/ng15/resources/antimicrobial-stewardship-systems-and-processes-for-effective-antimicrobial-medicine-use-pdf-1837273110469> [15 July 2020]
- Ghosh S, Panarese A, Parker AJ, Bull PD. Quinolone ear drops for chronic otitis media. *BMJ* 2000;**321**:126–7
- Chen YA, Chan KC, Chen CK, Chin-Kuo C, Che-Ming W. Differential diagnosis and treatments of necrotizing otitis externa: a report of 19 cases. *Auris Nasus Larynx* 2011;**38**:666–70
- NHS England. Anti-Microbial Resistance (AMR) CQUIN. In: <https://www.england.nhs.uk/nhs-standard-contract/cquin/cquin-16-17/amr-cquin/> [15 July 2020]
- NICE. Clinical Knowledge Summaries. Otitis externa. In: <https://cks.nice.org.uk/otitis-externa> [15 December 2019]
- Public Health England. Summary of antimicrobial prescribing guidance: managing common infections. In: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/994444/Common_Infect_PHE_context_references_and_rationale_May_2021_Bites_and_Eczema_1_.pdf [15 December 2019]
- East London Health and Care Partnership. North East London (NEL) management of infection guidance for primary care. In: http://www.cityandhackneyccg.nhs.uk/Downloads/gp/Formulary/ClinicalGuidelines/NEL_mangementofInfectionguide.pdf [15 December 2019]
- Area Prescribing Committee. North of Tyne/Gateshead guidelines for management of common ENT conditions in primary care. In: <http://www.northoftyneapc.nhs.uk/guidance/> [15 December 2019]
- Northern Lincolnshire and Goole NHS Foundation Trust. North Lincolnshire prescribing guidelines in primary care. In: <https://www.nlg.nhs.uk/content/uploads/2019/11/Ratified-Antimicrobial-Guidance-April-2014-v4.pdf> [15 December 2019]