

On a class of nonlinear anisotropic parabolic problems

M. H. Abdou

Département de Mathématiques et Informatique, Faculté des Sciences
Dhar-Mahraz, Université Sidi Mohamed Ben Abdellah,
B.P. 1796 Fès-Atlas, 30003 Morocco (mohamedhousseine@yahoo.fr)

M. Chrif

Département de Mathématiques et Informatique, Faculté des Sciences
Dhar-Mahraz, Université Sidi Mohamed Ben Abdellah,
B.P. 1796 Fès-Atlas, 30003 Morocco (moussachrif@yahoo.fr)

S. El Manouni

Department of Mathematics, Faculty of Sciences, Al Imam Mohammad
Ibn Saud Islamic University (IMSIU), PO Box 90950, Riyadh 11623,
Saudi Arabia (samanouni@imamu.edu.sa)

H. Hjjaj

Département de Mathématiques et Informatique, Faculté des Sciences
Dhar-Mahraz, Université Sidi Mohamed Ben Abdellah,
B.P. 1796 Fès-Atlas, 30003 Morocco (hjjajhassane@yahoo.fr)

(MS received 16 September 2013; accepted 2 December 2014)

We prove the existence of weak solutions for the strongly nonlinear parabolic problem

$$u_t + Au + g(x, t, u) + \gamma|u|^{p_0-2}u = f$$

in the anisotropic Sobolev space $L^{\mathcal{P}}(0, T; W_0^{1, \mathcal{P}}(\Omega))$, where the data f are assumed to be in the dual, and the nonlinear term $g(x, t, s)$ has growth and sign conditions on s .

Keywords: anisotropic Sobolev space; strongly nonlinear parabolic problems;
weak solution

2010 *Mathematics subject classification:* Primary 35K55; 35K58; 35K59

1. Introduction

This paper is motivated by the study of the parabolic problem associated with the following equations:

$$\begin{aligned} u_t + Au + g(x, t, u) + \gamma|u|^{p_0-2}u &= f && \text{in } Q_T = \Omega \times (0, T), \\ u(x, t) &= 0 && \text{on } S_T = \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x) && \text{in } \Omega, \end{aligned}$$

where $u_0 \in L^2(\Omega)$, $\gamma > 0$ and $f \in L^{\mathcal{P}'}(0, T; W^{-1, \mathcal{P}'}(\Omega))$.

© 2015 The Royal Society of Edinburgh

Here A is an operator of Leray–Lions type acting from $L^p(0, T; W_0^{1,p}(\Omega))$ into its dual $L^{p'}(0, T; W^{-1,p'}(\Omega))$ and defined by

$$Au = - \sum_{i=1}^N D^i a_i(x, t, \nabla u),$$

while $(a_i(x, t, \xi))_{i=1, \dots, N}$ are Carathéodory functions.

In the isotropic case with $p > 2$, Boccardo *et al.* [10] studied the problem

$$\left. \begin{aligned} u'_t - \operatorname{div}(a(x, t, u, \nabla u)) &= \mu && \text{in } Q_T = \Omega \times (0, T), \\ u &= 0 && \text{on } S_T = \partial\Omega \times (0, T), \\ u(x, 0) &= 0 && \text{in } \Omega, \end{aligned} \right\} \quad (1.1)$$

where a is a Carathéodory function satisfying certain growth, coerciveness and monotonicity conditions and μ belongs to $\mathcal{M}(Q_T)$, the space of bounded Borel measure on Q_T . Boccardo *et al.* proved the existence of solutions for problem (1.1) and showed some existence and regularity results in specific cases.

In [8], Boccardo *et al.* considered the following nonlinear parabolic equation:

$$\left. \begin{aligned} u'_t + Au + \alpha_0 |u|^{s-1} u &= f && \text{in } Q_T = \Omega \times (0, T), \\ u &= 0 && \text{on } S_T = \partial\Omega \times (0, T), \\ u(x, 0) &= 0 && \text{in } \Omega, \end{aligned} \right\} \quad (1.2)$$

where Ω is a bounded open subset of \mathbb{R}^N , $T > 0$, $\alpha_0 > 0$ and $s > (p(N+1) - N)/N$. A is the so-called p -Laplacian operator, $A(v) = -\operatorname{div}(|\nabla v|^{p-2} \nabla v)$, with $p > 1 + N/(N+1)$. They established the regularity of solutions for problem (1.2) when the data f are assumed to be merely in $L^1(\Omega)$.

In the case of anisotropic spaces $W^{1,p}(\Omega)$, Bendahmane *et al.* studied in [7] (see also [9]) an approximation result to prove the existence of weak solutions for some strongly nonlinear boundary-value problems of the form

$$Au + g(x, u) = f \quad \text{in } \Omega.$$

In addition, we refer the reader to the interesting papers [2–5], which deal with anisotropic problems governed by a general class of anisotropic operators in both stationary and parabolic cases.

In our case if f belongs only in $L^1(Q_T)$ and $u_0 \in L^1(\Omega)$, it does not make sense to talk about weak solutions. However, it is interesting to study the existence result of entropy solutions or renormalized solutions using the truncation functions.

Finally, in the framework of variable exponent Sobolev spaces, Bendahmane *et al.* [6] studied the following nonlinear parabolic problem:

$$\left. \begin{aligned} u'_t - \operatorname{div}(|\nabla u|^{p(x)-2} \nabla u) &= f && \text{in } Q_T = \Omega \times (0, T), \\ u &= 0 && \text{on } S_T = \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x) && \text{in } \Omega, \end{aligned} \right\} \quad (1.3)$$

where Ω is a bounded open subset of \mathbb{R}^N and $f \in L^1(\Omega)$. They established the existence and uniqueness of renormalized solutions to (1.3). Moreover, they proved

that if $p(\cdot) > 2 - 1/(N + 1)$, then $u \in L^{q(\cdot)}(0, T; W_0^{1,q(\cdot)}(\Omega))$ for all continuous variable exponents $q(\cdot)$ on $\bar{\Omega}$ satisfying

$$1 \leq q(x) < \frac{N(p(x) - 1) + p(x)}{N + 1} \quad \text{for all } x \in \bar{\Omega}.$$

2. Preliminaries

In this section we list briefly some definitions and well-known facts about anisotropic Sobolev spaces and give a technical lemma.

Recall that anisotropic Sobolev spaces were introduced and studied by Nikol'skiĭ [13], Slobodeckii [14] and Troisi [15], and later by Trudinger [16] in the framework of Orlicz spaces. Let Ω be an open bounded domain in \mathbb{R}^N with boundary $\partial\Omega$, and let p_0, p_1, \dots, p_N be $N + 1$ real numbers with $1 < p_i < \infty, i = 0, 1, \dots, N$. We define

$$\mathbf{p} = (p_0, p_1, \dots, p_N), \quad \underline{p} = \min\{p_i, i = 0, 1, 2, \dots, N\}, \quad D^0u = u, \quad D^i u = \frac{\partial u}{\partial x_i},$$

for $i = 1, \dots, N$. With a slight abuse of notation, we introduce the anisotropic Sobolev space,

$$W^{1,\mathbf{p}}(\Omega) = \{u \in L^{p_0}(\Omega) \text{ and } D^i u \in L^{p_i}(\Omega), i = 1, 2, \dots, N\},$$

endowed with the norm

$$\|u\|_{1,\mathbf{p}} = \sum_{i=0}^N \|D^i u\|_{L^{p_i}(\Omega)}. \tag{2.1}$$

We also define $W_0^{1,\mathbf{p}}(\Omega)$ as the closure of $C_0^\infty(\Omega)$ in $W^{1,\mathbf{p}}(\Omega)$ with respect to the norm (2.1). The dual of $W_0^{1,\mathbf{p}}(\Omega)$ is denoted by $W^{-1,\mathbf{p}' }(\Omega)$, where

$$\mathbf{p}' = (p'_0, p'_1, \dots, p'_N), \quad \frac{1}{p'_i} + \frac{1}{p_i} = 1$$

(cf. [7]).

REMARK 2.1. Arguing as in [1], it can easily be seen that $W^{1,\mathbf{p}}(\Omega)$ is a separable Banach space, and reflexive if $1 < p_i < \infty$ for $i = 0, 1, \dots, N$.

We shall use the following Sobolev embeddings later.

LEMMA 2.2. *Let Ω be a bounded open set in \mathbb{R}^N . Then the following embeddings are compact:*

- (i) if $\underline{p} < N$, then $W_0^{1,\mathbf{p}}(\Omega) \hookrightarrow L^q(\Omega)$ for all $q \in [\underline{p}, \underline{p}^*]$, where

$$\frac{1}{\underline{p}^*} = \frac{1}{\underline{p}} - \frac{1}{N};$$

- (ii) if $\underline{p} = N$, then $W_0^{1,\mathbf{p}}(\Omega) \hookrightarrow L^q(\Omega)$ for all $q \in [\underline{p}, +\infty[$;
- (iii) if $\underline{p} > N$, then $W_0^{1,\mathbf{p}}(\Omega) \hookrightarrow L^\infty(\Omega) \cap C^0(\bar{\Omega})$.

The proof of this lemma follows from the fact that the embedding $W_0^{1,\mathbf{p}}(\Omega) \hookrightarrow W_0^{1,\mathbf{p}'}(\Omega)$ is continuous, and in view of the classical embedding theorems of the Sobolev spaces.

DEFINITION 2.3. We denote the dual of the Sobolev space $W_0^{1,\mathbf{p}}(\Omega)$ by $W^{-1,\mathbf{p}'}(\Omega)$, and for each $F \in W^{-1,\mathbf{p}'}(\Omega)$ there exists $f_i \in L^{p'_i}(\Omega)$ for $i = 0, \dots, N$, such that

$$F = f_0 - \sum_{i=1}^N D^i f_i.$$

Moreover, for all $u \in W_0^{1,\mathbf{p}}(\Omega)$ we have

$$\langle F, u \rangle = \sum_{i=0}^N \int_{\Omega} f_i D^i u \, dx.$$

Furthermore, we define a norm on the dual space by

$$\|F\|_{-1,\mathbf{p}'} = \sum_{i=0}^N \|f_i\|_{p'_i}.$$

2.1. Parabolic spaces

Let $Q_T = \Omega \times (0, T)$ with $0 < T < \infty$. We introduce the parabolic space $L^{\mathbf{p}}(0, T; W^{1,\mathbf{p}}(\Omega))$ by

$$L^{\mathbf{p}}(0, T; W^{1,\mathbf{p}(x)}(\Omega)) = \left\{ u \text{ is measurable} \mid \sum_{i=0}^N \int_0^T \|D^i u\|_{p_i}^{p_i} \, dt < \infty \right\}, \quad (2.2)$$

equipped with the norm

$$\|u\|_{L^{\mathbf{p}}(0,T;W^{1,\mathbf{p}}(\Omega))} := \sum_{i=0}^N \|D^i u\|_{L^{p_i}(Q_T)}.$$

We also introduce the functional space $L^{\mathbf{p}}(0, T; W_0^{1,\mathbf{p}}(\Omega))$ by

$$L^{\mathbf{p}}(0, T; W_0^{1,\mathbf{p}}(\Omega)) = \{u \in L^{\mathbf{p}}(0, T; W^{1,\mathbf{p}}(\Omega)) \mid u = 0 \text{ on } \partial\Omega \times [0, T]\}. \quad (2.3)$$

The spaces $L^{\mathbf{p}}(0, T; W^{1,\mathbf{p}}(\Omega))$ and $L^{\mathbf{p}}(0, T; W_0^{1,\mathbf{p}}(\Omega))$ are separable and reflexive Banach spaces.

The dual space of $L^{\mathbf{p}}(0, T; W_0^{1,\mathbf{p}}(\Omega))$ is defined as follows:

$$L^{\mathbf{p}'}(0, T; W^{-1,\mathbf{p}'}(\Omega)) = \left\{ F = f_0 - \sum_{i=1}^N D^i f_i \text{ with } f_0 \in L^{p'_0}(Q_T), f_i \in L^{p'_i}(Q_T) \right\}, \quad (2.4)$$

and normed by

$$\|F\|_{L^{\mathbf{p}'}(0,T;W^{-1,\mathbf{p}'}(\Omega))} = \sum_{i=0}^N \|f_i\|_{L^{p'_i}(Q_T)}.$$

Finally, the duality of the spaces

$$L^p(0, T; W_0^{1,p}(\Omega)) \quad \text{and} \quad L^{p'}(0, T; W^{-1,p'}(\Omega))$$

is given by

$$\langle F, v \rangle = \sum_{i=0}^N \int_{Q_T} f_i D^i v(x) \, dx \quad \text{for any } v \in L^p(0, T; W_0^{1,p}(\Omega)).$$

3. Essential assumptions

Let $Q_T = \Omega \times (0, T)$ with $0 < T < \infty$ and $2N/(N + 2) < p_i < \infty$. We consider a Leray–Lions operator A acting from $L^p(0, T; W_0^{1,p}(\Omega))$ into its dual $L^{p'}(0, T; W^{-1,p'}(\Omega))$ defined by

$$Au = - \sum_{i=1}^N D^i a_i(x, t, \nabla u).$$

The functions $(a_i(x, t, \xi))_{i=1, \dots, N}$ are Carathéodory functions (measurable with respect to (x, t) in Q_T for every ξ in \mathbb{R}^N and continuous with respect to ξ in \mathbb{R}^N for almost every (x, t) in Q_T) satisfying

$$|a_i(x, t, \xi)| \leq \beta(K_i(x, t) + |\xi_i|^{p_i-1}) \quad \text{for } i = 1, \dots, N, \tag{3.1}$$

$$a_i(x, t, \xi)\xi_i \geq \alpha|\xi_i|^{p_i} \quad \text{for } i = 1, \dots, N, \tag{3.2}$$

and for all $\xi = (\xi_1, \dots, \xi_N)$ and $\xi' = (\xi'_1, \dots, \xi'_N)$ we have

$$(a_i(x, t, \xi) - a_i(x, t, \xi'))(\xi_i - \xi'_i) > 0 \quad \text{for } \xi_i \neq \xi'_i, \tag{3.3}$$

for almost every $(x, t) \in Q_T$ and all $\xi \in \mathbb{R}^N$, where $K_i(x, t)$ is a non-negative function lying in $L^{p_i}(Q_T)$ and $\alpha, \beta > 0$.

The nonlinear operator g is a Carathéodory function that satisfies

$$|g(x, t, s)| \leq |s|^{p_0-1} + b(x, t), \tag{3.4}$$

$$g(x, t, s)s \geq 0, \tag{3.5}$$

for almost every $(x, t) \in Q_T$ and any $s \in \mathbb{R}$, with the function $b: \Omega \times (0, T) \rightarrow \mathbb{R}^+$ such that $b \in L^{p'_0}(Q_T)$.

3.1. Some technical lemmas

In this subsection, we state and prove the following technical lemmas that will be needed later.

LEMMA 3.1 (Hewitt and Stromberg [11, theorem 13.47]). *Let $(u_n)_n$ be a sequence in $L^1(\Omega)$ and let $u \in L^1(\Omega)$ such that $u_n \rightarrow u$ almost everywhere (a.e.) in Ω , $u_n, u \geq 0$ a.e. and*

$$\int_{\Omega} u_n \, dx \rightarrow \int_{\Omega} u \, dx.$$

Then $u_n \rightarrow u$ in $L^1(\Omega)$.

LEMMA 3.2. Let $g \in L^r(Q_T)$ and $g_n \in L^r(Q_T)$ with $\|g_n\|_{L^r(Q_T)} \leq C$ for $1 < r < \infty$. If $g_n(x, t) \rightarrow g(x, t)$ a.e. on Q_T , then $g_n \rightarrow g$ in $L^r(Q_T)$.

LEMMA 3.3. Assume that (3.1)–(3.3) hold, and let $(u_n)_n$ be a sequence in $L^p(0, T; W_0^{1,p}(\Omega))$ such that $u_n \rightarrow u$ in $L^p(0, T; W_0^{1,p}(\Omega))$ and

$$\int_0^T \langle Au_n - Au, u_n - u \rangle dt + \int_{Q_T} (|u_n|^{p_0-2}u_n - |u|^{p_0-2}u)(u_n - u) dx dt \rightarrow 0. \quad (3.6)$$

Then $u_n \rightarrow u$ in $L^p(0, T; W_0^{1,p}(Q_T))$ for a subsequence.

Proof. Let

$$D_n(x, t) = \sum_{i=1}^N (a_i(x, t, \nabla u_n) - a_i(x, t, \nabla u))(D^i u_n - D^i u) + (|u_n|^{p_0-2}u_n - |u|^{p_0-2}u)(u_n - u).$$

By (3.3), D_n is a positive function, and, in view of (3.6), $D_n \rightarrow 0$ in $L^1(Q_T)$ as $n \rightarrow \infty$.

We have $u_n \rightarrow u$ in $L^p(0, T; W_0^{1,p}(\Omega))$. Then $u_n \rightarrow u$ in $L^1(Q_T)$, and since $D_n \rightarrow 0$ a.e. in Q_T there exists a subset B in Q_T with measure zero such that for all $(x, t) \in Q_T \setminus B$ we have

$$|u(x, t)| < \infty, \quad |\nabla u(x, t)| < \infty, \quad K_i(x, t) < \infty, \quad u_n \rightarrow u \quad \text{and} \quad D_n \rightarrow 0.$$

We also have

$$\begin{aligned} D_n(x, t) &= \sum_{i=1}^N (a_i(x, t, \nabla u_n) - a_i(x, t, \nabla u))(D^i u_n - D^i u) \\ &\quad + (|u_n|^{p_0-2}u_n - |u|^{p_0-2}u)(u_n - u) \\ &= \sum_{i=1}^N (a_i(x, t, \nabla u_n)D^i u_n + a_i(x, t, \nabla u)D^i u \\ &\quad - a_i(x, t, \nabla u)D^i u_n - a_i(x, t, \nabla u_n)D^i u) \\ &\quad + |u_n|^{p_0} + |u|^{p_0} - |u_n|^{p_0-2}u_n u - |u|^{p_0-2}u u_n \\ &\geq \underline{\alpha} \sum_{i=0}^N |D^i u_n|^{p_i} + \underline{\alpha} \sum_{i=0}^N |D^i u|^{p_i} - \beta \sum_{i=1}^N (K_i(x, t) + |D^i u|^{p_i-1})|D^i u_n| \\ &\quad - \beta \sum_{i=1}^N (K_i(x, t) + |D^i u_n|^{p_i-1})|D^i u| - |u_n|^{p_0-1}|u| - |u|^{p_0-1}|u_n| \\ &\geq \underline{\alpha} \sum_{i=0}^N |D^i u_n|^{p_i} - C_{x,t} \sum_{i=0}^N (1 + |D^i u_n|^{p_i-1} + |D^i u_n|), \end{aligned}$$

with $\underline{\alpha} = \min(\alpha, 1)$ and $C_{x,t}$ depending on x and t , without dependence on n . Hence, we obtain

$$D_n(x, t) \geq \sum_{i=0}^N |D^i u_n|^{p_i} \left(\underline{\alpha} - \frac{C_{x,t}}{|D^i u_n|^{p_i}} - \frac{C_{x,t}}{|D^i u_n|} - \frac{C_{x,t}}{|D^i u_n|^{p_i-1}} \right).$$

By standard arguments, $(D^i u_n)_n$ is bounded a.e. in Q_T for $i = 0, 1, \dots, N$.¹

Letting ξ_i^* be an accumulation point of $(D^i u_n)_n$ for $i = 1, \dots, N$, we have $|\xi_i^*| < \infty$, and by the continuity of $a(x, t, \cdot)$ we obtain

$$[a_i(x, t, \xi^*) - a_i(x, t, \nabla u)](\xi_i^* - D^i u) = 0 \quad \text{for } i = 1, \dots, N.$$

By (3.3), we have $\xi^* = \nabla u$. The uniqueness of the accumulation point implies that $\nabla u_n \rightarrow \nabla u$ a.e in Ω .

Now, since $(a_i(x, t, \nabla u_n))_n$ is bounded in $L^{p'_i}(Q_T)$ and

$$a_i(x, t, \nabla u_n) \rightarrow a_i(x, t, \nabla u) \quad \text{a.e in } Q_T$$

by lemma 3.2, we can establish that

$$a_i(x, t, \nabla u_n) \rightharpoonup a_i(x, t, \nabla u) \text{ in } L^{p'_i}(Q_T) \quad \text{for } i = 1, \dots, N.$$

Using (3.6) and in view of lemma 3.1, we deduce that

$$|u_n|^{p_0} \rightarrow |u|^{p_0} \quad \text{in } L^1(Q_T). \tag{3.7}$$

Then $u_n \rightarrow u$ in $L^{p_0}(Q_T)$, and

$$a_i(x, t, \nabla u_n) D^i u_n \rightarrow a_i(x, t, \nabla u) D^i u \quad \text{in } L^1(Q_T). \tag{3.8}$$

According to the condition (3.2), we have

$$\alpha |D^i u_n|^{p_i} \leq a_i(x, t, \nabla u_n) D^i u_n \quad \text{for } i = 1, \dots, N.$$

Letting

$$y_n^i = \frac{1}{\alpha} a_i(x, t, \nabla u_n) D^i u_n \quad \text{and} \quad y^i = \frac{1}{\alpha} a_i(x, t, \nabla u) D^i u,$$

in view of Fatou's lemma we get

$$\int_{Q_T} 2y^i \, dx \, dt \leq \liminf_{n \rightarrow \infty} \int_{Q_T} \left(y_n^i + y^i - \frac{1}{2^{p_i-1}} |D^i u_n - D^i u|^{p_i} \right) \, dx \, dt.$$

Hence,

$$0 \leq - \limsup_{n \rightarrow \infty} \int_{Q_T} |D^i u_n - D^i u|^{p_i} \, dx \, dt.$$

¹ Indeed, if $|D^i u_n| \rightarrow \infty$ in a measurable subset $E \subset Q_T$, then

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_{\Omega} D_n(x, t) \, dx \, dt \\ & \geq \lim_{n \rightarrow \infty} \int_E |D^i u_n|^{p_i} \left(\underline{\alpha} - \frac{C_{x,t}}{|D^i u_n|^{p_i}} - \frac{C_{x,t}}{|D^i u_n|} - \frac{C_{x,t}}{|D^i u_n|^{p_i-1}} \right) \, dx \, dt \\ & = \infty, \end{aligned}$$

which is absurd since $D_n \rightarrow 0$ in $L^1(Q_T)$.

Thus, since

$$0 \leq \liminf_{n \rightarrow \infty} \int_{Q_T} |D^i u_n - D^i u|^{p_i} dx dt \leq \limsup_{n \rightarrow \infty} \int_{Q_T} |D^i u_n - D^i u|^{p_i} dx dt \leq 0,$$

it follows that

$$\int_{Q_T} |D^i u_n - D^i u|^{p_i} dx dt \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Then, we obtain

$$D^i u_n \rightarrow D^i u \quad \text{in } L^{p_i}(Q_T) \quad \text{for } i = 1, \dots, N.$$

Consequently, due to (3.7), we deduce that

$$u_n \rightarrow u \quad \text{in } L^p(0, T; W_0^{1,p}(Q_T)).$$

□

4. Main result

Now, we state the main result in our paper.

THEOREM 4.1. *Assume that (3.1)–(3.5) hold and let $f \in L^{p'}(0, T; W^{-1,p'}(\Omega))$. Then there exists at least one solution $u \in L^p(0, T; W_0^{1,p}(\Omega))$ of the problem*

$$\left. \begin{aligned} u_t + Au + g(x, t, u) + \gamma|u|^{p_0-2}u &= f && \text{in } Q_T = \Omega \times (0, T), \\ u(x, t) &= 0 && \text{on } S_T = \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x) && \text{in } \Omega, \end{aligned} \right\} \quad (4.1)$$

where $u_0 \in L^2(\Omega)$.

Proof of theorem 4.1.

STEP 1 (approximate problem). Consider the following problem:

$$\left. \begin{aligned} \frac{\partial u_n}{\partial t} + Au_n + g_n(x, t, u_n) + \gamma|u|^{p_0-2}u &= f(x, t) && \text{in } \Omega \times (0, T), \\ u_k(x, t) &= 0 && \text{on } \partial\Omega \times (0, T), \\ u_k(x, 0) &= u_0 && \text{in } \Omega, \end{aligned} \right\} \quad (4.2)$$

with

$$g_n(x, t, s) = \frac{g(x, t, s)}{1 + |g(x, t, s)|/n}.$$

Note that

$$g_n(x, t, s) \geq 0, \quad |g_n(x, t, s)| \leq |g(x, t, s)| \quad \text{and} \quad |g_n(x, t, s)| \leq n \quad \forall n \in \mathbb{N}^*.$$

We define the operator $G_n : L^p(0, T; W_0^{1,p}(\Omega)) \rightarrow L^{p'}(0, T; W^{-1,p'}(\Omega))$ by

$$\int_0^T \langle G_n u, v \rangle dt = \int_{Q_T} g_n(x, t, u)v dx dt + \gamma \int_{Q_T} |u|^{p_0-2}uv dx dt$$

$$\forall v \in L^p(0, T; W_0^{1,p}(\Omega)).$$

By the Hölder inequality, we have

$$\begin{aligned} \left| \int_0^T \langle G_n u, v \rangle dt \right| &\leq \int_{Q_T} |g_n(x, t, u)| |v| dx dt + \gamma \int_{Q_T} |u|^{p_0-1} |v| dx dt \\ &\leq \int_{Q_T} ((\gamma + 1)|u|^{p_0-1} + b(x, t)) |v| dx dt \\ &\leq ((\gamma + 1)\|u\|_{L^{p_0}(Q_T)}^{p_0-1} + \|b(x, t)\|_{L^{p'_0}(Q_T)}) \|v\|_{L^{p_0}(Q_T)} \end{aligned} \tag{4.3}$$

and

$$\begin{aligned} \left| \int_0^T \langle Au, v \rangle dt \right| &\leq \sum_{i=1}^N \int_{Q_T} |a_i(x, t, \nabla u)| |D^i v| dx dt \\ &\leq \sum_{i=1}^N \int_{Q_T} \beta(K_i(x, t) + |D^i u|^{p_i-1}) |D^i v| dx dt \\ &\leq \beta \sum_{i=1}^N (\|K_i(x, t)\|_{L^{p'_i}(Q_T)} + \|D^i u\|_{L^{p_i}(Q_T)}^{p_i-1}) \|D^i v\|_{L^{p_i}(Q_T)} \end{aligned} \tag{4.4}$$

for all $u, v \in L^p(0, T; W_0^{1,p}(\Omega))$.

LEMMA 4.2. *The operator*

$$B_n = A + G_n$$

is pseudo-monotone from $L^p(0, T; W_0^{1,p}(\Omega))$ into $L^{p'}(0, T; W^{-1,p'}(\Omega))$. Moreover, B_n is coercive in the following sense:

$$\left(\frac{1}{\|v\|_{L^p(0,T;W_0^{1,p}(\Omega))}} \right) \int_0^T \langle B_n v, v \rangle dt \rightarrow +\infty \quad \text{as } \|v\|_{L^p(0,T;W_0^{1,p}(\Omega))} \rightarrow +\infty,$$

for $v \in L^p(0, T; W_0^{1,p}(\Omega))$.

Proof of lemma 4.2. In view of inequalities (4.3) and (4.4), the operator B_n is bounded. For the coercivity, due to (3.2) and (3.5), we have

$$\begin{aligned} \int_0^T \langle B_n u, u \rangle dt &= \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u) D^i u dx dt \\ &\quad + \int_{Q_T} g(x, t, u) u dx dt + \gamma \int_{Q_T} |u|^{p_0} dx dt \\ &\geq \alpha \sum_{i=1}^N \int_{Q_T} |D^i u|^{p_i} dx dt + \gamma \int_{Q_T} |u|^{p_0} dx dt \\ &\geq \alpha \sum_{i=1}^N (\|D^i u\|_{L^{p_i}(Q_T)}^{p_i} - 1) + \gamma (\|u\|_{L^{p_0}(Q_T)}^{p_0} - 1) \\ &\geq \alpha' \|v\|_{L^p(0,T;W_0^{1,p}(\Omega))}^p - \alpha N - \gamma \end{aligned}$$

for all $u \in L^p(0, T; W_0^{1,p}(\Omega))$. It follows that

$$\left(\frac{1}{\|u\|_{L^p(0,T;W_0^{1,p}(\Omega))}} \right) \int_0^T \langle B_n u, u \rangle dt \rightarrow +\infty \quad \text{as } \|u\|_{L^p(0,T;W_0^{1,p}(\Omega))} \rightarrow +\infty.$$

It remains to show that B_n is pseudo-monotone. Indeed, let $(u_k)_k$ be a sequence in $L^p(0, T; W_0^{1,p}(\Omega))$ such that

$$\left. \begin{aligned} &u_k \rightharpoonup u \quad \text{in } L^p(0, T; W_0^{1,p}(\Omega)), \\ &B_n u_k \rightharpoonup \chi \quad \text{in } L^{p'}(0, T; W^{-1,p'}(\Omega)), \\ &\limsup_{k \rightarrow \infty} \int_0^T \langle B_n u_k, u_k \rangle dt \leq \int_0^T \langle \chi, u \rangle dt. \end{aligned} \right\} \quad (4.5)$$

We shall prove that

$$\chi = B_n u \quad \text{and} \quad \int_0^T \langle B_n u_k, u_k \rangle dt \rightarrow \int_0^T \langle \chi, u \rangle dt \quad \text{as } k \rightarrow +\infty.$$

First, since the embedding $L^p(0, T; W_0^{1,p}(\Omega)) \hookrightarrow L^1(Q_T)$ is compact, there exists a subsequence, still denoted (u_k) , such that $u_k \rightarrow u$ in $L^1(Q_T)$.

Second, we have that $(u_k)_k$ is a bounded sequence in $L^p(0, T; W_0^{1,p}(\Omega))$. Then, by (3.1), $(a_i(x, t, \nabla u_k))_k$ is bounded in $L^{p_i}(Q_T)$. Therefore, there exists a function $\varphi_i \in L^{p_i}(Q_T)$ such that

$$a_i(x, t, \nabla u_k) \rightharpoonup \varphi_i \quad \text{in } L^{p_i}(Q_T) \quad \text{for } i = 1, \dots, N. \quad (4.6)$$

We have $g_n(x, t, u_k) \rightarrow g_n(x, t, u)$ a.e. in Q_T and $|g_n(x, t, u_k)| \leq n \in L^{p_0}(Q_T)$. Then, by the Lebesgue dominated convergence theorem, we deduce that

$$g_n(x, t, u_k) \rightarrow g_n(x, t, u) \quad \text{in } L^{p_0}(Q_T). \quad (4.7)$$

We also have

$$|u_k|^{p_0-2} u_k \rightharpoonup |u|^{p_0-2} u \quad \text{in } L^{p_0}(Q_T). \quad (4.8)$$

Thus, for all $v \in L^p(0, T; W_0^{1,p}(\Omega))$, we get

$$\begin{aligned} \langle \chi, v \rangle &= \lim_{k \rightarrow \infty} \langle B_n u_k, v \rangle \\ &= \lim_{k \rightarrow \infty} \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i v \, dx \, dt \\ &\quad + \lim_{k \rightarrow \infty} \int_{Q_T} g_n(x, t, u_k) v \, dx \, dt + \lim_{k \rightarrow \infty} \gamma \int_{Q_T} |u_k|^{p_0-2} u_k v \, dx \, dt \\ &= \sum_{i=1}^N \int_{Q_T} \varphi_i D^i v \, dx \, dt + \int_{Q_T} g_n(x, t, u) v \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0-2} u v \, dx \, dt. \end{aligned} \quad (4.9)$$

By using (4.5) and (4.9), we obtain

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \langle B_n(u_k), u_k \rangle \\ &= \limsup_{k \rightarrow \infty} \left\{ \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i u_k \, dx \, dt \right. \\ & \quad \left. + \int_{Q_T} g_n(x, t, u_k) u_k \, dx \, dt + \gamma \int_{Q_T} |u_k|^{p_0} \, dx \, dt \right\} \\ & \leq \sum_{i=1}^N \int_{Q_T} \varphi_i D^i u \, dx \, dt + \int_{Q_T} g_n(x, t, u) u \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0} \, dx \, dt. \end{aligned} \tag{4.10}$$

By (4.7), we have

$$\int_{Q_T} g_n(x, t, u_k) u_k \, dx \, dt \rightarrow \int_{Q_T} g_n(x, t, u) u \, dx \, dt. \tag{4.11}$$

Therefore,

$$\begin{aligned} & \limsup_{k \rightarrow \infty} \left\{ \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i u_k \, dx \, dt + \int_{Q_T} |u_k|^{p_0} \, dx \, dt \right\} \\ & \leq \sum_{i=1}^N \int_{Q_T} \varphi_i D^i u \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0} \, dx \, dt. \end{aligned} \tag{4.12}$$

On the other hand, by using (3.3) we get

$$\begin{aligned} & \sum_{i=1}^N \int_{Q_T} (a_i(x, t, \nabla u_k) - a_i(x, t, \nabla u)) (D^i u_k - D^i u) \, dx \, dt \\ & \quad + \gamma \int_{Q_T} (|u_k|^{p_0-2} u_k - |u|^{p_0-2} u) (u_k - u) \, dx \, dt \geq 0. \end{aligned} \tag{4.13}$$

Then, we obtain

$$\begin{aligned} & \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i u_k \, dx \, dt + \gamma \int_{Q_T} |u_k|^{p_0} \, dx \, dt \\ & \geq - \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u) D^i u \, dx \, dt - \gamma \int_{Q_T} |u|^{p_0} \, dx \, dt \\ & \quad + \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u) D^i u_k \, dx \, dt + \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i u \, dx \, dt \\ & \quad + \gamma \int_{Q_T} |u_k|^{p_0-2} u_k u \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0-2} u u_k \, dx \, dt. \end{aligned}$$

Thus, due to (4.6) and (4.8), we have

$$\begin{aligned} \liminf_{k \rightarrow \infty} \left\{ \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i u_k \, dx \, dt + \gamma \int_{Q_T} |u_k|^{p_0} \, dx \, dt \right\} \\ \geq \sum_{i=1}^N \int_{Q_T} \varphi_i D^i u \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0} \, dx \, dt. \end{aligned}$$

This implies, by (4.12), that

$$\begin{aligned} \lim_{k \rightarrow \infty} \left\{ \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_k) D^i u_k \, dx \, dt + \gamma \int_{Q_T} |u_k|^{p_0} \, dx \, dt \right\} \\ = \sum_{i=1}^N \int_{Q_T} \varphi_i D^i u \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0} \, dx \, dt. \quad (4.14) \end{aligned}$$

By combining (4.9), (4.11) and (4.14), we deduce that

$$\int_0^T \langle B_n u_k, u_k \rangle \, dt \rightarrow \int_0^T \langle \chi, u \rangle \, dt \quad \text{as } k \rightarrow +\infty.$$

Now, by (4.14) we can obtain

$$\begin{aligned} \sum_{i=1}^N \int_{Q_T} [a_i(x, t, \nabla u_k) - a_i(x, t, \nabla u)] (D^i u_k - D^i u) \, dx \, dt \\ + \int_{Q_T} (|u_k|^{p_0-2} u_k - |u|^{p_0-2} u) (u_k - u) \, dx \, dt \rightarrow 0 \quad \text{as } k \rightarrow +\infty. \end{aligned}$$

By lemma 3.3, we get

$$u_k \rightarrow u \quad \text{in } L^{\mathcal{P}}(0, T; W_0^{1, \mathcal{P}}(Q_T)).$$

Then $D^i u_n \rightarrow D^i u$ a.e. in Q_T . Thus, it follows that $a_i(x, t, \nabla u_n) \rightarrow a_i(x, t, \nabla u)$ a.e. in Q_T . Hence, in view of (3.1) and lemma 3.2, we have

$$a_i(x, t, \nabla u_n) \rightharpoonup a_i(x, t, \nabla u) \quad \text{in } L^{p'_i}(Q_T) \quad \text{for } i = 1, \dots, N.$$

Therefore, using (4.7) and (4.8), we deduce that $\chi = B_n u$, and the proof of lemma 4.2 is completed. \square

As a consequence of lemma 4.2, we deduce that there exists at least one weak solution $u_n \in L^{\mathcal{P}}(0, T; W_0^{1, \mathcal{P}}(\Omega))$ of problem (4.2) (see [12]).

STEP 2 (*a priori* estimates). Taking u_n as a test function in (4.2), we obtain

$$\begin{aligned} \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n \right\rangle \, dt + \int_0^T \langle A u_n, u_n \rangle \, dt \\ + \int_{Q_T} g_n(x, t, u_n) u_n \, dx \, dt + \gamma \int_{Q_T} |u_n|^{p_0} \, dx \, dt = \int_0^T \langle f, u_n \rangle \, dt. \quad (4.15) \end{aligned}$$

By (3.2), we have

$$\begin{aligned} \int_0^T \langle Au_n, u_n \rangle dt &= \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_n) D^i u_n dx dt \\ &\geq \alpha \sum_{i=1}^N \int_{Q_T} |D^i u_n|^{p_i} dx dt. \end{aligned} \tag{4.16}$$

We also have

$$\begin{aligned} \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n \right\rangle dt &= \frac{1}{2} \int_{\Omega} \int_0^T \frac{\partial (u_n(t))^2}{\partial t} dt dx \\ &= \frac{1}{2} \int_{\Omega} |u_n(T)|^2 dx - \frac{1}{2} \int_{\Omega} |u(0)|^2 dx, \end{aligned} \tag{4.17}$$

As regards the term on the right-hand side of (4.15), since $f \in L^{p'}(0, T; W_0^{-1, p'}(\Omega))$ there exist $f_0 \in L^{p'_0}(Q_T)$ and $f_i \in L^{p'_i}(Q_T)$ for $i = 1, \dots, N$ such that

$$f = f_0 - \sum_{i=1}^N D^i f_i.$$

Then by using Young's inequality, we get

$$\begin{aligned} \int_0^T \langle f, u_n \rangle dt &= \sum_{i=0}^N \int_{Q_T} f_i D^i u_n dx dt \\ &\leq c_1 \sum_{i=0}^N \int_{Q_T} |f_i|^{p'_i} dx dt + \frac{1}{2} \alpha \sum_{i=1}^N \int_{Q_T} |D^i u_n|^{p_i} dx dt \\ &\quad + \frac{1}{2} \gamma \int_{Q_T} |u_n|^{p_0} dx dt. \end{aligned} \tag{4.18}$$

Combining (4.15)–(4.18), we obtain

$$\begin{aligned} \frac{1}{2} \int_{\Omega} |u_n(T)|^2 dx + \frac{1}{2} \alpha \sum_{i=1}^N \int_{Q_T} |D^i u_n|^{p_i} dx dt \\ + \int_{Q_T} g_n(x, t, u_n) u_n dx dt + \frac{1}{2} \gamma \int_{Q_T} |u_n|^{p_0} dx dt \leq c_2. \end{aligned} \tag{4.19}$$

Then there exists a constant c_3 that does not depend on n , such that

$$\sum_{i=0}^N \|D^i u_n\|_{L^{p_i}(Q_T)} \leq c_3. \tag{4.20}$$

Therefore, $(u_n)_n$ is bounded in $L^p(0, T; W_0^{1, p}(\Omega))$. Hence, there exists a subsequence, still denoted $(u_n)_n$, such that

$$\left. \begin{aligned} u_n &\rightharpoonup u \quad \text{in } L^p(0, T; W_0^{1, p}(\Omega)), \\ u_n &\rightarrow u \quad \text{in } L^1(Q_T) \text{ and a.e. in } Q_T. \end{aligned} \right\} \tag{4.21}$$

We deduce that

$$|u_n(T)|^2 \rightarrow |u(T)|^2 \quad \text{a.e. in } \Omega \quad \text{and} \quad g_n(x, t, u_n)u_n \rightarrow g(x, t, u)u \quad \text{a.e. in } Q_T.$$

Then, due to (4.19) and Fatou's lemma, we get

$$\int_{\Omega} |u(T)|^2 dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |u_n(T)|^2 dx \quad (4.22)$$

and

$$\int_{Q_T} g(x, t, u)u dx dt \leq \liminf_{n \rightarrow \infty} \int_{Q_T} g_n(x, t, u_n)u_n dx dt. \quad (4.23)$$

STEP 3 (weak convergence of $(u_n)_t$ in $L^{p'}(0, T; W^{-1, p'}(\Omega))$). Taking $v \in L^p(0, T; W_0^{1, p}(\Omega))$ as a test function in (4.2), we get

$$\begin{aligned} \int_0^T \left\langle \frac{\partial u_n}{\partial t}, v \right\rangle dt + \int_0^T \langle Au_n, v \rangle dt + \int_{Q_T} g_n(x, t, u_n)v dx dt \\ + \gamma \int_{Q_T} |u_n|^{p_0-2}u_n v dx dt = \int_0^T \langle f, v \rangle dt. \end{aligned}$$

Then,

$$\begin{aligned} \left| \int_0^T \left\langle \frac{\partial u_n}{\partial t}, v \right\rangle dt \right| \leq \left| \int_0^T \langle Au_n, v \rangle dt \right| + \int_{Q_T} |g_n(x, t, u_n)||v| dx dt \\ + \gamma \int_{Q_T} |u_n|^{p_0-1}|v| dx dt + \left| \int_0^T \langle f, v \rangle dt \right|. \quad (4.24) \end{aligned}$$

For the first term on the right-hand side of (4.24), using Hölder's inequality and (4.20), we get

$$\begin{aligned} \left| \int_0^T \langle Au_n, v \rangle dt \right| \\ \leq \sum_{i=1}^N \int_{Q_T} |a_i(x, t, \nabla u_n)| |D^i v| dx dt \\ \leq \sum_{i=1}^N \int_{Q_T} \beta(K_i(x, t) + |D^i u_n|^{p_i-1}) |D^i v| dx dt \\ \leq \beta \sum_{i=1}^N (\|K_i(x, t)\|_{L^{p'_i}(Q_T)} + \|D^i u_n\|_{L^{p_i}(Q_T)}^{p_i-1}) \|D^i v\|_{L^{p_i}(Q_T)} \\ \leq \beta \left(\sum_{i=1}^N \|K_i(x, t)\|_{L^{p'_i}(Q_T)} + \sum_{i=1}^N \|D^i u_n\|_{L^{p_i}(Q_T)}^{p_i-1} \right) \sum_{i=1}^N \|D^i v\|_{L^{p_i}(Q_T)} \\ \leq c_4 \|v\|_{L^p(0, T; W_0^{1, p}(\Omega))}. \quad (4.25) \end{aligned}$$

Regarding the second and third terms on the left-hand side of (4.24), due to (3.4) we have

$$\begin{aligned} & \int_{Q_T} |g(x, t, u_k)| |v| \, dx \, dt + \gamma \int_{Q_T} |u_n|^{p_0-1} |v| \, dx \, dt \\ & \leq (1 + \gamma) \int_{Q_T} |u_k|^{p_0-1} |v| \, dx \, dt + \int_{Q_T} b(x, t) |v| \, dx \, dt \\ & \leq ((1 + \gamma) \| |u_k|^{p_0-1} \|_{L^{p'_0}(Q_T)} + \| b(x, t) \|_{L^{p'_0}(Q_T)}) \| v \|_{L^{p_0}(Q_T)} \\ & \leq c_5 \| v \|_{L^{\mathbf{p}}(0, T; W_0^{1, \mathbf{p}}(\Omega))}. \end{aligned} \tag{4.26}$$

Concerning the last term on the right-hand side of (4.24), we have

$$\begin{aligned} \left| \int_0^T \langle f, v \rangle \, dt \right| & \leq \sum_{i=0}^N \int_{Q_T} |f_i| |D^i v| \, dx \, dt \leq \sum_{i=0}^N \| f_i \|_{L^{p'_i}(Q_T)} \| D^i v \|_{L^{p_i}(Q_T)} \\ & \leq \| f \|_{L^{\mathbf{p}'}(0, T; W^{1, \mathbf{p}'}(\Omega))} \| v \|_{L^{\mathbf{p}}(0, T; W_0^{1, \mathbf{p}}(\Omega))}. \end{aligned} \tag{4.27}$$

Consequently, by combining (4.24)–(4.27), we deduce that

$$\left| \int_0^T \left\langle \frac{\partial u_n}{\partial t}, v \right\rangle \, dt \right| \leq c_6 \| v \|_{L^{\mathbf{p}}(0, T; W_0^{1, \mathbf{p}}(\Omega))}, \tag{4.28}$$

where c_6 is a constant that does not depend on n . Then $\partial u_n / \partial t$ is bounded in $L^{\mathbf{p}'}(0, T; W^{-1, \mathbf{p}'}(\Omega))$ and

$$\frac{\partial u_k}{\partial t} \rightharpoonup \frac{\partial u}{\partial t} \quad \text{in } L^{\mathbf{p}'}(0, T; W^{-1, \mathbf{p}'}(\Omega)).$$

STEP 4 (convergence of the gradient). By taking $u_n - u$ as a test function in (4.2), we obtain

$$\begin{aligned} & \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n - u \right\rangle \, dt + \int_0^T \langle Au_n, u_n - u \rangle \, dt \\ & \quad + \int_{Q_T} g_n(x, t, u_n)(u_n - u) \, dx \, dt + \gamma \int_{Q_T} |u_n|^{p_0-2} u_n (u_n - u) \, dx \, dt \\ & \quad \quad \quad = \int_0^T \langle f, u_n - u \rangle \, dt. \end{aligned}$$

Hence,

$$\begin{aligned} & \int_0^T \langle Au_n - Au, u_n - u \rangle \, dt + \gamma \int_{Q_T} (|u_n|^{p_0-2} u_n - |u|^{p_0-2} u)(u_n - u) \, dx \, dt \\ & = - \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n - u \right\rangle \, dt - \int_{Q_T} g_n(x, t, u_n)(u_n - u) \, dx \, dt \\ & \quad + \int_0^T \langle f, u_n - u \rangle \, dt - \int_0^T \langle Au, u_n - u \rangle \, dt - \gamma \int_{Q_T} |u|^{p_0-2} u (u_n - u) \, dx \, dt. \end{aligned} \tag{4.29}$$

Now, we shall study the first and second terms on the right-hand side of (4.29). Due to the fact that

$$\frac{\partial u_n}{\partial t} \rightharpoonup \frac{\partial u}{\partial t} \quad \text{in } L^{p'}(0, T; W^{-1, p'}(\Omega))$$

and due to (4.22), we obtain

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n - u \right\rangle dt \\ &= \liminf_{n \rightarrow \infty} \int_{\Omega} \int_0^T \frac{\partial u_n}{\partial t} (u_n - u) dt dx \\ &= \liminf_{n \rightarrow \infty} \int_{\Omega} \int_0^T \frac{\partial u_n}{\partial t} u_n dt dx - \int_{\Omega} \int_0^T \frac{\partial u}{\partial t} u dt dx \\ &= \liminf_{n \rightarrow \infty} \int_{\Omega} \left[\frac{u_n^2(T) - u_n^2(0)}{2} \right] dx - \int_{\Omega} \left[\frac{u^2(T) - u^2(0)}{2} \right] dx \\ &= \liminf_{n \rightarrow \infty} \int_{\Omega} \frac{u_n^2(T)}{2} dx - \int_{\Omega} \frac{u^2(T)}{2} dx \\ &\geq 0, \end{aligned}$$

since $u_n(0) = u(0)$. Thus,

$$\limsup_{n \rightarrow \infty} - \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n - u \right\rangle dt = - \liminf_{n \rightarrow \infty} \int_0^T \left\langle \frac{\partial u_n}{\partial t}, u_n - u \right\rangle dt \leq 0. \quad (4.30)$$

On the other hand, we get that $g_n(x, t, u_n)$ is bounded in $L^{p_0'}(Q_T)$ and

$$g_n(x, t, u_n) \rightarrow g(x, t, u) \quad \text{a.e. in } Q_T.$$

Hence, in view of lemma 3.2, we get

$$g_n(x, t, u_n) \rightharpoonup g(x, t, u) \quad \text{in } L^{p_0'}(Q_T).$$

Then, by (4.23), we also get

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \int_{Q_T} g_n(x, t, u_n)(u_n - u) dx dt \\ &= \liminf_{n \rightarrow \infty} \int_{Q_T} g_n(x, t, u_n)u_n dx dt - \int_{Q_T} g(x, t, u)u dx dt \\ &\geq 0. \end{aligned}$$

Consequently,

$$\limsup_{n \rightarrow \infty} - \int_{Q_T} g_n(x, t, u_n)(u_n - u) dx dt \leq 0. \quad (4.31)$$

By combining (4.29)–(4.31), we deduce that

$$\begin{aligned}
 0 &\leq \limsup_{n \rightarrow \infty} \left(\int_0^T \langle Au_n - Au, u_n - u \rangle dt \right. \\
 &\quad \left. + \gamma \int_{Q_T} (|u_n|^{p_0-2}u_n - |u|^{p_0-2}u)(u_n - u) dx dt \right) \\
 &\leq \lim_{n \rightarrow \infty} \int_0^T \langle f, u_n - u \rangle dt - \lim_{n \rightarrow \infty} \int_0^T \langle Au, u_n - u \rangle dt \\
 &\quad - \gamma \lim_{n \rightarrow \infty} \int_{Q_T} |u|^{p_0-2}u(u_n - u) dx dt. \tag{4.32}
 \end{aligned}$$

Now, in view of (4.21), it is clear that, as $n \rightarrow \infty$,

$$\int_0^T \langle f, u_n - u \rangle dt \rightarrow 0, \tag{4.33}$$

$$\int_0^T \langle Au, u_n - u \rangle dt = \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u)(D^i u_n - D^i u) dx dt \rightarrow 0, \tag{4.34}$$

$$\int_{Q_T} |u|^{p_0-2}u(u_n - u) dx dt \rightarrow 0. \tag{4.35}$$

Hence, it follows that

$$\begin{aligned}
 &\int_0^T \langle Au_n - Au, u_n - u \rangle dt \\
 &\quad + \gamma \int_{Q_T} (|u_n|^{p_0-2}u_n - |u|^{p_0-2}u)(u_n - u) dx dt \rightarrow 0 \quad \text{as } n \rightarrow \infty. \tag{4.36}
 \end{aligned}$$

Thus, due to lemma 3.3, we deduce that

$$u_n \rightarrow u \quad \text{in } L^p(0, T; W_0^{1,p}(\Omega)). \tag{4.37}$$

STEP 5 (equi-integrability of $g_n(x, t, u_n)$). In order to prove that

$$g_n(x, t, u_n) \rightarrow g(x, t, u) \quad \text{strongly in } L^1(Q_T)$$

using Vitali’s theorem, it is sufficient to prove that $g_n(x, t, u_n)$ is uniformly equi-integrable. Indeed, let $m > 0$ and let E be a measurable subset of Q_T . Then, we have

$$\begin{aligned}
 &\int_E |g_n(x, t, u_n)| dx dt \\
 &\leq \int_{E \cap \{|u_n| \leq m\}} |g_n(x, t, u_n)| dx dt + \frac{1}{m} \int_{E \cap \{|u_n| > m\}} g_n(x, t, u_n)u_n dx dt \\
 &\leq \int_{E \cap \{|u_n| \leq m\}} (|u_n|^{p_0-1} + b(x, t)) dx dt + \frac{1}{m} \int_{Q_T} g_n(x, t, u_n)u_n dx dt \\
 &\leq |m|^{p_0-1}|E| + \int_E b(x, t) dx dt + \frac{c_2}{m}, \tag{4.38}
 \end{aligned}$$

where c_2 is the constant of (4.19), which is independent of n . Then, for all $\varepsilon > 0$, there exist m large enough such that $c_2/m < \varepsilon/2$ and $|E|$ sufficiently small to obtain

$$|m|^{p_0-1}|E| + \int_E b(x, t) \, dx \, dt < \frac{1}{2}\varepsilon.$$

Thus,

$$\int_E |g_n(x, t, u_n)| \, dx \, dt \leq \varepsilon.$$

Using Vitali's theorem, we obtain

$$g_n(x, t, u_n) \rightarrow g(x, t, u) \quad \text{in } L^1(Q_T). \quad (4.39)$$

Moreover, in view of (4.19), we have

$$\int_{Q_T} g(x, t, u)u \, dx \, dt \leq \liminf_{n \rightarrow \infty} \int_{Q_T} g_n(x, t, u_n)u_n \, dx \, dt \leq c_2.$$

Therefore, $g(x, t, u)u \in L^1(Q_T)$.

STEP 6 (passing to the limit). Let us take $v \in L^p(0, T; W_0^{1,p}(\Omega))$ as a test function in (4.2). We obtain

$$\begin{aligned} \int_0^T \left\langle \frac{\partial u_n}{\partial t}, v \right\rangle dt + \int_0^T \langle Au_n, v \rangle dt \\ + \int_{Q_T} g_n(x, t, u_n)v \, dx \, dt + \gamma \int_{Q_T} |u_n|^{p_0-2}u_n v \, dx \, dt = \int_0^T \langle f, v \rangle dt. \end{aligned} \quad (4.40)$$

It is clear to see that

$$\int_0^T \left\langle \frac{\partial u_n}{\partial t}, v \right\rangle dt \rightarrow \int_0^T \left\langle \frac{\partial u}{\partial t}, v \right\rangle dt \quad \text{as } n \rightarrow \infty. \quad (4.41)$$

By (4.37), we have $a_i(x, t, \nabla u_n) \rightarrow a_i(x, t, \nabla u)$ a.e. in Q_T , and $a_i(x, t, \nabla u_n)$ is bounded in $L^{p'_i}(Q_T)$. Then, in view of lemma 3.2, we get

$$a_i(x, t, \nabla u_n) \rightharpoonup a_i(x, t, \nabla u) \quad \text{in } L^{p'_i}(Q_T) \quad \text{for } i = 1, \dots, N.$$

Consequently, as $n \rightarrow \infty$,

$$\begin{aligned} \int_0^T \langle Au_n, v \rangle dt &= \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u_n) D^i v \, dx \, dt \\ &\rightarrow \sum_{i=1}^N \int_{Q_T} a_i(x, t, \nabla u) D^i v \, dx \, dt = \int_0^T \langle Au, v \rangle dt, \end{aligned} \quad (4.42)$$

$$\int_{Q_T} g_n(x, t, u_n)v \, dx \, dt \rightarrow \int_{Q_T} g(x, t, u)v \, dx \, dt, \quad (4.43)$$

and

$$\int_{Q_T} |u_n|^{p_0-2}u_n v \, dx \, dt \rightarrow \int_{Q_T} |u|^{p_0-2}uv \, dx \, dt. \quad (4.44)$$

Then, passing to the limit in (4.40), we obtain

$$\int_0^T \left\langle \frac{\partial u}{\partial t}, v \right\rangle dt + \int_0^T \langle Au, v \rangle dt + \int_{Q_T} g(x, t, u)v \, dx \, dt + \gamma \int_{Q_T} |u|^{p_0-2}uv \, dx \, dt = \int_0^T \langle f, v \rangle dt \quad (4.45)$$

for all $u, v \in L^p(0, T; W_0^{1,p}(\Omega))$.

This completes the proof. □

PROPOSITION 4.3. *Assume, furthermore, that*

$$(g(x, t, s) - g(x, t, r))(s - r) \geq 0 \text{ a.e. in } \Omega \text{ for any } s, r \in \mathbb{R}. \quad (4.46)$$

Then, the weak solution of problem (4.1) is unique.

Proof. Let $u_1, u_2 \in L^p(0, T; W_0^{1,p}(\Omega))$ be two weak solutions of (4.1). Then, we have

$$\begin{aligned} \int_0^T \left\langle \frac{\partial u_1}{\partial t} - \frac{\partial u_2}{\partial t}, v \right\rangle dt + \sum_{i=1}^N \int_{Q_T} (a_i(x, t, \nabla u_1) - a_i(x, t, \nabla u_2))D^i v \, dx \, dt \\ + \int_{Q_T} (g(x, t, u_1) - g(x, t, u_2))v \, dx \, dt \\ + \gamma \int_{Q_T} (|u_1|^{p_0-2}u_1 - |u_2|^{p_0-2}u_2)v \, dx \, dt = 0 \end{aligned}$$

for any $v \in L^p(0, T; W_0^{1,p}(\Omega))$. In particular, by taking $v = u_1 - u_2$ we obtain

$$\begin{aligned} 0 &\leq \sum_{i=1}^N \int_{Q_T} (a_i(x, t, \nabla u_1) - a_i(x, t, \nabla u_2))(D^i u_1 - D^i u_2) \, dx \, dt \\ &\quad + \gamma \int_{Q_T} (|u_1|^{p_0-2}u_1 - |u_2|^{p_0-2}u_2)(u_1 - u_2) \, dx \, dt \\ &= - \int_{Q_T} (g(x, t, u_1) - g(x, t, u_2))(u_1 - u_2) \, dx \, dt - \int_0^T \left\langle \frac{\partial u_1}{\partial t} - \frac{\partial u_2}{\partial t}, u_1 - u_2 \right\rangle dt. \end{aligned} \quad (4.47)$$

By (4.46), we get

$$\int_{Q_T} (g(x, t, u_1) - g(x, t, u_2))(u_1 - u_2) \, dx \, dt \geq 0.$$

Now, since $u_1(x, 0) = u_2(x, 0) = u_0(x)$,

$$\begin{aligned} \int_0^T \left\langle \frac{\partial u_1}{\partial t} - \frac{\partial u_2}{\partial t}, u_1 - u_2 \right\rangle dt &= \int_{\Omega} \int_0^T \frac{\partial (u_1 - u_2)^2}{\partial t} \, dt \, dx \\ &= \int_{\Omega} (u_1(T) - u_2(T))^2 \, dx \\ &\geq 0. \end{aligned}$$

Therefore, due to (4.47), we conclude that

$$\sum_{i=1}^N \int_{Q_T} (a_i(x, t, \nabla u_1) - a_i(x, t, \nabla u_2))(D^i u_1 - D^i u_2) \, dx \, dt + \gamma \int_{Q_T} (|u_1|^{p_0-2} u_1 - |u_2|^{p_0-2} u_2)(u_1 - u_2) \, dx \, dt = 0. \quad (4.48)$$

Then $u_1 = u_2$ a.e. in Q_T , which concludes our proof. \square

REMARK 4.4. If the hypothesis (4.46) is not supported, the uniqueness of the weak solution of the problem (4.1) is not guaranteed.

PROPOSITION 4.5. *Let $f \in L^{p'}(0, T; W^{-1, p'}(\Omega))$ and assume that (3.1)–(3.5) hold. Moreover, assume that*

$$g(x, t, s) \geq \delta |s|^{p_0} \quad (4.49)$$

with $\delta > 0$. Then, the problem

$$\left. \begin{aligned} u_t + Au + g(x, t, u) &= f && \text{in } Q_T = \Omega \times (0, T), \\ u(x, t) &= 0 && \text{on } S_T = \partial\Omega \times (0, T), \\ u(x, 0) &= u_0(x) && \text{in } \Omega, \end{aligned} \right\} \quad (4.50)$$

has at least one weak solution $u \in L^p(0, T; W_0^{1, p}(\Omega))$.

Indeed, we set $g(x, t, s) = h(x, t, s) + \delta |s|^{p_0-2} s$. Then

$$h(x, t, s) = g(x, t, s) - \delta |s|^{p_0-2} s \geq 0 \quad \text{and} \quad |h(x, t, s)| \leq |g(x, t, s)|.$$

Therefore, the existence of weak solutions follows immediately from theorem 4.1.

Acknowledgements

This work was done while S.E.M. was on sabbatical leave at Technische Universität Berlin (TU-Berlin), Germany. He is grateful to the Department of Mathematics at TU-Berlin, in particular to Professor Karl-Heinz Förster for his hospitality.

References

- 1 R. Adams. *Sobolev spaces* (New York: Academic, 1975).
- 2 S. Antontsev and S. Shmarev. Anisotropic parabolic equations with variable nonlinearity. *Publ. Mat.* **53** (2009), 355–399.
- 3 S. Antontsev and S. Shmarev. Vanishing solutions of anisotropic parabolic equations with variable nonlinearity. *J. Math. Analysis Applic.* **361** (2010), 371–391.
- 4 M. Bendahmane and K. H. Karlsen. Anisotropic nonlinear elliptic systems with measure data and anisotropic harmonic maps into spheres. *Electron. J. Diff. Eqns* **2006**(46) (2006), 1–30.
- 5 M. Bendahmane and M. Saad. Entropy solution for anisotropic reaction–diffusion–advection systems with L^1 data. *Rev. Mat. Complut.* **18** (2005), 49–67.
- 6 M. Bendahmane, P. Wittbold and A. Zimmermann. Renormalized solutions for a nonlinear parabolic equation with variable exponents and L^1 -data. *J. Diff. Eqns* **249** (2010), 1483–1515.
- 7 M. Bendahmane, M. Chrif and S. El Manouni. An approximation result in generalized anisotropic Sobolev spaces and application. *Z. Analysis Anwend.* **30** (2011), 341–353.

- 8 L. Boccardo, T. Gallouët and J. L. Vázquez. Some regularity results for some nonlinear parabolic equations in L^1 . *Rend. Sem. Mat. Univ. Polit. Torino* **49** (Special Issue) (1991), 69–74.
- 9 L. Boccardo, T. Gallouët and P. Marcellini. Anisotropic equations in L^1 . *Diff. Integ. Eqns* **9** (1996), 209–212.
- 10 L. Boccardo, A. Dall’Aglio, T. Gallouët and L. Orsina. Nonlinear parabolic equations with measure data. *J. Funct. Analysis* **147** (1997), 237–258.
- 11 E. Hewitt and K. Stromberg. *Real and abstract analysis* (Springer, 1965).
- 12 J. L. Lions. *Quelque méthodes de résolution des problèmes aux limites non linéaires* (Paris: Dunod/Gauthier-Villars, 1969).
- 13 S. M. Nikolskiĭ. An imbedding theorem for function with partial derivatives considered in different metric. *Am. Math. Soc. Transl. 2* **90** (1970), 27–44.
- 14 L. N. Slobodeckii. Generalized Sobolev spaces and their application to boundary problems for partial differential equation. *Am. Math. Soc. Transl. 2* **57** (1966), 207–275.
- 15 M. Troisi. Teoremi di inclusioine per spazi di Sobolev non isotropi. *Ricerche. Mat.* **18** (1969), 3–24.
- 16 N. S. Trudinger. An imbedding theorem for $H_0(G, \Omega)$ spaces. *Studia. Math.* **50** (1974), 17–30.