

THE LOGIC OF RESOURCES AND CAPABILITIES

MARTA BÍLKOVÁ

Department of Logic, Faculty of Arts, Charles University

GIUSEPPE GRECO

Department of Languages, Literature and Communication, University of Utrecht

ALESSANDRA PALMIGIANO

Faculty of Technology, Policy and Management, Delft University of Technology;
Department of Pure and Applied Mathematics, University of Johannesburg

APOSTOLOS TZIMOULIS

Faculty of Technology, Policy and Management, Delft University of Technology
and

NACHOEM WIJNBERG

Faculty of Economics and Business, University of Amsterdam; College of Business and
Economics, University of Johannesburg

Abstract. We introduce the logic LRC, designed to describe and reason about agents' abilities and capabilities in using resources. The proposed framework bridges two—up to now—mutually independent strands of literature: the one on logics of abilities and capabilities, developed within the theory of agency, and the one on logics of resources, motivated by program semantics. The logic LRC is suitable to describe and reason about key aspects of social behaviour in organizations. We prove a number of properties enjoyed by LRC (soundness, completeness, canonicity, and disjunction property) and its associated analytic calculus (conservativity, cut elimination, and subformula property). These results lay at the intersection of the algebraic theory of unified correspondence and the theory of multitype calculi in structural proof theory. Case studies are discussed which showcase several ways in which this framework can be extended and enriched while retaining its basic properties, so as to model an array of issues, both practically and theoretically relevant, spanning from planning problems to the logical foundations of the theory of organizations.

§1. Introduction. Organizations are social units of agents structured and managed to meet a need, or pursue collective goals. In economics and social science, organizations are studied in terms of agency, goals, capabilities, and inter-agent coordination [40,66,70]. In strategic management, the dominant approach in the study of organizational performances is the so-called *resource-based view* [2,55,73], which has recognized that a central role in determining the success of an organization in market competition is played by the acquisition, management, and transformation of *resources* within that organization. In order to capture this insight and create the building blocks of the logical foundations of

Received: August 7, 2016.

2010 *Mathematics Subject Classification*: 03B42, 03B20, 03B60, 03B45, 03F03, 03G10, 03A99.

Key words and phrases: display calculus, logics for organizations, multitype calculus, algebraic proof theory.

the theory of organizations, a formal framework is needed in which it is possible to express and reason about agents' abilities and capabilities to use resources for achieving goals, to transform resources into other resources, and to coordinate the use of resources with other agents; i.e., a formal framework is needed for capturing and reasoning about the *resource flow within organizations*. The present article aims at introducing such a framework.

There is extensive literature in philosophical logic and formal AI accounting for agents' abilities (cf. e.g., [8,27]) and capabilities (cf. e.g., [28,29,71]) and their interaction, embedding in the wider context of the logics of agency (cf. e.g., [4,5,9,10,30,67]); some of these frameworks (viz. [28,29]) have been used to formalize some aspects of the theory of organizations. There is also literature in theoretical computer science on the logic of resources (cf. e.g., [63,64]), motivated by the build-up of mathematical models of computational systems. However, these two strands of research have been pursued independently, and in particular, the *interaction* between abilities, capabilities, and resources has not been explored before.

The present article introduces a logical framework, the *logic of resources and capabilities* (LRC), designed as an environment for the logical modelling of the behaviour of agents motivated and mediated by the use and transformation of resources. In this framework, agents' capabilities are not captured via primitive actions, as is done, e.g., in [71], but rather via dedicated modalities, similarly to the frameworks adopting the STIT logic approach [5,28,29]. However, LRC differs from these logics in two main respects: the first is the focus on *resources*, discussed above; the second is that, as a modal extension of intuitionistic logic, LRC inherits its *constructive* character: it comes equipped with a constructive proof theory which provides an explicit computational content brought out by the cut elimination theorem. This guarantees that each LRC-theorem (prediction) translates into an effective procedure, thus allowing for a greater amenability to concrete applications in planning, and paving the way for implementations in constructive programming environments. In particular, LRC enjoys the disjunction property, proof of which we have included in §2.4.

In the present article, the basic mathematical theory of the logic of resources and capabilities is developed in an algebraic and proof-theoretic environment. Specifically, the most important technical tool we introduce for LRC is the *proof calculus* D.LRC (cf. §3). This calculus is designed according to the *multitype* methodology, introduced in [32–34], and further developed in [36,43,45,46]. This methodology exploits facts and insights coming from various semantic theories: from the coalgebraic semantics of dynamic epistemic logics (cf. [42]), to the algebraic dual of the team semantics for inquisitive logic (cf. [36]), the representation theorems for lattices (cf. [45]), and the recently developed algebraic theory of unified correspondence [16,19,20,23–25,60], in the context of which, systematic connections have been developed (cf. [44,54]) between Sahlqvist-type correspondence results and the theory of analytic rules for proper display calculi (cf. [72]) and Gentzen calculi.

Multitype languages make it possible to express constituents such as actions, agents, or resources not as *parameters* in the generation of formulas, but as *terms* in their own right. They thus are regarded as first-class citizens of the multitype framework and are endowed with their corresponding structural connectives and rules. In this rich environment, it is possible to encode certain key interactions *within the language*, by means of structural analytic rules. This approach has made it possible to develop analytic calculi for logics notoriously impervious to the standard proof-theoretic treatment, such as Public Announcement Logic [62], Dynamic Epistemic Logic [1], their nonclassical counterparts [51,53], and PDL [48].

One of the most important benefits of multitype calculi is the degree of *modularity* for which they allow. When applied to the present setting, the metatheory of multitype

calculi makes it possible to add (resp. remove) analytic structural rules to (resp. from) the basic calculus D.LRC, and obtain variants endowed with a package of basic properties (soundness, completeness, cut-elimination, subformula property, and conservativity) as immediate consequences of general results. This feature is illustrated and exploited in §5, where we specialize D.LRC to various situations by adding certain analytic structural rules to it. More in general, an infinite class of axiomatic extensions and combinatoric variants of LRC can be captured in a systematic way within this framework. Hence, LRC can be regarded not just as one single logic, but as a *class* of interconnected logical systems. Besides being of theoretical interest, this feature is of great usefulness in practice, since this class of logics forms a coherent framework which can be adapted to very different concrete settings with minimum effort. The combined strengths of this class of logics make the resulting LRC framework into a viable proposal for capturing and reasoning about the resource flow within organizations.

Finally, LRC is the first example of a logical system designed from first principles according to the multitype methodology. As this example shows, multitype calculi can serve not only to provide existing logics with well-performing calculi, but also as a methodological platform for the analysis and the meta-design of new logical frameworks.

Structure of the article. In §2.1, the logic LRC is introduced by means of a Hilbert-style presentation, which is shown to be complete w.r.t. certain algebraic models (cf. §2.2), canonical (cf. §2.3) and to enjoy the disjunction property (cf. §2.4). Then, in §3, the multitype calculus D.LRC is introduced, and is shown to be sound w.r.t. the algebraic models (cf. §4.1), complete (cf. §4.2), and conservative (cf. §4.4) w.r.t. the Hilbert-style presentation introduced in §2.1. In §4.3, we prove that the calculus D.LRC satisfies the assumptions of the cut-elimination metatheorem proven in [33], and hence enjoys cut-elimination and subformula property. In §5, we start exploring various ways in which D.LRC can be modified and adapted to different contexts so that the resulting systems retain all the properties enjoyed by the basic system. Specifically, §5.1 illustrates how *coordination* among agents helps optimizing capabilities towards a goal; §5.2 explores the solution of a planning problem which requires the suitable concatenation of *reusable* and *nonreusable* resources; §5.3 focuses on a situation in which the possibility of resources to be used in different *roles* becomes relevant; §5.4 illustrates how the *resilience* of a fragment of a system can propagate to the system as a whole.

§2. The logic of resources and capabilities and its algebraic semantics.

2.1. Hilbert-style presentation of LRC. As mentioned in the introduction, the key idea is to introduce a language in which resources are not accounted for as *parameters* indexing the capability connectives, but as *logical terms* in their own right. Accordingly, we start by defining a *multitype language* in which the different types interact via special connectives. The present setting consists of the types **Res** for resources and **Fm** for formulas (describing states of affairs). We stipulate that **Res** and **Fm** are disjoint.

Similarly to the binary connectives introduced in [34], the connectives \triangleright , \diamond , and \triangleright (referred to as *heterogeneous connectives*) facilitate the interaction between resources and formulas:¹

¹ As discussed below, these modal operators intend to capture agents' abilities and capabilities vis-à-vis resources; in this section, for the sake of a simpler exposition, we present the single-agent version of LRC, where any explicit mention of the agent is omitted.

$$\begin{aligned} \triangleright & : \text{Res} \times \text{Fm} \rightarrow \text{Fm} & \triangleright & : \text{Res} \times \text{Res} \rightarrow \text{Fm} \\ \diamond & : \text{Fm} \rightarrow \text{Fm} & \diamond & : \text{Res} \rightarrow \text{Fm}. \end{aligned}$$

As discussed in the next section, the mathematical environment of heterogeneous LRC-algebras provides a natural interpretation for all these connectives. Let us introduce the language of the logic of resources and capabilities. Let **AtProp** and **AtRes** be countable and disjoint sets of atomic propositions and atomic resources, respectively. The set $\mathcal{R} = \mathcal{R}(\text{AtRes})$ of the *resource-terms* α over **AtRes**, and the set $\mathcal{L} = \mathcal{L}(\mathcal{R}, \text{AtProp})$ of the *formula-terms* A over \mathcal{R} and **AtProp** of the Logic of Resources and Capabilities (LRC) are defined as follows:

$$\alpha ::= a \in \text{AtRes} \mid 1 \mid 0 \mid \alpha \cdot \alpha \mid \alpha \sqcup \alpha \mid \alpha \sqcap \alpha,$$

$$A ::= p \in \text{AtProp} \mid \top \mid \perp \mid A \vee A \mid A \wedge A \mid A \rightarrow A \mid \alpha \triangleright A \mid \diamond A \mid \diamond \alpha \mid \alpha \triangleright \alpha.$$

When writing formulas, we will omit brackets whenever the functional type of the connectives allows for a unique reading. Hence, for instance, we will write $\alpha \triangleright (\diamond A)$ as $\alpha \triangleright \diamond A$ and $(\alpha \cdot \beta) \triangleright A$ as $\alpha \cdot \beta \triangleright A$. We will also abide by the convention that $\vee, \wedge, \diamond, \diamond, \triangleright$, and \triangleright bind more strongly than \rightarrow , that $\diamond, \diamond, \triangleright$, and \triangleright bind more strongly than \vee and \wedge , and that \leftrightarrow is a weaker binder than any other connective. With this convention, for instance, $\alpha \triangleright A \wedge B$ has the same reading as $(\alpha \triangleright A) \wedge B$.

The (single-agent version of the) logic of resources and capabilities LRC, in its Hilbert-style presentation H.LRC, is defined as the smallest set of formulas containing the axioms and rules of intuitionistic propositional logic² plus the following axiom schemas:

Pure-resource entailment schemas:

- R1. \sqcup and \sqcap are commutative, associative, idempotent, and distribute over each other;
- R2. \cdot is associative with unit 1;
- R3. $\alpha \vdash 1$ and $0 \vdash \alpha$;
- R4. $\alpha \cdot (\beta \sqcup \gamma) \vdash (\alpha \cdot \beta) \sqcup (\alpha \cdot \gamma)$ and $(\beta \sqcup \gamma) \cdot \alpha \vdash (\beta \cdot \alpha) \sqcup (\gamma \cdot \alpha)$.

Axiom schemas for \diamond and \diamond :

- D1. $\diamond(A \vee B) \leftrightarrow \diamond A \vee \diamond B$;
- D2. $\diamond \perp \leftrightarrow \perp$;
- D3. $\diamond(\alpha \sqcup \beta) \leftrightarrow \diamond \alpha \vee \diamond \beta$;
- D4. $\diamond 0 \leftrightarrow \perp$.

Axiom schemas for \triangleright and \triangleright :

- B1. $(\alpha \sqcup \beta) \triangleright A \leftrightarrow \alpha \triangleright A \wedge \beta \triangleright A$;
- B2. $0 \triangleright A$;
- B3. $\alpha \triangleright \beta \triangleright A \rightarrow \alpha \cdot \beta \triangleright A$;
- B4. $(\alpha \sqcup \beta) \triangleright \gamma \leftrightarrow \alpha \triangleright \gamma \wedge \beta \triangleright \gamma$;
- B5. $0 \triangleright \alpha$;
- B6. $\alpha \triangleright (\beta \sqcap \gamma) \leftrightarrow \alpha \triangleright \beta \wedge \alpha \triangleright \gamma$;
- B7. $\alpha \triangleright 1$.

Interaction axiom schemas:

- BD1. $\diamond \alpha \wedge \alpha \triangleright A \rightarrow \diamond A$;
- BD2. $\alpha \triangleright \beta \rightarrow \alpha \triangleright \diamond \beta$.

and closed under modus ponens, uniform substitution and the following rules:

² The classical propositional logic counterpart of LRC can be obtained as usual by adding, e.g., excluded middle to the present axiomatization. Notice that classical propositional base is not needed in any of the case studies of §5.

$$\frac{\alpha \vdash \beta}{\alpha \cdot \gamma \vdash \beta \cdot \gamma} \text{MF} \quad \frac{A \vdash B}{\alpha \triangleright A \vdash \alpha \triangleright B} \text{MB} \quad \frac{A \vdash B}{\diamond A \vdash \diamond B} \text{MD} \quad \frac{\alpha \vdash \beta}{\gamma \triangleright \alpha \vdash \gamma \triangleright \beta} \text{MB}'$$

$$\frac{\alpha \vdash \beta}{\gamma \cdot \alpha \vdash \gamma \cdot \beta} \text{MF}' \quad \frac{\alpha \vdash \beta}{\beta \triangleright A \vdash \alpha \triangleright A} \text{AB} \quad \frac{\alpha \vdash \beta}{\diamond \alpha \vdash \diamond \beta} \text{MD}' \quad \frac{\alpha \vdash \beta}{\beta \triangleright \gamma \vdash \alpha \triangleright \gamma} \text{AB}'.$$

Finally, for all $A, B \in \mathcal{L}$, we let $A \vdash_{\text{LRC}} B$ iff a proof of B exists in H.LRC which possibly uses A .

Let us expand on the intuitive meaning of the connectives, axioms and rules introduced above, and their formal properties.

The pure-resource fragment of LRC. The pure-resource fragment of the logic LRC is inspired by (distributive) linear logic.³ Indeed, as is witnessed by conditions R1–R4 and rules MF and MF', the algebraic behaviour of \sqcap (with unit 1), \sqcup (with unit 0), and \cdot (with unit 1) is that of the additive conjunction, additive disjunction, and multiplicative conjunction in (distributive) linear logic, respectively. The intuitive understanding of the difference between $\alpha \cdot \beta$ and $\alpha \sqcap \beta$ is also borrowed from linear logic (cf. [41, §1.1.2]): indeed, $\alpha \cdot \beta$ can be intuitively understood as the resource obtained by *putting* α and β *together*. This 'putting resources together' can be interpreted in many ways in different contexts: one of them is, e.g., when α (water) and β (flour) are mixed together to obtain $\alpha \cdot \beta$ (dough); another is, e.g., when α (water) and β (flour), juxtaposed in separate jars, are used at the same time so to form the counterweight $\alpha \cdot \beta$ to keep something in balance. Notice that under both interpretations, $\alpha \cdot \alpha$ is distinct from α . We understand $\alpha \sqcap \beta$ as the resource which is as powerful as α and β taken *separately*. In other words, if we identify any resource γ with the (upward-closed) set of the states of affairs which can be brought about using γ (for brevity let us call such set the *power* of γ), then the resource $\alpha \sqcap \beta$ is uniquely identified by the *union* of the power of α and the power of β . Finally, we understand $\alpha \sqcup \beta$ as the resource the power of which is the intersection of the power of α and the power of β . More in general, the intended meaning of the resource-type entailment $\alpha \vdash \beta$ (namely ' α is at least as powerful a resource as β '), together with the identification of the lattice of resources with the lattice of their powers (which is a lattice of sets closed under union and intersection and hence distributive), explain intuitively the validity of resource-type entailments such as $\alpha \sqcap \alpha \dashv\vdash \alpha$, $\alpha \sqcup \alpha \dashv\vdash \alpha$, $\alpha \vdash \alpha \sqcup \beta$, and $\beta \vdash \alpha \sqcup \beta$, as well as $\alpha \sqcap (\beta \sqcup \gamma) \vdash (\alpha \sqcap \beta) \sqcup (\alpha \sqcap \gamma)$ and $(\alpha \sqcup \beta) \sqcap (\alpha \sqcup \gamma) \vdash \alpha \sqcup (\beta \sqcap \gamma)$. Moreover, under this reading of \vdash , by R3, the bottom 0 and top 1 of the lattice of resources can, respectively, be understood as the resource that is at least as powerful as any other resource (hence 0 is impossibly powerful), and the resource any other resource, no matter how weak, is at least as powerful as (hence 1 is the resource with no power, or the *empty resource*). This intuition, together with the uniqueness of the neutral element, also justifies one of the main differences between this setting and general linear logic; namely, the fact that the unit of \cdot is the unit of \sqcap . Indeed, it seems intuitively plausible that, under the most common interpretations of \cdot , putting together (e.g., mixing or juxtaposing) the empty resource and

³ However, the conceptual distinction is worth being stressed that, while formulas in linear logic *behave like* resources, pure-resource terms of LRC *literally denote* resources. In this respect, the pure-resource fragment of LRC is similar to the logic of resources introduced in [63,64].

any resource α yields α as outcome.⁴ Our inability to distinguish between the units of \sqcap and of \cdot yields as a consequence that the following entailments hold, which are also valid in linear affine logic [49,50]

$$\alpha \cdot \beta \vdash \alpha \quad \text{and} \quad \alpha \cdot \beta \vdash \beta. \quad (2.1)$$

Indeed, by R3, R2 and MF', $\alpha \cdot \beta \vdash \alpha \cdot 1 \vdash \alpha$, and the second entailment goes likewise. This restricts the scope of applications of the present setting: for instance, the fact that the compound resource $\alpha \cdot \beta$ must be at least as powerful as its two components rules out the general examples of, e.g., those chemical reactions in which the compound and its components are resources of incomparable power. On the other hand, it includes the case of all resources which can be quantified: two 50 euros bills are at least as powerful a resource than each 50 euros bill; two hours of time are at least as powerful a resource than one hour time, and so on. Moreover, this restriction does not rule out the possibility that the power of $\alpha \cdot \beta$ be strictly *greater* than the union of the separate powers of α and β (which is the power of $\alpha \sqcap \beta$). This is the case for instance when a critical mass of fuel is needed for reaching a certain temperature, or a certain outcome (e.g., a nuclear chain reaction). Another difference between the pure-resource fragment of LRC and linear logic is that, in LRC, the connective \cdot is not necessarily commutative.

The modal operators. The intended meaning of the formulas $\diamond A$ and $\diamond \alpha$ is 'the agent is able to bring about state of affairs A ' and 'the agent is in possession of resource α ', respectively. By axioms D1 and D2 (resp. D3 and D4), the connective \diamond (resp. \diamond) is a normal diamond-type connective (i.e., its algebraic interpretation is finitely join-preserving). Axiom D1 expresses that being able to bring about $A \vee B$ is tantamount to either being able to bring about A or being able to bring about B . Axiom D2 encodes the fact that the agent can never bring about logical contradictions. Analogously, Axiom D3 says that the agent is in possession of $\alpha \sqcup \beta$ exactly in case is in possession of α or is in possession of β . Axiom D4 encodes the fact that the agent is never in possession of the 'impossibly powerful resource' 0.

The intended meaning of the formula $\alpha \triangleright A$ is 'whenever resource α is in possession of the agent, using α the agent is capable to bring about A '. By axioms B1 and B2, the connective \triangleright is an antitone normal box-type operator in the first coordinate (i.e., its algebraic interpretation is finitely join-reversing in that coordinate). Axiom B1 says that the agent is capable of bringing about A whenever in possession of $\alpha \sqcup \beta$ iff the agent is capable of bringing about A *both* whenever in possession of α *and* whenever in possession of β . Axiom B2 means that if the agent were in possession of the impossibly powerful resource (which is never the case by D4), the agent could bring about any state of affairs. The justification of axiom B3 is connected with the constraint, encoded in (2.1), that the fusion $\alpha \cdot \beta$ of two resources is at least as powerful as each of its components. Taking this fact into account, let us assume that the agent is in possession of $\alpha \cdot \beta$. Hence, by (2.1), the resource in its possession is at least as powerful as the resources α and β taken in isolation. If $\alpha \triangleright \beta \triangleright A$ is the case, then by using $\alpha \cdot \beta$ up to α , the agent can bring about $\beta \triangleright A$, and by using the remainder of $\alpha \cdot \beta$, the agent can bring about A , which motivates B3. However, the converse direction is arguably not valid. Indeed, let $\alpha \cdot \beta \triangleright A$ express the fact that a

⁴ This is one of the main differences between actions and resources: the idle action *skip*, represented as the identity relation, is the unit of the product operation on actions, and is clearly different from the top element in the lattice of actions (the total relation).

certain temperature is reached by burning a critical mass $\alpha \cdot \beta$ of fuel. However, burning α and then β in sequence might not be enough to reach the same temperature.⁵

The intended meaning of the formula $\alpha \triangleright \beta$ is ‘the agent is capable of getting β from α , whenever in possession of α ’. By axioms B4 and B5, the connective \triangleright is an antitone normal box-type operator in the first coordinate (i.e., its algebraic interpretation is finitely join-reversing in that coordinate). Axiom B4 says that the agent is capable of getting resource γ whenever in possession of $\alpha \sqcup \beta$ iff the agent is capable of getting resource γ *both* whenever in possession of α *and* whenever in possession of β . Axiom B5 says that if the agent were in possession of the impossibly powerful resource (which is never the case by D4), the agent could get any resource. By axioms B6 and B7, the connective \triangleright is a monotone normal box-type operator in the second coordinate (i.e., its algebraic interpretation is finitely meet-preserving in that coordinate). Axiom B6 says that the agent is capable of getting resource $\beta \sqcap \gamma$ whenever in possession of α iff the agent is capable of getting *both* β and γ whenever in possession of α . Axiom B7 says that any agent is capable to get the empty resource whenever in possession of any resource.

Axiom BD1 encodes the link between the agent’s capabilities and abilities: indeed, it expresses the fact that if the agent is capable to bring about A whenever in possession of α ($\alpha \triangleright A$), and moreover the agent is actually in possession of α ($\Diamond \alpha$), then the agent is able to bring about A ($\Diamond A$). Notice also the analogy between this axiom and the intuitionistic axiom $A \wedge (A \rightarrow B) \leftrightarrow A \wedge B$. Axiom BD2 establishes a link between \triangleright and \triangleright , via \Diamond ; indeed, it says that the agent’s being capable to get β implies that the agent is capable to bring about a state of affairs in which the agent is in possession of β .

The rules MB and AB (resp. MB’ and AB’) encode the fact that \triangleright (resp. \triangleright) is monotone in its second coordinate and antitone in its first. In fact, AB, MB’, and AB’ can be derived using B1, B4, and B6. The monotonicity of \triangleright in its second coordinate expresses the intuition that if the agent is capable, whenever in possession of α , to bring about A , then is capable to bring about any state of affairs which is logically implied by A . The remaining rules encode the monotonicity of \Diamond , \Diamond , and \cdot .

Some additional axioms. We conclude the present discussion by mentioning some analytic axioms which might perhaps be interesting for different settings. We start mentioning $\Diamond \top$, $\Diamond 1$, and $\alpha \triangleright \top$, respectively, stating that the agent is able to bring about what is always the case, such as logical tautologies; the agent is in possession of the empty resource; the agent is capable of using any resource (hence also the empty one) to bring about what is always the case. We also mention $\alpha \triangleright \alpha$, stating that the agent is capable to get any resource already in the possession of the agent; $\Diamond \alpha \wedge \alpha \triangleright \beta \rightarrow \Diamond \beta$, and $\Diamond \alpha \wedge \alpha \triangleright \beta \rightarrow \Diamond \Diamond \beta$. The latter is a consequence of BD1 and BD2, while the former is used in the case study in §5.4. For the sake of achieving greater generality we chose not to include it in the general system. Axioms which might also be considered in special settings are $\alpha \triangleright (A \vee B) \rightarrow \alpha \triangleright A \vee \alpha \triangleright B$, and $\alpha \triangleright A \wedge \alpha \triangleright B \rightarrow \alpha \triangleright (A \wedge B)$. The first one would imply the distributivity of \triangleright over disjunction in its second coordinate. The axiom $\alpha \triangleright A \wedge \alpha \triangleright B \rightarrow \alpha \triangleright (A \wedge B)$ is not applicable in general, given that the consequence would require the duplication of the resource α . More generally applicable variants are

⁵ There is a surface similarity between B3 and Axiom Ac4 of [71, §4], which captures the interaction between the capabilities of agents to perform actions and composition of actions; however, as remarked in Footnote 4, composition of actions behaves differently from composition of resources, which is why B3 is an implication and not a bi-implication.

$\alpha \triangleright A \wedge \alpha \triangleright B \rightarrow \alpha \cdot \alpha \triangleright (A \wedge B)$ and $\alpha \triangleright \beta \wedge \alpha \triangleright \gamma \rightarrow \alpha \cdot \alpha \triangleright (\beta \cdot \gamma)$. The latter encodes the behaviour of scalable resources, and will be used in the case study of §5.2 and §5.4. Another interesting axiom is the converse of B3, which we have discussed above.

2.2. Algebraic completeness. In the present section we outline the completeness of LRC w.r.t. the heterogeneous LRC-algebras⁶ defined below, via a Lindenbaum–Tarski type construction.

DEFINITION 2.1. A heterogeneous LRC-algebra is a tuple $F = (\mathbb{A}, \mathbb{Q}, \triangleright, \diamond, \triangleright, \diamond)$ such that \mathbb{A} is a Heyting algebra, $\mathbb{Q} = (Q, \sqcup, \sqcap, \cdot, 0, 1)$ is a bounded distributive lattice with binary operator \cdot which preserves finite joins in each coordinate and the unit of which is 1,⁷ and $\triangleright : \mathbb{Q} \times \mathbb{A} \rightarrow \mathbb{A}$, $\diamond : \mathbb{A} \rightarrow \mathbb{A}$, $\triangleright : \mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{A}$, $\diamond : \mathbb{Q} \rightarrow \mathbb{A}$ verify the (quasi-)inequalities corresponding to the axioms and rules of LRC as presented in the previous section. A heterogeneous LRC-algebra is perfect if both \mathbb{A} and \mathbb{Q} are perfect,⁸ and the operations \triangleright , \diamond , \triangleright , and \diamond satisfy the infinitary versions of the join- and meet-preservation properties satisfied by definition in any heterogeneous LRC-algebra. An algebraic LRC-model is a tuple $\mathbb{M} := (F, v_{\text{Fm}}, v_{\text{Res}})$ such that F is a heterogeneous LRC-algebra, $v_{\text{Fm}} : \text{AtProp} \rightarrow \mathbb{A}$ and $v_{\text{Res}} : \text{AtRes} \rightarrow \mathbb{Q}$. Clearly, for every algebraic LRC-model \mathbb{M} , the assignments v_{Fm} and v_{Res} have unique homomorphic extensions which we identify with v_{Fm} and v_{Res} , respectively. For each $\text{T} \in \{\text{Fm}, \text{Res}\}$ and all terms a, b of type T , we let $a \models_{\text{LRC}} b$ iff $v_{\text{T}}(a) \leq v_{\text{T}}(b)$ for every model \mathbb{M} .

Given AtProp and AtRes , the Lindenbaum–Tarski heterogeneous LRC-algebra over AtProp and AtRes is defined to be the following structure:

$$F^* := (\mathbb{A}^*, \mathbb{Q}^*, \triangleright^*, \diamond^*, \triangleright^*, \diamond^*),$$

where:

1. \mathbb{A}^* is the quotient algebra $\text{Fm}/\dashv\vdash$, where Fm is the formula algebra corresponding to the language \mathcal{L} defined in the previous subsection, and $\dashv\vdash$ is the equivalence relation on Fm defined as $A \dashv\vdash A'$ iff $A \vdash A'$ and $A' \vdash A$. Notice that the rules MD, MB, AB, MD', MB', and AB' guarantee that $\dashv\vdash$ is compatible with \diamond , \triangleright , \diamond , and \triangleright , hence the quotient algebra construction is well defined. The elements of \mathbb{A}^* will be typically denoted $[B]$ for some formula $B \in \mathcal{L}$;
2. \mathbb{Q}^* is the quotient algebra $\text{Res}/\dashv\vdash$, where Res is the resource algebra corresponding to the language \mathcal{R} defined in the previous subsection, and $\dashv\vdash$ is the equivalence relation on Res defined as $\alpha \dashv\vdash \alpha'$ iff $\alpha \vdash \alpha'$ and $\alpha' \vdash \alpha$. Notice that the rules MF and MF' guarantee that $\dashv\vdash$ is compatible with \cdot , hence the quotient algebra construction is well defined. The elements of \mathbb{Q}^* will be typically denoted $[\alpha]$ for some resource $\alpha \in \mathcal{R}$;

⁶ This notion specializes the more general notion of heterogeneous algebras introduced in [6] to the setting of interest of the present article.

⁷ It immediately follows from the definition that $\alpha \cdot \beta \leq \alpha$ and $\alpha \cdot \beta \leq \beta$ for all $\alpha, \beta \in Q$.

⁸ A bounded distributive lattice (BDL) is *perfect* if it is complete, completely distributive and completely join-generated by its completely join-irreducible elements. A BDL is perfect iff it is isomorphic to the lattice of the upward-closed subsets of some poset. A Heyting algebra is *perfect* if its lattice reduct is a perfect BDL. A bounded distributive lattice with operators (abbreviated DLO. Operators are additional operations which are finitely join-preserving in each coordinate) is *perfect* if its lattice reduct is a perfect BDL, and each operator is completely join-preserving in each coordinate.

3. $\triangleright^* : Q^* \times A^* \rightarrow A^*$ is defined as $[a] \triangleright^*[B] := [a \triangleright B]$;
4. $\diamond^* : A^* \rightarrow A^*$ is defined as $\diamond^*[B] := [\diamond B]$;
5. $\bowtie^* : Q^* \times Q^* \rightarrow A^*$ is defined as $[a_1] \bowtie^*[a_2] := [a_1 \bowtie a_2]$;
6. $\diamondotimes^* : Q^* \rightarrow A^*$ is defined as $\diamondotimes^*[a] := [\diamondotimes a]$.

LEMMA 2.2. *For any AtProp and AtRes, F^* is a heterogeneous LRC-algebra.*

Proof. It is a standard verification that A^* is a Heyting algebra and that Q^* is a bounded distributive lattice with binary operator \cdot which preserves finite joins in each coordinate and the unit of which is 1. It is also an easy verification that \triangleright^* , \diamond^* , \bowtie^* , and \diamondotimes^* are well-defined, and verify the additional conditions by construction. □

The canonical assignments can be defined as usual, i.e., mapping atomic propositions and resources to their canonical value in F^* . Let \mathbb{M}^* be the resulting LRC-algebraic model. With this definition, the proof of the following proposition is routine, and is omitted.

PROPOSITION 2.3. *For all $X \subseteq \mathcal{L}$ and $A \in \mathcal{L}$, if $X \not\vdash_{\text{LRC}} A$, then $X \not\vdash_{\text{LRC}} A$.*

2.3. Algebraic canonicity. The present subsection is aimed at showing that LRC is strongly complete w.r.t. perfect heterogeneous LRC-algebras. This will be a key ingredient in the conservativity proof of §4.4.

DEFINITION 2.4. *Let $F = (A, Q, \triangleright, \diamond, \bowtie, \diamondotimes)$ be a heterogeneous LRC-algebra. The canonical extension of F is*

$$F^\delta = (A^\delta, Q^\delta, \triangleright^\delta, \diamond^\delta, \bowtie^\delta, \diamondotimes^\delta),$$

where A^δ and Q^δ are the canonical extensions of A and Q , respectively,⁹ the operations $\diamond^\sigma : Q^\delta \rightarrow A^\delta$ and $\bowtie^\sigma : Q^\delta \times Q^\delta \rightarrow A^\delta$ and $\diamond^\sigma : A^\delta \rightarrow A^\delta$ and $\triangleright^\sigma : Q^\delta \times A^\delta \rightarrow A^\delta$ are defined as follows: for any $k \in K(A^\delta)$, $\kappa \in K(Q^\delta)$ and $o \in O(A^\delta)$, $\omega \in O(Q^\delta)$,¹⁰

$$\diamond^\sigma \kappa := \bigwedge \{ \diamond \alpha \mid \alpha \in Q \text{ and } \kappa \leq \alpha \} \quad \kappa \bowtie^\sigma \omega := \bigvee \{ \alpha \bowtie \beta \mid \beta \in Q, \beta \leq \omega, \alpha \in Q \text{ and } \kappa \leq \alpha \},$$

⁹ The canonical extension of a BDL L is a complete distributive lattice L^δ containing L as a sublattice, such that

1. (*denseness*) every element of L^δ can be expressed both as a join of meets and as a meet of joins of elements from L ;
2. (*compactness*) for all $S, T \subseteq L$, if $\bigwedge S \leq \bigvee T$ in L^δ , then $\bigwedge F \leq \bigvee G$ for some finite sets $F \subseteq S$ and $G \subseteq T$.

It is well known that the canonical extension of a BDL is a perfect BDL (cf. Footnote 8). Completeness and complete distributivity imply that each perfect BDL is naturally endowed with a Heyting algebra structure, and hence each perfect BDL is also a perfect Heyting algebra. Moreover, if L is the lattice reduct of some Heyting algebra A , then A is a subalgebra of L^δ , seen as a perfect Heyting algebra. The canonical extension A^δ of a Heyting algebra A is defined as the canonical extension of the lattice reduct of A endowed with its natural Heyting algebra structure. The canonical extension Q^δ of a DLO Q is defined as the canonical extension of the lattice reduct of Q endowed with the σ -extension of each additional operator. It is well known that the canonical extension of a Heyting algebra (resp. DLO) is a perfect Heyting algebra (resp. DLO).

¹⁰ For any BDL L , an element $k \in L^\delta$ (resp. $o \in L^\delta$) is closed (resp. open) if it is the meet (resp. join) of some subset of L . The set of closed (resp. open) elements of L^δ is $K(L^\delta)$ (resp. $O(L^\delta)$). We will slightly abuse notation and write $K(A^\delta)$ (resp. $O(A^\delta)$) and $K(Q^\delta)$ (resp. $O(Q^\delta)$) to refer to the sets of closed and open elements of their lattice reducts.

$$\diamond^\sigma k := \bigwedge \{ \diamond a \mid a \in \mathbb{A} \text{ and } k \leq a \} \quad \kappa \triangleright^\sigma o := \bigvee \{ \alpha \triangleright a \mid a \in \mathbb{A}, a \leq o, \alpha \in \mathbb{Q} \text{ and } \kappa \leq \alpha \},$$

and for any $u \in \mathbb{A}^\delta$ and $q, w \in \mathbb{Q}^\delta$,

$$\diamond^\sigma q := \bigvee \{ \diamond^\sigma \kappa \mid \kappa \in K(\mathbb{Q}^\delta) \text{ and } \kappa \leq q \}$$

$$q \bowtie^\sigma w := \bigwedge \{ \kappa \bowtie^\sigma \omega \mid \omega \in O(\mathbb{Q}^\delta), w \leq \omega, \kappa \in K(\mathbb{Q}^\delta) \text{ and } \kappa \leq q \}$$

$$\diamond^\sigma u := \bigvee \{ \diamond^\sigma k \mid k \in K(\mathbb{A}^\delta) \text{ and } k \leq u \}$$

$$q \triangleright^\sigma u := \bigwedge \{ \kappa \triangleright^\sigma o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa \in K(\mathbb{Q}^\delta) \text{ and } \kappa \leq q \}.$$

Below we also report the definition of \cdot^σ for the reader's convenience: For $\kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta)$

$$\kappa_1 \cdot^\sigma \kappa_2 = \bigwedge \{ \alpha \cdot \beta \mid \alpha, \beta \in \mathbb{Q} \text{ and } \kappa_1 \leq \alpha, \kappa_2 \leq \beta \},$$

and for any $q_1, q_2 \in \mathbb{Q}^\delta$

$$q_1 \cdot^\sigma q_2 = \bigvee \{ \kappa_1 \cdot^\sigma \kappa_2 \mid \kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta) \text{ and } \kappa_1 \leq q_1, \kappa_2 \leq q_2 \}.$$

In what follows, for the sake of readability, we will write \cdot^σ without the superscript. This will not create ambiguities, since we use different variables to denote the elements of $\mathbb{Q}, K(\mathbb{Q}^\delta), O(\mathbb{Q}^\delta)$ and \mathbb{Q}^δ , and since \cdot and \cdot^σ coincide over \mathbb{Q} .

LEMMA 2.5. For any heterogeneous LRC-algebra F , the canonical extension F^δ is a perfect heterogeneous LRC-algebra.

Proof. As discussed in Footnote 9, \mathbb{A}^δ is a perfect Heyting algebra and \mathbb{Q}^δ is a perfect DLO, so to finish the proof it is enough to show that the validity of all axioms and rules of LRC transfers from F to F^δ , and moreover, the join-and meet-preservation properties of the operations of F hold in their infinitary versions in F^δ . Conditions R2 hold in \mathbb{Q}^δ as consequences of the general theory of canonicity of terms purely built on operators (cf. [38, Theorem 4.6]). As to D1 and D2, by assumption the operation \diamond preserves finite joins. Hence, by a well known fact of the theory of the σ -extensions of finitely join-preserving maps, \diamond^σ preserves arbitrary joins (cf. [38, Theorem 3.2]). The same argument applies to D3, D4, B4, B5, B6, and B7. Furthermore, by [39, Lemma 2.22] it follows that \triangleright^σ turns arbitrary joins into arbitrary meets in the first coordinate, which is the infinitary version of B1.

As to axiom B2, it is enough to show that for every $u \in \mathbb{A}^\delta$,

$$0 \triangleright^\sigma u = \top.$$

Let us preliminarily show the identity above for $o \in O(\mathbb{A}^\delta)$. Notice that the set $\{a \mid a \in \mathbb{A}, a \leq o\}$ is always nonempty since \perp belongs to it. Hence,

$$\begin{aligned} 0 \triangleright^\sigma o &= \bigvee \{ 0 \triangleright a \mid a \in \mathbb{A}, a \leq o \} \\ &= \bigvee \{ \top \mid a \in \mathbb{A}, a \leq o \} \\ &= \top. \end{aligned}$$

Hence, for arbitrary $u \in \mathbb{A}^\delta$

$$\begin{aligned} 0 \triangleright^\sigma u &= \bigwedge \{ 0 \triangleright^\sigma o \mid o \in O(\mathbb{A}^\delta), u \leq o \} \\ &= \bigwedge \{ \top \mid o \in O(\mathbb{A}^\delta), u \leq o \} \\ &= \top. \end{aligned}$$

As to B3, let us show that for all $q, w \in \mathbb{Q}^\delta$ and $u \in \mathbb{A}^\delta$,

$$q \triangleright^\pi w \triangleright^\pi u \leq q \cdot w \triangleright^\pi u.$$

Let us preliminarily show that the inequality above is true for any $o \in O(\mathbb{A}^\delta)$, $\kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta)$. By definition, if $o \in O(\mathbb{A}^\delta)$ and $\kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta)$ then $\kappa_2 \triangleright^\pi o \in O(\mathbb{A}^\delta)$ and $\kappa_1 \cdot \kappa_2 \in K(\mathbb{Q}^\delta)$. Therefore

$$\begin{aligned} & \kappa_1 \triangleright^\pi \kappa_2 \triangleright^\pi o \\ &= \bigvee \{ \alpha \triangleright a \mid a \in \mathbb{A}, \alpha \in \mathbb{Q}, \kappa_1 \leq \alpha, \alpha \leq \kappa_2 \triangleright^\pi o \} && \text{(by definition)} \\ &= \bigvee \{ \alpha \triangleright a \mid a \in \mathbb{A}, \alpha \in \mathbb{Q}, \kappa_1 \leq \alpha, \alpha \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \} \} && \text{(by definition)} \\ &= \bigvee \{ \alpha \triangleright \beta \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \alpha, \beta \in \mathbb{Q}, \kappa_1 \leq \alpha, \kappa_2 \leq \beta \} && (*) \\ &\leq \bigvee \{ \alpha \cdot \beta \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \alpha, \beta \in \mathbb{Q}, \kappa_1 \leq \alpha, \kappa_2 \leq \beta \} && \text{(B3 holds in } \mathbb{A} \text{)} \\ &\leq \bigvee \{ \gamma \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \gamma \in \mathbb{Q}, \kappa_1 \cdot \kappa_2 \leq \gamma \} && (**) \\ &= \kappa_1 \cdot \kappa_2 \triangleright^\pi o. && \text{(by definition)} \end{aligned}$$

Let us prove the equality marked with (*). If $a \in \mathbb{A}$, $\beta \in \mathbb{Q}$, $a \leq o$ and $\kappa \leq \beta$, then $\beta \triangleright a \in \mathbb{A}$ and $\beta \triangleright a \in \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \}$, hence $\beta \triangleright a \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \}$. This, in turn, implies that

$$\alpha \triangleright \beta \triangleright a \in \{ \alpha \triangleright a \mid a \in \mathbb{A}, \alpha \in \mathbb{Q}, \kappa_1 \leq \alpha, \alpha \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \} \}.$$

Therefore

$$\begin{aligned} & \{ \alpha \triangleright \beta \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \alpha, \beta \in \mathbb{Q}, \kappa_1 \leq \alpha, \kappa_2 \leq \beta \} \\ & \subseteq \{ \alpha \triangleright a \mid a \in \mathbb{A}, \alpha \in \mathbb{Q}, \kappa_1 \leq \alpha, \alpha \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \} \} \end{aligned}$$

and thus

$$\begin{aligned} & \bigvee \{ \alpha \triangleright \beta \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \alpha, \beta \in \mathbb{Q}, \kappa_1 \leq \alpha, \kappa_2 \leq \beta \} \\ & \leq \bigvee \{ \alpha \triangleright a \mid a \in \mathbb{A}, \alpha \in \mathbb{Q}, \kappa_1 \leq \alpha, \alpha \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \} \}. \end{aligned}$$

To prove the converse inequality, it is enough to show that if $a \in \mathbb{A}$ and $a \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \}$, then $\alpha \triangleright a \leq \alpha \triangleright \beta \triangleright b$ for some $b \in \mathbb{A}$ such that $b \leq o$ and some $\beta \in \mathbb{Q}$ such that $\kappa_2 \leq \beta$. By compactness (cf. Footnote 9), $a \leq \bigvee \{ \beta \triangleright b \mid b \in \mathbb{A}, b \leq o, \beta \in \mathbb{Q}, \kappa_2 \leq \beta \}$ implies that $a \leq \bigvee \{ \beta_i \triangleright b_i \mid 1 \leq i \leq n \}$ for some $b_i \in \mathbb{A}$, $\beta_i \in \mathbb{Q}$ such that $b_i \leq o$, $\kappa_2 \leq \beta_i$ for every $1 \leq i \leq n$. Since \triangleright is monotone in its second coordinate and antitone in its first, this implies that

$$a \leq \beta_1 \triangleright b_1 \vee \dots \vee \beta_n \triangleright b_n \leq (\beta_1 \sqcap \dots \sqcap \beta_n) \triangleright (b_1 \vee \dots \vee b_n).$$

Let $b := b_1 \vee \dots \vee b_n$ and $\beta = \beta_1 \sqcap \dots \sqcap \beta_n$. By definition, $b \in \mathbb{A}$, $\beta \in \mathbb{Q}$ and $b \leq o$, $\kappa_2 \leq \beta$. Moreover, again by monotonicity, the displayed inequality implies that $\alpha \triangleright a \leq \alpha \triangleright \beta \triangleright b$, as required. This finishes the proof of (*). The inequality marked with (**) holds since if $\kappa_1 \leq \alpha$ and $\kappa_2 \leq \beta$ then $\kappa_1 \cdot \kappa_2 \leq \alpha \cdot \beta$, so $\alpha \cdot \beta \triangleright a \in \{ \gamma \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \gamma \in \mathbb{Q}, \kappa_1 \cdot \kappa_2 \leq \gamma \}$ and therefore

$$\{ \alpha \cdot \beta \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \alpha, \beta \in \mathbb{Q}, \kappa_1 \leq \alpha, \kappa_2 \leq \beta \} \subseteq \{ \gamma \triangleright a \mid a \in \mathbb{A}, \alpha \leq o, \gamma \in \mathbb{Q}, \kappa_1 \cdot \kappa_2 \leq \gamma \}.$$

Let us show that B3 holds for arbitrary $u \in \mathbb{A}^\delta$ and $q, w \in \mathbb{Q}^\delta$.

$$\begin{aligned} & q \triangleright^\pi w \triangleright^\pi u \\ &= \bigwedge \{ \kappa_1 \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), \kappa_1 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, w \triangleright^\pi u \leq o \} && \text{(by definition)} \\ &= \bigwedge \{ \kappa_1 \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), \kappa_1 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \bigwedge \{ \kappa_2 \triangleright^\pi o' \mid u \leq o', \kappa_2 \leq w \} \leq o \} && \text{(by definition)} \\ &\leq \bigwedge \{ \kappa_1 \triangleright^\pi \kappa_2 \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \kappa_2 \leq w \} && (***) \\ &\leq \bigwedge \{ \kappa_1 \cdot \kappa_2 \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \kappa_2 \leq w \} && (\dagger) \\ &= \bigwedge \left\{ \left(\bigvee \{ \kappa_1 \cdot \kappa_2 \mid \kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \kappa_2 \leq w \} \right) \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), u \leq o \right\} && (\ddagger) \\ &= \bigwedge \{ q \cdot w \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), u \leq o \} && \text{(by definition)} \\ &= q \cdot w \triangleright^\pi u. && \text{(by definition)} \end{aligned}$$

The inequality marked with $(***)$ holds since, for any $o \in O(\mathbb{A}^\delta)$ and $\kappa \in K(\mathbb{Q}^\delta)$, if $u \leq o$ and $\kappa \leq w$ then $\kappa \triangleright^x o \in O(\mathbb{A}^\delta)$ and $\kappa \triangleright^x o \in \{\kappa_2 \triangleright^x o' \mid u \leq o', \kappa_2 \leq w\}$, hence $\bigwedge \{\kappa_2 \triangleright^x o' \mid u \leq o', \kappa_2 \leq w\} \leq \kappa \triangleright^x o$. This implies that

$$\kappa_1 \triangleright^x \kappa_2 \triangleright^x o \in \{\kappa_1 \triangleright^x o \mid o \in O(\mathbb{A}^\delta), \kappa_1 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \bigwedge \{\kappa_2 \triangleright^x o' \mid u \leq o', \kappa_2 \leq w\} \leq o\}$$

and therefore

$$\begin{aligned} & \{\kappa_1 \triangleright^x \kappa_2 \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \kappa_2 \leq w\} \\ & \subseteq \{\kappa_1 \triangleright^x o \mid o \in O(\mathbb{A}^\delta), \kappa_1 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \bigwedge \{\kappa_2 \triangleright^x o' \mid u \leq o', \kappa_2 \leq w\} \leq o\}, \end{aligned}$$

which implies that

$$\begin{aligned} & \bigwedge \{\kappa_1 \triangleright^x o \mid o \in O(\mathbb{A}^\delta), \kappa_1 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \bigwedge \{\kappa_2 \triangleright^x o' \mid u \leq o', \kappa_2 \leq w\} \leq o\} \\ & \leq \bigwedge \{\kappa_1 \triangleright^x \kappa_2 \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta), \kappa_1 \leq q, \kappa_2 \leq w\}. \end{aligned}$$

The inequality marked with (\dagger) holds since as we showed above B3 holds for any $o \in O(\mathbb{A}^\delta)$, $\kappa_1, \kappa_2 \in K(\mathbb{Q}^\delta)$. The equality marked with (\ddagger) holds because \triangleright^x is completely join reversing in the first coordinate.

As to axiom BD1, let us show that for any $q \in \mathbb{Q}^\delta$ and $u \in \mathbb{A}^\delta$,

$$\diamond^\sigma q \wedge q \triangleright^x u \leq \diamond^\sigma u.$$

Let us preliminarily show that the inequality above is true for any $o \in O(\mathbb{A}^\delta)$ and $\kappa \in K(\mathbb{Q}^\delta)$:

$$\begin{aligned} & \diamond^\sigma \kappa \wedge \kappa \triangleright^x o \\ & = \bigwedge \{\diamond \beta \mid \beta \in \mathbb{Q}, \kappa \leq \beta\} \wedge \bigvee \{\alpha \triangleright a \mid a \in \mathbb{A}, a \leq o, \alpha \in \mathbb{Q}, \kappa \leq \alpha\} \quad (\text{by definition}) \\ & = \bigvee \{(\bigwedge \{\diamond \beta \mid \beta \in \mathbb{Q}, \kappa \leq \beta\}) \wedge \alpha \triangleright a \mid a \in \mathbb{A}, a \leq o\} \quad (\text{distributivity}) \\ & \leq \bigvee \{\diamond \alpha \wedge \alpha \triangleright a \mid a \in \mathbb{A}, a \leq o\} \quad (*) \\ & \leq \bigvee \{\diamond a \mid a \in \mathbb{A}, a \leq o\} \quad (\text{BD2 holds in } \mathbb{A}) \\ & = \diamond^\sigma \bigvee \{a \mid a \in \mathbb{A}, a \leq o\} \quad (\diamond^\sigma \text{ is completely join-preserving}) \\ & = \diamond^\sigma o. \quad (\text{by definition}) \end{aligned}$$

The inequality marked with $(*)$ holds because if $\kappa \leq \alpha$, then $\diamond \alpha \in \{\diamond \beta \mid \beta \in \mathbb{Q}, \kappa \leq \beta\}$ and therefore $\bigwedge \{\diamond \beta \mid \beta \in \mathbb{Q}, \kappa \leq \beta\} \leq \diamond \alpha$. Let us show the inequality for arbitrary $u \in \mathbb{A}^\delta$ and $q \in \mathbb{Q}^\delta$. In what follows, let \blacksquare denote the right adjoint of \diamond^σ . It is well known (cf. [21, Lemma 10.3.3]) that $\blacksquare o \in O(\mathbb{A}^\delta)$ for any $o \in O(\mathbb{A}^\delta)$.

$$\begin{aligned} & \diamond^\sigma q \wedge q \triangleright^x u \\ & = \bigvee \{\diamond^\sigma \kappa \mid \kappa \in K(\mathbb{Q}^\delta), \kappa \leq q\} \wedge \bigwedge \{\kappa' \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa' \in K(\mathbb{Q}^\delta), \kappa' \leq q\} \quad (\text{by definition}) \\ & = \bigvee \{\diamond^\sigma \kappa \wedge \bigwedge \{\kappa' \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa' \in K(\mathbb{Q}^\delta), \kappa' \leq q\} \mid \kappa \in K(\mathbb{Q}^\delta), \kappa \leq q\} \quad (\text{distributivity}) \\ & \leq \bigvee \{\diamond^\sigma \kappa \wedge \bigwedge \{\kappa \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o\} \mid \kappa \in K(\mathbb{Q}^\delta), \kappa \leq q\} \quad (*) \\ & \leq \bigvee \{\diamond^\sigma \kappa \wedge \bigwedge \{\kappa \triangleright^x \blacksquare o \mid o \in O(\mathbb{A}^\delta), u \leq \blacksquare o\} \mid \kappa \in K(\mathbb{Q}^\delta), \kappa \leq q\} \quad (\blacksquare o \in O(\mathbb{A}^\delta)) \\ & \leq \bigvee \{\bigwedge \{\diamond^\sigma \kappa \wedge \kappa \triangleright^x \blacksquare o \mid o \in O(\mathbb{A}^\delta), u \leq \blacksquare o\} \mid \kappa \in K(\mathbb{Q}^\delta), \kappa \leq q\} \quad (\text{distributivity}) \\ & \leq \bigvee \{\bigwedge \{\diamond^\sigma \blacksquare o \mid o \in O(\mathbb{A}^\delta), u \leq \blacksquare o\} \mid \kappa \in K(\mathbb{Q}^\delta), \kappa \leq q\} \quad (\text{BD2 holds in } O(\mathbb{A}^\delta)) \\ & = \bigwedge \{\diamond^\sigma \blacksquare o \mid o \in O(\mathbb{A}^\delta), u \leq \blacksquare o\} \quad (\diamond^\sigma \blacksquare o \text{ does not contain } \kappa) \\ & \leq \bigwedge \{o \in O(\mathbb{A}^\delta) \mid u \leq \blacksquare o\} \quad (\diamond^\sigma \blacksquare o \leq o) \\ & = \bigwedge \{o \in O(\mathbb{A}^\delta) \mid \diamond^\sigma u \leq o\} \quad (\text{by adjunction}) \\ & = \diamond^\sigma u. \quad (\text{by denseness}) \end{aligned}$$

The inequality marked with $(*)$ holds because if $\kappa \leq q$ and $u \leq o$, then

$$\kappa \triangleright^x o \in \{\kappa' \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa' \in K(\mathbb{Q}^\delta), \kappa' \leq q\}$$

and therefore

$$\{\kappa \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o\} \subseteq \{\kappa' \triangleright^x o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa' \in K(\mathbb{Q}^\delta), \kappa' \leq q\},$$

which yields

$$\bigwedge \{ \kappa' \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), u \leq o, \kappa' \in K(\mathbb{Q}^\delta), \kappa' \leq q \} \leq \bigwedge \{ \kappa \triangleright^\pi o \mid o \in O(\mathbb{A}^\delta), u \leq o \}.$$

Finally for axiom BD2 let us show that for any $q, w \in \mathbb{Q}^\delta$,

$$q \vDash^\pi w \leq q \triangleright^\pi \Diamond^\sigma w.$$

Let us preliminarily show that the inequality above is true for any $\omega \in O(\mathbb{Q}^\delta)$ and $\kappa \in K(\mathbb{Q}^\delta)$. Notice that since \Diamond^σ is completely join preserving, if $\omega \in O(\mathbb{Q}^\delta)$ then $\Diamond^\sigma \omega \in O(\mathbb{A}^\delta)$.

$$\begin{aligned} & \kappa \vDash^\pi \omega \\ &= \bigvee \{ \alpha \triangleright \beta \mid \alpha, \beta \in \mathbb{Q}, \kappa \leq \alpha, \beta \leq \omega \} && \text{(by definition)} \\ &= \bigvee \{ \alpha \triangleright \Diamond \beta \mid \alpha, \beta \in \mathbb{Q}, \kappa \leq \alpha, \beta \leq \omega \} && \text{(BD2 holds in } \mathbb{A} \text{)} \\ &\leq \bigvee \{ \alpha \triangleright a \mid \alpha \in \mathbb{Q}, a \in \mathbb{A}, \kappa \leq \alpha, a \leq \Diamond^\sigma \omega \} && (*) \\ &= \kappa \triangleright^\pi \Diamond^\sigma \omega. && \text{(by definition)} \end{aligned}$$

The inequality marked with (*) holds because if $\beta \leq \omega$ then $\Diamond \beta \leq \Diamond^\sigma \omega$, thus if $\kappa \leq \alpha$ we have

$$\alpha \triangleright \Diamond \beta \in \{ \alpha \triangleright a \mid \alpha \in \mathbb{Q}, a \in \mathbb{A}, \kappa \leq \alpha, a \leq \Diamond^\sigma \omega \}$$

and therefore

$$\{ \alpha \triangleright \Diamond \beta \mid \alpha, \beta \in \mathbb{Q}, \kappa \leq \alpha, \beta \leq \omega \} \subseteq \{ \alpha \triangleright a \mid \alpha \in \mathbb{Q}, a \in \mathbb{A}, \kappa \leq \alpha, a \leq \Diamond^\sigma \omega \}.$$

Let us show the inequality for arbitrary $q, w \in \mathbb{Q}^\delta$. In what follows, let $\blacksquare : \mathbb{A}^\delta \rightarrow \mathbb{Q}^\delta$ denote the right adjoint of \Diamond^σ .

$$\begin{aligned} & q \vDash^\pi w \\ &= \bigwedge \{ \kappa \vDash^\pi \omega \mid \kappa \in K(\mathbb{Q}^\delta), \omega \in O(\mathbb{Q}^\delta), \kappa \leq q, w \leq \omega \} && \text{(by definition)} \\ &\leq \bigwedge \{ \kappa \vDash^\pi \blacksquare o \mid \kappa \in K(\mathbb{Q}^\delta), o \in O(\mathbb{A}^\delta), \kappa \leq q, w \leq \blacksquare o \} && (\blacksquare o \in O(\mathbb{Q}^\delta)) \\ &\leq \bigwedge \{ \kappa \triangleright^\pi \Diamond^\sigma \blacksquare o \mid \kappa \in K(\mathbb{Q}^\delta), o \in O(\mathbb{A}^\delta), \kappa \leq q, w \leq \blacksquare o \} && \text{(BD2 holds for } \omega \in O(\mathbb{Q}^\delta) \text{ and } \kappa \in K(\mathbb{Q}^\delta) \text{)} \\ &\leq \bigwedge \{ \kappa \triangleright^\pi o \mid \kappa \in K(\mathbb{Q}^\delta), o \in O(\mathbb{A}^\delta), \kappa \leq q, w \leq \blacksquare o \} && (\Diamond^\sigma \blacksquare o \leq o) \\ &= \bigwedge \{ \kappa \triangleright^\pi o \mid \kappa \in K(\mathbb{Q}^\delta), o \in O(\mathbb{A}^\delta), \kappa \leq q, \Diamond^\sigma w \leq o \} && \text{(by adjunction)} \\ &= q \triangleright^\pi \Diamond^\sigma w. && \text{(by definition)} \end{aligned}$$

□

As an immediate consequence of Proposition 2.3 and Lemma 2.5 we get the following:

COROLLARY 2.6. *The logic LRC is strongly complete w.r.t. the class of perfect heterogeneous LRC-algebras.*

2.4. Disjunction property. In the present section, we show that the disjunction property holds for LRC, by adapting the standard argument to the setting of heterogeneous LRC-algebras. For any heterogeneous LRC-algebra $F = (\mathbb{A}, \mathbb{Q}, \triangleright, \Diamond, \vDash, \Diamond)$, we let $F^* := (\mathbb{A}^*, \mathbb{Q}, \triangleright^*, \Diamond^*, \vDash^*, \Diamond^*)$, where:

1. \mathbb{A}^* is the Heyting algebra obtained by adding a new top element \top^* to \mathbb{A} (we let $\top_{\mathbb{A}}$ denote the top element of \mathbb{A}). Joins and meets in \mathbb{A}^* are defined as expected. The implication \rightarrow^* of \mathbb{A}^* maps any $(u, w) \in \mathbb{A}^* \times \mathbb{A}^*$ to \top^* if $u \leq w$, to w if $u = \top^*$, and to $u \rightarrow w$ in any other case.
2. $\triangleright^* : \mathbb{Q} \times \mathbb{A}^* \rightarrow \mathbb{A}^*$ maps any (α, u) to \top^* if $\alpha = 0$ or $\Diamond^* 1 \leq u$, and to $\alpha \triangleright u$ otherwise.
3. $\Diamond^* : \mathbb{A}^* \rightarrow \mathbb{A}^*$ maps any u to $\Diamond u$ if $u \neq \top^*$, and to $\Diamond \top_{\mathbb{A}}$ if $u = \top^*$.

4. $\triangleright^* : \mathbb{Q} \times \mathbb{Q} \rightarrow \mathbb{A}^*$ maps any (α, β) to \top^* if $\alpha = 0$ or $\beta = 1$, and to $\alpha \triangleright \beta$ otherwise.
5. $\diamond^* : \mathbb{Q} \rightarrow \mathbb{A}^*$ maps any α to $\diamond \alpha$.

LEMMA 2.7. *F* is a heterogeneous LRC-algebra.*

Proof. It can be easily verified that the maps $\triangleright^*, \diamond^*, \triangleright^*, \diamond^*$ satisfy by definition all the monotonicity (resp. antitonicity) properties that yield the validity of the rules of LRC. Let us verify that F^* validates all the axioms of LRC. By construction, \top^* is join-irreducible, i.e., if $u \vee w = \top^*$ then either $u = \top^*$ or $w = \top^*$. Hence, $\diamond^*(u \vee w) = \diamond^*\top^* = \diamond\top_{\mathbb{A}} = \diamond^*u \vee \diamond^*w$. All the remaining cases follow from the assumptions on \diamond . This finishes the verification of the validity of $D1$. The validity of axioms $D2, D3$, and $D4$ immediately follows from their validity in F . The validity of axiom $B1$ can be shown using the identities $\alpha \sqcup 0 = \alpha$ and $0 \sqcup \beta = \beta$. The validity of $B2$ follows immediately from the definition of \triangleright^* . As to $B3$, if $\alpha = 0$ or $\beta = 0$, the assumption that \cdot preserves finite joins in each coordinate yields $\alpha \cdot \beta = 0$, and hence $\alpha \cdot \beta \triangleright^* A = \top^*$, which implies that the inequality holds. The remaining cases follow from the definition of \triangleright^* and the assumption that $B3$ is valid in F . Axiom $B4$ is argued similarly to $B1$. The validity of axioms $B5$ and $B7$ follows immediately from the definition of \triangleright , and the validity of $B6$ can be shown using the identities $\alpha \sqcap 1 = \alpha = 1 \sqcap \alpha$.

As to $BD2$, if $\alpha = 0$ or $\beta = 1$ then $\alpha \triangleright^* \diamond^* \beta = \top^*$, therefore the inequality holds. All the remaining cases follow from the assumption that $BD2$ is valid in F .

As to $BD1$, if $\alpha = 0$ then $\diamond^* \alpha \wedge \alpha \triangleright^* u = \perp$ for any u , therefore the inequality holds. If $\diamond^* 1 \leq u$ then, by definition, $\alpha \triangleright^* u = \top^*$, hence it is enough to show that $\diamond^* \alpha \leq \diamond^* u$. We proceed by cases: (a) if $u = \top^*$, then $\diamond^* \alpha = \diamond \alpha \leq \top_{\mathbb{A}} = \diamond^* u$, as required; (b) if $u \in \mathbb{A}$, then, by the assumption that $B7, BD2$ and MB hold in F ,

$$\top_{\mathbb{A}} \leq \alpha \triangleright 1 \leq \alpha \triangleright \diamond 1 \leq \alpha \triangleright u.$$

Since $BD1$ holds in F , this implies that $\diamond^* \alpha = \diamond \alpha \leq \diamond u = \diamond^* u$, as required. All the remaining cases follow from the assumption that $BD1$ is valid in F . □

For every algebraic LRC-model $\mathbb{M} = (F, v_{Fm}, v_{Res})$, we let $\mathbb{M}^* := (F^*, v_{Fm}^*, v_{Res})$, where v_{Fm}^* is defined by composing v_{Fm} with the natural injection $\mathbb{A} \hookrightarrow \mathbb{A}^*$. Henceforth, we let $\llbracket a \rrbracket$ denote the interpretation of any \top -term a in \mathbb{M} and $\llbracket a \rrbracket_*$ the interpretation of a in \mathbb{M}^* .

LEMMA 2.8. *For every formula A,*

1. *If $\llbracket A \rrbracket_* \neq \top^*$ then $\llbracket A \rrbracket_* = \llbracket A \rrbracket$.*
2. *If $\llbracket A \rrbracket_* = \top^*$ then $\llbracket A \rrbracket = \top_{\mathbb{A}}$.*

Proof. We prove the two statements simultaneously by induction on A . The cases of constants and atomic variables are straightforward. The case of $A = B \wedge C$ immediately follows from the induction hypothesis. The case of $A = B \vee C$ immediately follows from the induction hypothesis using the join-irreducibility of \top^* . If $A = B \rightarrow C$, then $\llbracket A \rrbracket_* = \llbracket B \rrbracket_* \rightarrow^* \llbracket C \rrbracket_*$. By definition of \rightarrow^* , if $\llbracket A \rrbracket_* \neq \top^*$ then either (a) $\llbracket B \rrbracket_* \not\leq \llbracket C \rrbracket_*$ and $\llbracket B \rrbracket_* \neq \top^*$, which implies that $\llbracket B \rrbracket_* \neq \top^* \neq \llbracket C \rrbracket_*$ in which case item 1 follows by induction hypothesis; or (b) $\llbracket B \rrbracket_* \leq \llbracket C \rrbracket_*$ and $\llbracket C \rrbracket_* \neq \top^*$, which implies that $\llbracket C \rrbracket_* = \llbracket C \rrbracket$ by induction hypothesis. Then either (b1) $\llbracket B \rrbracket_* = \top^*$, hence by induction hypothesis $\llbracket B \rrbracket = \top_{\mathbb{A}}$ and $\llbracket A \rrbracket_* = \llbracket C \rrbracket_* = \llbracket C \rrbracket = \llbracket A \rrbracket$, as required; or (b2) $\llbracket B \rrbracket_* \neq \top^*$,

hence by induction hypothesis $\llbracket B \rrbracket_* = \llbracket B \rrbracket$ and we finish the proof as in case (a). If $\llbracket A \rrbracket_* = \top^*$, then either (c) $\llbracket B \rrbracket_* = \top^* = \llbracket C \rrbracket_*$, which implies by induction hypothesis that $\llbracket B \rrbracket = \top^\Delta = \llbracket C \rrbracket$, which yields $\llbracket A \rrbracket = \top^\Delta$, as required; or (d) $\llbracket B \rrbracket_* \leq \llbracket C \rrbracket_*$, which implies $\llbracket B \rrbracket \leq \llbracket C \rrbracket$ and hence $\llbracket A \rrbracket = \top^\Delta$, as required.

If $A = \diamond B$, then $\llbracket A \rrbracket_* = \diamond^* \llbracket B \rrbracket_*$. The definition of \diamond^* implies that $\llbracket A \rrbracket_* \neq \top^*$, hence to finish the proof of this case we need to show that $\llbracket A \rrbracket_* = \llbracket A \rrbracket$. If $\llbracket B \rrbracket_* \neq \top^*$, then by induction hypothesis $\llbracket B \rrbracket_* = \llbracket B \rrbracket$, hence $\llbracket A \rrbracket_* = \diamond^* \llbracket B \rrbracket_* = \diamond \llbracket B \rrbracket = \llbracket \diamond B \rrbracket = \llbracket A \rrbracket$, as required. If $\llbracket B \rrbracket_* = \top^*$, then by induction hypothesis $\llbracket B \rrbracket = \top^\Delta$, hence $\llbracket A \rrbracket_* = \diamond^* \llbracket B \rrbracket_* = \diamond^* \top^* = \diamond \top^\Delta = \diamond \llbracket B \rrbracket = \llbracket \diamond B \rrbracket = \llbracket A \rrbracket$, as required.

If $A = \diamond \alpha$, item 2 is again vacuously true, and item 1 immediately follows from the definition of \diamond^* .

If $A = \alpha \triangleright \beta$, then $\llbracket A \rrbracket_* = \llbracket \alpha \rrbracket_* \triangleright^* \llbracket \beta \rrbracket_* = \llbracket \alpha \rrbracket \triangleright^* \llbracket \beta \rrbracket$. Then by definition of \triangleright^* , if $\llbracket A \rrbracket_* \neq \top^*$, then $\llbracket A \rrbracket_* = \llbracket A \rrbracket$, as required, and if $\llbracket A \rrbracket_* = \top^*$, then either $\llbracket \alpha \rrbracket = 0$ or $\llbracket \beta \rrbracket = 1$; since axioms B5 and B7 hold in F , each case yields $\llbracket A \rrbracket = \top^\Delta$, as required.

Finally, if $A = \alpha \triangleright B$, then $\llbracket A \rrbracket_* = \llbracket \alpha \rrbracket_* \triangleright^* \llbracket B \rrbracket_* = \llbracket \alpha \rrbracket \triangleright^* \llbracket B \rrbracket_*$. By definition of \triangleright^* , if $\llbracket A \rrbracket_* \neq \top^*$, then $\llbracket \alpha \rrbracket \neq 0$, $\llbracket A \rrbracket_* = \llbracket \alpha \rrbracket \triangleright \llbracket B \rrbracket_*$, and $\diamond^* 1 \not\leq \llbracket B \rrbracket_*$. The latter condition implies that $\llbracket B \rrbracket_* \neq \top^*$, hence, by induction hypothesis, $\llbracket B \rrbracket_* = \llbracket B \rrbracket$, and so $\llbracket A \rrbracket_* = \llbracket \alpha \rrbracket \triangleright \llbracket B \rrbracket = \llbracket A \rrbracket$, as required. If $\llbracket A \rrbracket_* = \top^*$, then either (a) $\llbracket \alpha \rrbracket = 0$, which implies by B2 that $\llbracket A \rrbracket = \llbracket \alpha \rrbracket \triangleright \llbracket B \rrbracket = \top^\Delta$, as required; or (b) $\diamond^* 1 \leq \llbracket B \rrbracket_*$, which implies by induction hypothesis that $\diamond^* 1 \leq \llbracket B \rrbracket$. Hence, by BD2 and monotonicity of \triangleright ,

$$\top^\Delta \leq \llbracket \alpha \rrbracket \triangleright 1 \leq \llbracket \alpha \rrbracket \triangleright \diamond 1 \leq \llbracket \alpha \rrbracket \triangleright \llbracket B \rrbracket,$$

which finishes the proof that $\llbracket A \rrbracket = \llbracket \alpha \triangleright B \rrbracket = \top^\Delta$, as required. □

The *product* $F_1 \times F_2$ of the heterogeneous LRC-algebras F_1 and F_2 is defined in the expected way, based on the product algebras $\mathbb{A}_1 \times \mathbb{A}_2$ and $\mathbb{Q}_1 \times \mathbb{Q}_2$, and defining all (i.e., both internal and external) operations component-wise. It can be readily verified that the resulting construction is a heterogeneous LRC-algebra. The product construction can be extended to algebraic LRC-models in the expected way, i.e., by pairing the valuations. Such valuations extend as usual to T-terms, and it can be proved by a straightforward induction that $\llbracket a \rrbracket_\times = (\llbracket a \rrbracket_1, \llbracket a \rrbracket_2)$.

PROPOSITION 2.9. *The disjunction property holds for the logic LRC.*

Proof. If B and C are not LRC-theorems, by completeness, algebraic LRC-models \mathbb{M}_1 and \mathbb{M}_2 exist such that $\llbracket B \rrbracket_1 \neq \top_1$ and $\llbracket C \rrbracket_2 \neq \top_2$. Consider the product model $\mathbb{M} := \mathbb{M}_1 \times \mathbb{M}_2$ as described above. Notice that $\llbracket B \rrbracket \neq (\top_1, \top_2)$ and likewise for C . The model \mathbb{M}^* does not satisfy $B \vee C$. Indeed, since \top^* is join-irreducible, if $\llbracket B \vee C \rrbracket_* = \top^*$ then either $\llbracket B \rrbracket_* = \top^*$ or $\llbracket C \rrbracket_* = \top^*$. By Lemma 2.8 this implies that either $\llbracket B \rrbracket = (\top_1, \top_2)$ or $\llbracket C \rrbracket = (\top_1, \top_2)$, contradicting the assumptions. □

§3. The calculus D.LRC. In the present section, we introduce the multitype calculus D.LRC for the logic of resources and capabilities. As is typical of similar existing calculi, the language manipulated by this calculus is built up from structural and operational term constructors. In the tables below, each structural symbol in the upper rows corresponds to one or two logical (aka operational) symbols in the lower rows. The idea, which will be made precise in §4.1, is that each structural connective is interpreted as the corresponding logical connective on the left-hand (resp. right-hand) side (if it exists) when occurring in antecedent (resp. consequent) position.

As discussed in the previous section, the mathematical environment of heterogeneous LRC-algebras provides natural interpretations for all connectives of the basic language of LRC. In particular, on *perfect* heterogeneous LRC-algebras, these interpretations have the following extra properties: the interpretations of \diamond and \heartsuit are completely join-preserving, that of \triangleright is completely join-reversing in its first coordinate and order preserving in its second coordinate, and \triangleright is completely join-reversing in its first coordinate and completely meet-preserving in its second coordinate. This implies that, in each perfect heterogeneous LRC-algebra,

- \diamond and \heartsuit have right adjoints, denoted \blacksquare and \blacksuit , respectively;
- \triangleright has a Galois-adjoint \blacktriangleright in its first coordinate, and \blacktriangleright has a Galois-adjoint \blacktriangleright in its first coordinate and a left adjoint \blacktriangleleft in its second coordinate.

Hence, the following connectives have a natural interpretation on perfect heterogeneous LRC-algebras:

$$\blacksquare : Fm \rightarrow Fm, \tag{3.1}$$

$$\blacksuit : Fm \rightarrow Res, \tag{3.2}$$

$$\blacktriangleright : Fm \times Fm \rightarrow Res, \tag{3.3}$$

$$\blacktriangleright : Fm \times Res \rightarrow Res, \tag{3.4}$$

$$\blacktriangleleft : Res \times Fm \rightarrow Res. \tag{3.5}$$

- Structural and operational symbols for pure **Fm**-connectives:

Structural symbols	\mathbf{I}	$\mathbf{;}$	$\mathbf{>}$	$\mathbf{(<)}$				
Operational symbols	\top	\perp	\wedge	\vee	(\triangleright)	\rightarrow	(\blacktriangleleft)	(\blackleftarrow)

- Structural and operational symbols for pure **Res**-connectives:

Structural symbols	Φ	\odot	$\mathbf{,}$	$\mathbf{>}$	$\mathbf{<}$	$\mathbf{\sqsupset}$	$\mathbf{(\sqsubset)}$				
Operational symbols	1	0	\cdot	\sqcap	\sqcup	(\backslash)	$(/)$	(\cup)	(\cap)	(\cup)	(\cap)

- Structural and operational symbols for the modal operators:

Structural symbols	\circ	$\cdot \triangleright \cdot$	Φ	$\cdot \triangleright \cdot$
Operational symbols	\diamond		\heartsuit	\triangleright

- Structural and operational symbols for the adjoints and residuals of the modal operators:

Structural symbols	\bullet	$\cdot \blacktriangleright \cdot$	Φ	$\cdot \blacktriangleleft \cdot$	$\cdot \blacktriangleright \cdot$	
Operational symbols		(\blacksquare)		(\blacksuit)	(\blacktriangleleft)	(\blacktriangleright)

The display-type calculus **D.LRC** consists of the following display postulates, structural rules, and operational rules:

1. Identity and cut rules:

$$\begin{array}{c}
 p \vdash p \\
 \\
 \frac{(X \vdash Y)[A]^{succ} \quad A \vdash Z}{(X \vdash Y)[Z/A]^{succ}}
 \end{array}
 \qquad
 \begin{array}{c}
 a \vdash a \\
 \\
 \frac{\Gamma \vdash \alpha \quad \alpha \vdash \Delta}{\Gamma \vdash \Delta}.
 \end{array}$$

2. Display postulates for pure **Fm**-connectives:

$$\frac{X; Y \vdash Z}{Y \vdash X > Z} \quad \frac{Z \vdash X; Y}{X > Z \vdash Y} \quad \frac{X; Y \vdash Z}{X \vdash Z < Y} \quad \frac{Z \vdash X; Y}{Z < Y \vdash X}$$

3. Display postulates for pure **Res**-connectives:

$$\frac{\Gamma, \Delta \vdash \Sigma}{\Delta \vdash \Gamma \sqsupset \Sigma} \quad \frac{\Gamma, \Delta \vdash \Sigma}{\Gamma \vdash \Sigma \sqsubset \Delta} \quad \frac{\Gamma \vdash \Delta, \Sigma}{\Delta \sqsupset \Gamma \vdash \Sigma} \quad \frac{\Gamma \vdash \Delta, \Sigma}{\Gamma \sqsubset \Sigma \vdash \Delta}$$

$$\frac{\Gamma \odot \Delta \vdash \Sigma}{\Delta \vdash \Gamma \triangleright \Sigma} \quad \frac{\Gamma \odot \Delta \vdash \Sigma}{\Gamma \vdash \Sigma \triangleleft \Delta}$$

4. Display postulates for the modal operators:

$$\frac{\circ X \vdash Y}{X \vdash \bullet Y} \quad \frac{\circ \Gamma \vdash X}{\Gamma \vdash \bullet X} \quad \frac{X \vdash \Gamma \cdot \triangleright \cdot Y}{\Gamma \vdash X \cdot \blacktriangleright \cdot Y} \quad \frac{X \vdash \Gamma \cdot \blacktriangleright \cdot \Delta}{\Gamma \cdot \blacktriangleleft \cdot X \vdash \Delta} \quad \frac{X \vdash \Gamma \cdot \blacktriangleright \cdot \Delta}{\Gamma \vdash X \cdot \blacktriangleright \cdot \Delta}$$

5. Pure **Fm**-type structural rules:

$$I_L \frac{X \vdash Y}{I; X \vdash Y} \quad \frac{Y \vdash X}{Y \vdash X; I} I_R \quad E_L \frac{Y; X \vdash Z}{X; Y \vdash Z} \quad \frac{Z \vdash X; Y}{Z \vdash Y; X} E_R$$

$$W_L \frac{Y \vdash Z}{X; Y \vdash Z} \quad \frac{Z \vdash Y}{Z \vdash Y; X} W_R \quad C_L \frac{X; X \vdash Y}{X \vdash Y} \quad \frac{Y \vdash X; X}{Y \vdash X} C_R$$

$$A_L \frac{X; (Y; Z) \vdash W}{(X; Y); Z \vdash W} \quad \frac{W \vdash (Z; Y); X}{W \vdash Z; (Y; X)} A_R$$

6. Pure **Res**-type structural rules:

$$\Phi_{L1} \frac{\Gamma \odot \Phi \vdash \Delta}{\Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta, \Phi} \Phi_R \quad A_L \frac{\Gamma \odot (\Delta \odot \Sigma) \vdash \Pi}{(\Gamma \odot \Delta) \odot \Sigma \vdash \Pi} \quad W_\Phi \frac{\Phi \vdash \Delta}{\Gamma \vdash \Delta}$$

$$W_L \frac{\Gamma \vdash \Delta}{\Gamma, \Sigma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta, \Sigma} W_R \quad C_L \frac{\Gamma, \Gamma \vdash \Delta}{\Gamma \vdash \Delta} \quad \frac{\Gamma \vdash \Delta, \Delta}{\Gamma \vdash \Delta} C_R$$

$$E_L \frac{\Gamma, \Delta \vdash \Sigma}{\Delta, \Gamma \vdash \Sigma} \quad \frac{\Sigma \vdash \Gamma, \Delta}{\Sigma \vdash \Delta, \Gamma} E_R \quad A_L \frac{\Sigma, (\Delta, \Gamma) \vdash \Pi}{(\Sigma, \Delta), \Gamma \vdash \Pi} \quad \frac{\Pi \vdash \Sigma, (\Delta, \Gamma)}{\Pi \vdash (\Sigma, \Delta), \Gamma} A_R$$

$$\frac{\Pi \vdash (\Gamma \triangleright \Delta), (\Gamma \triangleright \Sigma)}{\Pi \vdash \Gamma \triangleright (\Delta, \Sigma)} dis.$$

7. Structural rules corresponding to the D-axioms:

$$\frac{X \vdash \bullet Y; \bullet Z}{X \vdash \bullet (Y; Z)} D1 \quad \frac{\Gamma \vdash \bullet X, \bullet Y}{\Gamma \vdash \bullet (X; Y)} D3 \quad \frac{X \vdash I}{X \vdash \bullet I} D2 \quad \frac{\Gamma \vdash \Phi}{\Gamma \vdash \bullet I} D4$$

8. Structural rules corresponding to the B-axioms:

$$\frac{\Gamma \vdash (Y \cdot \triangleright \cdot \Delta), (Z \cdot \triangleright \cdot \Delta)}{\Gamma \vdash (Y; Z) \cdot \triangleright \cdot \Delta} \text{B4} \quad \frac{\Gamma \vdash (Y \cdot \triangleright \cdot W), (Z \cdot \triangleright \cdot W)}{\Gamma \vdash (Y; Z) \cdot \triangleright \cdot W} \text{B1}$$

$$\text{B6} \frac{(\Gamma \cdot \blacktriangleleft \cdot X), (\Gamma \cdot \blacktriangleleft \cdot Y) \vdash \Delta}{\Gamma \cdot \blacktriangleleft \cdot (X; Y) \vdash \Delta} \quad \frac{X \vdash \Gamma \cdot \triangleright \cdot (\Delta \cdot \triangleright \cdot Y)}{X \vdash \Gamma \cdot \odot \cdot \Delta \cdot \triangleright \cdot Y} \text{B3} \quad \text{B7} \frac{\Phi \vdash \Delta}{\Gamma \cdot \blacktriangleleft \cdot I \vdash \Delta}$$

9. Structural rules corresponding to the BD-axioms:

$$\frac{X \vdash \Gamma \cdot \triangleright \cdot \bullet Y}{X \vdash \circ \Gamma > Y} \text{BD1} \quad \frac{X \vdash \Gamma \cdot \triangleright \cdot \bullet Y}{X \vdash \Gamma \cdot \triangleright \cdot Y} \text{BD2}$$

10. Introduction rules for pure Fm-connectives (in the presence of the exchange rules E_L and E_R , the structural connective $<$ and the corresponding operational connectives \prec and \leftarrow are redundant and they are omitted):

$$\perp_L \frac{}{\perp \vdash I} \quad \frac{X \vdash I}{X \vdash \perp} \perp_R \quad \top_L \frac{I \vdash X}{\top \vdash X} \quad \frac{}{I \vdash \top} \top_R$$

$$\wedge_L \frac{A; B \vdash X}{A \wedge B \vdash X} \quad \frac{X \vdash A \quad Y \vdash B}{X; Y \vdash A \wedge B} \wedge_R \quad \vee_L \frac{A \vdash X \quad B \vdash Y}{A \vee B \vdash X; Y} \quad \frac{X \vdash A; B}{X \vdash A \vee B} \vee_R$$

$$\rightarrow_L \frac{X \vdash A \quad B \vdash Y}{A \rightarrow B \vdash X > Y} \quad \frac{X \vdash A > B}{X \vdash A \rightarrow B} \rightarrow_R$$

11. Introduction rules for pure Res-connectives:

$$0_L \frac{}{0 \vdash \Phi} \quad \frac{\Gamma \vdash \Phi}{\Gamma \vdash 0} 0_R \quad 1_L \frac{\Phi \vdash \Gamma}{1 \vdash \Gamma} \quad \frac{}{\Phi \vdash 1} 1_R$$

$$\cdot_L \frac{\alpha \odot \beta \vdash \Gamma}{\alpha \cdot \beta \vdash \Gamma} \quad \frac{\Gamma \vdash \alpha \quad \Delta \vdash \beta}{\Gamma \odot \Delta \vdash \alpha \cdot \beta} \cdot_R \quad \sqcup_L \frac{\alpha \vdash \Gamma \quad \beta \vdash \Delta}{\alpha \sqcup \beta \vdash \Gamma, \Delta} \quad \frac{\Gamma \vdash \alpha, \beta}{\Gamma \vdash \alpha \sqcup \beta} \sqcup_R$$

12. Introduction rules for the modal operators:

$$\diamond_L \frac{\circ A \vdash X}{\diamond A \vdash X} \quad \frac{X \vdash A}{\circ X \vdash \diamond A} \diamond_R \quad \frac{\Gamma \vdash \alpha \quad A \vdash X}{\alpha \triangleright A \vdash \Gamma \cdot \triangleright \cdot X} \triangleright_L \quad \frac{X \vdash \alpha \cdot \triangleright \cdot A}{X \vdash \alpha \triangleright A} \triangleright_R$$

$$\diamond_L \frac{\circ \alpha \vdash X}{\diamond \alpha \vdash X} \quad \frac{\Gamma \vdash \alpha}{\circ \Gamma \vdash \diamond \alpha} \diamond_R \quad \frac{\Gamma \vdash \alpha \quad \beta \vdash \Delta}{\alpha \triangleright \alpha \vdash \Gamma \cdot \triangleright \cdot \Delta} \triangleright_L \quad \frac{\Gamma \vdash \alpha \cdot \triangleright \cdot \alpha}{\Gamma \vdash \alpha \triangleright \alpha} \triangleright_R$$

We conclude the present section by listing some observations about D.LRC. First, notice that, although very similar in spirit to a display calculus [3,72], D.LRC does not enjoy the display property, the reason being that a display rule for displaying substructures in the scope of the second coordinate of $\cdot \triangleright \cdot$ occurring in consequent position would not be sound. This is the reason why a more general form of cut rule, sometimes referred to as *surgical cut*, has been included than the standard one in display calculi where both cut formulas occur in display. However, as discussed in [33], calculi without display property can still verify the assumptions of some Belnap-style cut elimination metatheorem. In §4.3, we will verify that this is the case of D.LRC. Second, as usual, the version of D.LRC on a classical propositional base can be obtained by adding, e.g., the following *Grishin rules*:

$$\frac{X > (Y; Z) \vdash W}{(X > Y); Z \vdash W} \quad \frac{X \vdash Y > (Z; W)}{X \vdash (Y > Z); W}$$

Third, the rule W_Φ encodes (and is used to derive) $\alpha \cdot \beta \vdash \alpha$, $\alpha \cdot \beta \vdash \beta$, $\alpha \vdash 1$, B2, and B5.

§4. Basic properties of D.LRC. In the present section, we verify that the calculus D.LRC is sound w.r.t. the semantics of perfect heterogeneous LRC-algebras (cf. Definition 2.1), is syntactically complete w.r.t. the Hilbert calculus for LRC introduced in §2.1, enjoys cut-elimination and subformula property, and conservatively extends the Hilbert calculus of §2.1.

4.1. Soundness. In the present subsection, we outline the verification of the soundness of the rules of D.LRC w.r.t. the semantics of perfect heterogeneous LRC-algebras (cf. Definition 2.1). The first step consists in interpreting structural symbols as logical symbols according to their (precedent or consequent) position,¹¹ as indicated in the synoptic tables at the beginning of §3. This makes it possible to interpret sequents as inequalities, and rules as quasi-inequalities. For example, the rules on the left-hand side below are interpreted as the quasi-inequalities on the right-hand side:

$$\frac{X \vdash \Gamma \cdot \triangleright \cdot \bullet Y}{X \vdash \circ \Gamma > Y} \text{BD1} \rightsquigarrow \forall \gamma \forall x \forall y [x \leq \gamma \triangleright \blacksquare y \Rightarrow x \leq \diamond \gamma \rightarrow y]$$

$$\frac{X \vdash \Gamma \cdot \triangleright \bullet Y}{X \vdash \Gamma \cdot \triangleright \cdot Y} \text{BD2} \rightsquigarrow \forall x \forall \gamma \forall y [x \leq \gamma \triangleright \blacksquare \blacksquare y \Rightarrow x \leq \gamma \triangleright y].$$

The verification that the rules of D.LRC are sound on perfect LRC-algebras then consists in verifying the validity of their corresponding quasi-inequalities in perfect LRC-algebras. The validity of these quasi-inequalities follows straightforwardly from two observations. The first observation is that the quasi-inequality corresponding to each rule is obtained by running the algorithm ALBA on the axiom of the Hilbert-style presentation of §2.1 bearing the same name as the rule. Below we perform the ALBA reduction on the axiom BD1:

$$\begin{aligned} & \forall a \forall p [\diamond a \wedge a \triangleright p \leq \diamond p] \\ \text{iff } & \forall a \forall p \forall \gamma \forall x \forall y [(\gamma \leq a \ \& \ x \leq a \triangleright p \ \& \ \diamond p \leq y) \Rightarrow \diamond \gamma \wedge x \leq y] \\ \text{iff } & \forall a \forall p \forall \gamma \forall x \forall y [(\gamma \leq a \ \& \ x \leq a \triangleright p \ \& \ p \leq \blacksquare y) \Rightarrow \diamond \gamma \wedge x \leq y] \\ \text{iff } & \forall \gamma \forall x \forall y [x \leq \gamma \triangleright \blacksquare y \Rightarrow \diamond \gamma \wedge x \leq y] \\ \text{iff } & \forall \gamma \forall x \forall y [x \leq \gamma \triangleright \blacksquare y \Rightarrow x \leq \diamond \gamma \rightarrow y]. \end{aligned}$$

It can be readily checked that the ALBA manipulation rules applied in the computation above (adjunction rules and Ackermann rules) are sound on perfect LRC-algebras. As discussed in [44], the soundness of these rules only depends on the order-theoretic properties of the interpretation of the logical connectives and their adjoints and residuals. The fact that some of these maps are not internal operations but have different domains and codomains does not make any substantial difference. A more substantial difference with the setting of [44] might be in principle the fact that the connective \triangleright is only monotone—rather than normal—in its second coordinate. However, notice that each manipulation in the chain of equivalences above involving that coordinate is an application of the Ackermann rule of ALBA, which relies on no more than monotonicity. The second observation is that the axioms of the Hilbert-style presentation of §2.1 are valid by definition on perfect

¹¹ For any (formula or resource) sequent $x \vdash y$ in the language of D.LRC, we define the signed generation trees $+x$ and $-y$ by labelling the root of the generation tree of x (resp. y) with the sign $+$ (resp. $-$), and then propagating the sign to all nodes according to the polarity of the coordinate of the connective assigned to each node. Positive (resp. negative) coordinates propagate the same (resp. opposite) sign to the corresponding child node. The only negative coordinates are the first coordinates of $>$, $\triangleright \cdot$ and $\bullet \triangleright \cdot$. Then, a substructure z in $x \vdash y$ is in *precedent* (resp. *consequent*) *position* if the sign of its root node as a subtree of $+x$ or $-y$ is $+$ (resp. $-$).

LRC-algebras. We conclude the present subsection reporting the ALBA-reduction of (the condition expressing the validity of) axiom BD2.

$$\begin{aligned} & \forall \alpha \forall \beta [\alpha \triangleright \beta \leq \alpha \triangleright \diamond \beta] \\ \text{iff } & \forall \alpha \forall \beta \forall x \forall \gamma \forall y [(x \leq \alpha \triangleright \beta \ \& \ \gamma \leq \alpha \ \& \ \diamond \beta \leq y) \Rightarrow x \leq \gamma \triangleright y] \\ \text{iff } & \forall \alpha \forall \beta \forall x \forall \gamma \forall y [(x \leq \alpha \triangleright \beta \ \& \ \gamma \leq \alpha \ \& \ \beta \leq \blacksquare y) \Rightarrow x \leq \gamma \triangleright y] \\ \text{iff } & \forall x \forall \gamma \forall y [x \leq \gamma \triangleright \blacksquare y \Rightarrow x \leq \gamma \triangleright y]. \end{aligned}$$

4.2. Completeness. In the present subsection, we show that the axioms of the Hilbert-style calculus H.LRC introduced in §2.1 are derivable sequents of D.LRC, and that the rules of H.LRC are derivable rules of D.LRC. Since H.LRC is complete w.r.t. the semantics of perfect heterogeneous LRC-algebras (cf. Definition 2.1), we obtain as a corollary that D.LRC is also complete w.r.t. the semantics of perfect heterogeneous LRC-algebras. The derivations of the axioms R1-R3 of H.LRC are standard and we omit them.

R4. $\alpha \cdot (\beta \sqcup \gamma) \leftrightarrow (\alpha \cdot \beta) \sqcup (\alpha \cdot \gamma)$

$$\frac{\frac{\frac{\alpha \vdash \alpha}{\alpha \odot \beta \vdash \alpha \cdot \beta} \quad \frac{\beta \vdash \beta}{\beta \vdash \alpha \triangleright \alpha \cdot \beta}}{\beta \vdash \alpha \triangleright \alpha \cdot \beta} \quad \frac{\frac{\alpha \vdash \alpha}{\alpha \odot \gamma \vdash \alpha \cdot \gamma} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \alpha \triangleright \alpha \cdot \gamma}}{\gamma \vdash \alpha \triangleright \alpha \cdot \gamma}}{\beta \sqcup \gamma \vdash (\alpha \triangleright \alpha \cdot \beta), (\alpha \triangleright \alpha \cdot \gamma)} \text{ dis} \quad \frac{\frac{\beta \vdash \beta}{\beta \vdash \beta, \gamma} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \beta, \gamma}}{\beta \vdash \beta \sqcup \gamma} \quad \frac{\frac{\alpha \vdash \alpha}{\alpha \odot \gamma \vdash \alpha \cdot (\beta \sqcup \gamma)} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \beta \sqcup \gamma}}{\alpha \odot \gamma \vdash \alpha \cdot (\beta \sqcup \gamma)} \\ \frac{\frac{\beta \sqcup \gamma \vdash \alpha \triangleright (\alpha \cdot \beta), (\alpha \cdot \gamma)}{\beta \sqcup \gamma \vdash \alpha \triangleright (\alpha \cdot \beta, \alpha \cdot \gamma)} \quad \frac{\alpha \odot \beta \vdash \alpha \cdot (\beta \sqcup \gamma)}{\alpha \cdot \beta \vdash \alpha \cdot (\beta \sqcup \gamma)}}{\frac{\alpha \odot (\beta \sqcup \gamma) \vdash \alpha \cdot \beta, \alpha \cdot \gamma}{\alpha \cdot (\beta \sqcup \gamma) \vdash \alpha \cdot \beta, \alpha \cdot \gamma}} \quad \frac{\frac{\alpha \odot \gamma \vdash \alpha \cdot (\beta \sqcup \gamma)}{\alpha \cdot \gamma \vdash \alpha \cdot (\beta \sqcup \gamma)} \quad \frac{\alpha \odot \beta \vdash \alpha \cdot (\beta \sqcup \gamma)}{\alpha \cdot \beta \vdash \alpha \cdot (\beta \sqcup \gamma)}}{\frac{(\alpha \cdot \beta) \sqcup (\alpha \cdot \gamma) \vdash \alpha \cdot (\beta \sqcup \gamma), \alpha \cdot (\beta \sqcup \gamma)}{(\alpha \cdot \beta) \sqcup (\alpha \cdot \gamma) \vdash \alpha \cdot (\beta \sqcup \gamma)}}$$

The proof of $(\beta \sqcup \gamma) \cdot \alpha \leftrightarrow (\beta \cdot \alpha) \sqcup (\gamma \cdot \alpha)$ is analogous and we omit it.

D1. $\diamond(A \vee B) \leftrightarrow \diamond A \vee \diamond B$

$$\frac{\frac{\frac{A \vdash A}{\circ A \vdash \diamond A} \quad \frac{B \vdash B}{\circ B \vdash \diamond B}}{A \vdash \bullet \diamond A \quad B \vdash \bullet \diamond B}}{\frac{A \vee B \vdash \bullet \diamond A; \bullet \diamond B}{A \vee B \vdash \bullet (\diamond A; \diamond B)}} \text{ D1} \quad \frac{\frac{A \vdash A}{A \vdash A; B} \quad \frac{B \vdash B}{B \vdash A; B}}{A \vdash A \vee B \quad B \vdash A \vee B}}{\frac{\circ A \vdash \diamond(A \vee B) \quad \circ B \vdash \diamond(A \vee B)}{\diamond A \vdash \diamond(A \vee B) \quad \diamond B \vdash \diamond(A \vee B)}} \\ \frac{\frac{\circ A \vee B \vdash \diamond A; \diamond B}{\diamond(A \vee B) \vdash \diamond A; \diamond B}}{\diamond(A \vee B) \vdash \diamond A \vee \diamond B} \quad \frac{\frac{\diamond A \vee \diamond B \vdash \diamond(A \vee B); \diamond(A \vee B)}{\diamond A \vee \diamond B \vdash \diamond(A \vee B)}}$$

D3. $\diamond(\alpha \sqcup \beta) \leftrightarrow \diamond \alpha \vee \diamond \beta$

$$\frac{\frac{\frac{\alpha \vdash \alpha}{\circ \alpha \vdash \diamond \alpha} \quad \frac{\beta \vdash \beta}{\circ \beta \vdash \diamond \beta}}{\alpha \vdash \bullet \diamond \alpha \quad \beta \vdash \bullet \diamond \beta}}{\frac{\alpha \sqcup \beta \vdash \bullet \diamond \alpha, \bullet \diamond \beta}{\alpha \sqcup \beta \vdash \bullet (\diamond \alpha; \diamond \beta)}} \text{ D3} \quad \frac{\frac{\alpha \vdash \alpha}{\alpha \vdash \alpha, \beta} \quad \frac{\beta \vdash \beta}{\beta \vdash \alpha, \beta}}{\alpha \vdash \alpha \sqcup \beta \quad \beta \vdash \alpha \sqcup \beta}}{\frac{\circ \alpha \vdash \diamond(\alpha \sqcup \beta) \quad \circ \beta \vdash \diamond(\alpha \sqcup \beta)}{\diamond \alpha \vdash \diamond(\alpha \sqcup \beta) \quad \diamond \beta \vdash \diamond(\alpha \sqcup \beta)}} \\ \frac{\frac{\circ \alpha \sqcup \beta \vdash \diamond \alpha; \diamond \beta}{\diamond(\alpha \sqcup \beta) \vdash \diamond \alpha; \diamond \beta}}{\diamond(\alpha \sqcup \beta) \vdash \diamond \alpha \vee \diamond \beta} \quad \frac{\frac{\diamond \alpha \sqcup \diamond \beta \vdash \diamond(\alpha \sqcup \beta), \diamond(\alpha \sqcup \beta)}{\diamond \alpha \sqcup \diamond \beta \vdash \diamond(\alpha \sqcup \beta)}}$$

D2. $\diamond \perp \leftrightarrow \perp$

$$\frac{\frac{\frac{\perp \vdash \perp}{\perp \vdash \bullet \perp}}{\circ \perp \vdash \perp} \quad \frac{\perp \vdash \perp}{\perp \vdash \perp}}{\diamond \perp \vdash \perp} \text{ D2} \quad \frac{\frac{\perp \vdash \perp}{\perp \vdash \diamond \perp; \perp}}{\perp \vdash \diamond \perp}}$$

D4. $\diamond 0 \leftrightarrow \perp$

$$\frac{\frac{\frac{0 \vdash \Phi}{0 \vdash \bullet I}}{\bullet 0 \vdash I}}{\bullet 0 \vdash \perp}}{\diamond 0 \vdash \perp} \text{D4} \quad \frac{0 \vdash \Phi}{0 \vdash \diamond 0, \Phi}}{0 \vdash \diamond 0}$$

B1. $\alpha \sqcup \beta \triangleright A \leftrightarrow (\alpha \triangleright A) \wedge (\beta \triangleright A)$

$$\frac{\frac{\frac{\alpha \vdash \alpha}{\alpha \vdash \alpha, \beta}}{\alpha \vdash \alpha \sqcup \beta} \quad \frac{A \vdash A}{A \vdash A}}{\alpha \sqcup \beta \triangleright A \vdash \alpha \cdot \triangleright \cdot A} \quad \frac{\frac{\beta \vdash \beta}{\beta \vdash \alpha, \beta}}{\beta \vdash \alpha \sqcup \beta} \quad \frac{A \vdash A}{A \vdash A}}{\alpha \sqcup \beta \triangleright A \vdash \beta \cdot \triangleright \cdot A} \quad \frac{\frac{\alpha \vdash \alpha}{\alpha \triangleright A \vdash \alpha \cdot \triangleright \cdot A} \quad \frac{A \vdash A}{A \vdash A}}{\alpha \vdash \alpha \triangleright A \cdot \blacktriangleright \cdot A} \quad \frac{\frac{\beta \vdash \beta}{\beta \triangleright A \vdash \beta \cdot \triangleright \cdot A} \quad \frac{A \vdash A}{A \vdash A}}{\beta \vdash \beta \triangleright A \cdot \blacktriangleright \cdot A}}{\frac{\alpha \sqcup \beta \vdash (\alpha \triangleright A \cdot \blacktriangleright \cdot A), (\beta \triangleright A \cdot \blacktriangleright \cdot A)}{\alpha \sqcup \beta \vdash (\alpha \triangleright A; \beta \triangleright A) \cdot \blacktriangleright \cdot A} \text{B1}}{\frac{\alpha \triangleright A; \beta \triangleright A \vdash \alpha \sqcup \beta \cdot \triangleright \cdot A}{\alpha \triangleright A; \beta \triangleright A \vdash \alpha \sqcup \beta \triangleright A}}{(\alpha \triangleright A) \wedge (\beta \triangleright A) \vdash \alpha \sqcup \beta \triangleright A}}$$

B4. $\alpha \sqcup \beta \triangleright \gamma \leftrightarrow (\alpha \triangleright \gamma) \wedge (\beta \triangleright \gamma)$

$$\frac{\frac{\frac{\alpha \vdash \alpha}{\alpha \vdash \alpha, \beta}}{\alpha \vdash \alpha \sqcup \beta} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \gamma}}{\alpha \sqcup \beta \triangleright \gamma \vdash \alpha \cdot \triangleright \cdot \gamma} \quad \frac{\frac{\beta \vdash \beta}{\beta \vdash \alpha, \beta}}{\beta \vdash \alpha \sqcup \beta} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \gamma}}{\alpha \sqcup \beta \triangleright \gamma \vdash \beta \cdot \triangleright \cdot \gamma} \quad \frac{\frac{\alpha \vdash \alpha}{\alpha \triangleright \gamma \vdash \alpha \cdot \triangleright \cdot \gamma} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \gamma}}{\alpha \vdash \alpha \triangleright \gamma \cdot \blacktriangleright \cdot \gamma} \quad \frac{\frac{\beta \vdash \beta}{\beta \triangleright \gamma \vdash \beta \cdot \triangleright \cdot \gamma} \quad \frac{\gamma \vdash \gamma}{\gamma \vdash \gamma}}{\beta \vdash \beta \triangleright \gamma \cdot \blacktriangleright \cdot \gamma}}{\frac{\alpha \sqcup \beta \vdash (\alpha \triangleright \gamma \cdot \blacktriangleright \cdot \gamma), (\beta \triangleright \gamma \cdot \blacktriangleright \cdot \gamma)}{\alpha \sqcup \beta \vdash (\alpha \triangleright \gamma; \beta \triangleright \gamma) \cdot \blacktriangleright \cdot \gamma} \text{B4}}{\frac{\alpha \triangleright \gamma; \beta \triangleright \gamma \vdash \alpha \sqcup \beta \cdot \triangleright \cdot \gamma}{\alpha \triangleright \gamma; \beta \triangleright \gamma \vdash \alpha \sqcup \beta \triangleright \gamma}}{(\alpha \triangleright \gamma) \wedge (\beta \triangleright \gamma) \vdash \alpha \sqcup \beta \triangleright \gamma}}$$

B2. $0 \triangleright A$

$$\frac{\frac{0 \vdash \Phi}{0 \vdash I \cdot \blacktriangleright \cdot A, \Phi}}{0 \vdash I \cdot \blacktriangleright \cdot A}}{I \vdash 0 \cdot \triangleright \cdot A}}{I \vdash 0 \triangleright A}$$

B5. $0 \triangleright \alpha$

$$\frac{\frac{0 \vdash \Phi}{0 \vdash I \cdot \blacktriangleright \cdot \alpha, \Phi}}{0 \vdash I \cdot \blacktriangleright \cdot \alpha}}{I \vdash 0 \cdot \triangleright \cdot \alpha}}{I \vdash 0 \triangleright \alpha}$$

B3. $\alpha \triangleright (\beta \triangleright A) \rightarrow (\alpha \cdot \beta \triangleright A)$

$$\frac{\frac{\frac{\beta \vdash \beta}{\beta \triangleright A \vdash \beta \cdot \triangleright \cdot A} \quad \frac{A \vdash A}{A \vdash A}}{\alpha \triangleright (\beta \triangleright A) \vdash \alpha \cdot \triangleright \cdot (\beta \cdot \triangleright \cdot A)} \text{B3}}{\frac{\alpha \triangleright (\beta \triangleright A) \vdash (\alpha \odot \beta) \cdot \triangleright \cdot A}}{\frac{\alpha \odot \beta \vdash (\alpha \triangleright (\beta \triangleright A)) \cdot \blacktriangleright \cdot A}{\alpha \cdot \beta \vdash (\alpha \triangleright (\beta \triangleright A)) \cdot \blacktriangleright \cdot A}}{\frac{\alpha \triangleright (\beta \triangleright A) \vdash (\alpha \cdot \beta) \cdot \triangleright \cdot A}{\alpha \triangleright (\beta \triangleright A) \vdash (\alpha \cdot \beta) \triangleright A}}$$

B6. $\alpha \triangleright (\beta \sqcap \gamma) \leftrightarrow \alpha \triangleright \beta \wedge \alpha \triangleright \gamma$

$$\begin{array}{c}
 \frac{\beta \vdash \beta}{\beta, \gamma \vdash \beta} \\
 \frac{\alpha \vdash \alpha \quad \beta \sqcap \gamma \vdash \beta}{\alpha \triangleright \beta \sqcap \gamma \vdash \alpha \cdot \triangleright \cdot \beta} \\
 \frac{\alpha \triangleright \beta \sqcap \gamma \vdash \alpha \triangleright \beta}{\alpha \triangleright \beta \sqcap \gamma \vdash \alpha \triangleright \beta} \\
 \frac{\alpha \triangleright \beta \sqcap \gamma; \alpha \triangleright \beta \sqcap \gamma \vdash (\alpha \triangleright \beta) \wedge (\alpha \triangleright \gamma)}{\alpha \triangleright \beta \sqcap \gamma \vdash (\alpha \triangleright \beta) \wedge (\alpha \triangleright \gamma)} \\
 \\
 \frac{\gamma \vdash \gamma}{\gamma, \beta \vdash \gamma} \\
 \frac{\beta, \gamma \vdash \gamma}{\beta \sqcap \gamma \vdash \gamma} \\
 \frac{\alpha \vdash \alpha \quad \beta \sqcap \gamma \vdash \gamma}{\alpha \triangleright \beta \sqcap \gamma \vdash \alpha \cdot \triangleright \cdot \gamma} \\
 \frac{\alpha \triangleright \beta \sqcap \gamma \vdash \alpha \triangleright \beta \quad \alpha \triangleright \beta \sqcap \gamma \vdash \alpha \triangleright \gamma}{\alpha \triangleright \beta \sqcap \gamma \vdash \alpha \triangleright \beta} \\
 \\
 \frac{\alpha \vdash \alpha \quad \beta \vdash \beta}{\alpha \triangleright \beta \vdash \alpha \cdot \triangleright \cdot \beta} \\
 \frac{\alpha \vdash \alpha \quad \gamma \vdash \gamma}{\alpha \triangleright \gamma \vdash \alpha \cdot \triangleright \cdot \gamma} \\
 \frac{\alpha \cdot \blacktriangleleft \cdot \alpha \triangleright \beta \vdash \beta \quad \alpha \cdot \blacktriangleleft \cdot \alpha \triangleright \gamma \vdash \gamma}{(\alpha \cdot \blacktriangleleft \cdot \alpha \triangleright \beta), (\alpha \cdot \blacktriangleleft \cdot \alpha \triangleright \gamma) \vdash \beta \sqcap \gamma} \\
 \frac{(\alpha \cdot \blacktriangleleft \cdot \alpha \triangleright \beta), (\alpha \cdot \blacktriangleleft \cdot \alpha \triangleright \gamma) \vdash \beta \sqcap \gamma}{\alpha \cdot \blacktriangleleft \cdot (\alpha \triangleright \beta; \alpha \triangleright \gamma) \vdash \beta \sqcap \gamma} \\
 \frac{\alpha \triangleright \beta; \alpha \triangleright \gamma \vdash \alpha \cdot \triangleright \cdot \beta \sqcap \gamma}{\alpha \triangleright \beta; \alpha \triangleright \gamma \vdash \alpha \triangleright \beta \sqcap \gamma} \\
 \frac{\alpha \triangleright \beta; \alpha \triangleright \gamma \vdash \alpha \triangleright \beta \sqcap \gamma}{(\alpha \triangleright \beta) \wedge (\alpha \triangleright \gamma) \vdash \alpha \triangleright \beta \sqcap \gamma}
 \end{array}$$

B7. $\alpha \triangleright 1$

$$\frac{\Phi \vdash 1}{\alpha \cdot \blacktriangleleft \cdot I, \Phi \vdash 1} \\
 \frac{\alpha \cdot \blacktriangleleft \cdot I \vdash 1}{I \vdash \alpha \cdot \triangleright \cdot 1} \\
 I \vdash \alpha \triangleright 1$$

BD1. $\diamond \alpha \wedge \alpha \triangleright A \rightarrow \diamond A$

$$\frac{A \vdash A}{\circ A \vdash \diamond A} \\
 \frac{\alpha \vdash \alpha \quad A \vdash \bullet \diamond A}{\alpha \triangleright A \vdash \alpha \cdot \triangleright \cdot \bullet \diamond A} \\
 \frac{\alpha \triangleright A \vdash \alpha \cdot \triangleright \cdot \bullet \diamond A}{\alpha \triangleright A \vdash \phi \alpha > \diamond A} \text{BD1} \\
 \frac{\phi \alpha; \alpha \triangleright A \vdash \diamond A}{\phi \alpha \vdash \diamond A < \alpha \triangleright A} \\
 \frac{\phi \alpha \vdash \diamond A < \alpha \triangleright A}{\diamond \alpha; \alpha \triangleright A \vdash \diamond A} \\
 \diamond \alpha \wedge \alpha \triangleright A \vdash \diamond A$$

BD2. $\alpha \triangleright \beta \rightarrow \alpha \triangleright \diamond \beta$

$$\frac{\beta \vdash \beta}{\phi \beta \vdash \diamond \beta} \\
 \frac{\alpha \vdash \alpha \quad \phi \beta \vdash \diamond \beta}{\beta \vdash \bullet \diamond \beta} \\
 \frac{\alpha \triangleright \beta \vdash \alpha \cdot \triangleright \cdot \bullet \diamond \beta}{\alpha \triangleright \beta \vdash \alpha \cdot \triangleright \cdot \diamond \beta} \text{BD2} \\
 \alpha \triangleright \beta \vdash \alpha \triangleright \diamond \beta$$

The rules of H.LRC immediately follow from applications of the introduction rules of the corresponding logical connectives in the usual way and we omit their derivations.

4.3. Cut-elimination and subformula property. In the present subsection, we sketch the verification that the D.LRC is a proper multitype calculus (cf. §7). By Theorem 7.3, this is enough to establish that the calculus enjoys cut elimination and subformula property. With the exception of C'_8 , all conditions are straightforwardly verified by inspecting the rules, and this verification is left to the reader.

As to the verification of condition C'_8 , the only interesting case is the one in which the cut formula is of the form $\alpha \triangleright A$, since the connective \triangleright is monotone rather than normal in its second coordinate, which is the reason why not even a weak form of display property holds for D.LRC. This case is treated below. Notice that, since all principal formulas are in display, no surgical cuts need to be eliminated in the principal stage.

$$\frac{\frac{\vdots \pi_1}{X \vdash \alpha \cdot \triangleright \cdot A} \quad \frac{\vdots \pi_2}{\Gamma \vdash \alpha} \quad \frac{\vdots \pi_3}{A \vdash Y}}{X \vdash \alpha \triangleright A} \quad \frac{\vdots \pi_2}{\Gamma \vdash \alpha} \quad \frac{\vdots \pi_3}{A \vdash Y}}{\alpha \triangleright A \vdash \Gamma \cdot \triangleright \cdot Y} \rightsquigarrow \frac{\frac{\frac{\vdots \pi_1}{X \vdash \alpha \cdot \triangleright \cdot A} \quad \frac{\vdots \pi_2}{\Gamma \vdash \alpha}}{\alpha \vdash X \cdot \triangleright \cdot A} \quad \frac{\vdots \pi_3}{A \vdash Y}}{\Gamma \vdash X \cdot \triangleright \cdot A} \quad \frac{\vdots \pi_3}{A \vdash Y}}{X \vdash \Gamma \cdot \triangleright \cdot Y}$$

4.4. Semantic conservativity. To argue that the calculus D.LRC adequately captures LRC, we follow the standard proof strategy discussed in [44]. Recall that \vdash_{LRC} denotes the syntactic consequence relation arising from the Hilbert system for LRC introduced in §2.1. We need to show that, for all LRC-formulas A and B , if $A \vdash B$ is a provable sequent in the calculus D.LRC, then $A \vdash_{\text{LRC}} B$. This fact can be verified using the following standard argument and facts: (a) the rules of D.LRC are sound w.r.t. perfect heterogeneous LRC-algebras (cf. §4.1), and (b) LRC is strongly complete w.r.t. perfect heterogeneous LRC-algebras (cf. Corollary 2.6). Then, let A, B be LRC-formulas such that $A \vdash B$ is a derivable sequent in D.LRC. By (a), this implies that $A \models_{\text{LRC}} B$, which implies, by (b), that $A \vdash_{\text{LRC}} B$, as required.

§5. Case studies. In this section, we present a number of case studies, with the purpose of highlighting various aspects of the basic framework and also various ways in which it can be adapted to different settings. The most common adaptations performed in the case studies below consist in adding analytic structural rules to the basic calculus. Interestingly, the resulting calculi still enjoy the same package of basic properties (soundness, completeness, cut-elimination, subformula property, and conservativity) which hold of D.LRC as an immediate consequence of general results. Indeed, it can be readily verified that the axioms corresponding to each of the rules introduced below are analytic inductive (cf. [44, Definition 55]), and hence are canonical (cf. [44, Theorem 19]). Therefore, the axiomatic extensions of LRC corresponding to these axioms is *sound* and *complete* w.r.t. the corresponding subclass of LRC-models. *Conservativity* can be argued by repeating verbatim the same argument given in §4.4 which uses the soundness of the augmented calculus w.r.t. the corresponding class of perfect LRC-models, and the completeness of the Hilbert-style presentation of the axiomatic extension which holds because the additional axioms are canonical. Finally, *cut-elimination* and *subformula property* follow from the general cut-elimination metatheorem.

In what follows, we will sometimes abuse terminology and speak of a formula A being derived from certain assumptions $A_1; \dots; A_n$ meaning that the sequent $A_1; \dots; A_n \vdash A$ is derivable in the calculus.

5.1. Pooling capabilities (correcting a homework assignment). Two teaching assistants, Carl (c) and Dan (d), are assigned the task of grading a set of homework assignments consisting of two exercises, a model-theoretic one (M) and a proof-theoretic one (P). Carl is only capable of correcting exercise P , while Dan is only capable of correcting exercise M . None of the two teaching assistants can individually complete the task they have been assigned. However, they can if they *pool* their capabilities. One way in which they can complete the task is by implementing the following plan: they split the set of homework assignments into two sets α and β . Initially, Carl grades the solutions to exercise P in α and Dan those of M in β . Then they switch sets and each of them grades the solutions to the same exercise in the other set.

To capture this case study in (a multi-agent version of) D.LRC, we introduce atomic propositions such as P_α (resp. M_β), the intended meaning of which is that all solutions to exercise P (resp. M) in α (resp. β) have been graded. We also treat α and β as resources. The following table contains formulas expressing the assumptions about agents' capabilities, the initial state of affairs (which resources are initially in possession of which agent), and the plan of switching after completing the correction of one exercise in a given set:

Capabilities	initial state	planning
$\alpha \triangleright_c P_\alpha \quad \beta \triangleright_c P_\beta$	$\Diamond_c \alpha$	$M_\beta \rightarrow \Diamond_c \beta$
$\alpha \triangleright_d M_\alpha \quad \beta \triangleright_d M_\beta$	$\Diamond_d \beta$	$P_\alpha \rightarrow \Diamond_d \alpha$

In the present setting we also assume that, whenever an agent is *able* to bring about a certain state of affairs, the agent will. Formally, this corresponds to the validity of the axioms $\Diamond_i A \rightarrow A$ for every agent i and formula A . This axiom does not follow from the logic H.LRC, and in many settings it would not be sound. However, for the sake of the present case study, we will assume that this axiom holds. In fact, this axiom corresponds to the following rules 'Ex_i' ('Ex' stands for Execution), for each $i \in \{c, d\}$:

$$\text{Ex}_i \frac{X \vdash Y}{\circ_i X \vdash Y}.$$

Notice that these rules are analytic (cf. §7). Hence, by Theorem 7.3, when adding these rules to the basic calculus D.LRC, the resulting calculus (which we refer to as D.LRC + Ex) enjoys cut elimination and subformula property.

We aim at deriving the formula $(P_\alpha \wedge M_\beta) \wedge (P_\beta \wedge M_\alpha)$ from the assumptions above in the calculus D.LRC + Ex. This will provide the formal verification that executing the plan yields the completion of the task. Let us start by considering the following derivations:

π_1 $\text{Cut} \frac{\begin{array}{c} \text{: proof for} \\ \text{: BD1} \\ \hline \Phi_c \alpha; \alpha \triangleright_c P_\alpha \vdash \circ_c P_\alpha \end{array} \quad \text{Ex} \frac{P_\alpha \vdash P_\alpha}{\circ_c P_\alpha \vdash P_\alpha}}{\Phi_c \alpha; \alpha \triangleright_c P_\alpha \vdash P_\alpha}$	π_2 $\text{Cut} \frac{\begin{array}{c} \text{: proof for} \\ \text{: BD1} \\ \hline \Phi_d \beta; \alpha \triangleright_d M_\beta \vdash \circ_d M_\beta \end{array} \quad \text{Ex} \frac{M_\beta \vdash M_\beta}{\circ_d M_\beta \vdash M_\beta}}{\Phi_d \beta; \beta \triangleright_d M_\beta \vdash M_\beta}$
π_3 $\text{Cut} \frac{\begin{array}{c} \text{: proof for} \\ \text{: BD1} \\ \hline \Phi_c \beta; \beta \triangleright_c P_\beta \vdash \circ_c P_\beta \end{array} \quad \text{Ex} \frac{P_\beta \vdash P_\beta}{\circ_c P_\beta \vdash P_\beta}}{\Phi_c \beta; \beta \triangleright_c P_\beta \vdash P_\beta}$	π_4 $\text{Cut} \frac{\begin{array}{c} \text{: proof for} \\ \text{: BD1} \\ \hline \Phi_d \alpha; \alpha \triangleright_d M_\alpha \vdash \circ_d M_\alpha \end{array} \quad \text{Ex} \frac{M_\alpha \vdash M_\alpha}{\circ_d M_\alpha \vdash M_\alpha}}{\Phi_d \alpha; \alpha \triangleright_d M_\alpha \vdash M_\alpha}$

These derivations follow one and the same pattern, and each derives one piece of the desired conclusion. Hence, one would want to suitably prolong these derivations by applying \wedge_R to reach the conclusion. However, while the conclusions of π_1 and π_2 contain only formulas which are assumptions in our case study as reported in the table above, the formulas $\Diamond_c \beta$ and $\Diamond_d \alpha$, occurring in the conclusions of π_3 and π_4 , respectively, are not assumptions. However, they are provable from the assumptions. Indeed, they encode states of affairs which hold after c and d have switched the sets α and β .

Notice that the following sequents are provable (their derivations are straightforward and are omitted):

$$M_\beta; M_\beta \rightarrow \Diamond_c \beta \vdash \Diamond_c \beta \quad P_\alpha; P_\alpha \rightarrow \Diamond_d \alpha \vdash \Diamond_d \alpha.$$

These sequents say that the formulas $\Diamond_c \beta$ and $\Diamond_d \alpha$ are provable from the 'planning assumptions' (cf. table above) using the formulas M_β and P_α which have been derived purely

from the assumptions by π_1 and π_2 . Hence, the atoms P_β and M_α can be derived from the original assumptions via cut. Then, applying \wedge_R and possibly contraction, one can derive the desired sequent.

5.2. Conjoining capabilities (the wisdom of the crow). A BBC documentary program shows a problem-solving test conducted on a crow. In the present subsection we formalize an adapted version of this test. There is food (ϕ) positioned deep in a narrow box, out of the reach of the crow’s beak. There is a short stick (σ) directly available to the crow, two stones (ρ_1, ρ_2) each inside a cage, and a long stick (λ) inside a transparent box which releases the stick if enough weight (that of two stones or more) lays inside the box. The stick σ is too short for the crow to reach the food using it. However, previous tests have shown that the crow is capable of performing the following individual steps: (a) reaching the food using the long stick; (b) retrieving the stones from the cages using the short stick; (c) retrieving the long stick by dropping stones into a slot in the box. The crow succeeded in executing these individual steps in the right order and got to the food.

An interesting feature of this case study is the interplay of different kinds of resources. Specifically, σ is a *reusable* resource (indeed, the crow uses the same stick to reach the two stones), which fact can be expressed by the sequent $\sigma \vdash \sigma \cdot \sigma$. Also, the following formula holds of all resources relevant to the present case study: $\alpha \triangleright \gamma \wedge \beta \triangleright \delta \rightarrow \alpha \cdot \beta \triangleright \gamma \cdot \delta$. This formula implies a form of *scalability* of resources,¹² which is not a property holding in general, and hence has not been added to the general calculus. The crow passing the test shows to be able to conjoin the separate capabilities together. This is expressed by the following *transitivity*-type axiom: $\alpha \triangleright \beta \wedge \beta \triangleright \gamma \rightarrow \alpha \triangleright \gamma$. The crow’s achievement is remarkable precisely because this axiom cannot be expected to hold of any agent. These conditions translate into the following analytic rules:

$$\text{Contr} \frac{\Sigma \odot \Sigma \vdash \Omega}{\Sigma \vdash \Omega} \quad \text{Scalab} \frac{(\Gamma \cdot \blacktriangle \cdot X) \odot (\Pi \cdot \blacktriangle \cdot Y) \vdash \Delta}{(\Gamma \odot \Pi) \cdot \blacktriangle \cdot (X; Y) \vdash \Delta} \quad \text{Trans} \frac{(\Gamma \cdot \blacktriangle \cdot X) \cdot \blacktriangle \cdot Y \vdash \Delta}{\Gamma \cdot \blacktriangle \cdot (X; Y) \vdash \Delta}.$$

In order for the rule Contr to satisfy C₆ and C₉, we need to work with a version of D.LRC which admits *two* types of resources: the *reusable* ones (for which the contraction rule is sound) and the general ones for which contraction is not sound. Hence, the contraction would be introduced only for the reusable type. Once the new type has been introduced, the language and calculus of LRC need to be expanded with copies of each original connective, so as to account for the fact that each copy takes in input and outputs exactly one type unambiguously. Correspondingly, copies of each original rule have to be added so that each copy accounts for exactly one reading of the original rule. This is a tedious but entirely safe procedure that guarantees that a proper multitype calculus (cf. Definition 7.2) can be introduced which admits *all* the rules above. The reader is referred to [32,34] for examples of such a disambiguation procedure.

The following table shows the assumptions of the present case study:

Initial state	Capabilities
$\diamond \sigma$	$\sigma \triangleright \rho$ $\rho \cdot \rho \triangleright \lambda$ $\lambda \triangleright \phi$

¹² That is, if the agent is capable of getting one (measure of) β from one (measure of) α , then is also capable to get two or n (measures of) β from two or n (measures of) α .

We aim at proving the following sequent:

$$\sigma \triangleright \rho; \rho \cdot \rho \triangleright \lambda; \lambda \triangleright \phi; \Diamond \sigma \vdash \Diamond \phi.$$

We do it in several steps: first, in the following derivation π_1 , we prove that for any reusable resource σ , if $\sigma \triangleright \rho$ then $\sigma \triangleright \rho \cdot \rho$:

$$\begin{array}{c} \frac{\frac{\frac{\sigma \vdash \sigma \quad \rho \vdash \rho}{\sigma \triangleright \rho \vdash \sigma \cdot \triangleright \cdot \rho}}{\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho \vdash \rho} \quad \frac{\frac{\sigma \vdash \sigma \quad \rho \vdash \rho}{\sigma \triangleright \rho \vdash \sigma \cdot \triangleright \cdot \rho}}{\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho \vdash \rho}}{(\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho) \odot (\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho) \vdash \rho \cdot \rho} \\ \text{Scalab} \frac{(\sigma \odot \sigma) \cdot \blacktriangleleft \cdot (\sigma \triangleright \rho; \sigma \triangleright \rho) \vdash \rho \cdot \rho}{\sigma \triangleright \rho; \sigma \triangleright \rho \vdash \sigma \odot \sigma \cdot \triangleright \cdot \rho \cdot \rho} \\ \text{Contr} \frac{\sigma \odot \sigma \vdash (\sigma \triangleright \rho; \sigma \triangleright \rho) \cdot \triangleright \cdot \rho \cdot \rho}{\sigma \vdash (\sigma \triangleright \rho; \sigma \triangleright \rho) \cdot \triangleright \cdot \rho \cdot \rho} \\ \frac{\sigma \triangleright \rho; \sigma \triangleright \rho \vdash \sigma \cdot \triangleright \cdot \rho \cdot \rho}{\sigma \triangleright \rho \vdash \sigma \cdot \triangleright \cdot \rho \cdot \rho} \\ \frac{\sigma \triangleright \rho \vdash \sigma \cdot \triangleright \cdot \rho \cdot \rho}{\sigma \triangleright \rho \vdash \sigma \triangleright \rho \cdot \rho} \end{array}$$

Second, in the following derivation π_2 , we prove an instance of the transitivity axiom:

$$\begin{array}{c} \frac{\frac{\frac{\frac{\rho \vdash \rho \quad \rho \vdash \rho}{\rho \odot \rho \vdash \rho \cdot \rho}}{\sigma \vdash \sigma \quad \rho \cdot \rho \vdash \rho \cdot \rho}}{\sigma \triangleright \rho \cdot \rho \vdash \sigma \cdot \triangleright \cdot \rho \cdot \rho}}{\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho \cdot \rho \vdash \rho \cdot \rho} \quad \lambda \vdash \lambda \\ \frac{\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho \cdot \rho \vdash \rho \cdot \rho \quad \lambda \vdash \lambda}{\rho \cdot \rho \triangleright \lambda \vdash \sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho \cdot \rho \cdot \triangleright \cdot \lambda} \\ \text{Trans} \frac{(\sigma \cdot \blacktriangleleft \cdot \sigma \triangleright \rho \cdot \rho) \cdot \blacktriangleleft \cdot \rho \cdot \rho \triangleright \lambda \vdash \lambda}{\sigma \cdot \blacktriangleleft \cdot (\sigma \triangleright \rho \cdot \rho; \rho \cdot \rho \triangleright \lambda) \vdash \lambda} \\ \frac{\sigma \cdot \blacktriangleleft \cdot (\sigma \triangleright \rho \cdot \rho; \rho \cdot \rho \triangleright \lambda) \vdash \lambda}{\sigma \triangleright \rho \cdot \rho; \rho \cdot \rho \triangleright \lambda \vdash \sigma \cdot \triangleright \cdot \lambda} \\ \frac{\sigma \triangleright \rho \cdot \rho; \rho \cdot \rho \triangleright \lambda \vdash \sigma \cdot \triangleright \cdot \lambda}{\sigma \triangleright \rho \cdot \rho; \rho \cdot \rho \triangleright \lambda \vdash \sigma \triangleright \lambda} \end{array}$$

Similarly, a derivation π_3 can be given of the following instance of the transitivity axiom:

$$\sigma \triangleright \lambda; \lambda \triangleright \phi \vdash \sigma \triangleright \phi.$$

Finally, the following derivation π_4 is the missing piece:

$$\begin{array}{c} \text{proof for} \\ \vdots \\ \text{BD2} \quad \frac{\sigma \triangleright \Diamond \phi; \Diamond \sigma \vdash \Diamond \phi}{\sigma \triangleright \Diamond \phi \vdash \Diamond \phi < \Diamond \sigma} \\ \text{Cut} \frac{\sigma \triangleright \phi \vdash \sigma \triangleright \Diamond \phi \quad \sigma \triangleright \Diamond \phi \vdash \Diamond \phi < \Diamond \sigma}{\sigma \triangleright \phi \vdash \Diamond \phi < \Diamond \sigma} \\ \frac{\sigma \triangleright \phi \vdash \Diamond \phi < \Diamond \sigma}{\sigma \triangleright \phi; \Diamond \sigma \vdash \Diamond \phi} \end{array}$$

The requested sequent can be then derived using π_1 - π_4 via cuts and display postulates.

5.3. Resources having different roles (The Gift of the Magi). *The Gift of the Magi* is a short story, written by O. Henry and first appeared in 1905, about a young married couple of very modest means, Jim (j) and Della (d), who have only two possessions between them which are of value (both monetarily and in the sense that they take pride in them): Della's

unusually long hair (η), and Jim’s family gold watch (ω). On Christmas Eve, Della sells her hair to buy a chain (γ) for Jim’s watch, and Jim sells his watch to buy an ivory brush (β) for Della.

Jim and Della are materially worse off at the end of the story than at the beginning, since, while the resources ω and η could be used/enjoyed on their own, γ and β can only be used when coupled with ω and η , respectively. In fact, the very choice of γ and β as presents is a direct consequence of the fact that—besides being used by their respective owners as a means to get the money to buy a present for the other—the resources ω and η are used by the *partner* of their respective owners as *beacons* guiding them in their choice of a present. For instance, their final situation would not have been as bad if Della had bought Jim a new overcoat or a pair of gloves, or if Jim had bought Della replacements for her old brown jacket or hat, the need for which is indicated in the short story. However, each wants to make their present as meaningful as possible to the other one, and hence each *targets* his/her present at the one possession the other takes pride in.

Finally, the uniqueness of the meaningful resource of each agent is the reason why “the whole affair has something of the dark inevitability of Greek tragedy” (cit. P. G. Wodehouse, *Thank you, Jeeves*): indeed, ω (resp. η) is both the only target for a meaningful present for Jim (resp. Della), and also the only means he (resp. she) has to acquire such a present for her (resp. him).

To formalize the observations above, we will need a modification of the language of LRC capturing the fact, which is sometimes relevant, that resources might have different *roles*, e.g., in the generation or the acquisition of a given resource. For instance, in the production of bread, the oven has a different role as a resource than water and flour; in shooting sports, the shooter uses a shooting device, projectiles and a target in different roles, etc. Roles cannot be reduced to how resources are combined irrespective of agency (this aspect is modelled by the pure-resource connectives \sqcap and \cdot); rather, assigning roles to resources is a facet of agency. Accordingly, we consider the following ternary connective for each agent:

$$[-, -] \triangleright - : \text{Res} \times \text{Res} \times \text{Res} \rightarrow \text{Fm},$$

the intended meaning of which is ‘the agent is capable of obtaining the resource in the third coordinate, whenever in possession of the resources in the first two coordinates *in their respective roles*’. Algebraically (and axiomatically), this connective is finitely join-reversing in the first two coordinates and finitely meet-preserving in the third one. Its introduction rules and display postulates are as expected:

$$\frac{\Gamma \vdash \alpha \quad \Theta \vdash \beta \quad \gamma \vdash \Sigma}{[\alpha, \beta] \triangleright \gamma \vdash [\Gamma, \Theta] \cdot \triangleright \cdot \Sigma} \quad \frac{X \vdash [\alpha, \beta] \cdot \triangleright \cdot \gamma}{X \vdash [\alpha, \beta] \triangleright \gamma}$$

$$\frac{X \vdash [\Gamma, \Theta] \cdot \triangleright \cdot \Sigma}{[\Gamma, \Theta] \cdot \blacktriangleleft \cdot X \vdash \Sigma} \quad \frac{X \vdash [\Gamma, \Theta] \cdot \triangleright \cdot \Sigma}{\Gamma \vdash [X, \Theta] \cdot \blacktriangleright \cdot^1 \Sigma} \quad \frac{X \vdash [\Gamma, \Theta] \cdot \triangleright \cdot \Sigma}{\Theta \vdash [\Gamma, X] \cdot \blacktriangleright \cdot^2 \Sigma}.$$

In addition, we need two unary diamond operators $\diamond^1, \diamond^2 : \text{Res} \rightarrow \text{Fm}$ for each agent, the intended meaning of which is ‘the agent is in possession of the resource (in the argument) *in the first* (resp. *second*) *role*’. The basic algebraic and axiomatic behaviour of \diamond^1 and \diamond^2 coincides with that of \diamond , hence the introduction and display rules relative to these connectives are like those given for \diamond . The various roles and their differences

can be understood and formalized in different ways relative to different settings. In the specific situation of the short story, we stipulate that \diamond^2 has the meaning usually attributed to \diamond , and understand $\diamond^1\sigma$ as ‘the agent has resource σ available in the role of target (or beacon)’.

The interaction of these connectives, and the difference in meaning between \diamond^1 and \diamond^2 , are captured by the following axiom:

$$\diamond^1\sigma \wedge \diamond^2\xi \wedge [\sigma, \xi] \triangleright \chi \rightarrow \diamond \diamond^2\chi, \tag{5.1}$$

which is equivalent on perfect LRC-algebras to the following analytic rule:

$$\frac{\circ\omega^2[\Sigma, \Xi] \cdot \blacktriangle \cdot X \vdash Y}{\diamond^1\Sigma ; \diamond^2\Xi ; X \vdash Y} \text{RR.}$$

Finally, in the specific case at hand, we will use the rules corresponding to the following slightly modified multi-agent versions of axiom (5.1):

$$\diamond_j^1\sigma \wedge \diamond_j^2\xi \wedge [\sigma, \xi] \triangleright_j \chi \rightarrow \diamond_j \diamond_a^2\chi \quad \text{and} \quad \diamond_a^1\sigma \wedge \diamond_a^2\xi \wedge [\sigma, \xi] \triangleright_a \chi \rightarrow \diamond_a \diamond_j^2\chi.$$

The following table shows the assumptions of the present case study:

	Initial state	Capabilities	Abilities
Jim j	$\diamond_j^1\eta \ \diamond_j^2\omega$	$[\eta, \omega] \triangleright_j \beta$	$\diamond_j \diamond_a^2\beta \rightarrow \diamond_j \neg \diamond_j^2\omega$
Della d	$\diamond_d^1\omega \ \diamond_d^2\eta$	$[\omega, \eta] \triangleright_d \gamma$	$\diamond_a \diamond_j^2\gamma \rightarrow \diamond_a \neg \diamond_d^2\eta$

Let H be the structural conjunction of the assumptions above. We aim at deriving the following sequent in the calculus D.LRC to which the analytic rules introduced above have been added:

$$H \vdash \diamond_j \neg \diamond_j^2\omega \wedge \diamond_j \diamond_a^2\beta \wedge \diamond_a \neg \diamond_a^2\eta \wedge \diamond_a \diamond_j^2\gamma.$$

We do it in several steps: first, the following derivation π_1 :

$$\frac{\frac{\frac{\eta \vdash \eta \quad \omega \vdash \omega \quad \beta \vdash \beta}{[\eta, \omega] \triangleright_j \beta \vdash [\eta, \omega] \cdot \triangleright_j \beta}}{[\eta, \omega] \cdot \blacktriangle \cdot_j [\eta, \omega] \triangleright_j \beta \vdash \beta}}{\diamond_a^2([\eta, \omega] \cdot \blacktriangle \cdot_j [\eta, \omega] \triangleright_j \beta) \vdash \diamond_a^2\beta}}{\circ_j \diamond_a^2([\eta, \omega] \cdot \blacktriangle \cdot_j [\eta, \omega] \triangleright_j \beta) \vdash \diamond_j \diamond_a^2\beta} \text{RR}_{ja}$$

$$\frac{(\diamond_j^1\eta ; \diamond_j^2\omega) ; [\eta, \omega] \triangleright_j \beta \vdash \diamond_j \diamond_a^2\beta}{(\diamond_j^1\eta ; \diamond_j^2\omega) ; [\eta, \omega] \triangleright_j \beta \vdash \diamond_j \diamond_a^2\beta}$$

With an analogous derivation π_2 we can prove that

$$\diamond_a^1\omega ; \diamond_a^2\eta ; [\omega, \eta] \triangleright_a \gamma \vdash \diamond_a \diamond_j^2\gamma.$$

Next, let π_3 be the following derivation:

$$\begin{array}{c}
 \frac{\omega \vdash \omega}{\Phi_j^2 \omega \vdash \Phi_j^2 \omega} \\
 \frac{\Phi_j^2 \omega \vdash \Phi_j^2 \omega \quad \perp \vdash \perp}{\Phi_j^2 \omega \rightarrow \perp \vdash \Phi_j^2 \omega > \perp} \\
 \frac{\Phi_j^2 \omega \rightarrow \perp \vdash \Phi_j^2 \omega \rightarrow \perp}{\neg \Phi_j^2 \omega \vdash \neg \Phi_j^2 \omega} \text{ def} \\
 \frac{\beta \vdash \beta}{\Phi_a^2 \beta \vdash \Phi_a^2 \beta} \\
 \frac{\Phi_a^2 \beta \vdash \Phi_a^2 \beta}{\diamond_j \Phi_a^2 \beta \vdash \diamond_j \Phi_a^2 \beta} \\
 \frac{\diamond_j \Phi_a^2 \beta \vdash \diamond_j \Phi_a^2 \beta}{\diamond_j \Phi_a^2 \beta \rightarrow \diamond_j \neg \Phi_j^2 \omega \vdash \diamond_j \Phi_a^2 \beta > \diamond_j \neg \Phi_j^2 \omega} \\
 \hline
 \diamond_j \Phi_a^2 \beta ; \diamond_j \Phi_a^2 \beta \rightarrow \diamond_j \neg \Phi_j^2 \omega \vdash \diamond_j \neg \Phi_j^2 \omega
 \end{array}$$

With an analogous derivation π_4 we can prove that

$$\diamond_a \Phi_j^2 \gamma ; \diamond_a \Phi_j^2 \gamma \rightarrow \diamond_a \neg \Phi_a^2 \eta \vdash \diamond_a \neg \Phi_a^2 \eta.$$

Then, by applying cut (and left weakening) on π_1 and π_3 one derives the following:

$$\diamond_j^1 \eta ; \Phi_j^2 \omega ; [\eta, \omega] \triangleright_j \beta ; \diamond_j \Phi_a^2 \beta \rightarrow \diamond_j \neg \Phi_j^2 \omega \vdash \diamond_j \neg \Phi_j^2 \omega.$$

Likewise, by applying cut (and left weakening) on π_2 and π_4 one derives the following:

$$\diamond_a^1 \omega ; \Phi_a^2 \eta ; [\omega, \eta] \triangleright_a \gamma ; \diamond_a \Phi_j^2 \gamma \rightarrow \diamond_a \neg \Phi_a^2 \eta \vdash \diamond_a \neg \Phi_a^2 \eta.$$

The derivation is concluded with applications of right-introduction of \wedge and left contraction rules.

5.4. From local to global resilience (two production lines). Resilience is the ability of an agent or an organization to realize their goals notwithstanding unexpected changes and disruptions. The language of LRC provides a natural way to understand resilience as the capability to realize one’s goal(s) in a range of situations characterized by the reduced availability of key resources. Consider for example a factory with two production lines for products γ_1 and γ_2 . Product γ_1 is of higher quality than γ_2 and can only be produced using resource α , the availability of which is subject to fluctuations. Product γ_2 can be produced using either resource α or β , and the availability of β is not subject to fluctuations. It is interesting to note that the ‘local’ resilience in the production of γ_2 (namely, the fact that any shortage in α can be dealt with by switching to β) results in the resilience of both production lines. Indeed, when α is available for only one of the two production lines, all of it can be employed in the production line for γ_1 , and the production of γ_2 is switched to β . In the formal treatment that follows, we notice that the axioms $\Phi \sigma \wedge \sigma \triangleright \pi \rightarrow \Phi \pi$ and $\sigma \triangleright \chi \wedge \pi \triangleright \xi \rightarrow \sigma \cdot \pi \triangleright \chi \cdot \xi$ hold for the setting described above. These axioms are analytic and are equivalent on perfect LRC-algebras to the following rules:

$$\text{BDR} \frac{X \vdash \Gamma \cdot \triangleright \cdot \bullet Y}{X \vdash \bullet \Gamma > Y} \quad \frac{(\Gamma \cdot \blacktriangle \cdot X) \odot (\Pi \cdot \blacktriangle \cdot Y) \vdash \Delta}{(\Gamma \odot \Pi) \cdot \blacktriangle \cdot (X ; Y) \vdash \Delta} \text{ Scalab}$$

Resources	Capabilities
$\diamond(((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta)$	$\alpha \triangleright \gamma_1$ $\alpha \sqcup \beta \triangleright \gamma_2$

We aim at showing that the assumptions above are enough to conclude that the factory is able to realize the production of both γ_1 and γ_2 :

$$\Diamond(((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta); \alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash \Diamond(\gamma_1 \cdot \gamma_2).$$

Notice that the following is an instance of $\Diamond\sigma \wedge \sigma \triangleright \pi \rightarrow \Diamond\pi$, and hence is derivable using the rule BDR:

$$\Diamond(((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta); ((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta \triangleright \gamma_1 \cdot \gamma_2 \vdash \Diamond(\gamma_1 \cdot \gamma_2).$$

Hence, modulo cut and left weakening, it is enough to show that

$$\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash ((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta \triangleright \gamma_1 \cdot \gamma_2.$$

Notice that:

$$\frac{\begin{array}{c} \vdots \text{ proof for } \\ \vdots \text{ R4} \end{array} \quad \frac{\gamma_1 \vdash \gamma_1 \quad \gamma_2 \vdash \gamma_2}{\gamma_1 \odot \gamma_2 \vdash \gamma_1 \cdot \gamma_2}}{\frac{((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta \vdash (\alpha \cdot \alpha) \cdot \beta \sqcup \alpha \cdot \beta}{(\alpha \cdot \alpha) \cdot \beta \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2 \vdash ((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta \triangleright \gamma_1 \cdot \gamma_2}}{(\alpha \cdot \alpha) \cdot \beta \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2 \vdash ((\alpha \cdot \alpha) \sqcup \alpha) \cdot \beta \triangleright \gamma_1 \cdot \gamma_2}}$$

Hence, modulo cut and left weakening, it is enough to show that

$$\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash (\alpha \cdot \alpha) \cdot \beta \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2.$$

Indeed, a derivation for the sequent above is

$$\frac{\begin{array}{c} \vdots \pi_1 \\ \alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash \alpha \cdot (\alpha \sqcup \beta) \triangleright \gamma_1 \cdot \gamma_2 \end{array} \quad \begin{array}{c} \vdots \pi_2 \\ \alpha \cdot (\alpha \sqcup \beta) \triangleright \gamma_1 \cdot \gamma_2 \vdash (\alpha \cdot \alpha) \cdot \beta \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2 \end{array}}{\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash (\alpha \cdot \alpha) \cdot \beta \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2}$$

where π_1 is the following derivation:

$$\text{Scalab} \frac{\frac{\frac{\alpha \vdash \alpha \quad \gamma_1 \vdash \gamma_1}{\alpha \triangleright \gamma_1 \vdash \alpha \cdot \triangleright \cdot \gamma_1} \quad \frac{\frac{\alpha \vdash \alpha \quad \beta \vdash \beta}{\alpha \sqcup \beta \vdash \alpha, \beta} \quad \frac{\alpha \sqcup \beta \vdash \alpha \sqcup \beta \quad \gamma_2 \vdash \gamma_2}{\alpha \sqcup \beta \triangleright \gamma_2 \vdash \alpha \sqcup \beta \cdot \triangleright \cdot \gamma_2}}{\frac{\alpha \cdot \blacktriangle \cdot \alpha \triangleright \gamma_1 \vdash \gamma_1 \quad (\alpha \sqcup \beta) \cdot \blacktriangle \cdot \alpha \sqcup \beta \triangleright \gamma_2 \vdash \gamma_2}{(\alpha \cdot \blacktriangle \cdot \alpha \triangleright \gamma_1) \odot (\alpha \sqcup \beta) \cdot \blacktriangle \cdot \alpha \sqcup \beta \triangleright \gamma_2 \vdash \gamma_1 \cdot \gamma_2}}{\frac{\alpha \odot (\alpha \sqcup \beta) \cdot \blacktriangle \cdot (\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2) \vdash \gamma_1 \cdot \gamma_2}{\frac{\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash \alpha \odot (\alpha \sqcup \beta) \cdot \triangleright \cdot \gamma_1 \cdot \gamma_2}{\frac{\alpha \odot (\alpha \sqcup \beta) \vdash \alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \cdot \triangleright \cdot \gamma_1 \cdot \gamma_2}{\frac{\alpha \cdot (\alpha \sqcup \beta) \vdash \alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \cdot \triangleright \cdot \gamma_1 \cdot \gamma_2}{\frac{\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash \alpha \cdot (\alpha \sqcup \beta) \cdot \triangleright \cdot \gamma_1 \cdot \gamma_2}{\alpha \triangleright \gamma_1; \alpha \sqcup \beta \triangleright \gamma_2 \vdash \alpha \cdot (\alpha \sqcup \beta) \triangleright \gamma_1 \cdot \gamma_2}}}}}}}$$

and π_2 is the following derivation:

$$\begin{array}{c}
 \frac{\alpha \vdash \alpha}{\alpha \odot \Phi \vdash \alpha} \\
 \frac{\alpha \odot \Phi \vdash \alpha}{\Phi \vdash \alpha \triangleright \alpha} \\
 \frac{\Phi \vdash \alpha \triangleright \alpha}{\alpha \vdash \alpha \triangleright \alpha} \\
 \frac{\alpha \vdash \alpha \triangleright \alpha}{\alpha \odot \alpha \vdash \alpha} \\
 \frac{\alpha \odot \alpha \vdash \alpha}{\alpha \cdot \alpha \vdash \alpha} \\
 \frac{\alpha \cdot \alpha \vdash \alpha}{\alpha \cdot \alpha \odot \beta \vdash \alpha} \\
 \frac{\alpha \cdot \alpha \odot \beta \vdash \alpha}{(\alpha \cdot \alpha \cdot \beta \vdash \alpha)} \\
 \frac{(\alpha \cdot \alpha \cdot \beta) \sqcup (\alpha \cdot \beta) \vdash \alpha \cdot (\alpha \sqcup \beta)}{\alpha \cdot (\alpha \sqcup \beta) \triangleright \gamma_1 \cdot \gamma_2 \vdash (\alpha \cdot \alpha \cdot \beta) \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2} \\
 \frac{\alpha \cdot (\alpha \sqcup \beta) \triangleright \gamma_1 \cdot \gamma_2 \vdash (\alpha \cdot \alpha \cdot \beta) \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2}{\alpha \cdot (\alpha \sqcup \beta) \triangleright \gamma_1 \cdot \gamma_2 \vdash (\alpha \cdot \alpha \cdot \beta) \sqcup \alpha \cdot \beta \triangleright \gamma_1 \cdot \gamma_2}
 \end{array}
 \qquad
 \begin{array}{c}
 \frac{\alpha \vdash \alpha}{\alpha \odot \Phi \vdash \alpha} \\
 \frac{\alpha \odot \Phi \vdash \alpha}{\Phi \vdash \alpha \triangleright \alpha} \\
 \frac{\Phi \vdash \alpha \triangleright \alpha}{\alpha \vdash \alpha \triangleright \alpha} \\
 \frac{\alpha \vdash \alpha \triangleright \alpha}{\alpha \odot \alpha \vdash \alpha} \\
 \frac{\alpha \odot \alpha \vdash \alpha}{\alpha \odot \beta \vdash \alpha} \\
 \frac{\alpha \odot \beta \vdash \alpha}{\alpha \cdot \beta \vdash \alpha} \\
 \frac{\alpha \cdot \beta \vdash \alpha}{\alpha \cdot \beta \vdash \alpha, \beta} \\
 \frac{\alpha \cdot \beta \vdash \alpha, \beta}{\alpha \cdot \beta \vdash \alpha \sqcup \beta} \\
 \frac{\gamma_1 \vdash \gamma_1 \quad \gamma_2 \vdash \gamma_2}{\gamma_1 \odot \gamma_2 \vdash \gamma_1 \cdot \gamma_2} \\
 \frac{\gamma_1 \odot \gamma_2 \vdash \gamma_1 \cdot \gamma_2}{\gamma_1 \cdot \gamma_2 \vdash \gamma_1 \cdot \gamma_2}
 \end{array}$$

§6. Conclusions and further directions.

Resources and capabilities. In the present article, a logical framework is introduced aimed at capturing and reasoning about resource flow within organizations. This framework contributes to the line of investigation of the logics of agency (cf. e.g., [4,10,28–30]) by focusing specifically on the *resource*-dimension of agents’ (cap)abilities (e.g., to use resources to achieve goals, to transform resources into other resources, and to coordinate the use of resources with other agents). Formally, the logic of resources and capabilities (LRC) has been introduced in a language consisting of formula-terms and resource-terms. Besides pure-formula and pure-resource connectives, the language of LRC includes connectives bridging the two types in various ways. Although action-terms are not included in LRC, perhaps the logical system of which LRC is most reminiscent is the logic of capabilities introduced in [71], which formalizes the capabilities of agents to perform actions. Indeed, looking past the differences between the two formalisms deriving from the inherent differences between actions and resources, the focus of both axiomatizations is *interaction*, between (cap)abilities and actions in [71], and between (cap)abilities and resources in the present article. Precisely its focus on interaction makes it worthwhile to recast the logical framework of [71] in a multitype environment.

A study in algebraic proof theory. The main technical contribution of the article is the introduction of the multitype calculus D.LRC. The definition of this calculus and the proofs of its basic properties hinge on the integration of two theories in algebraic logic and structural proof theory—namely, *unified correspondence* and *multitype calculi*—which originated independently of each other. This integration contributes to the research program of *algebraic proof theory* [11,13], to which the results of the present article pertain. Specifically, the rules of D.LRC are introduced, and their soundness proved, by applying (and adapting) the ALBA-based methodology of [44] (cf. also [12] for a purely proof-theoretic perspective on the same methodology); cut elimination is proved ‘Belnap-style’, by verifying that D.LRC satisfies the assumptions of the cut elimination metatheorem for multitype calculi of [33]; conservativity is proved following the general proof strategy for conservativity illustrated in [44], to which the canonicity of the axioms of the Hilbert-style presentation of LRC is key.

It is perhaps worth stressing that the theory of proper display calculi developed in [44] cannot be applied directly to the Hilbert-style presentation of LRC, for two reasons. First,

the setting of [44] is a pure-formula setting, while the setting of the present article is multitype. However, the results of [44] can be ported to the multitype setting (as done also in [36,45,46]); indeed, the algorithm ALBA and the definition of analytic inductive inequalities are grounded in the order-theoretic properties of the algebraic interpretations of the logical connectives, and remain fundamentally unchanged when applied to maps with the required order-theoretic properties, irrespective of whether these maps are operations on one algebra or between different algebras. The second, more serious reason is that the algebraic interpretation of the capability connective \triangleright is a map which reverses finite joins in its first coordinate but is only *monotone* (rather than finitely meet-preserving) in its second coordinate. Hence, (the multitype version of) the definition of (analytic) inductive inequalities given in [44] does not apply to many axioms of the Hilbert-style presentation of LRC, and hence some results (e.g., the canonicity results of §2.3) could not be immediately inferred by directly applying the general theory. However, as we saw in §4.1, the algorithm ALBA is successful on the LRC axiomatization, which suggests the possibility of generalizing these results to arbitrary multitype signatures in which operations are allowed to be only monotone or antitone in some coordinates. Moreover, unified correspondence theory covers various settings, from general lattice-based propositional logics [17,18,21,22], to regular [61] and monotone modal logics [37], (distributive) lattice-based mu-calculi [14–16], hybrid logic [26] and many-valued logic [52]. It would be interesting to investigate whether structural proof calculi for each of these settings (or for multitype logics based on them) could be defined by suitably extending the techniques employed in the design of D.LRC.

Proof-theoretic formalizations of social behaviour. In §5, we have discussed the formalization of situations revolving around some instances of resource flow. These situations have been captured as inferences or sequents in the language of LRC, and derived in the basic calculus D.LRC or in some of its analytic extensions. This proof-theoretic analysis makes it possible to single out the steps and assumptions which are *essential* to a given situation. For instance, thanks to this analysis, it is clear that the full power of classical logic is *not essential* to any case study we treated. In fact, as can be readily verified by inspection, many derivations treated in §5 need less than the full power of intuitionistic logic, which is the propositional base of LRC. Also, reasoning from assumptions in a given proof-theoretic environment corresponds semantically to reasoning on *all* the models of that environment satisfying those assumptions. This is a *safer* practice than, e.g., starting out with an ad-hoc model, since it makes it impossible to rely on some implicit assumption or other extra feature of a chosen model.

The pure-resource fragment. In §2.1 we mentioned that the fact that 1 coincides with the weakest resource entails (and is in fact equivalent to) the validity of the sequents $\alpha \cdot \beta \vdash \alpha$ and $\alpha \cdot \beta \vdash \beta$, which in some contexts seems too restrictive. How to relax this restriction is current work in progress. However, this restriction brings also some advantages. Indeed, as discussed earlier on in §2.1, this restriction makes the pure resource fragment of LRC very similar to (the exponential-free fragment of) linear affine logic, which, unlike general linear logic, is decidable [50,58]. Hence, this leaves open the question of the decidability of LRC (see also below).

Agents as first-class citizens. In the present article, we focused on the basic setting of LRC, and for the sake of not overloading notation and machinery, we have treated agents as parameters. However, a fully multitype treatment would include terms of type **Ag** (agents)

in the language, as done, e.g., in [34]. This will be particularly relevant to the formalization of organization theory, where terms of type \mathbf{Ag} will represent members of an organization, and \mathbf{Ag} might be endowed with additional structure: for instance it can be a graph (capturing networks of agents), or a partial order (capturing hierarchies), or partitioned in coalitions or teams. Having agents as first-class citizens of the language will also make possible to attribute *roles* to them, analogously to the way roles are attributed to resources in §5.3. Roles in turn could provide concrete handles towards the modelling of agent coordination.

Group capabilities. Closely related to the issue of the previous paragraph is the formalization of various forms of group capabilities. This theme is particularly relevant to organization theory, since it might help to capture, e.g., the contribution of leadership to the results of an organization, versus the advantages of self-organization. Another interesting notion in organization theory which could benefit from a formal theory of group capabilities is *tacit group knowledge* [68], emerging from the individual capabilities to adapt, often implicitly, to the behaviour of others.

Different types of resources. Key to the analysis of the case study of §5.2 was the interplay between reusable and nonreusable resources. The treatment of this case study suggests that analytic extensions of D.LRC can be used to develop a formal theory of resource flow that also captures other differences between resources (e.g., storable vs. non storable, scalable vs. non scalable), their interaction, direct or mediated by agents, in the production process, or in facilitating more generally the competitive success of the organization [57].

Pre-orderings on resources. In §5.3, we mentioned that the resources the agents possess at the end of the story cannot be used without those they possess at the beginning, while these can be used on their own. This observation suggest that alternative or additional orderings of resources can be considered and studied, such as the ‘dependence’ preorder between resources, which might be relevant to the analysis of some situations.

Comparing capabilities. The logic LRC provides a formal environment where to explore the consequences for organizations of some agent’s being *more* capable than some other agent at bringing about a certain state of affairs. In this environment, we can express that agent a is *at least as capable* than agent b at bringing about A , e.g., when $\alpha \triangleright_a A$ and $\beta \triangleright_b A$, and $\beta \vdash \alpha$ (i.e., to bring about the same state of affairs, b uses a resource which is at least as powerful as, possibly more powerful than, the resource used by a). Ricardo’s economic theory of comparative advantage with regard to the division of labour in organizations [65] can be formalized on the basis of capabilities differentials.

Algebraic canonicity and relational semantics. The theory of canonical extensions provides a way to extract relational semantics from the algebraic semantics via algebraic canonicity. In §2.3, we have shown that the logic LRC is complete w.r.t. *perfect* LRC-algebraic models. Via standard discrete Stone-type duality, perfect LRC-algebraic models can be associated with set-based structures similar to Kripke models, thus providing complete relational semantics for LRC. The specification of this relational semantics and its properties is part of future work.

Semantics of Petri nets. We are currently studying Petri nets as an alternative semantic framework for LRC. In particular, the reachability problem for finite Petri nets is equivalent to the deducibility problem for sequents in finitely axiomatized theory in the pure-tensor

fragment of linear logic [56,69]. More recently, [31] proved completeness for several versions of linear logic w.r.t. Petri nets. We are investigating similar issues in the setting of LRC.

Decidability, finite model property, complexity. The computational properties of LRC such as decidability and complexity are certainly of interest. In particular, two, in general distinct, problems are to be considered: the decidability of the set of theorems, and the decidability of the (finite) consequence relation.¹³

A standard argument establishing decidability is via the so-called finite model property (FMP), i.e., proving that any nontheorem can be refuted in a finite structure. Together with finite axiomatizability and completeness of the underlying logic, FMP entails the decidability of the set of theorems. For the second problem a stronger property is needed: the finite embeddability property, which can be seen as the finite model property for quasi-identities and, together with finite axiomatizability and completeness, entails the decidability of the finite consequence relation of the underlying logic.

We wish to stress that the decidability problems for LRC subsume the complexity and decidability of certain substructural logics. Indeed, as mentioned earlier, the pure-resource fragment of LRC is similar to (propositional, exponential-free) linear affine logic, which essentially coincides with the distributive Full Lambek calculus with weakening, a logic for which the finite consequence relation, and hence the set of theorems, are known to be decidable (see [58,59]); FEP for integral residuated groupoids has been proved in [7], for a simple proof of FEP in the distributive setting see also [47], where a coNEXP upper bound is obtained. We hope we can use the algebraic semantics of LRC to investigate, and hopefully establish decidability of LRC and its variants using FMP or FEP.

Syntactic decidability. An alternative path towards decidability for LRC consists in adapting the techniques developed in [50], where a syntactic proof is given of the decidability of full propositional affine linear logic, by showing that it is enough to consider sequents in a suitable normal form. An encouraging hint is the fact that the full Lambek calculus with weakening is decidable [58,59]. However, it is also known that, for certain substructural logics, distributivity is problematic for decidability.

§7. Appendix: Proper multitype calculi and their cut elimination. In the present section, we report on the Belnap-style meta-theorem that we appeal to in order to show that the calculus introduced in §3 enjoys cut elimination. This meta-theorem was proven in [33] for the so-called *proper multitype calculi*. In order to make the exposition self-contained, in what follows we will report the definition of proper multitype calculi and the statement of the meta-theorem.

DEFINITION 7.1. *A sequent $x \vdash y$ is type-uniform if x and y are of the same type \top (cf. [34, Definition 3.1]).*

DEFINITION 7.2. *A proper multitype calculus is any calculus in a multitype language satisfying the following list of conditions:¹⁴*

¹³ The two problems coincide in presence of deduction theorem, which is available in intuitionistic logic and for the formula-fragment of LRC, but not for the pure-resource fragment of LRC.

¹⁴ See [35] for a discussion on C'_5 and C''_5 .

C₁: Preservation of operational terms. Each operational term occurring in a premise of an inference rule *inf* is a subterm of some operational term in the conclusion of *inf*.

C₂: Shape-alikeness of parameters. Congruent parameters (i.e., nonactive terms in the application of a rule) are occurrences of the same structure.

C₂: Type-alikeness of parameters. Congruent parameters have exactly the same type. This condition bans the possibility that a parameter changes type along its history.

C₃: Nonproliferation of parameters. Each parameter in an inference rule *inf* is congruent to at most one constituent in the conclusion of *inf*.

C₄: Position-alikeness of parameters. Congruent parameters are either all precedent or all succedent parts of their respective sequents. In the case of calculi enjoying the display property, precedent and succedent parts are defined in the usual way (see [3]). Otherwise, these notions can still be defined by induction on the shape of the structures, by relying on the polarity of each coordinate of the structural connectives.

C₅: Quasi-display of principal constituents. If an operational term *a* is principal in the conclusion sequent *s* of a derivation π , then *a* is in display, unless π consists only of its conclusion sequent *s* (i.e., *s* is an axiom).

C₅'': Display-invariance of axioms. If *a* is principal in an axiom *s*, then *a* can be isolated by applying Display Postulates and the new sequent is still an axiom.

C₅''': Closure of axioms under surgical cut. If $(x \vdash y)([a]^{pre}, [a]^{suc})$, $a \vdash z[a]^{suc}$ and $v[a]^{pre} \vdash a$ are axioms, then $(x \vdash y)([a]^{pre}, [z/a]^{suc})$ and $(x \vdash y)([v/a]^{pre}, [a]^{suc})$ are again axioms.

C₆'': Closure under substitution for succedent parts within each type. Each rule is closed under simultaneous substitution of arbitrary structures for congruent operational terms occurring in succedent position, within each type.

C₇'': Closure under substitution for precedent parts within each type. Each rule is closed under simultaneous substitution of arbitrary structures for congruent operational terms occurring in precedent position, within each type.

Condition C₆' (and likewise C₇') ensures, for instance, that if the following inference is an application of the rule *R*:

$$\frac{(x \vdash y)([a]_i^{suc} \mid i \in I)}{(x' \vdash y')[a]^{suc}} R,$$

and $([a]_i^{suc} \mid i \in I)$ represents all and only the occurrences of *a* in the premiss which are congruent to the occurrence of *a* in the conclusion (if $I = \emptyset$, then the occurrence of *a* in the conclusion is congruent to itself), then also the following inference is an application of the same rule *R*:

$$\frac{(x \vdash y)([z/a]_i^{suc} \mid i \in I)}{(x' \vdash y')[z/a]^{suc}} R,$$

where the structure *z* is substituted for *a*.

This condition caters for the step in the cut elimination procedure in which the cut needs to be “pushed up” over rules in which the cut-formula in succedent position is parametric (cf. [33, §4]).

C₈: Eliminability of matching principal constituents. This condition requests a standard Gentzen-style checking, which is now limited to the case in which both cut formulas are principal, i.e., each of them has been introduced with the last rule application of each corresponding subdeduction. In this case, analogously to the proof Gentzen-style, condition C₈ requires being able to transform the given deduction into a deduction with the same

conclusion in which either the cut is eliminated altogether, or is transformed in one or more applications of the cut rule, involving proper subterms of the original operational cut-term. In addition to this, specific to the multitype setting is the requirement that the new application(s) of the cut rule be also strongly type-uniform (cf. condition C_{10} below).

C_9 : **Type-uniformity of derivable sequents.** Each derivable sequent is type-uniform.

C_{10} : **Preservation of type-uniformity of cut rules.** All cut rules preserve type-uniformity.

In the context of proper multitype calculi we say that a rule is *analytic* if it satisfies conditions C_1 – C'_7 of the list above. Analytic rules can be added to a given proper multitype calculus, and the resulting calculus enjoys cut elimination and subformula property.

We state the cut-elimination metatheorem which we appeal to when establishing the cut elimination for the calculus introduced in §3.

THEOREM 7.3. *Any calculus satisfying $C_2, C'_2, C'_3, C_4, C'''_5, C''''_5, C'_6, C'_7, C'_8, C''_8, C_9$, and C_{10} is cut-admissible. If also C_1 is satisfied, then the calculus enjoys the subformula property.*

§8. Acknowledgments. The research of the first author has been supported by the project SEGA: From Shared Evidence to Group Agency, of the Czech Science Foundation, and DFG no. 16-07954J. The research of the second author was also supported by the Values4Water project, subsidised by the Responsible Innovation research programme, which is partly financed by the Netherlands Organisation for Scientific Research (NWO) under Grant Number 313-99-316. The research of the second, third, and fourth author has been made possible by the NWO Vidi grant 016.138.314, the NWO Aspasia grant 015.008.054, and a Delft Technology Fellowship awarded in 2013.

BIBLIOGRAPHY

- [1] Baltag, A., Moss, L. S., & Solecki, S. The logic of public announcements, common knowledge and private suspicious. Technical Report SEN-R9922, CWI, Amsterdam, 1999.
- [2] Barney, J. (1991). Firm resources and sustained competitive advantage. *Journal of Management*, **17**(1), 99–120.
- [3] Belnap, N. (1982). Display logic. *Journal of Philosophical Logic*, **11**(4), 375–417.
- [4] Belnap, N. (1991) Backwards and forwards in the modal logic of agency. *Philosophy and Phenomenological Research*, **51**(4), 777–807.
- [5] Belnap, N. & Perloff, M. (1988). Seeing to it that: A canonical form for agentives. *Theoria*, **54**(3), 175–199.
- [6] Birkhoff, G. & Lipson, J. D. (1970). Heterogeneous algebras. *Journal of Combinatorial Theory*, **8**(1), 115–133.
- [7] Blok, W. & Van Alten, C. (2005). On the finite embeddability property for residuated ordered groupoids. *Transactions of the American Mathematical Society*, **357**(10), 4141–4157.
- [8] Brown, M. A. (1988). On the logic of ability. *Journal of Philosophical Logic*, **17**(1), 1–26.
- [9] Chellas, B. F. (1969). The Logical form of Imperatives. Ph.D. Thesis, Stanford University.
- [10] Chellas, B. F. (1995). On bringing it about. *Journal of Philosophical Logic*, **24**(6), 563–571.

- [11] Ciabatonni, A., Galatos, N., & Terui, K. (2012). Algebraic proof theory for substructural logics: Cut-elimination and completions. *Annals of Pure and Applied Logic*, **163**(3), 266–290.
- [12] Ciabatonni, A. & Ramanayake, R. (2016). Power and limits of structural display rules. *ACM Transactions on Computational Logic (TOCL)*, **17**(3), 17.
- [13] Ciabatonni, A., Straßburger, L., & Terui, K. (2009). Expanding the realm of systematic proof theory. In Grädel, E., and Kahle, R., editors. *International Workshop on Computer Science Logic*. Berlin: Springer, pp. 163–178.
- [14] Conradie, W. & Craig, A. (2017). Canonicity results for mu-calculi: An algorithmic approach. *Journal of Logic and Computation*, **27**(3), 705–748.
- [15] Conradie, W., Craig, A., Palmigiano, A., & Zhao, Z. (2017). Constructive canonicity for lattice-based fixed point logics. In Kennedy, J., and de Queiroz, R. J. G. B., editors. *Logic, Language, Information, and Computation*. LNCS 10388. Berlin: Springer-Verlag, pp. 92–109.
- [16] Conradie, W., Fomatati, Y., Palmigiano, A., & Sourabh, S. (2015). Algorithmic correspondence for intuitionistic modal mu-calculus. *Theoretical Computer Science*, **564**, 30–62.
- [17] Conradie, W., Frittella, S., Palmigiano, A., Piazzai, M., Tzimoulis, A., & Wijnberg, N. (2016). Categories: How I learned to stop worrying and love two sorts. In Väänänen, J., Hirvonen, A., and de Queiroz, R., editors. *Logic, Language, Information, and Computation*. LNCS 9803. Berlin: Springer-Verlag, pp. 145–164.
- [18] Conradie, W., Frittella, S., Palmigiano, A., Piazzai, M., Tzimoulis, A., & Wijnberg, N. (2017). Towards an epistemic-logical theory of categorization. *Electronic Proceedings in Theoretical Computer Science*, **251**, 167–186. ArXiv:1707.08743.
- [19] Conradie, W., Ghilardi, S., & Palmigiano, A. (2014). Unified correspondence. In Baltag, A. and Smets, S., editors. *Johan van Benthem on Logic and Information Dynamics*. Outstanding Contributions to Logic, Vol. 5. Dordrecht: Springer International Publishing, pp. 933–975.
- [20] Conradie, W. & Palmigiano, A. (2012). Algorithmic correspondence and canonicity for distributive modal logic. *Annals of Pure and Applied Logic*, **163**(3), 338–376.
- [21] Conradie, W. & Palmigiano, A. Algorithmic correspondence and canonicity for non-distributive logics, preprint, arXiv: 1603.08515.
- [22] Conradie, W. & Palmigiano, A. Constructive canonicity of inductive inequalities, preprint, arXiv: 1603.08341.
- [23] Conradie, W., Palmigiano, A., & Sourabh, S. (2017). Algebraic modal correspondence: Sahlqvist and beyond. *Journal of Logical and Algebraic Methods in Programming*, **91**, 60–84.
- [24] Conradie, W., Palmigiano, A., Sourabh, S., & Zhao, Z. Canonicity and relativized canonicity via pseudo-correspondence: An application of ALBA, preprint, arXiv: 1511.04271.
- [25] Conradie, W., Palmigiano, A., & Zhao, Z. Sahlqvist via translation, preprint, arXiv: 1603.08220.
- [26] Conradie, W. & Robinson, C. (2017). On Sahlqvist theory for hybrid logics. *Journal of Logic and Computation*, **27**(3), 867–900.
- [27] Cross, C. B. (1986). ‘Can’ and the logic of ability. *Philosophical Studies*, **50**(1), 53–64.
- [28] Dignum, V. (2003). A Model for Organizational Interaction: Based on Agents, Founded in Logic. Ph.D. Thesis, The Netherlands: University of Utrecht.

[29] Dignum, V. & Dignum, F. (2009). A logic of agent organizations. *Logic Journal of IGPL*, **20**(1), 283–316.

[30] Elgesem, D. (1997). The modal logic of agency. *Nordic Journal of Philosophical Logic*, **2**, 1–46.

[31] Engberg, U. H. & Winskel, G. (1993). Linear logic on Petri nets. In de Bakker, J. W., de Roever, W. P., and Rozenberg, G., editors. *REX School/Symposium: A Decade of Concurrency, Reflections and Perspective*. LNCS 3580, Vol. 803. Berlin: Springer, pp. 176–229.

[32] Frittella, S., Greco, G., Kurz, A., & Palmigiano, A. (2016). Multi-type display calculus for propositional dynamic logic. *Journal of Logic and Computation*, **26**(6), 2067–2104.

[33] Frittella, S., Greco, G., Kurz, A., Palmigiano, A., & Sikimić, V. (2014). Multi-type sequent calculi. In Indrzejczak, A., Kaczmarek, J., and Zawidzki, M., editors. *Proceedings of Trends in Logic XIII*. Lodz: Lodz University Press, pp. 81–93.

[34] Frittella, S., Greco, G., Kurz, A., Palmigiano, A., & Sikimić, V. (2016a). Multi-type display calculus for dynamic epistemic logic. *Journal of Logic and Computation*, **26**(6), 2017–2065.

[35] Frittella, S., Greco, G., Kurz, A., Palmigiano, A., & Sikimić, V. (2016b). A proof-theoretic semantic analysis of dynamic epistemic logic. *Journal of Logic and Computation*, **26**(6), 1961–2015.

[36] Frittella, S., Greco, G., Palmigiano, A., & Yang, F. (2016). A multi-type calculus for inquisitive logic. In Väänänen, J., Hirvonen, Å., and de Queiroz, R., editors. *Logic, Language, Information, and Computation*. LNCS 9803. Berlin: Springer, pp. 215–233.

[37] Frittella, S., Palmigiano, A., & Santocanale, L. (2017). Dual characterizations for finite lattices via correspondence theory for monotone modal logic. *Journal of Logic and Computation*, **27**(3), 639–678.

[38] Gehrke, M. & Jónsson, B. (1994). Bounded distributive lattices with operators. *Mathematica Japonica*, **40**(2), 207–215.

[39] Gehrke, M. & Jónsson, B. (2004). Bounded distributive lattice expansions. *Mathematica Scandinavica*, **94**, 13–45.

[40] Gibbons, R. & Roberts, J. (2013). *The Handbook of Organizational Economics*. Princeton, NJ: Princeton University Press.

[41] Girard, J.-Y. (1995). Linear logic: Its syntax and semantics. In Girard, J.-Y., Lafont, Y., and Regnier, L., editors. *Advances in Linear Logic*. London Mathematical Society Lecture Note Series 222. Cambridge: Cambridge University Press, pp. 1–42.

[42] Greco, G., Kurz, A., & Palmigiano, A. (2013). Dynamic epistemic logic displayed. In Huang, H., Grossi, D., and Roy, O., editors. *Proceedings of the 4th International Workshop on Logic, Rationality and Interaction (LORI-4)*. LNCS, Vol. 8196. Berlin: Springer, pp. 135–148.

[43] Greco, G., Liang, F., Moshier, A., & Palmigiano, A. (2017). Multi-type display calculus for semi De Morgan logic. In Kennedy, J. and de Queiroz, R. J. G. B., editors. *Logic, Language, Information, and Computation*. LNCS 10388. Berlin: Springer-Verlag, pp. 199–215.

[44] Greco, G., Ma, M., Palmigiano, A., Tzimoulis, A., & Zhao, Z. (2016). Unified correspondence as a proof-theoretic tool. *Journal of Logic and Computation*, <https://doi.org/10.1093/logcom/exw022>.

[45] Greco, G. & Palmigiano, A. (2017). Lattice logic properly displayed. In Kennedy, J. and de Queiroz, R. J. G. B., editors. *Logic, Language, Information, and Computation*. LNCS 10388. Berlin: Springer-Verlag, pp. 153–169.

- [46] Greco, G. & Palmigiano, A. Linear logic properly displayed, preprint, arXiv: 1611.04181.
- [47] Haniková, Z. & Horčík, R. (2014). The finite embeddability property for residuated groupoids. *Algebra Universalis*, **72**(1), 1–13.
- [48] Harel, D., Kozen, D., & Tiuryn, J. (2000). *Dynamic Logic*. Cambridge, MA: MIT Press.
- [49] Hiroakira, O. & Komori, Y. (1985). Logics without the contraction rule. *The Journal of Symbolic Logic*, **50**(1), 169–201.
- [50] Kopylov, A. P. (2001). Decidability of linear affine logic. *Information and Computation*, **164**(1), 173–198.
- [51] Kurz, A. & Palmigiano, A. (2013). Epistemic updates on algebras. *Logical Methods in Computer Science*, **9**(4:17), 1–28.
- [52] le Roux, C. (2016). Correspondence Theory in Many-Valued Modal Logics. Master's Thesis. South Africa: University of Johannesburg.
- [53] Ma, M., Palmigiano, A., & Sadrzadeh, M. (2014). Algebraic semantics and model completeness for intuitionistic public announcement logic. *Annals of Pure and Applied Logic*, **165**(4), 963–995.
- [54] Ma, M. & Zhao, Z. (2017). Unified correspondence and proof theory for strict implication. *Journal of Logic and Computation*, **27**(3), 921–960.
- [55] Mahoney, J. T. (1995). The management of resources and the resource of management. *Journal of Business Research*, **33**(2), 91–101.
- [56] Marté-Oliet, N. & Meseguer, J. (1991). From Petri nets to linear logic. *Mathematical Structures in Computer Science*, **1**(1), 69–101.
- [57] Mol, J. M. & Wijnberg, N. M. (2011). From resources to value and back: Competition between and within organizations. *British Journal of Management*, **22**(1), 77–95.
- [58] Okada, M. & Terui, K. (1999). The finite model property for various fragments of intuitionistic linear logic. *The Journal of Symbolic Logic*, **64**(2), 790–802.
- [59] Ono, H. (1998). Decidability and finite model property of substructural logics. In Ginzburg, J., Khasidashvili, Z., Vogel, C.-W., Lévy, J.-J., and Vallduví, E., editors. *The Tbilisi Symposium on Logic, Language and Computation*. Stanford, CA: CSLI Publications, pp. 263–274.
- [60] Palmigiano, A., Sourabh, S., & Zhao, Z. (2017). Jónsson-style canonicity for ALBA-inequalities. *Journal of Logic and Computation*, **27**(3), 817–865.
- [61] Palmigiano, A., Sourabh, S., & Zhao, Z. (2017). Sahlqvist theory for impossible worlds. *Journal of Logic and Computation*, **27**(3), 775–816.
- [62] Plaza, J. (2007). Logics of public communications. *Synthese*, **158**(2), 165–179.
- [63] Pym, D. & Tofts, C. (2006). A calculus and logic of resources and processes. *Formal Aspects of Computing*, **18**(4), 495–517.
- [64] Pym, D. J., O'Hearn, P. W., & Yang, H. (2004). Possible worlds and resources: The semantics of BI. *Theoretical Computer Science*, **315**(1), 257–305.
- [65] Ricardo, D. (1891). *Principles of Political Economy and Taxation*. London: G. Bell and Sons.
- [66] Scott, W. G. (1961). Organization theory: An overview and an appraisal. *The Journal of the Academy of Management*, **4**(1), 7–26.
- [67] Segerberg, K. (1982). The logic of deliberate action. *Journal of Philosophical Logic*, **11**(2), 233–254.
- [68] Shamsie, J. & Mannor, M. J. (2013). Looking inside the dream team: Probing into the contributions of tacit knowledge as an organizational resource. *Organization Science*, **24**(2), 513–529.

[69] Troelstra, A. S. (1992). *Lectures on Linear Logic*. Lecture Notes, No. 29. Stanford, CA: CSLI Publications.

[70] Tsoukas, H. & Knudsen, C. (2005). *The Oxford Handbook of Organization Theory*. New York: Oxford University Press.

[71] van der Hoek, W., van Linder, B., & Meyer, J. J. C. (1994). A logic of capabilities. In Nerode, A., and Matiyasevich, Y. V., editors. *International Symposium on Logical Foundations of Computer Science*. Lecture Notes in Computer Science, Vol. 813. Berlin: Springer, pp. 366–378.

[72] Wansing, H. (1998). *Displaying Modal Logic*. Norwell, MA: Kluwer.

[73] Wernerfelt, B. (1984). A resource-based view of the firm. *Strategic Management Journal*, 5(2), 171–180.

DEPARTMENT OF LOGIC, FACULTY OF ARTS

CHARLES UNIVERSITY

PRAGUE, CZECH REPUBLIC

E-mail: marta.bilkova@ff.cuni.cz

DEPARTMENT OF LANGUAGES, LITERATURE AND COMMUNICATION

UNIVERSITY OF UTRECHT

UTRECHT, THE NETHERLANDS

E-mail: G.Greco@uu.nl

FACULTY OF TECHNOLOGY, POLICY AND MANAGEMENT

DELFT UNIVERSITY OF TECHNOLOGY

DELFT, THE NETHERLANDS

and

DEPARTMENT OF PURE AND APPLIED MATHEMATICS

UNIVERSITY OF JOHANNESBURG

JOHANNESBURG, SOUTH AFRICA

E-mail: A.Palmigiano@tudelft.nl

FACULTY OF TECHNOLOGY, POLICY AND MANAGEMENT

DELFT UNIVERSITY OF TECHNOLOGY

DELFT, THE NETHERLANDS

E-mail: A.Tzimoulis-1@tudelft.nl

FACULTY OF ECONOMICS AND BUSINESS

UNIVERSITY OF AMSTERDAM

AMSTERDAM, THE NETHERLANDS

and

COLLEGE OF BUSINESS AND ECONOMICS

UNIVERSITY OF JOHANNESBURG

JOHANNESBURG, SOUTH AFRICA

E-mail: N.M.Wijnberg@uva.nl