

EXCEPTIONAL SIMPLE REAL LIE ALGEBRAS \mathfrak{f}_4 AND \mathfrak{e}_6 VIA CONTACTIFICATIONS

PAWEŁ NUROWSKI 

Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668
Warszawa, Poland
(nurowski@fuw.edu.pl)

(Received 11 April 2023; revised 7 March 2024; accepted 7 March 2024;
first published online 3 July 2024)

Abstract In Cartan’s PhD thesis, there is a formula defining a certain rank 8 vector distribution in dimension 15, whose algebra of automorphism is the split real form of the simple exceptional complex Lie algebra \mathfrak{f}_4 . Cartan’s formula is written in the standard Cartesian coordinates in \mathbb{R}^{15} . In the present paper, we explain how to find analogous formulae for the flat models of any bracket generating distribution \mathcal{D} whose symbol algebra $\mathfrak{n}(\mathcal{D})$ is constant and 2-step graded, $\mathfrak{n}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$.

The formula is given in terms of a solution to a certain system of linear algebraic equations determined by two representations $(\rho, \mathfrak{n}_{-1})$ and $(\tau, \mathfrak{n}_{-2})$ of a Lie algebra \mathfrak{n}_{00} contained in the 0th order Tanaka prolongation \mathfrak{n}_0 of $\mathfrak{n}(\mathcal{D})$.

Numerous examples are provided, with particular emphasis put on the distributions with symmetries being real forms of simple exceptional Lie algebras \mathfrak{f}_4 and \mathfrak{e}_6 .

Contents

1	Introduction: the notion of a contactification	158
2	Magical equation for a contactification	161
3	Two-step filtered manifolds	163
4	Construction of contactifications with nice symmetries	166
5	Majorana spinor representations of $\mathfrak{so}(p, q)$	167
6	Application: Obtaining the flat model for (3,6) distributions	170
7	Application: Obtaining Biquard’s 7-dimensional flat quaternionic contact manifold via contactification using spin representations of $\mathfrak{so}(1, 2)$ and $\mathfrak{so}(3, 0)$	171
8	Application: Obtaining the exceptionals from contactifications of spin representations; the \mathfrak{f}_4 case	173
	8.1. Cartan’s realisation of \mathfrak{f}_I	174

The research was funded from the Norwegian Financial Mechanism 2014–2021 with project registration number 2019/34/H/ST1/00636.

Key words and phrases: contactifications; exceptional simple Lie algebras; real forms

2020 Mathematics subject classification: 17B05; 17B10; 17B60; 17B70; 53A35; 53A55; 57M50

© The Author(s), 2024. Published by Cambridge University Press.

8.2. Realisation of \mathfrak{f}_{II}	180
9 Spinorial representations in dimension 8	183
9.1. Triality representations of $\mathfrak{so}(4,4)$	184
9.2. Triality representations of $\mathfrak{so}(8,0)$	187
10 Application: 2-step graded realisations of real forms of the exceptional Lie algebra \mathfrak{e}_6	190
10.1. Realisations of \mathfrak{e}_I and \mathfrak{e}_{IV} : generalities	191
10.2. An explicit realisation of \mathfrak{e}_I in dimension 24	193
10.3. An explicit realisation of \mathfrak{e}_{IV} in dimension 24	196
11 Application: one more realisation of \mathfrak{e}_6 and a realisation of \mathfrak{b}_6	197
11.1. Realisation of \mathfrak{e}_I in dimension 25	198
11.2. A realisation of $\mathfrak{so}(7,6)$ in dimension 21	199

1. Introduction: the notion of a contactification

A *contact structure* (M, \mathcal{D}) on a $(2n + 1)$ dimensional real manifold M is usually defined in terms of a 1-form λ on M such that

$$\underbrace{d\lambda \wedge d\lambda \wedge \cdots \wedge d\lambda}_{n \text{ times}} \wedge \lambda \neq 0$$

at each point $x \in M$. Given such a 1-form, the contact structure (M, \mathcal{D}) on M is the rank $s = 2n$ *vector distribution*

$$\mathcal{D} = \{X \in TM \text{ s.t. } X \lrcorner \lambda = 0\}.$$

Note that any $\lambda' = a\lambda$, with a being a nonvanishing function on M , defines the same contact structure (M, \mathcal{D}) . We also note that given a contact structure (M, \mathcal{D}) , we additionally have a family of 2-forms on M

$$\omega' = a\omega + \mu \wedge \lambda, \quad \text{with } \omega = d\lambda,$$

where $a \neq 0$ is a function, and μ is a 1-form on M . This, in particular, means that given a contact structure (M, \mathcal{D}) , we have a rank $s = 2n$ (bracket generating) distribution \mathcal{D} , and a *line* of a closed 2-form ω *in the distribution* \mathcal{D} , with

$$d\omega = 0 \quad \& \quad \underbrace{\omega \wedge \omega \wedge \cdots \wedge \omega}_{n \text{ times}} \neq 0.$$

This can be compared with the notion of a *symplectic* structure $(N, [\omega])$ on a $s = 2n$ dimensional real manifold N . Such a structure is defined in terms of a line $\omega' = h\omega$ of a nowhere vanishing 2-form ω on N , such that

$$d\omega = 0 \quad \& \quad \underbrace{\omega \wedge \omega \wedge \cdots \wedge \omega}_{n \text{ times}} \neq 0.$$

Here, contrary to the contact case, we have a *line* of a closed 2-form ω *in the tangent space* TN rather than in the proper vector subbundle $\mathcal{D} \subsetneq TN$.

By the *Poincaré lemma*, locally, in an open set $\mathcal{O} \subset N$, the form ω defines a 1-form Λ on N such $d\Lambda = \omega$. Therefore, given a symplectic structure $(N, [\omega])$, we can locally *contactify* it by considering a $(2n + 1)$ dimensional manifold

$$\mathcal{U} = \mathbb{R} \times \mathcal{O} \xrightarrow{\pi} \mathcal{O},$$

with a 1-form

$$\lambda = du + \pi^*(\Lambda)$$

on \mathcal{U} ; here, the real variable u is a coordinate along the \mathbb{R} factor in $\mathcal{U} = \mathbb{R} \times \mathcal{O}$. As a result, the structure $(M, \mathcal{D}) = (\mathcal{U}, \ker(\lambda))$ is a *contact structure*, called a *contact structure associated with the symplectic structure* $(N, [\omega])$.

We introduce the notion of a *contactification* as a generalisation of the above considerations.

Definition 1.1. Let N be an s -dimensional manifold and let $d\mathcal{D}^\perp := \text{Span}(\omega^1, \omega^2, \dots, \omega^r)$ be a rank r subbundle of $\bigwedge^2 N$. Consider an $(s+r)$ -dimensional fiber bundle $F \rightarrow M \xrightarrow{\pi} N$ over N . Let (X_1, X_2, \dots, X_r) be a coframe of *vertical vectors* in M . In particular, we have $\pi_*(X_i) = 0$ for all $i = 1, 2, \dots, r$.

Let us assume that on M there exist r one-forms λ^i , $i = 1, 2, \dots, r$, such that $\det(X_i \lrcorner \lambda^j) \neq 0$ on M , and that $d\lambda^i = \sum_{j=1}^r a^i_j \pi^*(\omega^j) + \sum_{j=1}^r \mu^i_j \wedge \lambda^j$ for all $i = 1, 2, \dots, r$, with some 1-forms μ^i_j and some functions a^i_j on M satisfying $\det(a^i_j) \neq 0$. Consider the corresponding rank s distribution $\mathcal{D} = \{TM \ni X \mid X \lrcorner \lambda^i = 0, i = 1, 2, \dots, r\}$ on M .

Then the pair (M, \mathcal{D}) is called a *contactification* of the pair $(N, d\mathcal{D}^\perp)$.

Definition 1.2. A real Lie algebra \mathfrak{g} spanned over \mathbb{R} by the vector fields Y on M of the contactification (M, \mathcal{D}) satisfying

$$\mathcal{L}_Y \lambda^i \wedge \lambda^1 \wedge \dots \wedge \lambda^r = 0, \quad \forall i = 1, 2, \dots, r \tag{1.1}$$

is called the Lie algebra of infinitesimal symmetries of the contactification (M, \mathcal{D}) . By definition, it is the same as the Lie algebra of infinitesimal symmetries of the distribution \mathcal{D} on M . The vector fields Y on (M, \mathcal{D}) satisfying (1.1) are called infinitesimal symmetries of (M, \mathcal{D}) , or of \mathcal{D} , for short.

Below, we give a nontrivial example of the notions included in Definitions 1.1 and 1.2.

Example 1.3. Consider $N = \mathbb{R}^8$ with Cartesian coordinates $(x^1, x^2, x^3, x^4, x^5, x^6, x^7, x^8)$ and a space $d\mathcal{D}^\perp = \text{Span}(\omega^1, \omega^2, \omega^3, \omega^4, \omega^5, \omega^6, \omega^7) \subset \bigwedge^2 N$, which is spanned by the following seven 2-forms on N :

$$\begin{aligned} \omega^1 &= dx^1 \wedge dx^8 + dx^2 \wedge dx^5 + dx^3 \wedge dx^7 + dx^4 \wedge dx^6 \\ \omega^2 &= -dx^1 \wedge dx^5 + dx^2 \wedge dx^8 + dx^3 \wedge dx^6 - dx^4 \wedge dx^7 \\ \omega^3 &= -dx^1 \wedge dx^7 - dx^2 \wedge dx^6 + dx^3 \wedge dx^8 + dx^4 \wedge dx^5 \end{aligned}$$

$$\begin{aligned} \omega^4 &= dx^1 \wedge dx^2 + dx^3 \wedge dx^4 + dx^5 \wedge dx^8 + dx^6 \wedge dx^7 \\ \omega^5 &= -dx^1 \wedge dx^6 + dx^2 \wedge dx^7 - dx^3 \wedge dx^5 + dx^4 \wedge dx^8 \\ \omega^6 &= dx^1 \wedge dx^4 + dx^2 \wedge dx^3 - dx^5 \wedge dx^7 + dx^6 \wedge dx^8 \\ \omega^7 &= dx^1 \wedge dx^3 - dx^2 \wedge dx^4 + dx^5 \wedge dx^6 + dx^7 \wedge dx^8. \end{aligned}$$

As the bundle N , take $M = \mathbb{R}^7 \times \mathbb{R}^8 \rightarrow N$ with coordinates $(x^1, \dots, x^8, x^9, \dots, x^{15})$, and take seven 1-forms

$$\begin{aligned} \lambda^1 &= dx^9 + x^1 dx^8 + x^2 dx^5 + x^3 dx^7 + x^4 dx^6 \\ \lambda^2 &= dx^{10} - x^1 dx^5 + x^2 dx^8 + x^3 dx^6 - x^4 dx^7 \\ \lambda^3 &= dx^{11} - x^1 dx^7 - x^2 dx^6 + x^3 dx^8 + x^4 dx^5 \\ \lambda^4 &= dx^{12} + x^1 dx^2 + x^3 dx^4 + x^5 dx^8 + x^6 dx^7 \\ \lambda^5 &= dx^{13} - x^1 dx^6 + x^2 dx^7 - x^3 dx^5 + x^4 dx^8 \\ \lambda^6 &= dx^{14} + x^1 dx^4 + x^2 dx^3 - x^5 dx^7 + x^6 dx^8 \\ \lambda^7 &= dx^{15} + x^1 dx^3 - x^2 dx^4 + x^5 dx^6 + x^7 dx^8. \end{aligned}$$

This defines a rank 8 distribution $\mathcal{D} = \{TM \ni X \mid X \lrcorner \lambda^i = 0, i = 1, 2, \dots, 7\}$ on M . The pair (M, \mathcal{D}) is a contactification of $(N, d\mathcal{D}^\perp)$ since $X_i = \partial_{i+8}$, $\det(X_i \lrcorner \lambda^j) = 1$, and $d\lambda^i = \omega^i$ for all $i = 1, \dots, 7$. In particular, in this example, the rank 8 distribution \mathcal{D} gives a 2-step filtration $\mathcal{D}_{-1} \subset \mathcal{D}_{-2} = TM$, where $\mathcal{D}_{-1} = \mathcal{D}$ and $\mathcal{D}_{-2} = [\mathcal{D}_{-1}, \mathcal{D}_{-1}] = TM$.

This example is essentially taken from Élie Cartan’s PhD thesis [8] – actually its German version. We took it as our example inspired by the following quote from Sigurdur Helgason [13]:

Cartan represented [the simple exceptional Lie group] \mathbf{F}_4 (...) by the Pfaffian system in \mathbb{R}^{15} (...). Similar results for \mathbf{E}_6 in \mathbb{R}^{16} , \mathbf{E}_7 in \mathbb{R}^{27} and \mathbf{E}_8 in \mathbb{R}^{29} are indicated in [8]. Unfortunately, detailed proofs of these remarkable representations of the exceptional groups do not seem to be available.

The 15-dimensional contactification (M, \mathcal{D}) from our Example 1.3 is obtained in terms of the seven 1-forms λ^i , which are equivalent to the seven forms from the Cartan Pfaffian system in dimension 15 mentioned by Helgason. In particular, it follows that the distribution structure (M, \mathcal{D}) has the simple exceptional Lie group \mathbf{F}_4 , actually its real form F_I in the terminology of [11], as a group of automorphism.

In this paper, we will explain how one gets this realisation of the exceptional Lie group \mathbf{F}_4 , a realisation of its real form F_{II} , and realisations of the two (out of 5) real forms E_I and E_{IV} of the complex simple exceptional Lie group \mathbf{E}_6 . For this explanation, we need some preparations consisting of recalling few notions associated with vector distributions on manifolds and spinorial representations of the orthogonal groups in space of real spinors.

Finally, we note that our approach in this paper is purely utilitarian. We answer the question: *How to get the explicit formulas in Cartesian coordinates for Pfaffian forms $(\lambda^1, \dots, \lambda^r)$, which have simple Lie algebras as symmetries?* One can study more general problems related to this on purely Lie theoretical ground. For example, one can ask when a 2-step graded nilpotent Lie algebra $\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$ has a given Lie algebra

\mathfrak{n}_{00} as a part of its Lie algebra \mathfrak{n}_0 of derivations preserving the strata, or a question as to when the Tanaka prolongation of such \mathfrak{n}_- with $\mathfrak{n}_{00} \subset \mathfrak{n}_0$ is finite, or simple. This is beyond the scope of our paper. A reader interested in such problems may consult, for example, [3, 4, 16]. We also mention that since in this paper we *only* consider distributions \mathcal{D} with 2-step filtrations $\mathcal{D} = \mathcal{D}_{-1} \subset \mathcal{D}_{-2} = TM$ and without any additional geometric structure intentionally imposed on \mathcal{D} , the realisations of the real forms of the exceptional simple Lie group G_2 will *not* appear. These can be found, for example, in [1, 6, 10, 17].

2. Magical equation for a contactification

The purpose of this section is to prove the following crucial lemma about a certain algebraic equation, which we call a *magical equation*. It is the boxed equation (2.1) below.

Lemma 2.1. *Let $(\mathfrak{n}_{00}, [\cdot, \cdot]_0)$ be a finite dimensional Lie algebra, and let $\rho : \mathfrak{n}_{00} \xrightarrow{\text{hom}} \text{End}(S)$ be its finite dimensional representation in a real vector space S of dimension s . In addition, let R be an r -dimensional real vector space, and $\tau : \mathfrak{n}_{00} \rightarrow \text{End}(R)$, be a linear map. Finally, let ω be a linear map $\omega : \bigwedge^2 S \rightarrow R$, or what is the same, let $\omega \in \text{Hom}(\bigwedge^2 S, R)$.*

Suppose now that the triple (ρ, ω, τ) satisfies the following equation:

$$\boxed{\omega(\rho(A)X, Y) + \omega(X, \rho(A)Y) = \tau(A)\omega(X, Y),} \tag{2.1}$$

for all $A \in \mathfrak{n}_{00}$ and all $X, Y \in S$. Then we have the following:

(1) *The map τ satisfies*

$$(\tau([A, B]_0) - [\tau(A), \tau(B)]_{\text{End}(R)})\omega = 0 \quad \forall A, B \in \mathfrak{n}_{00}.$$

(2) *If the map $\tau : \mathfrak{n}_{00} \rightarrow \text{End}(R)$ is a representation of \mathfrak{n}_{00} , that is, if*

$$\tau([A, B]_0) = [\tau(A), \tau(B)]_{\text{End}(R)},$$

then the real vector space $\mathfrak{g}_0 := R \oplus S \oplus \mathfrak{n}_{00}$ is a graded Lie algebra

$$\mathfrak{g}_0 = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_{00},$$

with the graded components

$$\mathfrak{n}_{-2} = R, \quad \mathfrak{n}_{-1} = S, \quad \text{with } \mathfrak{n}_{00} \text{ as the 0 grade,}$$

and with the Lie bracket $[\cdot, \cdot]$ given by

- (a) *if $X, Y \in \mathfrak{n}_{00}$, then $[X, Y] = [X, Y]_0$,*
- (b) *if $A \in \mathfrak{n}_{00}$, $X \in \mathfrak{n}_{-1}$, then $[A, X] = \rho(A)X$,*
- (c) *if $A \in \mathfrak{n}_{00}$, $X \in \mathfrak{n}_{-2}$, then $[A, X] = \tau(A)X$,*
- (d) $[\mathfrak{n}_{-1}, \mathfrak{n}_{-2}] = [\mathfrak{n}_{-2}, \mathfrak{n}_{-2}] = \{0\}$,
- (e) *and, if $X, Y \in \mathfrak{n}_{-1}$, then $[X, Y] = \omega(X, Y)$.*

(3) *Moreover, in the case (2), the Lie subalgebra*

$$\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$$

of \mathfrak{g}_0 is a 2-step graded Lie algebra, and the algebra \mathfrak{n}_{00} is a Lie subalgebra of the Lie algebra

$$\mathfrak{n}_0 = \{ \text{Der}(\mathfrak{n}_-) \ni D \text{ s.t. } D\mathfrak{n}_j \subset \mathfrak{n}_j \text{ for } j = -1, -2 \}$$

of all derivations of \mathfrak{n}_- preserving its strata \mathfrak{n}_{-1} and \mathfrak{n}_{-2} .

Remark 2.2. Note that, in the respective bases $\{f_\mu\}_{\mu=1}^s$ in S and $\{e_i\}_{i=1}^r$ in R , the equation (2.1) is

$$\boxed{\rho(A)^\alpha_\mu \omega^i_{\alpha\nu} + \rho(A)^\alpha_\nu \omega^i_{\mu\alpha} = \tau(A)^i_j \omega^i_{\mu\nu}} \tag{2.2}$$

for all $A \in \mathfrak{n}_{00}$, all $i = 1, 2, \dots, r$ and all $\mu, \nu = 1, 2, \dots, s$. On this basis, the condition (1) is

$$(\tau([A, B]_0) - [\tau(A), \tau(B)]_{\text{End}(R)})^i_j \omega^j_{\mu\nu} = 0$$

for all $i = 1, 2, \dots, r, \mu, \nu = 1, 2, \dots, s$, and $A, B \in \mathfrak{n}_{00}$.

Proof of the lemma. The proof of part (1) is a pure calculation using the equation (2.1). We first rewrite it in the shorthand notation as

$$\rho(A)\omega + \omega\rho(A)^T = \tau(A)\omega, \quad \forall A \in \mathfrak{n}_{00}.$$

Then we have

$$\begin{aligned} \tau([A, B]_0)\omega &= \rho([A, B]_0)\omega + \omega\rho([A, B]_0)^T \\ &= \rho(A)\rho(B)\omega - \rho(B)\rho(A)\omega + \omega\rho(B)^T\rho(A)^T - \omega\rho(A)^T\rho(B)^T \\ &= \rho(A)\left(\tau(B)\omega - \omega\rho(B)^T\right) - \rho(B)\left(\tau(A)\omega - \omega\rho(A)^T\right) \\ &\quad + \left(\tau(B)\omega - \rho(B)\omega\right)\rho(A)^T - \left(\tau(A)\omega - \rho(A)\omega\right)\rho(B)^T \\ &= \rho(A)\left(\tau(B)\omega\right) - \rho(B)\left(\tau(A)\omega\right) + \left(\tau(B)\omega\right)\rho(A)^T - \left(\tau(A)\omega\right)\rho(B)^T \\ &= \tau(A)\tau(B)\omega - \tau(B)\omega\rho(A)^T - \left(\tau(B)\tau(A)\omega - \tau(A)\omega\rho(B)^T\right) \\ &\quad + \tau(B)\omega\rho(A)^T - \tau(A)\omega\rho(B)^T = \tau(A)\tau(B)\omega - \tau(B)\tau(A)\omega \\ &= ([\tau(A), \tau(B)]_{\text{End}(R)})\omega, \end{aligned}$$

which proves part (1).

The proof of parts (2) and (3) is as follows:

We need to check the Jacobi identity for the bracket $[\cdot, \cdot]$.

We first consider the representation

$$\sigma = \tau \oplus \rho \quad \text{of} \quad \mathfrak{n}_{00} \quad \text{in} \quad \mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1},$$

defined by

$$\sigma(A)(Y \oplus X) = \tau(A)Y \oplus \rho(A)X, \quad \forall A \in \mathfrak{n}_{00}, X \in \mathfrak{n}_{-1}, Y \in \mathfrak{n}_{-2}.$$

We then prove that the representation σ is a *strata preserving derivation* in \mathfrak{n}_- . This is implied by the definitions (a)–(e) of the bracket, and the fundamental equation (2.1) as follows:

The strata preserving property of $\sigma, \sigma(\mathfrak{n}_{-i}) \subset \mathfrak{n}_{-i}, i = 1, 2$, is obvious by the definitions of ρ and τ . However, we need to check that σ is a derivation – that is, that

$$\sigma(A)[X, Y] = [\sigma(A)X, Y] + [X, \sigma(A)Y] \tag{2.3}$$

for all $A \in \mathfrak{n}_{00}$ and for all $X, Y \in \mathfrak{n}_{-}$. Because of the strata preserving property of σ , which we have just established, and because of the point (d) of the definition of the bracket, the equation (2.3) is satisfied when both X and Y are in \mathfrak{n}_{-2} , or when X is in \mathfrak{n}_{-1} and Y is \mathfrak{n}_{-2} . The only thing to be checked is if (2.3) is also valid when both X and Y belong to \mathfrak{n}_{-1} . But this just follows directly from (2.1) since if $X, Y \in \mathfrak{n}_{-1}$, then

$$\begin{aligned} \sigma(A)[X, Y] &= \sigma(A)\omega(X, Y) = \tau(A)\omega(X, Y) = \\ &= \omega(\rho(A)X, Y) + \omega(X, \rho(A)Y) = [\rho(A)X, Y] + [X, \rho(A)Y] = \\ &= [\sigma(A)X, Y] + [X, \sigma(A)Y], \quad \forall A \in \mathfrak{n}_{00}. \end{aligned}$$

Now we return to checking the Jacobi identity for the bracket $[\cdot, \cdot]$ in \mathfrak{g}_0 :
On elements of the form $A, B \in \mathfrak{n}_{00}, Z \in \mathfrak{n}_{-}$, by (b)-(c), we have

$$[[A, B], Z] + [[Z, A], B] + [[B, Z], A] = \left(\sigma([A, B]) - [\sigma(A), \sigma(B)] \right) Z,$$

which vanishes due to the representation property of σ . However, on elements $A \in \mathfrak{n}_{00}$ and $Z_1, Z_2 \in \mathfrak{n}_{-}$, we have

$$[[A, Z_1], Z_2] + [[Z_2, A], Z_1] + [[Z_1, Z_2], A] = [\sigma(A)Z_1, Z_2] + [Z_1, \sigma(A)Z_2] - \sigma(A)[Z_1, Z_2],$$

which is again zero, on the ground of the derivation property 2.3 of σ . Obviously, the bracket satisfies the Jacobi identity when it is restricted to \mathfrak{n}_{00} ; it is the Lie bracket $[\cdot, \cdot]_o$ of the Lie algebra \mathfrak{n}_{00} . Finally, property (2) implies that $[[Z_1, Z_2], Z_3] = 0$ for all Z_1, Z_2, Z_3 in \mathfrak{n}_{-} ; hence, the Jacobi identity is trivially satisfied for $[\cdot, \cdot]$, when it is restricted to \mathfrak{n}_{-} . \square

In the following, we will use the map $\omega \in \text{Hom}(\wedge^2 S, R)$ satisfying the *magical equation* (2.1) to construct contactifications with nontrivial symmetry algebras \mathfrak{g} . The setting will include Cartan’s contactification with symmetry \mathbf{F}_4 mentioned in the Helgason’s quote. For this, however, we need few preparations.

3. Two-step filtered manifolds

A *2-step filtered structure* on an $(s+r)$ -dimensional manifold M is a pair (M, \mathcal{D}) , in which \mathcal{D} is a vector distribution of rank s on M , such that it is *bracket generating* in the quickest possible way. This means that its *derived distribution* $\mathcal{D}_{-2} := [\mathcal{D}_{-1}, \mathcal{D}_{-1}]$, with $\mathcal{D}_{-1} = \mathcal{D}$, is such that

$$\mathcal{D}_{-2} = TM.$$

It provides the simplest nontrivial *filtration*

$$TM = \mathcal{D}_{-2} \supset \mathcal{D}_{-1}$$

of the tangent bundle TM .

A (local) *automorphism* of a 2-step filtered manifold (M, \mathcal{D}) is a (local) diffeomorphism $\phi: M \rightarrow M$ such that $\phi_*\mathcal{D} \subset \mathcal{D}$. Since automorphisms can be composed and have inverses, they form a *group* G of (local) automorphisms of (M, \mathcal{D}) , also called a *group of (local) symmetries of \mathcal{D}* . Infinitesimally, the Lie group of automorphisms defines the *Lie algebra $\mathfrak{aut}(\mathcal{D})$ of symmetries*, which is the real span of all vector fields X on M such that $[X, Y] \subset \mathcal{D}$ for all $Y \in \mathcal{D}$.

Among all the 2-step filtered manifolds (M, \mathcal{D}) particularly simple are those which can be realised on a group manifold of a *2-step nilpotent* Lie group. These are related to the notion of the *nilpotent approximation* of a pair (M, \mathcal{D}) . This is defined as follows:

At every point $x \in M$ equipped with a 2-step filtration $\mathcal{D}_{-2} \supset \mathcal{D}_{-1}$, we have well-defined vector spaces $\mathfrak{n}_{-1}(x) = \mathcal{D}_{-1}(x)$ and $\mathfrak{n}_{-2}(x) = \mathcal{D}_{-2}(x)/\mathcal{D}_{-1}(x)$, which define a vector space

$$\mathfrak{n}(x) = \mathfrak{n}_{-2}(x) \oplus \mathfrak{n}_{-1}(x).$$

This vector space is naturally a *Lie algebra*, with a *Lie bracket* induced from the Lie bracket of vector fields in TM . Due to the 2-step property of the filtration defined by \mathcal{D} , this Lie algebra is *2-step nilpotent*,

$$[\mathfrak{n}_{-1}(x), \mathfrak{n}_{-1}(x)] = \mathfrak{n}_{-2}(x) \quad \& \quad [\mathfrak{n}_{-1}(x), \mathfrak{n}_{-2}(x)] = \{0\}.$$

This 2-step nilpotent Lie algebra is a *local invariant* of the structure (M, \mathcal{D}) , and it is called a *nilpotent approximation of the structure (M, \mathcal{D}) at $x \in M$* .

This enables for defining a class of particularly simple examples of 2-step filtered structures:

Consider a *2-step nilpotent Lie algebra $\mathfrak{n} = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$* , and let M be a Lie group, whose Lie algebra is \mathfrak{n} . The Lie algebra \mathfrak{n}_M of left invariant vector fields on M is isomorphic to \mathfrak{n} and mirrors its gradation, $\mathfrak{n}_M = \mathfrak{n}_{M-2} \oplus \mathfrak{n}_{M-1}$. Now, taking all linear combinations with *smooth functions* coefficients of all vector fields from the graded component \mathfrak{n}_{M-1} of \mathfrak{n}_M , one defines a *vector distribution $\mathcal{D} = \text{Span}_{\mathcal{F}(M)}(\mathfrak{n}_M)$* on M . The so constructed filtered structure (M, \mathcal{D}) is obviously 2-step graded and is the *simplest* filtered structure with nilpotent approximation being equal to \mathfrak{n} everywhere. We call this (M, \mathcal{D}) structure the *flat model* for all the 2-step filtered structures having the same constant nilpotent approximation \mathfrak{n} .

It is remarkable that the largest possible symmetry of all 2-step filtered structures (M, \mathcal{D}) is precisely the symmetry of the flat model. As such, it is *algebraically* determined by the nilpotent approximation \mathfrak{n} . This is the result of Noboru Tanaka [18]. To describe it, we recall the notion of *Tanaka prolongation*.

Definition 3.1. The *Tanaka prolongation* of a 2-step nilpotent Lie algebra \mathfrak{n} is a graded Lie algebra $\mathfrak{g}(\mathfrak{n})$ given by a direct sum

$$\mathfrak{g}(\mathfrak{n}) = \mathfrak{n} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \dots \oplus \mathfrak{n}_j \oplus \dots, \tag{3.1}$$

with

$$\mathfrak{n}_k = \left\{ \bigoplus_{j < 0} \mathfrak{n}_{k+j} \otimes \mathfrak{n}_j^* \ni A \text{ s.t. } A[X, Y] = [AX, Y] + [X, AY] \right\} \tag{3.2}$$

for each $k \geq 0$.

Furthermore, for each $j \geq 0$, the Lie algebra

$$\mathfrak{g}_j(\mathfrak{n}) = \mathfrak{n} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \cdots \oplus \mathfrak{n}_j$$

is called the Tanaka prolongations of \mathfrak{n} up to j^{th} order.

Setting $[A, X] = AX$ for all $A \in \mathfrak{n}_k$ with $k \geq 0$ and for all $X \in \mathfrak{n}$ makes the condition in (3.2) into the Jacobi identity. Moreover, if $A \in \mathfrak{n}_k$ and $B \in \mathfrak{n}_l$, $k, l \geq 0$, then their commutator $[A, B] \in \mathfrak{n}_{k+l}$ is defined on elements $X \in \mathfrak{n}$ inductively, according to the Jacobi identity. By this, we mean that it should satisfy

$$[A, B]X = [A, BX] - [B, AX],$$

which is sufficient enough to define $[A, B]$.

Remark 3.2. Note, in particular, that \mathfrak{n}_0 is the Lie algebra of all derivations of \mathfrak{n} preserving the two strata \mathfrak{n}_{-1} and \mathfrak{n}_{-2} of the direct sum $\mathfrak{n} = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$:

$$\mathfrak{n}_0 = \{ \text{Der}(\mathfrak{n}) \ni D \text{ s.t. } D\mathfrak{n}_j \subset \mathfrak{n}_j \text{ for } j = -1, -2 \}.$$

Although the Tanaka prolongation of a nilpotent Lie algebra \mathfrak{n} is in general infinite, in this paper, we will be interested in situations when the Tanaka prolongation

$$\mathfrak{g} = \mathfrak{g}(\mathfrak{n})$$

of the 2-step nilpotent part

$$\mathfrak{n} = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$$

is finite and symmetric, in the sense

$$\mathfrak{g}(\mathfrak{n}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with

$$\dim(\mathfrak{n}_{-k}) = \dim(\mathfrak{n}_k), \quad k = 1, 2.$$

Such situations are possible, and in them the so defined Lie algebra $\mathfrak{g}(\mathfrak{n})$ is simple. In such a case, the Tanaka prolongation $\mathfrak{g}(\mathfrak{n})$ is graded, and the subalgebra

$$\mathfrak{p} = \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2$$

in such $\mathfrak{g}(\mathfrak{n})$ is parabolic. Moreover, the Lie algebra

$$\mathfrak{p}_{opp} = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0$$

is also a parabolic subalgebra of this simple $\mathfrak{g}(\mathfrak{n})$. It is isomorphic to \mathfrak{p} , $\mathfrak{p} \simeq \mathfrak{p}_{opp}$.

Regardless of whether $\mathfrak{g}(\mathfrak{n})$ is finite or not, we have the following general theorem, which is a specialisation of a remarkable theorem by Noboru Tanaka [18]:

Theorem 3.3. Consider 2-step filtered structures (M, \mathcal{D}) , with distributions \mathcal{D} having the same constant nilpotent approximation \mathfrak{n} . Then

- The most symmetric of all of these distribution structures is the flat model (M, \mathcal{D}) , with M being a nilpotent Lie group associated of the nilpotent approximation algebra \mathfrak{n} , and with \mathcal{D} being the first component \mathcal{D}^{-1} of the natural filtration on M associated to the 2-step grading in \mathfrak{n} .
- The Lie algebra of automorphism $\text{aut}(\mathcal{D})$ of the flat model structure is isomorphic to the Tanaka prolongation $\mathfrak{g}(\mathfrak{n})$ of the nilpotent approximation \mathfrak{n} , $\text{aut}(\mathcal{D}) \simeq \mathfrak{g}(\mathfrak{n})$.

Remark 3.4. This theorem is of fundamental importance for explanation of the Cartan’s result about a realisation of \mathbf{F}_4 in \mathbb{R}^{15} . As we will see, Cartan’s \mathbb{R}^{15} is actually a domain of a chart (U, φ) on a certain 2-step nilpotent Lie group M , with a 2-step nilpotent Lie algebra \mathfrak{n} , and the equivalent description of \mathbf{F}_4 in terms of a symmetry group of the contactification (M, \mathcal{D}) from our Example 1.3 is valid because this contactification is just the flat model for the 2-step filtration (M, \mathcal{D}) with the nilpotent approximation \mathfrak{n} .

Using the information about the Tanaka prolongation of a nilpotent Lie algebra \mathfrak{n} , we can enlarge our Lemma 2.1 by changing its point (3) into the following more complete form:

Lemma 3.5. *With all the assumptions of Lemma 2.1, and with points (1) and (2) as in Lemma 2.1, its point (3) is equivalent to the following:*

(3) Moreover, in the case (2), the Lie subalgebra

$$\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$$

of

$$\mathfrak{g}_0 = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_{00}$$

is a 2-step graded nilpotent Lie algebra, and the algebra \mathfrak{n}_{00} is a Lie subalgebra of the Tanaka prolongation up to 0th order $\mathfrak{g}_0(\mathfrak{n}_-)$ of the Lie algebra $\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$.

Remark 3.6. The term ‘... \mathfrak{n}_{00} is a Lie subalgebra of the Tanaka prolongation up to 0th order $\mathfrak{g}_0(\mathfrak{n}_-)$ of the Lie algebra $\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1}$.’ in the above lemma means that \mathfrak{n}_{00} , although nontrivial, is in general only a subalgebra of the

$$\mathfrak{n}_0 = \{ \text{Der}(\mathfrak{n}_-) \ni D \text{ s.t. } D\mathfrak{n}_j \subset \mathfrak{n}_j \text{ for } j = -1, -2 \}, \quad \mathfrak{n}_{00} \subsetneq \mathfrak{n}_0,$$

which is the full 0 graded component of the Tanaka prolongation of \mathfrak{n}_- . So for applications, it is reasonable to choose \mathfrak{n}_{00} as large as possible.

4. Construction of contactifications with nice symmetries

Consider a Lie algebra $(\mathfrak{n}_{00}, [\cdot, \cdot]_0)$ and its two real representations (ρ, S) , (τ, R) , in the respective real s - and r -dimensional vector spaces S and R . Let $S = \mathbb{R}^s$, $R = \mathbb{R}^r$, and let $\{f_\mu\}_{\mu=1}^s$ and $\{e_i\}_{i=1}^r$ be respective bases in S and in R . Let $\{f^\mu\}_{\mu=1}^s$ be a basis in the vector space S^* dual to the basis $\{f_\mu\}_{\mu=1}^s$, $f_\nu \lrcorner f^\mu = \delta_\nu^\mu$. To be in a situation of Lemma 2.1, we also assume that we have the homomorphism $\omega \in \text{Hom}(\bigwedge^2 S, R)$ satisfying the magical equation (2.1).

Then the map ω is

$$\omega = \frac{1}{2}\omega^i{}_{\mu\nu}e_i \otimes f^\mu \wedge f^\nu,$$

and it defines the coefficients $\omega^i{}_{\mu\nu}$, $i = 1, \dots, r$, $\mu, \nu = 1, 2, \dots, s$, which satisfy $\omega^i{}_{\mu\nu} = -\omega^i{}_{\nu\mu}$.

Now, consider an s -dimensional manifold, which is an open set N of \mathbb{R}^s , $N \subset \mathbb{R}^s$, with coordinates $(x^\mu)_{\mu=1}^r$. Then we have r two-forms $(\omega^i)_{i=1}^r$ on N , defined by

$$\omega^i = \frac{1}{2}\omega^i{}_{\mu\nu}dx^\mu \wedge dx^\nu.$$

This produces an $(N, d\mathcal{D}^\perp)$ structure on N , with

$$d\mathcal{D}^\perp = \text{Span}_{\mathbb{R}}(\omega^1, \dots, \omega^r).$$

We contactify it. For this, we take a local $M = \mathbb{R}^r \times N$, with coordinates $(u^i, x^\mu)_{i=1}^r,_{\mu=1}^s$, and define the ‘contact forms’ on M by

$$\lambda^i = du^i + \omega^i{}_{\mu\nu}x^\mu dx^\nu.$$

Because of Lemmas 2.1 and 3.5, the distribution \mathcal{D} on M defined by this contactification as in Definition 1.1 equips M with a 2-step filtered structure having $\mathcal{D}_{-1} = \mathcal{D}$. This has rank s . Now using Lemmas 2.1 and 3.5, and Tanaka’s Theorem 3.3, we get the following corollary.

Corollary 4.1. *Let $M = \mathbb{R}^r \times \mathbb{R}^s$ and let*

$$\lambda^i = du^i + \omega^i{}_{\mu\nu}x^\mu dx^\nu, \quad i = 1, \dots, r,$$

with ω being a solution of the magical equation 2.1 such that $\text{Im}(\omega) = R$. Consider the distribution structure (M, \mathcal{D}) with a rank r distribution

$$\mathcal{D} = \{TM \ni X, \text{ s.t. } X \lrcorner \lambda^i = 0, i = 1, \dots, r\}$$

on M . Then the Lie algebra of automorphism $\text{aut}(\mathcal{D})$ of (M, \mathcal{D}) is isomorphic to the Tanaka prolongation of the 2-step nilpotent Lie algebra $\mathfrak{n}_- = R \oplus S$ defined in point (3) of Lemma 2.1 or 3.5. The Lie algebra $\mathfrak{g}_0 = R \oplus S \oplus \mathfrak{n}_{00}$ is nontrivially contained in the Tanaka prolongation up to the 0th order $\mathfrak{g}_0(\mathfrak{n}_-)$ of \mathfrak{n}_- , with $\{0\} \neq \mathfrak{n}_{00} \subset \mathfrak{n}_0$, and as such is a subalgebra of the algebra of $\text{aut}(\mathcal{D})$.

5. Majorana spinor representations of $\mathfrak{so}(p, q)$

In this section, we will explain how to construct the real spin representations of the Lie algebras $\mathfrak{so}(p, q)$, in cases when $p = n$, $q = n - 1$, or $p = q = n$, $n = 1, 2, \dots, n$. We will also give a construction of these representations for $\mathfrak{so}(0, n)$. We emphasise that we are only interested in *real* spin representations. They share a general name of *Majorana representations*. Our presentation of this material is adapted from [19].

We will need Pauli matrices

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \epsilon = -i\sigma_y = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \tag{5.1}$$

and the 2×2 identity matrix

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{5.2}$$

We have the following identities:

$$\begin{aligned} \sigma_x^2 = \sigma_z^2 = -\epsilon^2 = I \\ \sigma_x \epsilon = -\epsilon \sigma_x = \sigma_z, \quad \sigma_z \sigma_x = -\sigma_x \sigma_z = -\epsilon, \quad \epsilon \sigma_z = -\sigma_z \epsilon = \sigma_x. \end{aligned} \tag{5.3}$$

Now we quote [19]:

With this notation, *restricting to low dimensions* $p + q = 4, 5, 6$ and 7 , the real representations of the Clifford algebra $\mathcal{Cl}(0, p + q)$ are all in dimension $s = 8$, and are generated by the $p + q$ matrices $\rho_1, \dots, \rho_{(p+q)}$ given by:

$$\begin{aligned} \rho_1 &= \sigma_z \otimes I \otimes \epsilon \\ \rho_2 &= \sigma_z \otimes \epsilon \otimes \sigma_x \\ \rho_3 &= \sigma_z \otimes \epsilon \otimes \sigma_z \\ \rho_4 &= \sigma_x \otimes \epsilon \otimes I \\ \rho_5 &= \sigma_x \otimes \sigma_x \otimes \epsilon \\ \rho_6 &= \sigma_x \otimes \sigma_z \otimes \epsilon \\ \rho_7 &= \epsilon \otimes I \otimes I. \end{aligned} \tag{5.4}$$

The eight matrices $\theta_\mu = \sigma_x \otimes \rho_\mu, \mu = 1, \dots, 7$ and $\theta_8 = \epsilon \otimes I \otimes I \otimes I$, give the real representation of $\mathcal{Cl}(0, 8)$ in $S = \mathbb{R}^{16}$. Dropping the first factor in ρ_1, ρ_2, ρ_3 one obtains the matrices generating a representation of $\mathcal{Cl}(0, 3)$ in $S = \mathbb{R}^4$, etc.

Majorana representations of $\mathfrak{so}(n - 1, n)$ in dimension $s = 2^{n-1}$ are called *Pauli representations*, and Majorana representations of $\mathfrak{so}(n, n)$ in dimension $s = 2^n$ are called *Dirac representations*.

To construct them, we need generalisations of the *Pauli* σ matrices and *Dirac* γ matrices. The construction of those is *inductive*.

It starts with $p + q = 1$ with one matrix $\sigma_1 = 1$, and for every $n = 1, 2, \dots$, it alternates between $p + q = 2n - 1$ of Pauli matrices $\sigma_\mu, \mu = 1, \dots, 2n - 1$, and $p + q = 2n$ of Dirac matrices $\gamma_\mu, \mu = 1, \dots, 2n$.

Again quoting Trautman [19] we have the following:

- (1) In dimension $p + q = 1$, put $\sigma_1 = 1$.
- (2) Given $2^{n-1} \times 2^{n-1}$ matrices $\sigma_\mu, \mu = 1, \dots, 2n - 1$, define

$$\gamma_\mu = \begin{pmatrix} 0 & \sigma_\mu \\ \sigma_\mu & 0 \end{pmatrix} \text{ for } \mu = 1, \dots, 2n - 1,$$

and

$$\gamma_{2n} = \begin{pmatrix} 0 & -I \\ I & 0 \end{pmatrix},$$

where I is the identity $2^{n-1} \times 2^{n-1}$ matrix.

(3) Given $2^n \times 2^n$ matrices $\gamma_\mu, \mu = 1, \dots, 2n$, define $\sigma_\mu = \gamma_\mu$ for $\mu = 1, \dots, 2n$ and $\sigma_{2n+1} = \gamma_1 \dots \gamma_{2n}$, so that for $n > 0$,

$$\sigma_{2n+1} = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}.$$

In every dimension $p + q = 2n - 1, n \geq 1$, the Pauli matrices $\sigma_\mu, \mu = 1, \dots, 2n - 1$ satisfy

$$\sigma_\mu \sigma_\nu + \sigma_\nu \sigma_\mu = 2g_{\mu\nu} \underbrace{(I \otimes \dots \otimes I)}_{n-1 \text{ times}},$$

where the $(2n - 1) \times (2n - 1)$ symmetric matrix $(g_{\mu\nu})$ is *diagonal* and has the following diagonal elements:

$$(g_{\mu\nu}) = \text{diag} \underbrace{(1, -1, \dots, -1, 1)}_{(2n-1) \text{ times}}.$$

Likewise, in every dimension $p + q = 2n, n \geq 1$, the Dirac matrices $\gamma_\mu, \mu = 1, \dots, 2n$ satisfy

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2g_{\mu\nu} \underbrace{(I \otimes \dots \otimes I)}_{n \text{ times}},$$

where the $(2n) \times (2n)$ symmetric matrix $(g_{\mu\nu})$ is *diagonal* and has the following diagonal elements:

$$(g_{\mu\nu}) = \text{diag} \underbrace{(1, -1, \dots, 1, -1)}_{2n \text{ times}}.$$

Therefore, for each $n = 1, 2, \dots$, the set $\{\sigma_\mu\}_{\mu=1}^{2n-1}$ of Pauli matrices generates the elements of a real 2^{n-1} -dimensional representation of the Clifford algebra $\mathcal{Cl}(n - 1, n)$, and the set $\{\gamma_\mu\}_{\mu=1}^{2n}$ of Dirac matrices generates the elements of a real 2^n -dimensional representation of the Clifford algebra $\mathcal{Cl}(n, n)$.

Then, in turn, these real Clifford algebras representations can be further used to define the real spin representations of the Lie algebras $\mathfrak{so}(p + q, 0), \mathfrak{so}(n - 1, n)$ and $\mathfrak{so}(n, n)$ as follows. One obtains all the generators of the spin representation of $\mathfrak{so}(g)$ by spanning it by all the elements of the form

- $\frac{1}{2}\rho_\mu \rho_\nu$, with $1 \leq \mu < \nu \leq (p + q)$, in the case of $\mathfrak{so}(p + q, 0), p + q = 3, 5, 6, 7$;
- $\frac{1}{2}\theta_\mu \theta_\nu$, with $1 \leq \mu < \nu \leq 8$, in the case of $\mathfrak{so}(8, 0)$;
- $\frac{1}{2}\sigma_\mu \sigma_\nu$, with $1 \leq \mu < \nu \leq (p + q) = 2n - 1$, in the case of $\mathfrak{so}(n - 1, n)$;
- $\frac{1}{2}\gamma_\mu \gamma_\nu$, with $1 \leq \mu < \nu \leq (p + q) = 2n$, in the case of $\mathfrak{so}(n, n)$.

For further details, consult [19].

We will use all this information in next sections when we create examples.

6. Application: Obtaining the flat model for (3,6) distributions

Let (ρ, S) be the defining representation of $\mathfrak{so}(3)$ in $S = \mathbb{R}^3$. It can be generated by

$$\rho(A_1) = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \rho(A_2) = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \rho(A_3) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}. \tag{6.1}$$

And let (τ, R) be an equivalent 3-dimensional representation of $\mathfrak{so}(3)$ given by

$$\tau(A_1) = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \tau(A_2) = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tau(A_3) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}. \tag{6.2}$$

We claim that for these two representations of $\mathfrak{so}(3)$, in the standard bases in $S = \mathbb{R}^3$, $R = \mathbb{R}^3$, the magical equation (2.2) has the following solution:

$$\omega^1_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \omega^2_{\mu\nu} = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \omega^3_{\mu\nu} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Now using this solution (ρ, τ, ω) of the magical equation (2.1), we use the Corollary 4.1 with $\lambda^i = du^i + \omega^i_{\mu\nu} x^\mu dx^\nu$ and obtain the following theorem.

Theorem 6.1. *Let $M = \mathbb{R}^6$ with coordinates $(u^1, u^2, u^3, x^1, x^2, x^3)$ and consider three 1-forms*

$$\begin{aligned} \lambda^1 &= du^1 + x^2 dx^3 \\ \lambda^2 &= du^2 + x^1 dx^3 \\ \lambda^3 &= du^3 + x^1 dx^2 \end{aligned}$$

on M . Then the rank 3 distribution \mathcal{D} on M defined by $\mathcal{D} = \{\mathbb{T}\mathbb{R}^6 \ni X \mid X \lrcorner \lambda^i = 0, i = 1, 2, 3\}$ has its Lie algebra of infinitesimal symmetries $\mathfrak{aut}\mathcal{D}$ isomorphic to the Tanaka prolongation of $\mathfrak{n}_- = R \oplus S$, where $(\rho, S = \mathbb{R}^3)$ and $(\tau, R = \mathbb{R}^3)$ are the respective representations (6.1), (6.2) of $\mathfrak{n}_{00} = \mathfrak{so}(3)$.

The symmetry algebra $\mathfrak{aut}(\mathcal{D})$ is isomorphic to the simple graded Lie algebra $\mathfrak{so}(4, 3)$,

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{so}(4, 3),$$

with the following gradation:

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R, \mathfrak{n}_{-1} = S$,

$$\mathfrak{n}_0 = \mathfrak{gl}(3, \mathbb{R}) \supset \mathfrak{n}_{00}$$

$\mathfrak{n}_1 = S^, \mathfrak{n}_2 = R^*$, which is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in $\mathfrak{so}(4, 3)$.*

The contactification (M, \mathcal{D}) is locally a flat model for the parabolic geometry of type $(\mathbf{Spin}(4, 3), P)$ related to the following crossed Satake diagram: $\circ \text{---} \circ \rightleftharpoons \circ$.

Proof. Proof is by calculating the Tanaka prolongation of $\mathfrak{n}_- = R \oplus S$, which is $\mathfrak{gl}(3, \mathbb{R})$, naturally graded by the Tanaka prolongation algebraic procedure precisely as $\mathfrak{aut}(\mathcal{D})$ in the statement of the theorem. \square

7. Application: Obtaining Biquard’s 7-dimensional flat quaternionic contact manifold via contactification using spin representations of $\mathfrak{so}(1,2)$ and $\mathfrak{so}(3,0)$

According to Trautman’s procedure, [19] there is a real representation of $\mathcal{Cl}(0,3)$ in \mathbb{R}^4 . There also is an analogous representation of $\mathcal{Cl}(1,2)$. Both of them are generated by the σ matrices

$$\sigma_1 = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & 0 & 0 & -\varepsilon \\ 0 & 0 & -\varepsilon & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 0 & 0 & -\varepsilon & 0 \\ 0 & 0 & 0 & \varepsilon \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix},$$

where

$$\varepsilon = 1 \quad \text{for } \mathcal{Cl}(0,3),$$

and

$$\varepsilon = -1 \quad \text{for } \mathcal{Cl}(2,1).$$

One can check that these matrices¹ satisfy the (representation of) Clifford algebra relations:

$$\sigma_\mu \sigma_\nu + \sigma_\nu \sigma_\mu = 2g_{\mu\nu} (I \otimes I)$$

with all $g_{\mu\nu}$ being zero, except $g_{11} = -1, g_{22} = g_{33} = -\varepsilon$.

This leads to the following spinorial representation ρ of $\mathfrak{so}(0,3)$ or $\mathfrak{so}(1,2)$:

$$\rho(A_1) = -\frac{1}{2}\sigma_3, \quad \rho(A_2) = \frac{1}{2}\sigma_2, \quad \rho(A_3) = -\frac{1}{2}\varepsilon\sigma_1. \tag{7.1}$$

Here, (A_1, A_2, A_3) constitutes a basis for $\mathfrak{so}(0,3)$ when $\varepsilon = 1$ and for $\mathfrak{so}(2,1)$ when $\varepsilon = -1$. This can be extended to the representation of

$$\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right)$$

in $S = \mathbb{R}^4$ by setting the value of ρ on the generator $A_4 = \text{Id}$ as

$$\rho(A_4) = \frac{1}{2}(I \otimes I). \tag{7.2}$$

For this representation of $\mathbb{R} \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right)$, the magical equation (2.1) has a following solution:

¹In Trautman’s quote in the previous section, these matrices were denoted by ρ_1, ρ_2, ρ_3 , and they were only explicitly given for $\varepsilon = 1$.

$$\omega^1_{\mu\nu} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad \omega^2_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

$$\omega^1_{\mu\nu} = \begin{pmatrix} 0 & -\varepsilon & 0 & 0 \\ \varepsilon & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix},$$

with

$$\begin{aligned} \tau(A_1) &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \varepsilon \\ 0 & -1 & 0 \end{pmatrix}, \quad \tau(A_2) = \begin{pmatrix} 0 & 0 & -\varepsilon \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad \tau(A_3) = \begin{pmatrix} 0 & -\varepsilon & 0 \\ \varepsilon & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ \tau(A_4) &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned} \tag{7.3}$$

This in particular gives the vectorial representation τ of

$$\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right)$$

in $R = \mathbb{R}^3$.

Now, by using this solution for (ρ, ω, τ) and applying our Corollary 4.1, we have an $(s = 4)$ -dimensional manifold $N = \mathbb{R}^4$, equipped with $r = 3$ two-forms $\omega^i = \frac{1}{2}\omega_{\mu\nu}dx^\mu \wedge dx^\nu$, $i = 1, 2, 3$, which contactifies to an $(s + r) = 7$ -dimensional manifold $M = \mathbb{R}^7$ having a distribution structure (M, \mathcal{D}) defined as an annihilator of the $r = 3$ one-forms $\lambda^i = du^i + \omega^i_{\mu\nu}x^\mu dx^\nu$, $i = 1, 2, 3$. We have the following theorem.

Theorem 7.1. *Let $M = \mathbb{R}^7$ with coordinates $(u^1, u^2, u^3, x^1, x^2, x^3, x^4)$, and consider three 1-forms $\lambda^1, \lambda^2, \lambda^3$ on M given by*

$$\begin{aligned} \lambda^1 &= du^1 + x^1 dx^3 - x^2 dx^4, \\ \lambda^2 &= du^2 - x^1 dx^4 - x^2 dx^3, \quad \text{with} \quad \varepsilon = \pm 1. \\ \lambda^3 &= du^3 - \varepsilon x^1 dx^2 - x^3 dx^4, \end{aligned}$$

The rank 4 distribution \mathcal{D} on M defined as $\mathcal{D} = \{\mathbb{T}\mathbb{R}^7 \ni X \mid X \lrcorner \lambda^1 = X \lrcorner \lambda^2 = X \lrcorner \lambda^3 = 0\}$ has its Lie algebra of infinitesimal automorphism $\mathfrak{aut}(\mathcal{D})$ isomorphic to the Tanaka prolongation of $\mathfrak{n} = R \oplus S$, where $(\rho, S = \mathbb{R}^4)$ is the spinorial representation (7.1)-(7.2) of $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right)$, and $(\tau, R = \mathbb{R}^3)$ is the vectorial representation (7.3) of \mathfrak{n}_{00} .

The symmetry algebra $\mathfrak{aut}(\mathcal{D})$ is isomorphic to the simple Lie algebra $\mathfrak{sp}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right)$,

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{sp}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right),$$

having the following natural gradation:

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S$,

$$\begin{aligned} \mathfrak{n}_0 &= \mathfrak{n}_{00} \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right) = \\ &\mathbb{R} \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right) \oplus \mathfrak{so}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right), \end{aligned}$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$, which is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in $\mathfrak{sp}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right)$.

The contactification (M, \mathcal{D}) is locally a flat model for the parabolic geometry of type $(\mathbb{S}\mathfrak{p}\left(\frac{1-\varepsilon}{2}, \frac{5+\varepsilon}{2}\right), P)$ related to the following crossed satake diagrams:

- (1) $\bullet \rightleftarrows \bullet$ in the case of $\varepsilon = 1$, and
- (2) $\circ \rightleftarrows \circ$ in the case of $\varepsilon = -1$.

Remark 7.2. When $\varepsilon = 1$, the flat parabolic geometry described in the above theorem is the lowest dimensional example of the *quaternionic contact* geometry considered by Biquard [5] (see also [12]).

8. Application: Obtaining the exceptionals from contactifications of spin representations; the \mathfrak{f}_4 case

We will now explain the Cartan realisation of the simple exceptional Lie algebra \mathfrak{f}_4 in dimension \mathbb{R}^{15} mentioned in the introduction.

The Satake diagrams for the real forms of the complex simple exceptional Lie algebra \mathfrak{f}_4 are as follows:



The first diagram corresponds to the *compact* real form of \mathfrak{f}_4 and is not interesting for us.² The other two diagrams are interesting:

- (1) The last, $\circ \rightleftarrows \circ \rightleftarrows \circ$, corresponds to the *split* real form \mathfrak{f}_I , and
- (2) the middle one, $\bullet \rightleftarrows \bullet \rightleftarrows \circ$, denoted by \mathfrak{f}_{II} in [11], is also interesting, since similarly to \mathfrak{f}_I , it defines a *parabolic geometry* in dimension 15.

Crossing the last node on the right in the diagrams for \mathfrak{f}_I or \mathfrak{f}_{II} , as in $\circ \rightleftarrows \circ \rightleftarrows \circ$ or $\bullet \rightleftarrows \bullet \rightleftarrows \circ$, we see that in both algebras there exist *parabolic subalgebras* \mathfrak{p}_I or \mathfrak{p}_{II} , respectively, of dimension 37, $\dim(\mathfrak{p}_I) = \dim(\mathfrak{p}_{II}) = 37$. In both respective cases, these choices of parabolics define similar gradations in the corresponding real forms \mathfrak{f}_I , \mathfrak{f}_{II} , of the simple exceptional Lie \mathfrak{f}_4 :

$$\mathfrak{f}_A = \mathfrak{n}_{-2A} \oplus \mathfrak{n}_{-1A} \oplus \mathfrak{n}_{0A} \oplus \mathfrak{n}_{1A} \oplus \mathfrak{n}_{2A} \quad \text{for } A = I, II,$$

with

$$\mathfrak{n}_{-A} = \mathfrak{n}_{-2A} \oplus \mathfrak{n}_{-1A} \quad \text{for } A = I, II,$$

²For the compact realisation, see, for example, [2].

being 2-step nilpotent and having grading components \mathfrak{n}_{-2A} and \mathfrak{n}_{-1A} of respective dimension $r_A = 7$ and $s_A = 8$,

$$r_A = \dim(\mathfrak{n}_{-2A}) = 7, \quad s_A = \dim(\mathfrak{n}_{-1A}) = 8 \quad \text{for } A = I, II.$$

The Lie algebra \mathfrak{n}_{0A} in the Tanaka prolongation of \mathfrak{n}_{-A} up to 0^{th} order is

- (1) $\mathfrak{n}_{0I} = \mathbb{R} \oplus \mathfrak{so}(4,3)$ in the case of \mathfrak{f}_I , and
- (2) $\mathfrak{n}_{0II} = \mathbb{R} \oplus \mathfrak{so}(0,7)$ in the case of \mathfrak{f}_{II} .

Thus, from the analysis performed here, we see that there exist two different 2-step filtered structures (M_I, \mathcal{D}_I) and $(M_{II}, \mathcal{D}_{II})$, both in dimension 15, with the respective F_I -symmetric, or F_{II} -symmetric flat models, realised on $M_I = F_I/P_I$ or $M_{II} = F_{II}/P_{II}$. Here, F_I and F_{II} denote the real Lie groups whose Lie algebras are \mathfrak{f}_I and \mathfrak{f}_{II} , respectively. Similarly, P_I and P_{II} are parabolic subgroups of respective F_I and F_{II} , whose Lie algebras are \mathfrak{p}_I and \mathfrak{p}_{II} . Recalling that each of the real groups $\mathbf{SO}(4,3)$ and $\mathbf{SO}(0,7)$ has *two* real irreducible representations ρ in dimension $s = 8$ and τ in dimension $r = 7$, with the 8-dimensional representation ρ being the *spin* representation of either $\mathbf{SO}(4,3)$ or $\mathbf{SO}(0,7)$, we can now give the explicit realisations of the \mathbf{F}_4 -symmetric structures (M_A, \mathcal{D}_A) for $A = I, II$.

8.1. Cartan’s realisation of \mathfrak{f}_I

The plan is to start with the Lie algebra $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(4,3)$, as in the crossed Satake diagram $\circ \text{---} \circ \rightleftarrows \circ \text{---} \circ$ of \mathfrak{f}_I , and its two representations:

- a representaion $(\rho, S = \mathbb{R}^8)$, corresponding to the spin representation of $\mathbf{SO}(4,3)$ in $(s = 8)$ -dimensional space $\mathfrak{n}_{-1} = S$ of real Pauli spinors, and
- a representation $(\tau, R = \mathbb{R}^7)$, corresponding to the vectorial representation of $\mathbf{SO}(4,3)$ in $(r = 7)$ -dimensional space $\mathfrak{n}_{-2} = R$ of vectors in $\mathbb{R}^{(4,3)}$.

Having these two representations of $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(4,3)$ in the same basis, we will then solve the equations (2.1) for the map $\omega \in \text{Hom}(\wedge^2 S, R)$, which will give us the commutators between elements in \mathfrak{n}_{-1} . This via Corollary 4.1 will provide the explicit realisation of the 15-dimensional contactification (M, \mathcal{D}) with the exception of simple Lie algebra F_I as its symmetry.

Actually, the passage from ρ to τ in the above plan is a bit tricky, since we need to have these representations expressed in the same basis. To handle this obstacle, we will start with the spin representation ρ in the space of *Pauli spinors* S , and then we will use the fact that the skew representation $\rho \wedge \rho$ in the space of the bispinors $\wedge^2 S$ decomposes as

$$\wedge^2 S = \wedge_{21} \oplus \wedge_7,$$

where \wedge_{21} is the 21-dimensional *adjoint representation* of $\mathbf{SO}(4,3)$ and \wedge_7 is its 7-dimensional *vectorial representation* τ . In this way, we will have the two representations (ρ, S) and $(\tau, R = \wedge_7)$, expressed in the same basis $\{A_I\}$ of $\mathbb{R} \oplus \mathbf{SO}(4,3)$, and we will apply the Corollary 4.1 to get the desired F_I -symmetric contactification in dimension 15. On doing this, we will use notation from Section 5.

According to [19], the real 8-dimensional representation of the Clifford algebra $\mathcal{Cl}(4,3)$ is generated by the seven 8-dimensional Pauli matrices:

$$\begin{aligned} \sigma_1 &= \sigma_x \otimes \sigma_x \otimes \sigma_x \\ \sigma_2 &= \sigma_x \otimes \sigma_x \otimes \epsilon \\ \sigma_3 &= \sigma_x \otimes \sigma_x \otimes \sigma_z \\ \sigma_4 &= \sigma_x \otimes \epsilon \otimes I \\ \sigma_5 &= \sigma_x \otimes \sigma_z \otimes I \\ \sigma_6 &= \epsilon \otimes I \otimes I \\ \sigma_7 &= \sigma_z \otimes I \otimes I. \end{aligned}$$

Using the identities (5.3), especially the one saying that $\epsilon^2 = -I$, one easily finds that the seven Pauli matrices $\sigma_i, i = 1, 2, \dots, 7$, satisfy the Clifford algebra identity

$$\sigma_i \sigma_j + \sigma_j \sigma_i = 2g_{ij} (I \otimes I \otimes I), \quad i, j = 1, 2, \dots, 7,$$

with the coefficients g_{ij} forming a diagonal 7×7 matrix

$$(g_{ij}) = \text{diag}(1, -1, 1, -1, 1, -1, 1)$$

of signature (4,3). Thus, the 8-dimensional Pauli matrices $\sigma_i, i = 1, \dots, 7$ generate the Clifford algebra $\mathcal{Cl}(4,3)$, and in turn, by the general theory, as described in Section 5, they define the spin representation ρ of $\mathfrak{so}(4,3)$ in an 8-dimensional real vector space $S = \mathbb{R}^8$ of Pauli(-Majorana) spinors.

8.1.1. The spinorial representation of $\mathfrak{so}(4,3)$. To be more explicit, let (i, j) be such that $1 \leq i < j \leq 7$, and let I be a function

$$I(i, j) = 1 + i + \frac{1}{2}(j - 3)j \tag{8.1}$$

on such pairs. Note that the function I is a bijection between the 21 pairs (i, j) and the set of 21 natural numbers $I = 1, 2, \dots, 21$. Consider the twenty-one 8×8 real matrices $\sigma_i \sigma_j$ with $1 \leq i < j \leq 7$, and a basis $\{A_I\}_{I=1}^{21}$ in the Lie algebra $\mathfrak{so}(4,3)$. Then the spin representation ρ of $\mathfrak{so}(4,3)$ is given by

$$\rho(A_{I(i,j)}) = \frac{1}{2} \sigma_i \sigma_j \quad \text{with } 1 \leq i < j \leq 7.$$

Explicitly, we have

$$\begin{aligned} \rho(A_1) &= \frac{1}{2} I \otimes I \otimes \sigma_z, & \rho(A_8) &= \frac{1}{2} I \otimes \epsilon \otimes \epsilon, & \rho(A_{15}) &= \frac{1}{2} \sigma_z \otimes \sigma_z \otimes I, \\ \rho(A_2) &= \frac{1}{2} I \otimes I \otimes \epsilon, & \rho(A_9) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_z, & \rho(A_{16}) &= \frac{1}{2} \epsilon \otimes \sigma_x \otimes \sigma_x, \\ \rho(A_3) &= \frac{1}{2} I \otimes I \otimes \sigma_x, & \rho(A_{10}) &= \frac{1}{2} I \otimes \sigma_x \otimes I, & \rho(A_{17}) &= \frac{1}{2} \epsilon \otimes \sigma_x \otimes \epsilon, \\ \rho(A_4) &= \frac{1}{2} I \otimes I \otimes \sigma_x, & \rho(A_{11}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \sigma_x, & \rho(A_{18}) &= \frac{1}{2} \epsilon \otimes \sigma_x \otimes \sigma_z, \\ \rho(A_5) &= \frac{1}{2} I \otimes \sigma_z \otimes \epsilon, & \rho(A_{12}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \epsilon, & \rho(A_{19}) &= \frac{1}{2} \epsilon \otimes \epsilon \otimes I, \\ \rho(A_6) &= \frac{1}{2} I \otimes \sigma_z \otimes \sigma_z, & \rho(A_{13}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \sigma_z, & \rho(A_{20}) &= \frac{1}{2} \epsilon \otimes \sigma_z \otimes I, \\ \rho(A_7) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_x, & \rho(A_{14}) &= \frac{1}{2} \sigma_z \otimes \epsilon \otimes I, & \rho(A_{21}) &= \frac{1}{2} \sigma_x \otimes I \otimes I. \end{aligned} \tag{8.2}$$

The spin representation ρ of $V_0 = \mathbb{R} \oplus \mathfrak{so}(4,3)$ needs one generator more. Let us call it $\rho(A_{22})$. We have

$$\rho(A_{22}) = \frac{1}{2}I \otimes I \otimes I.$$

We determine the structure constants c^K_{IJ} of $\mathbb{R} \oplus \mathfrak{so}(4,3)$ in the basis A_I from

$$[\rho(A_I), \rho(A_J)] = c^K_{IJ} \rho(A_K). \tag{8.3}$$

8.1.2. Obtaining the vectorial representation of $\mathfrak{so}(4,3)$. Now, we take the space $\wedge^2 S$ and consider the skew symmetric representation

$$\overset{\circ}{\rho} = \rho \wedge \rho$$

in it. We will write it in the standard basis f_μ , $\mu = 1, \dots, 8$ in $S = \mathbb{R}^8$. We have $\rho(A_I)f_\mu = \rho I^\nu_\mu f_\nu$. Now, the components of the 28-dimensional representation $\overset{\circ}{\rho} = \rho \wedge \rho_1$ are

$$\overset{\circ}{\rho} I^{\mu\nu}{}_{\alpha\beta} = \rho I^\mu_\alpha \delta^\nu_\beta + \delta^\mu_\alpha \rho I^\nu_\beta - \rho I^\nu_\alpha \delta^\mu_\beta - \delta^\nu_\alpha \rho I^\mu_\beta,$$

and we have

$$(\overset{\circ}{\rho}(A_I)w)^{\mu\nu} = \overset{\circ}{\rho} I^{\mu\nu}{}_{\alpha\beta} w^{\alpha\beta}, \quad \forall w^{\alpha\beta} = w^{[\alpha\beta]}.$$

The Casimir operator for this representation is

$$\mathcal{C} = 10 K^{IJ} \overset{\circ}{\rho}(A_I) \overset{\circ}{\rho}(A_J),$$

where K^{IJ} is the inverse of the Killing form matrix $K_{IJ} = c^L_{IM} c^M_{JL}$ in the basis A_I . Since for the Killing form to be nondegenerate we must restrict to the semisimple part of $\mathbb{R} \oplus \mathfrak{so}(4,3)$, here the indices $I, J, K, L, M = 1, 2, \dots, 21$, and as always are summed over the repeated indices. One can check that in this basis of $\mathfrak{so}(4,3)$, the Killing form matrix is diagonal and reads

$$(K_{IJ}) = 10 \text{diag}(1, -1, 1, 1, -1, 1, -1, 1, -1, 1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, 1).$$

The Casimir \mathcal{C} defines the decomposition of the 28-dimensional reducible representation $\overset{\circ}{\rho} = \rho_1 \wedge \rho_1$ onto

$$\wedge^2 S = \wedge_{21} \oplus \wedge_7,$$

where the 7-dimensional irreducible representation space \wedge_7 is the eigenspace of the Casimir operator consisting of eigen-bispinors with eigenvalue equal to 6,

$$\mathcal{C}(\wedge_7) = 6 \wedge_7.$$

Explicitly, in the same basis A_I , $I = 1, 2, \dots, 21$ as before, this 7-dimensional representation $(\tau, R = \bigwedge_7)$ of the $\mathfrak{so}(4,3)$ Lie algebra is given by

$$\begin{aligned}
 \tau(A_1) &= E_{66} - E_{22}, \\
 \tau(A_2) &= \frac{1}{2}(E_{23} - E_{32} + E_{25} - E_{52} + E_{36} - E_{63} + E_{56} - E_{65}), \\
 \tau(A_3) &= \frac{1}{2}(E_{23} + E_{32} + E_{25} + E_{52} + E_{36} + E_{63} + E_{56} + E_{65}), \\
 \tau(A_4) &= \frac{1}{2}(E_{23} + E_{32} - E_{25} - E_{52} - E_{36} - E_{63} + E_{56} + E_{65}), \\
 \tau(A_5) &= \frac{1}{2}(E_{23} - E_{32} - E_{25} + E_{52} - E_{36} + E_{63} + E_{56} - E_{65}), \\
 \tau(A_6) &= E_{33} - E_{55}, \\
 \tau(A_7) &= \frac{1}{2}(E_{12} - E_{21} - E_{16} + E_{61} - E_{27} + E_{72} + E_{67} - E_{76}), \\
 \tau(A_8) &= \frac{1}{2}(-E_{12} - E_{21} - E_{16} - E_{61} - E_{27} - E_{72} - E_{67} - E_{76}), \\
 \tau(A_9) &= \frac{1}{2}(E_{13} - E_{31} + E_{15} - E_{51} - E_{37} + E_{73} - E_{57} + E_{75}), \\
 \tau(A_{10}) &= \frac{1}{2}(E_{13} + E_{31} - E_{15} - E_{51} + E_{37} + E_{73} - E_{57} - E_{75}), \\
 \tau(A_{11}) &= \frac{1}{2}(-E_{12} - E_{21} + E_{16} + E_{61} + E_{27} + E_{72} - E_{67} - E_{76}), \\
 \tau(A_{12}) &= \frac{1}{2}(E_{12} - E_{21} + E_{16} - E_{61} + E_{27} - E_{72} + E_{67} - E_{76}), \\
 \tau(A_{13}) &= \frac{1}{2}(-E_{13} - E_{31} - E_{15} - E_{51} + E_{37} + E_{73} + E_{57} + E_{75}), \\
 \tau(A_{14}) &= \frac{1}{2}(-E_{13} + E_{31} + E_{15} - E_{51} - E_{37} + E_{73} + E_{57} - E_{75}), \\
 \tau(A_{15}) &= E_{11} - E_{77}, \\
 \tau(A_{16}) &= \frac{1}{2}(2E_{24} - E_{42} + E_{46} - 2E_{64}), \\
 \tau(A_{17}) &= \frac{1}{2}(2E_{24} + E_{42} + E_{46} + 2E_{64}), \\
 \tau(A_{18}) &= \frac{1}{2}(2E_{34} - E_{43} - E_{45} + 2E_{54}), \\
 \tau(A_{19}) &= \frac{1}{2}(-2E_{34} - E_{43} + E_{45} + 2E_{54}), \\
 \tau(A_{20}) &= \frac{1}{2}(-2E_{14} + E_{41} + E_{47} - 2E_{74}), \\
 \tau(A_{21}) &= \frac{1}{2}(2E_{14} + E_{41} - E_{47} - 2E_{74}), \\
 \tau(A_{22}) &= E_{11} + E_{22} + E_{33} + E_{44} + E_{55} + E_{66} + E_{77},
 \end{aligned} \tag{8.4}$$

where E_{ij} , $i, j = 1, 2, \dots, 7$ denote 7×7 matrices with zeroes everywhere except the value 1 in the entry (i, j) seating at the crossing of the i th row and the j th column.

One can check that

$$[\tau(A_I), \tau(A_J)] = c^K_{IJ} \tau(A_K),$$

with the same structure constants as in (8.3).

8.1.3. A contactification with \mathfrak{f}_I symmetry. So now we are in the situation of having two representations (ρ, S) and $(\tau, R = \bigwedge_7)$ on $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(4,3)$, and we can try to solve the equation (2.1) for the map $\omega \in \text{Hom}(\bigwedge^2 S, R)$. Of course, if we started with some arbitrary ρ and τ , this equation would not have solutions other than 0, but here we expect to have solution since we know it from the Cartan’s PhD thesis [8] and the announcement in Helgason’s paper [13]. And indeed, there is a solution for a nonzero ω , which when written in the basis $\{f_\mu\}$ in S and $\{e_i\}$ in R is such that it gives the *seven*

2-forms $\omega^i = \frac{1}{2}\omega^i_{\mu\nu}dx^\mu \wedge dx^\nu$, $i = 1, \dots, 7$, in $N = \mathbb{R}^8$ given by

$$\begin{aligned} \omega^1 &= dx^1 \wedge dx^2 - dx^7 \wedge dx^8, \\ \omega^2 &= dx^2 \wedge dx^4 - dx^6 \wedge dx^8, \\ \omega^3 &= dx^1 \wedge dx^4 - dx^5 \wedge dx^8, \\ \omega^4 &= \frac{1}{2} (dx^1 \wedge dx^6 - dx^2 \wedge dx^5 - dx^3 \wedge dx^8 + dx^4 \wedge dx^7), \\ \omega^5 &= dx^2 \wedge dx^3 - dx^6 \wedge dx^7, \\ \omega^6 &= dx^1 \wedge dx^3 - dx^5 \wedge dx^7, \\ \omega^7 &= dx^3 \wedge dx^4 - dx^5 \wedge dx^6. \end{aligned} \tag{8.5}$$

These, via the contactification and the theory summarised in Corollary 4.1, lead to the following theorem.

Theorem 8.1. *Let $M = \mathbb{R}^{15}$ with coordinates $(u^1, \dots, u^7, x^1, \dots, x^8)$, and consider seven 1-forms $\lambda^i, \dots, \lambda^7$ on M given by*

$$\begin{aligned} \lambda^1 &= du^1 + x^1 dx^2 - x^7 dx^8, \\ \lambda^2 &= du^2 + x^2 dx^4 - x^6 dx^8, \\ \lambda^3 &= du^3 + x^1 dx^4 - x^5 dx^8, \\ \lambda^4 &= du^4 + \frac{1}{2} (x^1 dx^6 - x^2 dx^5 - x^3 dx^8 + x^4 dx^7), \\ \lambda^5 &= du^5 + x^2 dx^3 - x^6 dx^7, \\ \lambda^6 &= du^6 + x^1 dx^3 - x^5 dx^7, \\ \lambda^7 &= du^7 + x^3 dx^4 - x^5 dx^6. \end{aligned}$$

The rank 8 distribution \mathcal{D} on M defined as $\mathcal{D} = \{\text{TRR}^{15} \ni X \mid X \lrcorner \lambda^1 = \dots = X \lrcorner \lambda^7 = 0\}$ has its Lie algebra of infinitesimal automorphism $\text{aut}(\mathcal{D})$ isomorphic to the Tanaka prolongation of $\mathfrak{n}_- = R \oplus S$, where $(\rho, S = \mathbb{R}^8)$ is the spinorial representation (8.2) of $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(4,3)$ and $(\tau, R = \mathbb{R}^7)$ is the vectorial representation (8.4) of \mathfrak{n}_{00} .

The symmetry algebra $\text{aut}(\mathcal{D})$ is isomorphic to the simple Lie algebra \mathfrak{f}_I ,

$$\text{aut}(\mathcal{D}) = \mathfrak{f}_I,$$

having the following natural gradation:

$$\text{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S$,

$$\mathfrak{n}_0 = \mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(4,3),$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$, which is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in \mathfrak{f}_I .

The contactification (M, \mathcal{D}) is locally a flat model for the parabolic geometry of type (F_I, P_I) related to the following crossed Satake diagram $\circ \text{---} \circ \rightleftarrows \circ \text{---} \circ$ of \mathfrak{f}_I .

Remark 8.2. Please note that this is an example of an application of the magical equation (2.1) in which the starting algebra \mathfrak{n}_{00} was big enough, so that its Tanaka

prolongation \mathfrak{n}_0 counterpart is precisely equal to \mathfrak{n}_{00} . This was actually expected from the construction based on the crossed Satake diagram, $\circ \text{---} \circ \text{---} \circ \text{---} \circ$, which shows that \mathfrak{n}_0 of this parabolic geometry is precisely our $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(4,3)$.

Remark 8.3. One sees that the distribution \mathcal{D} in \mathbb{R}^{15} with \mathfrak{f}_I symmetry presented in Theorem 8.1 looks different than the distribution from our Example 1.3. It follows, however, that both these distributions are locally equivalent, and both have the same simple exceptional Lie algebra \mathfrak{f}_I as an algebra of their automorphisms.

8.1.4. Contactification for \mathfrak{f}_I : more algebra about $\mathfrak{so}(4,3)$. In our construction of the \mathfrak{f}_I symmetric distribution \mathcal{D} in Theorem 8.1, the crucial role was played by the 7-dimensional span of 2-forms $\omega^i, i = 1, 2, \dots, 7$. If we were given these seven 2-forms, we would produce the \mathfrak{f}_I symmetric distribution \mathcal{D} by the procedure of contactification.

It turns out that in $S = \mathbb{R}^8$, there is a particular 4-form³

$$\Phi = \frac{1}{4!} \Phi_{\mu\nu\rho\sigma} dx^\mu \wedge dx^\nu \wedge dx^\rho \wedge dx^\sigma$$

that is $\mathbb{R} \oplus \mathfrak{so}(4,3)$ invariant

$$\rho_I^\alpha{}_\mu \Phi_{\alpha\nu\rho\sigma} + \rho_I^\alpha{}_\nu \Phi_{\mu\alpha\rho\sigma} + \rho_I^\alpha{}_\rho \Phi_{\mu\nu\alpha\sigma} + \rho_I^\alpha{}_\sigma \Phi_{\mu\nu\rho\alpha} = S_I \Phi_{\mu\nu\rho\sigma}.$$

It may be represented by

$$\Phi = h_{ij} \omega^i \wedge \omega^j,$$

where ω^i are given by (8.5) and

$$(h_{ij}) = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

or in words⁴: $h_{ij}, i, j = 1, 2, \dots, 7$, are all zero except $h_{17} = h_{71} = -h_{26} = -h_{62} = h_{35} = h_{53} = h_{44} = 1$.

The form Φ in full beauty reads

$$\begin{aligned} \frac{2}{3} \Phi = 2 \Big(& dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^4 + dx^5 \wedge dx^6 \wedge dx^7 \wedge dx^8 \Big) - \\ & dx^1 \wedge dx^2 \wedge dx^5 \wedge dx^6 + dx^1 \wedge dx^3 \wedge dx^6 \wedge dx^8 - \\ & dx^1 \wedge dx^4 \wedge dx^6 \wedge dx^7 - dx^2 \wedge dx^3 \wedge dx^5 \wedge dx^8 + \\ & dx^2 \wedge dx^4 \wedge dx^5 \wedge dx^7 - dx^3 \wedge dx^4 \wedge dx^7 \wedge dx^8. \end{aligned} \tag{8.6}$$

³Compare this and other 4-forms appearing in the sequel with forms introduced in [7].

⁴Note that since (h_{ij}) is a symmetric matrix of signature (4,3), this fact alone shows that the span of seven 2-forms ω^i is a 7-dimensional representation space of $\mathbf{SO}(4,3)$. Actually, this fact easily leads to the construction of the double cover $\mathbb{Z}_2 \rightarrow \mathbf{Spin}(4,3) \rightarrow \mathbf{SO}(4,3)$.

Remark 8.4. It is remarkable that this 4-form alone encaptures all the features of the \mathfrak{f}_I symmetric contactification we discussed in the entire Section 8.1. By this, we mean following:

- (1) Consider $N = \mathbb{R}^8$ with coordinates (x^μ) , $\mu = 1, 2, \dots, 8$ and the 4-form

$$\Phi = \frac{1}{4!} \Phi_{\mu\nu\rho\sigma} dx^\mu \wedge dx^\nu \wedge dx^\rho \wedge dx^\sigma$$

given by (8.6).

- (2) Consider an equation

$$A^\alpha{}_\mu \Phi_{\alpha\nu\rho\sigma} + A^\alpha{}_\nu \Phi_{\mu\alpha\rho\sigma} + A^\alpha{}_\rho \Phi_{\mu\nu\alpha\sigma} + A^\alpha{}_\sigma \Phi_{\mu\nu\rho\alpha} = S \Phi_{\mu\nu\rho\sigma}$$

for the real 7×7 matrix $A = (A^\mu{}_\nu)$.

- (3) For simplicity, solve it in two steps:
 - First with $S = 0$. You obtain 21-dimensional solution space, which will be the *spin representation* ρ of $\mathfrak{so}(4,3)$. It is given $\rho(A) = A$.
 - Then prove that the only solution with $S \neq 0$ corresponds to $S = 4$, and that, modulo the addition of linear combinations of solutions with $S = 0$, it is given by $A = \text{Id}_{8 \times 8}$. Extend your possible A s with $S = 0$ to A s including $A = \text{Id}_{8 \times 8}$.
- (4) In this way, you will show that the stabiliser in $\mathfrak{gl}(8, \mathbb{R})$ of the 4-form Φ is the Lie algebra $\mathbb{R} \oplus \mathfrak{so}(4,3)$ in the *spin representation* ρ of Pauli spinors; $\rho(A) = A$.
- (5) Then search for a 7-dimensional space of 2-forms spanned say by the 2-forms $\omega^i = \frac{1}{2} \omega^i{}_{\mu\nu} dx^\mu \wedge dx^\nu$ satisfying

$$A^\alpha{}_\mu \omega^i{}_{\alpha\nu} + A^\alpha{}_\nu \omega^i{}_{\mu\alpha} = s^i{}_j \omega^j{}_{\mu\nu}$$

for all A s from the spin representation $\rho(A) = A$ of $\mathbb{R} \oplus \mathfrak{so}(4,3)$. Here, $s^i{}_j$ are auxiliary constants⁵.

- (6) This space is uniquely defined by these equations, and after solving them, you will get 7 linearly independent 2-forms $(\omega^1, \dots, \omega^7)$ in $N = \mathbb{R}^8$.
- (7) Contactifying the resulting structure $(N, \text{Span}(\omega^1, \dots, \omega^7))$, as we did, for example, in Theorem 8.1, you will get \mathfrak{f}_I symmetric distribution \mathcal{D} in $\mathbb{R}^7 \rightarrow (M = \mathbb{R}^{15}) \rightarrow (N = \mathbb{R}^8)$.

8.2. Realisation of \mathfrak{f}_{II}

It seems that Cartan was only interested in the explicit realisation of \mathfrak{f}_I . The realisation of \mathfrak{f}_{II} can be obtained in the same spirit as we have described in Section 8.1. Here, without much of the explanations since they parallel Section 8.1, we only display the main steps leading to this realisation.

We start with the representation of the Clifford algebra $\mathcal{C}\ell(0,7)$ generated by the seven ρ -matrices from (5.4). They satisfy

⁵Note, however, that although you look for $\omega^i{}_{\mu\nu}$ with *some* constants $s^i{}_j$, these constants have geometric meaning: comparing with our magical equation (2.2), we see that the 7×7 matrices $(s^i{}_j)$ constitute matrices of the defining representation τ of $\mathbb{R} \oplus \mathfrak{so}(4,3)$.

$$\rho_i \rho_j + \rho_j \rho_i = -2\delta_{ij} I_{8 \times 8}, \quad i, j = 1, \dots, 7.$$

They induce the 8-dimensional representation

$$\rho : \mathbb{R} \oplus \mathfrak{so}(0,7) \rightarrow \text{End}(S)$$

of $n_{00} = \mathbb{R} \oplus \mathfrak{so}(0,7)$ in the space $S = \mathbb{R}^8$ of real Pauli spinors, generated by the 22 real 8×8 matrices:

$$\begin{aligned} \rho(A_{I(i,j)}) &= \frac{1}{2} \rho_i \rho_j, \quad 1 \leq i < j \leq 7, \\ \rho(A_{22}) &= \frac{1}{2} (I \otimes I \otimes I), \end{aligned}$$

with the index $I = I(i, j)$ given by (8.1) and with $I, \sigma_x, \epsilon, \sigma_z$ given by (5.1)–(5.2). Explicitly, in terms of matrices $I, \sigma_x, \epsilon, \sigma_z$, the generators of this spinorial representation of $\mathfrak{so}(0,7)$ are

$$\begin{aligned} \rho(A_1) &= -\frac{1}{2} I \otimes \epsilon \otimes \sigma_z, & \rho(A_8) &= \frac{1}{2} \epsilon \otimes \sigma_z \otimes \sigma_z, & \rho(A_{15}) &= -\frac{1}{2} I \otimes \epsilon \otimes I, \\ \rho(A_2) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_x, & \rho(A_9) &= -\frac{1}{2} \epsilon \otimes \sigma_z \otimes \sigma_x, & \rho(A_{16}) &= -\frac{1}{2} \sigma_x \otimes I \otimes \epsilon, \\ \rho(A_3) &= -\frac{1}{2} I \otimes I \otimes \epsilon, & \rho(A_{10}) &= -\frac{1}{2} I \otimes \sigma_z \otimes \epsilon, & \rho(A_{17}) &= -\frac{1}{2} \sigma_x \otimes \epsilon \otimes \sigma_x, \\ \rho(A_4) &= -\frac{1}{2} \epsilon \otimes \epsilon \otimes \epsilon, & \rho(A_{11}) &= \frac{1}{2} \epsilon \otimes \sigma_z \otimes I, & \rho(A_{18}) &= -\frac{1}{2} \sigma_x \otimes \epsilon \otimes \sigma_z, \\ \rho(A_5) &= \frac{1}{2} \epsilon \otimes I \otimes \sigma_x, & \rho(A_{12}) &= -\frac{1}{2} \epsilon \otimes \sigma_x \otimes \sigma_z, & \rho(A_{19}) &= \frac{1}{2} \sigma_z \otimes \epsilon \otimes I, \\ \rho(A_6) &= \frac{1}{2} \epsilon \otimes I \otimes \sigma_z, & \rho(A_{13}) &= \frac{1}{2} \epsilon \otimes \sigma_x \otimes \sigma_x, & \rho(A_{20}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \epsilon, \\ \rho(A_7) &= \frac{1}{2} \epsilon \otimes \sigma_x \otimes I, & \rho(A_{14}) &= \frac{1}{2} I \otimes \sigma_x \otimes \epsilon, & \rho(A_{21}) &= \frac{1}{2} \sigma_z \otimes \sigma_z \otimes \epsilon. \end{aligned} \tag{8.7}$$

We also write down the corresponding generators of the vectorial representation τ , which is the 7-dimensional irreducible component Λ_7 of the representation $\rho \wedge \rho$, which decomposes as $\Lambda^2 S = \Lambda_{21} \oplus \Lambda_7$. These generators read

$$\begin{aligned} \tau(A_1) &= E_{31} - E_{13}, & \tau(A_8) &= E_{37} - E_{73}, & \tau(A_{15}) &= E_{75} - E_{57}, \\ \tau(A_2) &= E_{12} - E_{21}, & \tau(A_9) &= E_{72} - E_{27}, & \tau(A_{16}) &= E_{14} - E_{41}, \\ \tau(A_3) &= E_{32} - E_{23}, & \tau(A_{10}) &= E_{76} - E_{67}, & \tau(A_{17}) &= E_{34} - E_{43}, \\ \tau(A_4) &= E_{61} - E_{16}, & \tau(A_{11}) &= E_{51} - E_{15}, & \tau(A_{18}) &= E_{42} - E_{24}, \\ \tau(A_5) &= E_{36} - E_{63}, & \tau(A_{12}) &= E_{53} - E_{35}, & \tau(A_{19}) &= E_{46} - E_{64}, \\ \tau(A_6) &= E_{62} - E_{26}, & \tau(A_{13}) &= E_{25} - E_{52}, & \tau(A_{20}) &= E_{47} - E_{74}, \\ \tau(A_7) &= E_{17} - E_{71}, & \tau(A_{14}) &= E_{65} - E_{56}, & \tau(A_{21}) &= E_{54} - E_{45}, \end{aligned} \tag{8.8}$$

where E_{ij} are 7×7 matrices with all zero entries, except at the i th- j th entry, where 1 resides.

We are again in a position ready for application of our Lemma 2.1. Given the representations $(\rho, S = \mathbb{R}^8)$ and $(\tau, R = \Lambda_7)$ of $\mathfrak{so}(0,7)$, we solve the magical equation (2.1) for $\omega = \frac{1}{2} \omega^i{}_{\mu\nu} e_i \otimes f^\mu \wedge f^\nu$. In this way, we obtain the seven 2-forms $\omega^i = -\frac{1}{2} \omega^i{}_{\mu\nu} dx^\mu \wedge dx^\nu$

on $N = \mathbb{R}^8$, with coordinates $(x^\mu)_{\mu=1}^8$, which read as follows:

$$\begin{aligned}
 \omega^1 &= -dx^1 \wedge dx^2 - dx^3 \wedge dx^4 + dx^5 \wedge dx^6 + dx^7 \wedge dx^8, \\
 \omega^2 &= dx^1 \wedge dx^3 - dx^2 \wedge dx^4 - dx^5 \wedge dx^7 + dx^6 \wedge dx^8, \\
 \omega^3 &= -dx^1 \wedge dx^4 - dx^2 \wedge dx^3 + dx^5 \wedge dx^8 + dx^6 \wedge dx^7, \\
 \omega^4 &= dx^1 \wedge dx^5 + dx^2 \wedge dx^6 + dx^3 \wedge dx^7 + dx^4 \wedge dx^8, \\
 \omega^5 &= -dx^1 \wedge dx^6 + dx^2 \wedge dx^5 + dx^3 \wedge dx^8 - dx^4 \wedge dx^7, \\
 \omega^6 &= dx^1 \wedge dx^7 + dx^2 \wedge dx^8 - dx^3 \wedge dx^5 - dx^4 \wedge dx^6, \\
 \omega^7 &= dx^1 \wedge dx^8 - dx^2 \wedge dx^7 + dx^3 \wedge dx^6 - dx^4 \wedge dx^5.
 \end{aligned}
 \tag{8.9}$$

These, via the contactification, lead to the following theorem.

Theorem 8.5. *Let $M = \mathbb{R}^{15}$ with coordinates $(u^1, \dots, u^7, x^1, \dots, x^8)$, and consider seven 1-forms $\lambda^i, \dots, \lambda^7$ on M given by*

$$\begin{aligned}
 \lambda^1 &= du^1 - x^1 dx^2 - x^3 dx^4 + x^5 dx^6 + x^7 dx^8, \\
 \lambda^2 &= du^2 + x^1 dx^3 - x^2 dx^4 - x^5 dx^7 + x^6 dx^8, \\
 \lambda^3 &= du^3 - x^1 dx^4 - x^2 dx^3 + x^5 dx^8 + x^6 dx^7, \\
 \lambda^4 &= du^4 + x^1 dx^5 + x^2 dx^6 + x^3 dx^7 + x^4 dx^8, \\
 \lambda^5 &= du^5 - x^1 dx^6 + x^2 dx^5 + x^3 dx^8 - x^4 dx^7, \\
 \lambda^6 &= du^6 + x^1 dx^7 + x^2 dx^8 - x^3 dx^5 - x^4 dx^6, \\
 \lambda^7 &= du^7 + x^1 dx^8 - x^2 dx^7 + x^3 dx^6 - x^4 dx^5.
 \end{aligned}$$

The rank 8 distribution \mathcal{D} on M defined as $\mathcal{D} = \{T\mathbb{R}^{15} \ni X \mid X \lrcorner \lambda^1 = \dots = X \lrcorner \lambda^7 = 0\}$ has its Lie algebra of infinitesimal automorphisms $\mathfrak{aut}(\mathcal{D})$ isomorphic to the Tanaka prolongation of $\mathfrak{n}_- = R \oplus S$, where $(\rho, S = \mathbb{R}^8)$ is the spinorial representation (8.7) of $\mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(0,7)$, and $(\tau, R = \mathbb{R}^7)$ is the vectorial representation (8.8) of \mathfrak{n}_{00} .

The symmetry algebra $\mathfrak{aut}(\mathcal{D})$ is isomorphic to the simple Lie algebra \mathfrak{f}_{II} ,

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{f}_{II},$$

having the following natural gradation

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S$,

$$\mathfrak{n}_0 = \mathfrak{n}_{00} = \mathbb{R} \oplus \mathfrak{so}(0,7),$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$, which is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in \mathfrak{f}_{II} .

The contactification (M, \mathcal{D}) is locally a flat model for the parabolic geometry of type (F_{II}, P_{II}) related to the following crossed Satake diagram $\bullet \rightleftarrows \bullet \circ$ of \mathfrak{f}_{II} .

Remark 8.6. In this way, we realised the real form \mathfrak{f}_{II} of the simple exceptional complex Lie algebra \mathfrak{f}_4 in $M = \mathbb{R}^{15}$ as a symmetry algebra of the Pfaffian system $(\lambda^1, \dots, \lambda^7)$. This realisation does not appear in Cartan’s theses.

Remark 8.7. Our present case of f_{II} also admits description in terms of a certain $\mathbb{R} \oplus \mathfrak{so}(0,7)$ invariant 4-form Φ in $S = \mathbb{R}^8$, analogous to the 4-form Φ introduced in Section 8.1.4, when we discussed f_I . Skipping the details, we only mention that now Φ may be represented by

$$\Phi = h_{ij}\omega^i \wedge \omega^j,$$

where ω^i are given by (8.9) and

$$(h_{ij}) = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}.$$

Explicitly, the form Φ reads

$$\begin{aligned} -\frac{1}{6}\Phi = & dx^1 \wedge dx^2 \wedge dx^3 \wedge dx^4 - dx^1 \wedge dx^2 \wedge dx^5 \wedge dx^6 - \\ & dx^1 \wedge dx^2 \wedge dx^7 \wedge dx^8 - dx^1 \wedge dx^3 \wedge dx^5 \wedge dx^7 + \\ & dx^1 \wedge dx^3 \wedge dx^6 \wedge dx^8 - dx^1 \wedge dx^4 \wedge dx^5 \wedge dx^8 - \\ & dx^1 \wedge dx^4 \wedge dx^6 \wedge dx^7 - dx^2 \wedge dx^3 \wedge dx^5 \wedge dx^8 - \\ & dx^2 \wedge dx^3 \wedge dx^6 \wedge dx^7 + dx^2 \wedge dx^4 \wedge dx^5 \wedge dx^7 - \\ & dx^2 \wedge dx^4 \wedge dx^6 \wedge dx^8 - dx^3 \wedge dx^4 \wedge dx^5 \wedge dx^6 - \\ & dx^3 \wedge dx^4 \wedge dx^7 \wedge dx^8 + dx^5 \wedge dx^6 \wedge dx^7 \wedge dx^8. \end{aligned} \tag{8.10}$$

This 4-form alone encaptures all the features of the f_{II} symmetric contactification we discussed in the entire Section 8.2. In particular, analogous statements as in Remark 8.4, with now $\mathfrak{so}(4,3)$ replaced by $\mathfrak{so}(0,7)$, apply to the present 4-form Φ .


9. Spinorial representations in dimension 8

Dimension *eight* is quite exceptional, as, for example, 8 is the highest possible dimension for the existence of Euclidean Hurwitz algebras, gifting us with the algebra of *octonions*. From the perspective of our paper, which meanders through the realm of simple Lie algebras, eight is *very* special: among all the complex simple Lie algebras, the Dynkin diagram of $\mathfrak{d}_4 = \mathfrak{so}(8, \mathbb{C})$ which is *defined* in dimension *eight*, is the most symmetric:



Visibly it has a threefold symmetry S_3 .

The Lie algebra $\mathfrak{so}(8, \mathbb{C})$ has six real forms. These are: $\mathfrak{so}(8, 0)$, $\mathfrak{so}(7, 1)$, $\mathfrak{so}(6, 2)$, $\mathfrak{so}^*(8)$, $\mathfrak{so}(5, 3)$ and $\mathfrak{so}(4, 4)$, with the following respective Satake diagrams:



We see that among these Satake diagrams, the only ones that share the S_3 symmetry of the Dynkin diagram of the complex algebra \mathfrak{d}_4 are those of the compact real form $\mathfrak{so}(8, 0)$ and of the split real form $\mathfrak{so}(4, 4)$.

This S_3 symmetry of these two diagrams indicates that the lowest dimensional real representations of $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$ may have additional features when compared with spinorial representations of other $\mathfrak{so}(p, q)$ s. In particular, for both $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$, we have the following:

- Their Dirac representation (ρ, S) in the real vector space $S = \mathbb{R}^{16}$ is reducible over \mathbb{R} and is split into two real Weyl representations (ρ_+, S_+) and (ρ_-, S_-) in the respective vector spaces of Weyl spinors $S_+ = \mathbb{R}^8$ and $S_- = \mathbb{R}^8$, which have the same real dimension *eight*,

$$\rho = \rho_+ \oplus \rho_- \quad \text{in} \quad S = S_+ \oplus S_-, \quad \dim_{\mathbb{R}} S_{\pm} = 8.$$

- The real Weyl representations (ρ_{\pm}, S_{\pm}) are *faithful, irreducible and nonequivalent*.
- The defining representations (τ, R) of $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$, as the algebra of endomorphisms in the space $R = \mathbb{R}^8$ of vectors preserving the bilinear form of respective signatures $(4, 4)$ and $(8, 0)$ has the same dimension *eight* as the two Weyl representations (ρ_{\pm}, S_{\pm}) .
- The real defining representations (τ, R) are *irreducible* for both $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$.
- All three real 8-dimensional irreducible representations (ρ_+, S_+) , (ρ_-, S_-) and (τ, R) of, respectively, both $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$ are *pairwise nonequivalent*.

Thus, the Lie algebras $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$ have three real, irreducible and nonequivalent representations (ρ_+, ρ_-, τ) in the vector space \mathbb{R}^8 of the defining dimension $p + q = 8$. For all $\mathfrak{so}(p, q)$ Lie algebras, this is the only dimension $p + q$ that such situation with the irreducible representations occurs.

Below, we provide the explicit description of the *triatlity representations* (ρ_+, ρ_-, τ) separately for $\mathfrak{so}(4, 4)$ and $\mathfrak{so}(8, 0)$.

9.1. Triatlity representations of $\mathfrak{so}(4, 4)$

We recall from Section 5 that the Lie algebra $\mathfrak{so}(4, 4)$ admits a representation ρ in the 16-dimensional real vector space $S = \mathbb{R}^{16}$ of Dirac spinors. This is obtained by using the Dirac γ matrices generating the representation of the Clifford algebra $\mathcal{Cl}(4, 4)$. In terms of the 2-dimensional Pauli matrices $(\sigma_x, \epsilon, \sigma_z, I)$, these look as follows:

$$\begin{aligned}
 \gamma_1 &= \sigma_x \otimes \sigma_x \otimes \sigma_x \otimes \sigma_x \\
 \gamma_2 &= \sigma_x \otimes \sigma_x \otimes \sigma_x \otimes \epsilon \\
 \gamma_3 &= \sigma_x \otimes \sigma_x \otimes \sigma_x \otimes \sigma_z \\
 \gamma_4 &= \sigma_x \otimes \sigma_x \otimes \epsilon \otimes I \\
 \gamma_5 &= \sigma_x \otimes \sigma_x \otimes \sigma_z \otimes I \\
 \gamma_6 &= \sigma_x \otimes \epsilon \otimes I \otimes I \\
 \gamma_7 &= \sigma_x \otimes \sigma_z \otimes I \otimes I \\
 \gamma_8 &= \epsilon \otimes I \otimes I \otimes I.
 \end{aligned}
 \tag{9.1}$$

They satisfy the *Dirac identity*

$$\gamma_i \gamma_j + \gamma_j \gamma_i = 2g_{ij}(I \otimes I \otimes I \otimes I), \quad i, j = 1, \dots, 8,
 \tag{9.2}$$

with

$$(g_{ij}) = \text{diag}(1, -1, 1, -1, 1, -1, 1, -1).$$

The 28 generators of $\mathfrak{so}(4,4)$ in the Majorana-Dirac spinor representation ρ in the space of Dirac spinors $S = \mathbb{R}^{16}$ are given by

$$\rho(A_{I(i,j)}) = \frac{1}{2} \gamma_i \gamma_j, \quad 1 \leq i < j \leq 8,$$

where we again have used the function $I = I(i,j)$ defined in (8.1). Note that since now $i < j$ can run from 1 to 8, the function has a range from 1 to 28. We add to these generators the scaling generator, $\rho(A_{29})$,

$$\rho(A_{29}) = \frac{1}{2} I \otimes I \otimes I \otimes I.$$

This extends the Dirac representation ρ of the Lie algebra $\mathfrak{so}(4,4)$ to the representation of the *homothety* Lie algebra $\mathfrak{co}(4,4) = \mathbb{R} \oplus \mathfrak{so}(4,4)$.

In terms of the 2-dimensional Pauli matrices, these generators look like

$$\begin{aligned}
 \rho(A_1) &= \frac{1}{2} I \otimes I \otimes I \otimes \sigma_z, & \rho(A_{15}) &= \frac{1}{2} I \otimes \sigma_z \otimes \sigma_z \otimes I, \\
 \rho(A_2) &= \frac{1}{2} I \otimes I \otimes I \otimes \epsilon, & \rho(A_{16}) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_x \otimes \sigma_x, \\
 \rho(A_3) &= \frac{1}{2} I \otimes I \otimes I \otimes \sigma_x, & \rho(A_{17}) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_x \otimes \epsilon, \\
 \rho(A_4) &= \frac{1}{2} I \otimes I \otimes \sigma_z \otimes \sigma_x, & \rho(A_{18}) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_x \otimes \sigma_z, \\
 \rho(A_5) &= \frac{1}{2} I \otimes I \otimes \sigma_z \otimes \epsilon, & \rho(A_{19}) &= \frac{1}{2} I \otimes \epsilon \otimes \epsilon \otimes I, \\
 \rho(A_6) &= \frac{1}{2} I \otimes I \otimes \sigma_z \otimes \sigma_z, & \rho(A_{20}) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_z \otimes I, \\
 \rho(A_7) &= \frac{1}{2} I \otimes I \otimes \epsilon \otimes \sigma_x, & \rho(A_{21}) &= \frac{1}{2} I \otimes \sigma_x \otimes I \otimes I, \\
 \rho(A_8) &= \frac{1}{2} I \otimes I \otimes \epsilon \otimes \epsilon, & \rho(A_{22}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \sigma_x \otimes \sigma_x, \\
 \rho(A_9) &= \frac{1}{2} I \otimes I \otimes \epsilon \otimes \sigma_z, & \rho(A_{23}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \sigma_x \otimes \epsilon, \\
 \rho(A_{10}) &= \frac{1}{2} I \otimes I \otimes \sigma_x \otimes I, & \rho(A_{24}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \sigma_x \otimes \sigma_z, \\
 \rho(A_{11}) &= \frac{1}{2} I \otimes \sigma_z \otimes \sigma_x \otimes \sigma_x, & \rho(A_{25}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \epsilon \otimes I, \\
 \rho(A_{12}) &= \frac{1}{2} I \otimes \sigma_z \otimes \sigma_x \otimes \epsilon, & \rho(A_{26}) &= \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \sigma_z \otimes I, \\
 \rho(A_{13}) &= \frac{1}{2} I \otimes \sigma_z \otimes \sigma_x \otimes \sigma_z, & \rho(A_{27}) &= \frac{1}{2} \sigma_z \otimes \epsilon \otimes I \otimes I, \\
 \rho(A_{14}) &= \frac{1}{2} I \otimes \sigma_z \otimes \epsilon \otimes I, & \rho(A_{28}) &= \frac{1}{2} \sigma_z \otimes \sigma_z \otimes I \otimes I.
 \end{aligned}
 \tag{9.3}$$

Looking at the first factor in *all* of these generators, we observe that it is either I or σ_z , (i.e., it is *diagonal*). This means that this 16-dimensional representation of $\mathbb{R} \oplus \mathfrak{so}(4,4)$ is *reducible*. It *splits* onto two real 8-dimensional *Weyl representations*

$$\rho = \rho_+ \oplus \rho_- \quad \text{in} \quad S = S_+ \oplus S_-, \quad \dim_{\mathbb{R}} S_{\pm} = 8$$

in the spaces S_{\pm} of (Majorana)-Weyl spinors.

On generators of $\mathfrak{so}(4,4)$, these two 8-dimensional representations ρ_{\pm} are given by

$$\begin{aligned} \rho_{\pm}(A_1) &= \frac{1}{2}I \otimes I \otimes \sigma_z, & \rho_{\pm}(A_{15}) &= \frac{1}{2}\sigma_z \otimes \sigma_z \otimes I, \\ \rho_{\pm}(A_2) &= \frac{1}{2}I \otimes I \otimes \epsilon, & \rho_{\pm}(A_{16}) &= \frac{1}{2}\epsilon \otimes \sigma_x \otimes \sigma_x, \\ \rho_{\pm}(A_3) &= \frac{1}{2}I \otimes I \otimes \sigma_x, & \rho_{\pm}(A_{17}) &= \frac{1}{2}\epsilon \otimes \sigma_x \otimes \epsilon, \\ \rho_{\pm}(A_4) &= \frac{1}{2}I \otimes \sigma_z \otimes \sigma_x, & \rho_{\pm}(A_{18}) &= \frac{1}{2}\epsilon \otimes \sigma_x \otimes \sigma_z, \\ \rho_{\pm}(A_5) &= \frac{1}{2}I \otimes \sigma_z \otimes \epsilon, & \rho_{\pm}(A_{19}) &= \frac{1}{2}\epsilon \otimes \epsilon \otimes I, \\ \rho_{\pm}(A_6) &= \frac{1}{2}I \otimes \sigma_z \otimes \sigma_z, & \rho_{\pm}(A_{20}) &= \frac{1}{2}\epsilon \otimes \sigma_z \otimes I, \\ \rho_{\pm}(A_7) &= \frac{1}{2}I \otimes \epsilon \otimes \sigma_x, & \rho_{\pm}(A_{21}) &= \frac{1}{2}\sigma_x \otimes I \otimes I, \\ \rho_{\pm}(A_8) &= \frac{1}{2}I \otimes \epsilon \otimes \epsilon, & \rho_{\pm}(A_{22}) &= \pm \frac{1}{2}\sigma_x \otimes \sigma_x \otimes \sigma_x, \\ \rho_{\pm}(A_9) &= \frac{1}{2}I \otimes \epsilon \otimes \sigma_z, & \rho_{\pm}(A_{23}) &= \pm \frac{1}{2}\sigma_x \otimes \sigma_x \otimes \epsilon, \\ \rho_{\pm}(A_{10}) &= \frac{1}{2}I \otimes \sigma_x \otimes I, & \rho_{\pm}(A_{24}) &= \pm \frac{1}{2}\sigma_x \otimes \sigma_x \otimes \sigma_z, \\ \rho_{\pm}(A_{11}) &= \frac{1}{2}\sigma_z \otimes \sigma_x \otimes \sigma_x, & \rho_{\pm}(A_{25}) &= \pm \frac{1}{2}\sigma_x \otimes \epsilon \otimes I, \\ \rho_{\pm}(A_{12}) &= \frac{1}{2}\sigma_z \otimes \sigma_x \otimes \epsilon, & \rho_{\pm}(A_{26}) &= \pm \frac{1}{2}\sigma_x \otimes \sigma_z \otimes I, \\ \rho_{\pm}(A_{13}) &= \frac{1}{2}\sigma_z \otimes \sigma_x \otimes \sigma_z, & \rho_{\pm}(A_{27}) &= \pm \frac{1}{2}\epsilon \otimes I \otimes I, \\ \rho_{\pm}(A_{14}) &= \frac{1}{2}\sigma_z \otimes \epsilon \otimes I, & \rho_{\pm}(A_{28}) &= \pm \frac{1}{2}\sigma_z \otimes I \otimes I. \end{aligned} \tag{9.4}$$

We extend them to $\mathbb{R} \oplus \mathfrak{so}(4,4)$ by adding

$$\rho_{\pm}(A_{29}) = \frac{1}{2}I \otimes I \otimes I.$$

It follows that the Weyl representations (ρ_{\pm}, S_{\pm}) of $\mathfrak{so}(4,4)$ are *irreducible* and *nonequivalent*.

They can be used to find yet another real 8-dimensional representation of $\mathfrak{so}(4,4)$. For this, one considers the tensor product representation

$$\rho_+ \otimes \rho_-.$$

This 64-dimensional real representation of $\mathfrak{so}(4,4)$ is *reducible*. It decomposes as

$$\rho_+ \otimes \rho_- = \alpha \oplus \tau \quad \text{in} \quad S_+ \otimes S_- = T_{56} \oplus R, \quad \text{with} \quad \dim_{\mathbb{R}}(R) = 8, \dim_{\mathbb{R}}(T_{56}) = 56,$$

having irreducible components (α, T_{56}) and (τ, R) of respective dimensions 56 and 8. Explicitly, on generators of $\mathbb{R} \oplus \mathfrak{so}(4,4)$, the 8-dimensional representation τ reads

$$\begin{aligned} \tau(A_1) &= E_{66} - E_{22}, \\ \tau(A_2) &= \frac{1}{2}(E_{23} - E_{32} + E_{25} - E_{52} + E_{36} - E_{63} + E_{56} - E_{65}), \\ \tau(A_3) &= \frac{1}{2}(E_{23} + E_{32} + E_{25} + E_{52} + E_{36} + E_{63} + E_{56} + E_{65}), \end{aligned}$$

$$\begin{aligned}
 \tau(A_4) &= \frac{1}{2}(E_{23} + E_{32} - E_{25} - E_{52} - E_{36} - E_{63} + E_{56} + E_{65}), \\
 \tau(A_5) &= \frac{1}{2}(E_{23} - E_{32} - E_{25} + E_{52} - E_{36} + E_{63} + E_{56} - E_{65}), \\
 \tau(A_6) &= E_{33} - E_{55}, \\
 \tau(A_7) &= \frac{1}{2}(E_{12} - E_{21} - E_{16} + E_{61} - E_{27} + E_{72} + E_{67} - E_{76}), \\
 \tau(A_8) &= \frac{1}{2}(-E_{12} - E_{21} - E_{16} - E_{61} - E_{27} - E_{72} - E_{67} - E_{76}), \\
 \tau(A_9) &= \frac{1}{2}(E_{13} - E_{31} + E_{15} - E_{51} - E_{37} + E_{73} - E_{57} + E_{75}), \\
 \tau(A_{10}) &= \frac{1}{2}(E_{13} + E_{31} - E_{15} - E_{51} + E_{37} + E_{73} - E_{57} - E_{75}), \\
 \tau(A_{11}) &= \frac{1}{2}(-E_{12} - E_{21} + E_{16} + E_{61} + E_{27} + E_{72} - E_{67} - E_{76}), \\
 \tau(A_{12}) &= \frac{1}{2}(E_{12} - E_{21} + E_{16} - E_{61} + E_{27} - E_{72} + E_{67} - E_{76}), \\
 \tau(A_{13}) &= \frac{1}{2}(-E_{13} - E_{31} - E_{15} - E_{51} + E_{37} + E_{73} + E_{57} + E_{75}), \\
 \tau(A_{14}) &= \frac{1}{2}(-E_{13} + E_{31} + E_{15} - E_{51} - E_{37} + E_{73} + E_{57} - E_{75}), \\
 \tau(A_{15}) &= E_{11} - E_{77}, \\
 \tau(A_{16}) &= \frac{1}{2}(E_{24} - E_{42} + E_{28} - E_{82} + E_{46} - E_{64} - E_{68} + E_{86}), \\
 \tau(A_{17}) &= \frac{1}{2}(E_{24} + E_{42} + E_{28} + E_{82} + E_{46} + E_{64} + E_{68} + E_{86}), \\
 \tau(A_{18}) &= \frac{1}{2}(E_{34} - E_{43} + E_{38} - E_{83} - E_{45} + E_{54} + E_{58} - E_{85}), \\
 \tau(A_{19}) &= \frac{1}{2}(-E_{34} - E_{43} - E_{38} - E_{83} + E_{45} + E_{54} + E_{58} + E_{85}), \\
 \tau(A_{20}) &= \frac{1}{2}(-E_{14} + E_{41} - E_{18} + E_{81} + E_{47} - E_{74} - E_{78} + E_{87}), \\
 \tau(A_{21}) &= \frac{1}{2}(E_{14} + E_{41} + E_{18} + E_{81} - E_{47} - E_{74} - E_{78} - E_{87}), \\
 \tau(A_{22}) &= \frac{1}{2}(-E_{24} - E_{42} + E_{28} + E_{82} + E_{46} + E_{64} - E_{68} - E_{86}), \\
 \tau(A_{23}) &= \frac{1}{2}(-E_{24} + E_{42} + E_{28} - E_{82} + E_{46} - E_{64} + E_{68} - E_{86}), \\
 \tau(A_{24}) &= \frac{1}{2}(-E_{34} - E_{43} + E_{38} + E_{83} - E_{45} - E_{54} + E_{58} + E_{85}), \\
 \tau(A_{25}) &= \frac{1}{2}(E_{34} - E_{43} - E_{38} + E_{83} + E_{45} - E_{54} + E_{58} - E_{85}), \\
 \tau(A_{26}) &= \frac{1}{2}(E_{14} + E_{41} - E_{18} - E_{81} + E_{47} + E_{74} - E_{78} - E_{87}), \\
 \tau(A_{27}) &= \frac{1}{2}(-E_{14} + E_{41} + E_{18} - E_{81} - E_{47} + E_{74} - E_{78} + E_{87}), \\
 \tau(A_{28}) &= -E_{44} + E_{88}, \\
 \tau(A_{29}) &= E_{11} + E_{22} + E_{33} + E_{44} + E_{55} + E_{66} + E_{77} + E_{88},
 \end{aligned}
 \tag{9.5}$$

where E_{ij} , $i, j = 1, 2, \dots, 8$ denote 8×8 matrices with zeroes everywhere except the value 1 in the entry (i, j) seating at the crossing of the i th row and the j th column.

The three real, irreducible, pairwise nonequivalent representations (ρ_+, ρ_-, τ) of $\mathfrak{so}(4, 4)$, given by the formulas (9.4) and (9.5), constitute the set of the *triality representations* for $\mathfrak{so}(4, 4)$.

9.2. Triality representations of $\mathfrak{so}(8, 0)$

To get the real representation (ρ, S) of $\mathfrak{so}(8, 0)$ in the space $S = \mathbb{R}^{16}$ of Dirac spinors, we need the real Dirac γ matrices satisfying the *Dirac identity* (9.2), but now with

$$g_{ij} = \delta_{ij},$$

where δ_{ij} is the *Kronecker delta* in 8 dimensions.

Thus, we need to modify the Dirac matrices γ_i from (9.1) to have the proper signature of the metric. This is done in *two* steps [19]. *First*, one puts the *imaginary unit* i in front

of some of the Dirac matrices γ_i generating the Clifford algebra $\mathcal{Cl}(4,4)$ to get the proper signature of (g_{ij}) . Although this produces few *complex* generators, *in step two*, one uses them with the others and modifies them in an algorithmic fashion so that they become all real and still satisfy the *Dirac identity* with the proper signature of (g_{ij}) . Explicitly, it is done as follows:

By placing the *imaginary unit* i in front of $\gamma_2, \gamma_4, \gamma_6$ and γ_8 in (9.1), we obtain 8 matrices

$$\tilde{\gamma}_{2j-1} = \gamma_{2j-1}, \quad \tilde{\gamma}_{2j} = i\gamma_{2j}, \quad j = 1, 2, 3, 4,$$

with $\gamma_i, i, 1, \dots, 8$, in (9.1). These constitute generators of the *complex* 16-dimensional representation of the Clifford algebra $\mathcal{Cl}(8,0)$. We will also need the representation of this Clifford algebra, which is *complex conjugate* of $\tilde{\gamma}$. This is generated by

$$\bar{\tilde{\gamma}}_{2j-1} = \gamma_{2j-1}, \quad \bar{\tilde{\gamma}}_{2j} = -\gamma_{2j}, \quad j = 1, 2, 3, 4.$$

The Clifford algebra representations generated by the Dirac matrices $\tilde{\gamma}$ and $\bar{\tilde{\gamma}}$ are *real equivalent* – that is, there exists a real 16×16 matrix B such that

$$B\tilde{\gamma}_i = \bar{\tilde{\gamma}}_i B, \quad \forall i = 1, \dots, 8.$$

It can be chosen so that

$$B^2 = \text{Id},$$

where $\text{Id} = I \otimes I \otimes I \otimes I$.

Explicitly,

$$B = \sigma_z \otimes \epsilon \otimes \sigma_z \otimes \epsilon.$$

Using this matrix, we define a *new* set of eight γ matrices⁶ by

$$\gamma_i = (iB + \text{Id}) \tilde{\gamma}_i (iB + \text{Id})^{-1}, \quad \forall i = 1, \dots, 8.$$

One can check that these 8 matrices are *all real* and that they satisfy the desired Dirac identity:

$$\gamma_i \gamma_j + \gamma_j \gamma_i = 2\delta_{ij} (I \otimes I \otimes I \otimes I), \quad i, j = 1, \dots, 8.$$

Explicitly, we have

$$\begin{aligned} \gamma_1 &= \sigma_x \otimes \sigma_x \otimes \sigma_x \otimes \sigma_x \\ \gamma_2 &= -\epsilon \otimes \sigma_z \otimes \epsilon \otimes I \\ \gamma_3 &= \sigma_x \otimes \sigma_x \otimes \sigma_x \otimes \sigma_z \\ \gamma_4 &= \epsilon \otimes \sigma_z \otimes \sigma_x \otimes \epsilon \\ \gamma_5 &= \sigma_x \otimes \sigma_x \otimes \sigma_z \otimes I \\ \gamma_6 &= -\epsilon \otimes I \otimes \sigma_z \otimes \epsilon \\ \gamma_7 &= \sigma_x \otimes \sigma_z \otimes I \otimes I \\ \gamma_8 &= \sigma_x \otimes \epsilon \otimes \sigma_z \otimes \epsilon. \end{aligned}$$

⁶The γ -matrices used below should be considered as new symbols and should not be confused with the $\mathfrak{so}(4,4)$ γ -matrices in formulas defining $\tilde{\gamma}$ -matrices at the beginning of this section. One should forget about the definition of $\tilde{\gamma}_8$ in the formula below.

The 28 generators of $\mathfrak{so}(8,0)$ in the Majorana-Dirac spinor representation ρ in the space of Dirac spinors $S = \mathbb{R}^{16}$ are given by

$$\rho(A_{I(i,j)}) = \frac{1}{2}\gamma_i\gamma_j, \quad 1 \leq i < j \leq 8,$$

where again we have used the function $I = I(i,j)$ defined in (8.1). Note that since now $i < j$ can run from 1 to 8, the function has a range from 1 to 28. We add to this the scaling generator, $\rho(A_{29})$, extending the Lie algebra $\mathfrak{so}(4,4)$ to $\mathfrak{coa}(4,4)$, given by

$$\rho(A_{29}) = \frac{1}{2}I \otimes I \otimes I \otimes I.$$

In terms of the 2-dimensional Pauli matrices, these generators look like

$$\begin{aligned} \rho(A_1) &= -\frac{1}{2}\sigma_z \otimes \epsilon \otimes \sigma_z \otimes \sigma_x, & \rho(A_{15}) &= -\frac{1}{2}\sigma_z \otimes \sigma_x \otimes I \otimes \epsilon, \\ \rho(A_2) &= \frac{1}{2}I \otimes I \otimes I \otimes \epsilon, & \rho(A_{16}) &= \frac{1}{2}I \otimes \epsilon \otimes \sigma_x \otimes \sigma_x, \\ \rho(A_3) &= \frac{1}{2}\sigma_z \otimes \epsilon \otimes \sigma_z \otimes \sigma_z, & \rho(A_{17}) &= \frac{1}{2}\sigma_z \otimes I \otimes \epsilon \otimes I, \\ \rho(A_4) &= \frac{1}{2}\sigma_z \otimes \epsilon \otimes I \otimes \sigma_z, & \rho(A_{18}) &= \frac{1}{2}I \otimes \epsilon \otimes \sigma_x \otimes \sigma_z, \\ \rho(A_5) &= -\frac{1}{2}I \otimes I \otimes \sigma_z \otimes \epsilon, & \rho(A_{19}) &= -\frac{1}{2}\sigma_z \otimes I \otimes \sigma_x \otimes \epsilon, \\ \rho(A_6) &= -\frac{1}{2}\sigma_z \otimes \epsilon \otimes I \otimes \sigma_x, & \rho(A_{20}) &= \frac{1}{2}I \otimes \epsilon \otimes \sigma_z \otimes I, \\ \rho(A_7) &= \frac{1}{2}I \otimes I \otimes \epsilon \otimes \sigma_x, & \rho(A_{21}) &= \frac{1}{2}\sigma_z \otimes \sigma_z \otimes \sigma_z \otimes \epsilon, \\ \rho(A_8) &= -\frac{1}{2}\sigma_z \otimes \epsilon \otimes \sigma_x \otimes I, & \rho(A_{22}) &= \frac{1}{2}I \otimes \sigma_z \otimes \epsilon \otimes \sigma_z, \\ \rho(A_9) &= \frac{1}{2}I \otimes I \otimes \epsilon \otimes \sigma_z, & \rho(A_{23}) &= -\frac{1}{2}\sigma_z \otimes \sigma_x \otimes \sigma_x \otimes \epsilon, \\ \rho(A_{10}) &= \frac{1}{2}\sigma_z \otimes \epsilon \otimes \epsilon \otimes \epsilon, & \rho(A_{24}) &= -\frac{1}{2}I \otimes \sigma_z \otimes \epsilon \otimes \sigma_x, \\ \rho(A_{11}) &= -\frac{1}{2}\sigma_z \otimes \sigma_x \otimes \epsilon \otimes \sigma_z, & \rho(A_{25}) &= -\frac{1}{2}\sigma_z \otimes \sigma_x \otimes \epsilon \otimes I, \\ \rho(A_{12}) &= -\frac{1}{2}I \otimes \sigma_z \otimes \sigma_x \otimes \epsilon, & \rho(A_{26}) &= \frac{1}{2}I \otimes \sigma_z \otimes I \otimes \epsilon, \\ \rho(A_{13}) &= \frac{1}{2}\sigma_z \otimes \sigma_x \otimes \epsilon \otimes \sigma_x, & \rho(A_{27}) &= -\frac{1}{2}\sigma_z \otimes \epsilon \otimes I \otimes I, \\ \rho(A_{14}) &= -\frac{1}{2}I \otimes \sigma_z \otimes \epsilon \otimes I, & \rho(A_{28}) &= -\frac{1}{2}I \otimes \sigma_x \otimes \sigma_z \otimes \epsilon. \end{aligned} \tag{9.6}$$

Similarly, as in the case of $\mathfrak{so}(4,4)$, this 16-dimensional representation of $\mathbb{R} \oplus \mathfrak{so}(4,4)$ is *reducible*, again due to the appearance of I and σ_z only as the first factors in the above formulas. It *splits* onto two real 8-dimensional Weyl representations

$$\rho = \rho_+ \oplus \rho_- \quad \text{in} \quad S = S_+ \oplus S_-, \quad \dim_{\mathbb{R}} S_{\pm} = 8.$$

On generators of $\mathfrak{so}(8,0)$, these two 8-dimensional representations ρ_{\pm} , are given by

$$\begin{aligned} \rho_{\pm}(A_1) &= \mp \frac{1}{2}\epsilon \otimes \sigma_z \otimes \sigma_x, & \rho_{\pm}(A_{15}) &= \mp \frac{1}{2}\sigma_x \otimes I \otimes \epsilon, \\ \rho_{\pm}(A_2) &= \frac{1}{2}I \otimes I \otimes \epsilon, & \rho_{\pm}(A_{16}) &= \frac{1}{2}\epsilon \otimes \sigma_x \otimes \sigma_x, \\ \rho_{\pm}(A_3) &= \pm \frac{1}{2}\epsilon \otimes \sigma_z \otimes \sigma_z, & \rho_{\pm}(A_{17}) &= \pm \frac{1}{2}I \otimes \epsilon \otimes I, \end{aligned}$$

$$\begin{aligned}
 \rho_{\pm}(A_4) &= \pm \frac{1}{2} \epsilon \otimes I \otimes \sigma_z, & \rho_{\pm}(A_{18}) &= \frac{1}{2} \epsilon \otimes \sigma_x \otimes \sigma_z, \\
 \rho_{\pm}(A_5) &= -\frac{1}{2} I \otimes \sigma_z \otimes \epsilon, & \rho_{\pm}(A_{19}) &= \mp \frac{1}{2} I \otimes \sigma_x \otimes \epsilon, \\
 \rho_{\pm}(A_6) &= \mp \frac{1}{2} \epsilon \otimes I \otimes \sigma_x, & \rho_{\pm}(A_{20}) &= \frac{1}{2} \epsilon \otimes \sigma_z \otimes I, \\
 \rho_{\pm}(A_7) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_x, & \rho_{\pm}(A_{21}) &= \pm \frac{1}{2} \sigma_z \otimes \sigma_x \otimes \epsilon, \\
 \rho_{\pm}(A_8) &= \mp \frac{1}{2} \epsilon \otimes \sigma_x \otimes I, & \rho_{\pm}(A_{22}) &= \frac{1}{2} \sigma_z \otimes \epsilon \otimes \sigma_z, \\
 \rho_{\pm}(A_9) &= \frac{1}{2} I \otimes \epsilon \otimes \sigma_z, & \rho_{\pm}(A_{23}) &= \mp \frac{1}{2} \sigma_x \otimes \sigma_x \otimes \epsilon, \\
 \rho_{\pm}(A_{10}) &= \pm \frac{1}{2} \epsilon \otimes \epsilon \otimes \epsilon, & \rho_{\pm}(A_{24}) &= -\frac{1}{2} \sigma_z \otimes \epsilon \otimes \sigma_x, \\
 \rho_{\pm}(A_{11}) &= \mp \frac{1}{2} \sigma_x \otimes \epsilon \otimes \sigma_z, & \rho_{\pm}(A_{25}) &= \mp \frac{1}{2} \sigma_x \otimes \epsilon \otimes I, \\
 \rho_{\pm}(A_{12}) &= -\frac{1}{2} \sigma_z \otimes \sigma_x \otimes \epsilon, & \rho_{\pm}(A_{26}) &= \frac{1}{2} \sigma_z \otimes I \otimes \epsilon, \\
 \rho_{\pm}(A_{13}) &= \pm \frac{1}{2} \sigma_x \otimes \epsilon \otimes \sigma_x, & \rho_{\pm}(A_{27}) &= \mp \frac{1}{2} \epsilon \otimes I \otimes I, \\
 \rho_{\pm}(A_{14}) &= -\frac{1}{2} \sigma_z \otimes \epsilon \otimes I, & \rho_{\pm}(A_{28}) &= -\frac{1}{2} \sigma_x \otimes \sigma_z \otimes \epsilon.
 \end{aligned}
 \tag{9.7}$$

We extend them em to $\mathbb{R} \oplus \mathfrak{so}(8,0)$ by adding

$$\rho_{\pm}(A_{29}) = \frac{1}{2} I \otimes I \otimes I.$$

It follows that the Weyl representations (ρ_{\pm}, S_{\pm}) of $\mathfrak{so}(4,4)$ are *irreducible* and *nonequivalent*.

We use them to find the defining representation (τ, R) of $\mathfrak{so}(8,0)$ in the vector space $R = \mathbb{R}^8$ of vectors. We again consider the tensor product representation $\rho_+ \otimes \rho_-$. It decomposes as

$$\rho_+ \otimes \rho_- = \alpha \oplus \tau \quad \text{in} \quad S_+ \otimes S_- = T_{56} \oplus R, \quad \text{with} \quad \dim_{\mathbb{R}}(R) = 8, \dim_{\mathbb{R}}(T_{56}) = 56,$$

having irreducible components (α, T_{56}) and (τ, R) of respective dimensions 56 and 8. Explicitly, on generators A_I of $\mathbb{R} \oplus \mathfrak{so}(8,0)$, the 8-dimensional representation τ reads


$$\begin{aligned}
 \tau(A_1) &= E_{38} - E_{83}, & \tau(A_8) &= E_{35} - E_{53}, & \tau(A_{15}) &= E_{52} - E_{25}, & \tau(A_{22}) &= E_{68} - E_{86}, \\
 \tau(A_2) &= E_{78} - E_{87}, & \tau(A_9) &= E_{75} - E_{57}, & \tau(A_{16}) &= E_{18} - E_{81}, & \tau(A_{23}) &= E_{36} - E_{63}, \\
 \tau(A_3) &= E_{37} - E_{73}, & \tau(A_{10}) &= E_{54} - E_{45}, & \tau(A_{17}) &= E_{31} - E_{13}, & \tau(A_{24}) &= E_{76} - E_{67}, \\
 \tau(A_4) &= E_{84} - E_{48}, & \tau(A_{11}) &= E_{28} - E_{82}, & \tau(A_{18}) &= E_{71} - E_{17}, & \tau(A_{25}) &= E_{64} - E_{46}, \\
 \tau(A_5) &= E_{43} - E_{34}, & \tau(A_{12}) &= E_{32} - E_{23}, & \tau(A_{19}) &= E_{14} - E_{41}, & \tau(A_{26}) &= E_{56} - E_{65}, \\
 \tau(A_6) &= E_{47} - E_{74}, & \tau(A_{13}) &= E_{72} - E_{27}, & \tau(A_{20}) &= E_{51} - E_{15}, & \tau(A_{27}) &= E_{26} - E_{62}, \\
 \tau(A_7) &= E_{58} - E_{85}, & \tau(A_{14}) &= E_{24} - E_{42}, & \tau(A_{21}) &= E_{21} - E_{12}, & \tau(A_{28}) &= E_{16} - E_{61},
 \end{aligned}
 \tag{9.8}$$




where E_{ij} , $i, j = 1, 2, \dots, 8$ denote 8×8 matrices with zeroes everywhere except the value 1 in the entry (i, j) seating at the crossing of the i th row and the j th column.

The three real, irreducible, pairwise nonequivalent representations (ρ_+, ρ_-, τ) of $\mathfrak{so}(8,0)$ given by the formulas (9.7) and (9.8) constitute the set of the *triality representations* for $\mathfrak{so}(8,0)$.


10. Application: 2-step graded realisations of real forms of the exceptional Lie algebra \mathfrak{e}_6


The simple exceptional complex Lie algebra \mathfrak{e}_6 has the following *noncompact* real forms:

(1) \mathfrak{e}_I , with Satake diagram 

- (2) \mathfrak{e}_{II} , with Satake diagram ,
- (3) \mathfrak{e}_{III} , with Satake diagram , and
- (4) \mathfrak{e}_{IV} , with Satake diagram .



Elie Cartan in his theses [8, 9] mentioned realisation of the real form \mathfrak{e}_I in $N = \mathbb{R}^{16}$. In the modern language, Cartan’s realisation is such that \mathfrak{e}_I is the algebra of automorphisms of the flat model of a parabolic geometry of type (E_I, P) , where the choice of parabolic subgroup in the real form E_I of the exceptional Lie group \mathbf{E}_6 is indicated by the

following decoration of the Satake diagram for \mathfrak{e}_I : . The structure on the 16-dimensional manifold $N = E_I/P$, whose symmetry is E_I , is a Majorana-Weyl $\mathbb{R}\mathbf{Spin}(5,5)$ structure (i.e., the reduction of the structure group of the tangent bundle TN to the $\mathbb{R}\mathbf{Spin}(5,5) \subset \mathbf{GL}(16, \mathbb{R})$ in the irreducible 16-dimensional representation of Majorana-Weyl spinors [19]). This geometry, as 1-step graded, is quite different from 2-step graded geometries considered in our paper. We also mention that if we wanted to have a realisation of, say \mathfrak{e}_{II} or \mathfrak{e}_{III} , in the spirit of Cartan’s realisation of \mathfrak{e}_I (i.e., if we crossed one lateral node in the Satake diagram of \mathfrak{e}_{II} or \mathfrak{e}_{III}), we would be forced to cross the complex

conjugated lateral root, resulting in the Satake diagrams , which would give realisations of the respective \mathfrak{e}_{II} and \mathfrak{e}_{III} in dimension twenty four. This we did in [14], providing realisations of \mathfrak{e}_{II} and \mathfrak{e}_{III} as Lie algebras of CR-automorphisms of certain 24-dimensional CR manifolds of CR dimension 16, and CR (real) codimension 8. The important point of these realisations of these two real forms of \mathfrak{e}_6 was that these geometries were 2-step graded, as in the case of Cartan’s realisation of \mathfrak{f}_I , and they could have been also thought as realisations in terms of the symmetry algebras of the structure (M, \mathcal{D}) , where M is a certain 24-dimensional real manifold and \mathcal{D} is a real rank 16-distribution on M with $[\mathcal{D}, \mathcal{D}] = TM$. Thus, these two geometries described by us in [14] are 2-step graded geometries of distributions – very much like Cartan’s realisation of \mathfrak{f}_I .

In this section, we give the remaining similar realisations of the yet untreated cases of \mathfrak{e}_{II} and \mathfrak{e}_{III} .

10.1. Realisations of \mathfrak{e}_I and \mathfrak{e}_{IV} : generalities

To get realisations of \mathfrak{e}_I and \mathfrak{e}_{IV} in dimension 24, we decorate the Satake diagrams of these two Lie algebras as follows:  and . These choices of a parabolic subalgebra in the respective \mathfrak{e}_I and \mathfrak{e}_{IV} produces the following gradation in these algebras:

$$\mathfrak{e}_A = \mathfrak{n}_{-2A} \oplus \mathfrak{n}_{-1A} \oplus \mathfrak{n}_{0A} \oplus \mathfrak{n}_{1A} \oplus \mathfrak{n}_{2A} \quad \text{for } A = I, IV,$$

with

$$\mathfrak{n}_{-A} = \mathfrak{n}_{-2A} \oplus \mathfrak{n}_{-1A} \quad \text{for } A = I, IV,$$

being 2-step nilpotent and having grading components \mathfrak{n}_{-2A} and \mathfrak{n}_{-1A} of respective dimension $r_A = 8$ and $s_A = 16$,

$$r_A = \dim(\mathfrak{n}_{-2A}) = 8, \quad s_A = \dim(\mathfrak{n}_{-1A}) = 16 \quad \text{for } A = I, IV.$$

The Lie algebra \mathfrak{n}_{0A} in the Tanaka prolongation of \mathfrak{n}_{-A} up to 0th order is

- (1) $\mathfrak{n}_{0I} = 2\mathbb{R} \oplus \mathfrak{so}(4,4) = \mathbb{R} \oplus \mathfrak{co}(4,4)$ in the case of \mathfrak{e}_I , and
- (2) $\mathfrak{n}_{0IV} = 2\mathbb{R} \oplus \mathfrak{so}(8,0) = \mathbb{R} \oplus \mathfrak{co}(8,0)$ in the case of \mathfrak{e}_{IV} .

The last two statements, (1) and (2), get clear when one looks at the Satake diagrams we

have just decorated. If we strip off the crossed nodes from these diagrams, we get



and clearly the simple part of \mathfrak{n}_{0AS} above.

Because of the grading property $[\mathfrak{n}_{iA}, \mathfrak{n}_{jA}] \subset \mathfrak{n}_{(i+j)A}$ in the Lie algebras \mathfrak{e}_A , restricting to subalgebras \mathfrak{n}_{-A} , we see that we have representations $(\rho_A, \mathfrak{n}_{-1A})$ and $(\tau_A, \mathfrak{n}_{-2A})$ given by the adjoint action of $\mathfrak{co}(4,4)$ or $\mathfrak{co}(8,0)$ which naturally seat in \mathfrak{n}_{0A} , respectively.

There is no surprise that the representations $(\rho_A, \mathfrak{n}_{-1A})$ are the Dirac spinor representations (9.3) and (9.6) of the respective $\mathfrak{co}(4,4)$ and $\mathfrak{co}(8,0)$ parts of \mathfrak{n}_{0AS} in the 16-dimensional real vector spaces \mathfrak{n}_{-1A} . As such, these representations are *reducible*, and they split each \mathfrak{n}_{0A} , $A = I, IV$ onto two irreducible representations $(\rho_{A\pm}, \mathfrak{n}_{-1A\pm})$ in real 8-dimensional spaces $\mathfrak{n}_{-1A\pm}$ of Weyl spinors. This shows that the 2-step nilpotent Lie algebra \mathfrak{n}_{-A} is, for each $A = I, IV$, a natural representation space for the action of the three triality representations (ρ_+, ρ_-, τ) . We have

$$\begin{aligned} \mathfrak{n}_{-A} &= \mathfrak{n}_{-2A} \oplus \mathfrak{n}_{-1A} = \\ &= \mathfrak{n}_{-2A} \oplus \mathfrak{n}_{-1A+} \oplus \mathfrak{n}_{-1A-}, \end{aligned}$$

and the 8-dimensional real irreducible representations $(\tau_A, \rho_{A+}, \rho_{A-})$ of $\mathfrak{co}(4,4)$ or $\mathfrak{co}(8,0)$ acting in the respective \mathfrak{n}_{-2A} , \mathfrak{n}_{-1A+} and \mathfrak{n}_{-1A-} .

We summarise the considerations from this section in the following theorem,

Theorem 10.1. (Natural realisation of the triality representations)

- (1) *The $\mathfrak{so}(4,4)$ triality:* The real form \mathfrak{e}_I of the simple exceptional Lie algebra \mathfrak{e}_6 , when graded according to the following decoration of its Satake diagram has the \mathfrak{n}_- part as a real 24-dimensional vector space, naturally split onto the three real 8-dimensional components \mathfrak{n}_{-2} , \mathfrak{n}_{-1+} and \mathfrak{n}_{-1-} ,


$$\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1+} \oplus \mathfrak{n}_{-1-}.$$

This decomposition is $\mathfrak{so}(4,4)$ invariant and consists of components \mathfrak{n}_{-2} , \mathfrak{n}_{-1+} and \mathfrak{n}_{-1-} , on which the triality representation

$$\tau \oplus \rho_+ \oplus \rho_-$$

of $\mathfrak{so}(4,4)$ acts irreducibly.

- (2) The $\mathfrak{so}(8,0)$ triality: Likewise, the real form \mathfrak{e}_{IV} of the simple exceptional Lie algebra \mathfrak{e}_6 , when graded according to the following decoration of its Satake diagram



has the \mathfrak{n}_- part as a real 24-dimensional vector space, naturally split onto the three real 8-dimensional components \mathfrak{n}_{-2} , \mathfrak{n}_{-1+} and \mathfrak{n}_{-1-} ,

$$\mathfrak{n}_- = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1+} \oplus \mathfrak{n}_{-1-}.$$

This decomposition is $\mathfrak{so}(8,0)$ invariant and consists of components \mathfrak{n}_{-2} , \mathfrak{n}_{-1+} and \mathfrak{n}_{-1-} , on which the triality representation

$$\tau \oplus \rho_+ \oplus \rho_-$$

of $\mathfrak{so}(8,0)$ acts irreducibly.

10.2. An explicit realisation of \mathfrak{e}_I in dimension 24

Taking as (ρ, S) the Dirac spinors representation (9.3) of $\mathfrak{co}(4,4)$ in dimension 16, and as (τ, R) the vectorial representation (9.5) of $\mathfrak{co}(4,4)$ in dimension 8, we again are in the situation of a missing $\omega \in \text{Hom}(\bigwedge^2 S, R)$ from the triple (ρ, τ, ω) described by the magical equation (2.1). Solving this equation for ω , we obtain $\omega^i_{\mu\nu}$, $i = 1, \dots, 8$, $\mu, \nu = 1, \dots, 16$, which leads to the eight 2-forms $\omega^i = \frac{1}{2} \omega^i_{\mu\nu} dx^\mu \wedge dx^\nu$ on a 16-dimensional manifold $N = \mathbb{R}^{16}$, which read

$$\begin{aligned} \omega^1 &= -dx^1 \wedge dx^{10} + dx^2 \wedge dx^9 + dx^7 \wedge dx^{16} - dx^8 \wedge dx^{15} \\ \omega^2 &= -dx^2 \wedge dx^{12} + dx^4 \wedge dx^{10} + dx^6 \wedge dx^{16} - dx^8 \wedge dx^{14} \\ \omega^3 &= -dx^1 \wedge dx^{12} + dx^4 \wedge dx^9 + dx^5 \wedge dx^{16} - dx^8 \wedge dx^{13} \\ \omega^4 &= -dx^5 \wedge dx^{10} + dx^6 \wedge dx^9 + dx^7 \wedge dx^{12} - dx^8 \wedge dx^{11} \\ \omega^5 &= -dx^2 \wedge dx^{11} + dx^3 \wedge dx^{10} + dx^6 \wedge dx^{15} - dx^7 \wedge dx^{14} \\ \omega^6 &= -dx^1 \wedge dx^{11} + dx^3 \wedge dx^9 + dx^5 \wedge dx^{15} - dx^7 \wedge dx^{13} \\ \omega^7 &= -dx^3 \wedge dx^{12} + dx^4 \wedge dx^{11} + dx^5 \wedge dx^{14} - dx^6 \wedge dx^{13} \\ \omega^8 &= -dx^1 \wedge dx^{14} + dx^2 \wedge dx^{13} + dx^3 \wedge dx^{16} - dx^4 \wedge dx^{15}. \end{aligned} \tag{10.1}$$

The manifold $N = \mathbb{R}^{16}$ with these 2-forms, after contactification, gives the following Theorem.

Theorem 10.2. *Let $M = \mathbb{R}^{24}$ with coordinates $(u^1, \dots, u^8, x^1, \dots, x^{16})$, and consider eight 1-forms $\lambda^1, \dots, \lambda^8$ on M given by*

$$\begin{aligned} \lambda^1 &= du^1 - x^1 dx^{10} + x^2 dx^9 + x^7 dx^{16} - x^8 dx^{15} \\ \lambda^2 &= du^2 - x^2 dx^{12} + x^4 dx^{10} + x^6 dx^{16} - x^8 dx^{14} \\ \lambda^3 &= du^3 - x^1 dx^{12} + x^4 dx^9 + x^5 dx^{16} - x^8 dx^{13} \\ \lambda^4 &= du^4 - x^5 dx^{10} + x^6 dx^9 + x^7 dx^{12} - x^8 dx^{11} \\ \lambda^5 &= du^5 - x^2 dx^{11} + x^3 dx^{10} + x^6 dx^{15} - x^7 dx^{14} \\ \lambda^6 &= du^6 - x^1 dx^{11} + x^3 dx^9 + x^5 dx^{15} - x^7 dx^{13} \\ \lambda^7 &= du^7 - x^3 dx^{12} + x^4 dx^{11} + x^5 dx^{14} - x^6 dx^{13} \\ \lambda^8 &= du^8 - x^1 dx^{14} + x^2 dx^{13} + x^3 dx^{16} - x^4 dx^{15}. \end{aligned}$$

The rank 16 distribution \mathcal{D} on M defined as $\mathcal{D} = \{\text{TR}^{24} \ni X \mid X \lrcorner \lambda^1 = \dots = X \lrcorner \lambda^8 = 0\}$ has its Lie algebra of infinitesimal automorphisms $\text{aut}(\mathcal{D})$ isomorphic to the Tanaka prolongation of $\mathfrak{n} = R \oplus S$, where $(\rho, S = \mathbb{R}^{16})$ is the Dirac spinors representation (9.3) of $\mathfrak{n}_{00} = \mathfrak{co}(4,4)$, and $(\tau, R = \mathbb{R}^8)$ is the vectorial representation (9.5) of \mathfrak{n}_{00} .

The symmetry algebra $\text{aut}(\mathcal{D})$ is isomorphic to the simple exceptional Lie algebra \mathfrak{e}_I ,

$$\text{aut}(\mathcal{D}) = \mathfrak{e}_I,$$

having the following natural gradation

$$\text{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S = S_+ \oplus S_-$,

$$\mathfrak{n}_0 = \mathbb{R} \oplus \mathfrak{co}(4,4) \supset \mathfrak{n}_{00},$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$, and with the spaces S_{\pm} being the carrier spaces for the Weyl spinors representations ρ_{\pm} of $\mathfrak{co}(4,4)$. The gradation in \mathfrak{e}_I is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in \mathfrak{e}_I .

The contactification (M, \mathcal{D}) is locally the flat model for the parabolic geometry of type

(E_I, P_I) related to the following crossed Satake diagram:

Remark 10.3. Also the \mathfrak{e}_I case, considered in this section, admits a description in terms of an $\mathbb{R} \oplus \mathfrak{so}(4,4)$ invariant 4-form Φ in $S = \mathbb{R}^{16}$. Now Φ may be represented by

$$\Phi = h_{ij} \omega^i \wedge \omega^j,$$

where ω^i are given by (10.1) and

$$(h_{ij}) = \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Explicitly, the form Φ reads

$$\begin{aligned} \Phi = & 2dx^1 \wedge dx^2 \wedge dx^{11} \wedge dx^{12} - 2dx^1 \wedge dx^3 \wedge dx^{10} \wedge dx^{12} + \\ & 2dx^1 \wedge dx^4 \wedge dx^{10} \wedge dx^{11} + 2dx^1 \wedge dx^5 \wedge dx^{10} \wedge dx^{14} - \\ & dx^1 \wedge dx^6 \wedge dx^9 \wedge dx^{14} - dx^1 \wedge dx^6 \wedge dx^{10} \wedge dx^{13} - \\ & dx^1 \wedge dx^6 \wedge dx^{11} \wedge dx^{16} + dx^1 \wedge dx^6 \wedge dx^{12} \wedge dx^{15} - \\ & 2dx^1 \wedge dx^7 \wedge dx^{12} \wedge dx^{14} + 2dx^1 \wedge dx^8 \wedge dx^{11} \wedge dx^{14} + \\ & 2dx^2 \wedge dx^3 \wedge dx^9 \wedge dx^{12} - 2dx^2 \wedge dx^4 \wedge dx^9 \wedge dx^{11} - \\ & dx^2 \wedge dx^5 \wedge dx^9 \wedge dx^{14} - dx^2 \wedge dx^5 \wedge dx^{10} \wedge dx^{13} + \\ & dx^2 \wedge dx^5 \wedge dx^{11} \wedge dx^{16} - dx^2 \wedge dx^5 \wedge dx^{12} \wedge dx^{15} + \\ & 2dx^2 \wedge dx^6 \wedge dx^9 \wedge dx^{13} + 2dx^2 \wedge dx^7 \wedge dx^{12} \wedge dx^{13} - \\ & 2dx^2 \wedge dx^8 \wedge dx^{11} \wedge dx^{13} + 2dx^3 \wedge dx^4 \wedge dx^9 \wedge dx^{10} - \\ & 2dx^3 \wedge dx^5 \wedge dx^{10} \wedge dx^{16} + 2dx^3 \wedge dx^6 \wedge dx^9 \wedge dx^{16} + \\ & 2dx^3 \wedge dx^7 \wedge dx^{12} \wedge dx^{16} - dx^3 \wedge dx^8 \wedge dx^9 \wedge dx^{14} + \\ & dx^3 \wedge dx^8 \wedge dx^{10} \wedge dx^{13} - dx^3 \wedge dx^8 \wedge dx^{11} \wedge dx^{16} - \\ & dx^3 \wedge dx^8 \wedge dx^{12} \wedge dx^{15} + 2dx^4 \wedge dx^5 \wedge dx^{10} \wedge dx^{15} - \\ & 2dx^4 \wedge dx^6 \wedge dx^9 \wedge dx^{15} + dx^4 \wedge dx^7 \wedge dx^9 \wedge dx^{14} - \\ & dx^4 \wedge dx^7 \wedge dx^{10} \wedge dx^{13} - dx^4 \wedge dx^7 \wedge dx^{11} \wedge dx^{16} - \\ & dx^4 \wedge dx^7 \wedge dx^{12} \wedge dx^{15} + 2dx^4 \wedge dx^8 \wedge dx^{11} \wedge dx^{15} + \\ & 2dx^5 \wedge dx^6 \wedge dx^{15} \wedge dx^{16} - 2dx^5 \wedge dx^7 \wedge dx^{14} \wedge dx^{16} + \\ & 2dx^5 \wedge dx^8 \wedge dx^{14} \wedge dx^{15} + 2dx^6 \wedge dx^7 \wedge dx^{13} \wedge dx^{16} - \\ & 2dx^6 \wedge dx^8 \wedge dx^{13} \wedge dx^{15} + 2dx^7 \wedge dx^8 \wedge dx^{13} \wedge dx^{14}. \end{aligned} \tag{10.2}$$

This 4-form is such that its stabiliser in $\mathfrak{gl}(16, \mathbb{R})$ is $\mathfrak{n}_0 = \mathbb{R} \oplus \mathfrak{co}(4, 4)$. When restricted to $\mathfrak{n}_{00} = \mathfrak{co}(4, 4)$ this stabiliser is given precisely in the Majorana Dirac spinor representation

$$\rho = \rho_+ \oplus \rho_-$$

as in (9.3)–(9.4).

10.3. An explicit realisation of ϵ_{IV} in dimension 24

Similarly, as in the previous section, we take as (ρ, S) the Dirac spinors representation (9.6) of $\mathfrak{co}(8,0)$ in dimension 16, and as (τ, R) the vectorial representation (9.8) of $\mathfrak{co}(8,0)$ in dimension 8, and we search for $\omega \in \text{Hom}(\wedge^2 S, R)$, solving the magical equation (2.1). We obtain $\omega^i_{\mu\nu}$, $i = 1, \dots, 8$, $\mu, \nu = 1, \dots, 16$, which provides us with the eight 2-forms $\omega^i = \frac{1}{2}\omega^i_{\mu\nu}dx^\mu \wedge dx^\nu$ on a 16-dimensional manifold $N = \mathbb{R}^{16}$, which read

$$\begin{aligned}
 \omega^1 &= dx^1 \wedge dx^9 + dx^2 \wedge dx^{10} + dx^3 \wedge dx^{11} + dx^4 \wedge dx^{12} - \\
 &\quad dx^5 \wedge dx^{13} - dx^6 \wedge dx^{14} - dx^7 \wedge dx^{15} - dx^8 \wedge dx^{16} \\
 \omega^2 &= -dx^1 \wedge dx^{10} + dx^2 \wedge dx^9 + dx^3 \wedge dx^{12} - dx^4 \wedge dx^{11} - \\
 &\quad dx^5 \wedge dx^{14} + dx^6 \wedge dx^{13} + dx^7 \wedge dx^{16} - dx^8 \wedge dx^{15} \\
 \omega^3 &= -dx^1 \wedge dx^{11} - dx^2 \wedge dx^{12} + dx^3 \wedge dx^9 + dx^4 \wedge dx^{10} + \\
 &\quad dx^5 \wedge dx^{15} + dx^6 \wedge dx^{16} - dx^7 \wedge dx^{13} - dx^8 \wedge dx^{14} \\
 \omega^4 &= -dx^1 \wedge dx^{12} + dx^2 \wedge dx^{11} - dx^3 \wedge dx^{10} + dx^4 \wedge dx^9 + \\
 &\quad dx^5 \wedge dx^{16} - dx^6 \wedge dx^{15} + dx^7 \wedge dx^{14} - dx^8 \wedge dx^{13} \\
 \omega^5 &= dx^1 \wedge dx^{13} + dx^2 \wedge dx^{14} - dx^3 \wedge dx^{15} - dx^4 \wedge dx^{16} + \\
 &\quad dx^5 \wedge dx^9 + dx^6 \wedge dx^{10} - dx^7 \wedge dx^{11} - dx^8 \wedge dx^{12} \\
 \omega^6 &= dx^1 \wedge dx^{14} - dx^2 \wedge dx^{13} - dx^3 \wedge dx^{16} + dx^4 \wedge dx^{15} - \\
 &\quad dx^5 \wedge dx^{10} + dx^6 \wedge dx^9 + dx^7 \wedge dx^{12} - dx^8 \wedge dx^{11} \\
 \omega^7 &= dx^1 \wedge dx^{15} - dx^2 \wedge dx^{16} + dx^3 \wedge dx^{13} - dx^4 \wedge dx^{14} + \\
 &\quad dx^5 \wedge dx^{11} - dx^6 \wedge dx^{12} + dx^7 \wedge dx^9 - dx^8 \wedge dx^{10} \\
 \omega^8 &= -dx^1 \wedge dx^{16} - dx^2 \wedge dx^{15} - dx^3 \wedge dx^{14} - dx^4 \wedge dx^{13} - \\
 &\quad dx^5 \wedge dx^{12} - dx^6 \wedge dx^{11} - dx^7 \wedge dx^{10} - dx^8 \wedge dx^9.
 \end{aligned}
 \tag{10.3}$$

Contactifying, we have the following theorem:

Theorem 10.4. *Let $M = \mathbb{R}^{24}$ with coordinates $(u^1, \dots, u^8, x^1, \dots, x^{16})$, and consider eight 1-forms $\lambda^1, \dots, \lambda^8$ on M given by*

$$\begin{aligned}
 \lambda^1 &= du^1 + x^1 dx^9 + x^2 dx^{10} + x^3 dx^{11} + x^4 dx^{12} - x^5 dx^{13} - x^6 dx^{14} - x^7 dx^{15} - x^8 dx^{16} \\
 \lambda^2 &= du^2 - x^1 dx^{10} + x^2 dx^9 + x^3 dx^{12} - x^4 dx^{11} - x^5 dx^{14} + x^6 dx^{13} + x^7 dx^{16} - x^8 dx^{15} \\
 \lambda^3 &= du^3 - x^1 dx^{11} - x^2 dx^{12} + x^3 dx^9 + x^4 dx^{10} + x^5 dx^{15} + x^6 dx^{16} - x^7 dx^{13} - x^8 dx^{14} \\
 \lambda^4 &= du^4 - x^1 dx^{12} + x^2 dx^{11} - x^3 dx^{10} + x^4 dx^9 + x^5 dx^{16} - x^6 dx^{15} + x^7 dx^{14} - x^8 dx^{13} \\
 \lambda^5 &= du^5 + x^1 dx^{13} + x^2 dx^{14} - x^3 dx^{15} - x^4 dx^{16} + x^5 dx^9 + x^6 dx^{10} - x^7 dx^{11} - x^8 dx^{12} \\
 \lambda^6 &= du^6 + x^1 dx^{14} - x^2 dx^{13} - x^3 dx^{16} + x^4 dx^{15} - x^5 dx^{10} + x^6 dx^9 + x^7 dx^{12} - x^8 dx^{11} \\
 \lambda^7 &= du^7 + x^1 dx^{15} - x^2 dx^{16} + x^3 dx^{13} - x^4 dx^{14} + x^5 dx^{11} - x^6 dx^{12} + x^7 dx^9 - x^8 dx^{10} \\
 \lambda^8 &= du^8 - x^1 dx^{16} - x^2 dx^{15} - x^3 dx^{14} - x^4 dx^{13} - x^5 dx^{12} - x^6 dx^{11} - x^7 dx^{10} - x^8 dx^9.
 \end{aligned}$$

The rank 16 distribution \mathcal{D} on M defined as $\mathcal{D} = \{\text{TRR}^{24} \ni X \mid X \lrcorner \lambda^1 = \dots = X \lrcorner \lambda^8 = 0\}$ has its Lie algebra of infinitesimal automorphisms $\text{aut}(\mathcal{D})$ isomorphic to the Tanaka

prolongation of $\mathfrak{n}_- = R \oplus S$, where $(\rho, S = \mathbb{R}^{16})$ is the Dirac spinors representation (9.3) of $\mathfrak{n}_{00} = \mathfrak{co}(8,0)$, and $(\tau, R = \mathbb{R}^8)$ is the vectorial representation (9.5) of \mathfrak{n}_{00} .

The symmetry algebra $\mathfrak{aut}(\mathcal{D})$ is isomorphic to the simple exceptional Lie algebra \mathfrak{e}_{IV} ,

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{e}_{IV},$$


having the following natural gradation:

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S = S_+ \oplus S_-$,

$$\mathfrak{n}_0 = \mathbb{R} \oplus \mathfrak{co}(8,0) \supset \mathfrak{n}_{00},$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$, and with the spaces S_{\pm} being the Carrier spaces for the Weyl spinors representations ρ_{\pm} of $\mathfrak{co}(8,0)$. The gradation in \mathfrak{e}_{IV} is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in \mathfrak{e}_{IV} .

The contactification (M, \mathcal{D}) is locally the flat model for the parabolic geometry of type (E_{IV}, P_{IV}) related to the following crossed Satake diagram: 

Remark 10.5. Again, we have a description of the relevant representations in terms of an $\mathbb{R} \oplus \mathfrak{co}(8,0)$ invariant 4-form Φ in $S = \mathbb{R}^{16}$. Now Φ may be represented by

$$\Phi = h_{ij} \omega^i \wedge \omega^j,$$

where ω^i are given by (10.3) and


$$(h_{ij}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$


This 4-form is such that its stabiliser in $\mathfrak{gl}(16, \mathbb{R})$ is $\mathfrak{n}_0 = \mathbb{R} \oplus \mathfrak{co}(8,0)$. When restricted to $\mathfrak{n}_{00} = \mathfrak{co}(8,0)$, this stabiliser is given precisely in the Majorana Dirac spinor representation

$$\rho = \rho_+ \oplus \rho_-$$


as in (9.6)–(9.7).

11. Application: one more realisation of \mathfrak{e}_6 and a realisation of \mathfrak{b}_6

Between the 24-dimensional realisations of \mathfrak{e}_6 mentioned in this paper, and Cartan’s 16-dimensional realisation of \mathfrak{e}_6 associated with the grading , there are 21-dimensional realisations of this algebra \mathfrak{e}_6 associated with the following Dynkin diagram

crossing: . These define contact \mathfrak{e}_6 geometries and are described in [11] p. 425–426.

11.1. Realisation of \mathfrak{e}_I in dimension 25

Here, we will briefly discuss yet another realisation, now in dimension 25, corresponding to the following Dynkin diagram crossing:  of \mathfrak{e}_6 . This is, for example, mentioned in [15]. Looking at the Satake diagrams of real forms of \mathfrak{e}_6 , we see that this realisation is only possible for the real form \mathfrak{e}_I .

So we again use our Corollary 4.1 with now $\mathfrak{n}_{00} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(5, \mathbb{R})$ and with representations (ρ, S) and (τ, R) , as indicated in [15] Section 5.3, $S = \mathbb{R}^2 \otimes \bigwedge^2 \mathbb{R}^5$, $R = \bigwedge^2 \mathbb{R}^2 \otimes \bigwedge^4 \mathbb{R}^5$.

To be more explicit, we obtain this representations as follows:

- We start with the defining representations τ_2 of $\mathfrak{sl}(2, \mathbb{R})$ in \mathbb{R}^2 and τ_5 of $\mathfrak{sl}(5, \mathbb{R})$ in \mathbb{R}^5 , and we define the representation

$$\rho = \tau_2 \otimes (\tau_5 \wedge \tau_5) \quad \text{of} \quad \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(5, \mathbb{R}) \quad \text{in} \quad S = \mathbb{R}^2 \otimes \bigwedge^2 \mathbb{R}^5 = \mathbb{R}^{20}.$$

The representation (ρ, S) is an irreducible real 20-dimensional representation of

$$\mathfrak{n}_{00} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(5, \mathbb{R}).$$

- Then we decompose the 190-dimensional representation $\rho \wedge \rho$ onto the irreducibles:

$$\rho \wedge \rho = \alpha \oplus \tau \oplus \beta \quad \text{in} \quad \bigwedge_{50} \oplus R \oplus \bigwedge_{135},$$

with (α, \bigwedge_{50}) being 50-dimensional, (τ, R) being 5-dimensional and (β, \bigwedge_{135}) being 135-dimensional.

- We take the 20-dimensional representation (ρ, S) and the 5-dimensional representation (τ, R) of $\mathfrak{n}_{00} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(5, \mathbb{R})$ as above, and we apply our Corollary 4.1.

We obtain the following theorem.

Theorem 11.1. *Let $M = \mathbb{R}^{25}$ with coordinates $(u^1, \dots, u^5, x^1, \dots, x^{20})$, and consider five 1-forms $\lambda^1, \dots, \lambda^5$ on M given by*

$$\begin{aligned} \lambda^1 &= du^1 - x^3 dx^{20} + x^5 dx^{19} - x^6 dx^{18} - x^8 dx^{16} + x^9 dx^{15} - x^{10} dx^{13} \\ \lambda^2 &= du^2 - x^2 dx^{20} + x^4 dx^{19} - x^6 dx^{17} - x^7 dx^{16} + x^9 dx^{14} - x^{10} dx^{12} \\ \lambda^3 &= du^3 - x^1 dx^{20} + x^4 dx^{18} - x^5 dx^{17} - x^7 dx^{15} + x^8 dx^{14} - x^{10} dx^{11} \\ \lambda^4 &= du^4 - x^1 dx^{19} + x^2 dx^{18} - x^3 dx^{17} - x^7 dx^{13} + x^8 dx^{12} - x^9 dx^{11} \\ \lambda^5 &= du^5 - x^1 dx^{16} + x^2 dx^{15} - x^3 dx^{14} - x^4 dx^{13} + x^5 dx^{12} - x^6 dx^{11}. \end{aligned}$$

The rank 20 distribution \mathcal{D} on M defined as $\mathcal{D} = \{\text{TR}^{25} \ni X \mid X \lrcorner \lambda^1 = \dots = X \lrcorner \lambda^5 = 0\}$ has its Lie algebra of infinitesimal automorphisms $\text{aut}(\mathcal{D})$ isomorphic to the Tanaka prolongation of $\mathfrak{n}_- = R \oplus S$, where $(\rho, S = \mathbb{R}^{20})$ is the 20-dimensional irreducible representation of $\mathfrak{n}_{00} = \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(5, \mathbb{R})$, and $(\tau, R = \mathbb{R}^5)$ is the 5-dimensional irreducible subrepresentation $\tau \in (\rho \wedge \rho)$ of \mathfrak{n}_{00} .

The symmetry algebra $\mathfrak{aut}(\mathcal{D})$ is isomorphic to the simple exceptional Lie algebra \mathfrak{e}_I ,

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{e}_I,$$

having the following natural gradation:

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S$,

$$\mathfrak{n}_0 = \mathbb{R} \oplus \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{sl}(5, \mathbb{R}) \supset \mathfrak{n}_{00},$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$. The gradation in \mathfrak{e}_I is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in \mathfrak{e}_I .

The contactification (M, \mathcal{D}) is locally the flat model for the parabolic geometry of type (E_I, P_{I*}) related to the following crossed Satake diagram:

11.2. A realisation of $\mathfrak{so}(7,6)$ in dimension 21

We know from [11] that the crossed Satake diagram corresponds to the \mathfrak{e}_I -symmetric contact geometry in dimension 21. It corresponds to the grading

$$\mathfrak{e}_I = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\dim(\mathfrak{n}_{\pm 1}) = 20$, $\dim(\mathfrak{n}_{\pm 2}) = 1$ and $\mathfrak{n}_0 = \mathfrak{gl}(6, \mathbb{R})$.

Interestingly, dimension $n = 78$ is the dimension not only of the exceptional simple Lie algebra \mathfrak{e}_6 , but also for the simple Lie algebras \mathfrak{b}_6 and \mathfrak{c}_6 . For example, if we take the crossed Satake diagram

$$\mathfrak{so}(7,6) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\dim(\mathfrak{n}_{\pm 1}) = 6$, $\dim(\mathfrak{n}_{\pm 2}) = 15$ and $\mathfrak{n}_0 = \mathfrak{gl}(6, \mathbb{R})$, in the simple Lie algebra $\mathfrak{so}(7,6)$. Here, taking (ρ, S) as the defining representation $\rho(A) = A$ of $\mathbf{GL}(6, \mathbb{R})$ in $S = \mathbb{R}^6$, taking the representation (τ, R) to be $\tau = \rho \wedge \rho$ in $R = \bigwedge^2 \mathbb{R}^6 = \mathbb{R}^{15}$, and applying our Corollary 4.1, we get the following theorem.⁷

Theorem 11.2. Let $M = \mathbb{R}^{21}$ with coordinates $(u^1, \dots, u^{15}, x^1, \dots, x^6)$, and consider fifteen 1-forms $\lambda^1, \dots, \lambda^5$ on M given by

$$\lambda^{I(i,j)} = du^{I(i,j)} - x^i dx^j,$$

with

$$I(i,j) = 1 + i + \frac{1}{2}(j-3)j, \quad 1 \leq i < j \leq 6.$$

The rank 6 distribution \mathcal{D} on M defined as $\mathcal{D} = \{\mathbb{T}\mathbb{R}^{21} \ni X \mid X \lrcorner \lambda^1 = \dots = X \lrcorner \lambda^{15} = 0\}$ has its Lie algebra of infinitesimal automorphisms $\mathfrak{aut}(\mathcal{D})$ isomorphic to the Tanaka prolongation of $\mathfrak{n}_- = R \oplus S$, where $(\rho, S = \mathbb{R}^6)$ is the 6-dimensional defining representation

⁷We invoke it, just to show that we do not only use spin representations in this paper.

of $\mathfrak{n}_{00} = \mathfrak{gl}(6, \mathbb{R})$, and $(\tau, R = \wedge^2 \mathbb{R}^6)$ is the 15-dimensional irreducible subrepresentation $\tau = \rho \wedge \rho$ of \mathfrak{n}_{00} .

The symmetry algebra $\mathfrak{aut}(\mathcal{D})$ is isomorphic to the simple exceptional Lie algebra $\mathfrak{so}(7, 6)$,

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{so}(7, 6),$$

having the following natural gradation:

$$\mathfrak{aut}(\mathcal{D}) = \mathfrak{n}_{-2} \oplus \mathfrak{n}_{-1} \oplus \mathfrak{n}_0 \oplus \mathfrak{n}_1 \oplus \mathfrak{n}_2,$$

with $\mathfrak{n}_{-2} = R$, $\mathfrak{n}_{-1} = S$,

$$\mathfrak{n}_0 = \mathfrak{gl}(6, \mathbb{R}) = \mathfrak{n}_{00},$$

$\mathfrak{n}_1 = S^*$, $\mathfrak{n}_2 = R^*$. The gradation in $\mathfrak{so}(7, 6)$ is inherited from the distribution structure (M, \mathcal{D}) . The duality signs $*$ at R^* and S^* above are with respect to the Killing form in $\mathfrak{so}(7, 6)$.

The contactification (M, \mathcal{D}) is locally the flat model for the parabolic geometry of type $(\mathfrak{so}(7, 6), P)$ related to the following crossed Satake diagram: $\circ - \circ - \circ - \circ - \circ \rightleftharpoons \circ$.

Competing interests. The authors have no competing interest to declare.

References

- [1] I. ANDERSON, Z. NIE AND P. NUROWSKI (2015) Non-rigid parabolic geometries of Monge type. *Adv. Math.* **277**, 24–55. DOI: <https://doi.org/10.1016/j.aim.2015.01.021>.
- [2] A. MICHAEL AND J. BERNDT (2003) Projective planes, Severi varieties and spheres. In *Surveys in Differential Geometry*. Vol. VIII. 1–27.
- [3] DV. ALEKSEEVSKY AND V. CORTES (1997) Classification of N-(super)-extended Poincaré algebras and bilinear invariants of the spinor representation of $\text{Spin}(p, q)$. *Comm. Math. Phys.* **183**(3), 477–510.
- [4] A. ALTOMANI AND A. SANTI (2014) Tanaka structures modeled on extended Poincaré algebras. *Indiana Univ. Math. J.* **63**(1), 91–117.
- [5] O. BIQUARD, Quaternionic contact structures. In *Quaternionic Structures in Mathematics and Physics*. 23–30. DOI: [10.1142/9789812810038_0003](https://doi.org/10.1142/9789812810038_0003), available at https://www.worldscientific.com/doi/pdf/10.1142/9789812810038_0003.
- [6] O. BIQUARD (2015) From G2 geometry to quaternionic Kahler metrics. *J. Geom. Phys.* **91**, 101–107.
- [7] RB. BROWN AND A. GRAY (1972) Riemannian manifolds with holonomy group $\text{Spin}(9)$. In *Differential Geometry; in Honor of Kentaro Yano*. 41–59.
- [8] E. CARTAN (1893) Über die einfachen Transformationsgruppen. *Ber. Verh. k. Sachs. Ges. d. Wiss. Leipzig*, 395–420.
- [9] E. CARTAN (1894) Sur la structure des groupes de transformations finis et continus. *Oeuvres* **1**, 137–287.
- [10] E. CARTAN (1910) Les systèmes de Pfaff à cinq variables et les équations aux dérivées partielles du second ordre. *Ann. Sci. Éc. Norm. Supér.* **27**, 109–192.
- [11] A. ČAP AND J. SLOVÁK (2009) *Parabolic Geometries. I. Mathematical Surveys and Monographs*. Vol. 154. Providence, RI : American Mathematical Society. MR2532439

- [12] D. DUCHEMIN (2006) Quaternionic contact structures in dimension 7. *Ann. Inst. Fourier* **56**(4), 851–885 (eng).
- [13] S. HELGASON (1977) Invariant differential equations on homogeneous manifolds. *BAMS* **83**, 751–756.
- [14] CD. HILL, J. MERKER, Z. NIE AND P. NUROWSKI (2023) Accidental CR structures. arXiv preprint 2302.03119. Available at <https://arxiv.org/abs/2302.03119>.
- [15] W. KRAŚKIEWICZ AND J. WEYMAN (2012) Geometry of orbit closures for the representations associated to gradings of Lie algebras of types E6, F4 and G2. arXiv preprint 1201.1102. Available at <https://arxiv.org/abs/1201.1102>.
- [16] MG. MOLINA, B. KRUGLIKOV, I. MARKINA AND A. VASIL'EV (2018) Rigidity of 2-step Carnot groups. *J. Geom. Anal.* **28**, 1477–1501.
- [17] P. NUROWSKI (2005) Differential equations and conformal structures. *J. Geom. Phys.* **55**, 19–49. DOI: <https://doi.org/10.1016/j.geomphys.2004.11.006>.
- [18] N. TANAKA (1970) On differential systems, graded Lie algebras and pseudogroups. *Kyoto J. Math.* **10**, 1–82.
- [19] A. TRAUTMAN (2006) Clifford algebras and their representations. In GL. NABER, ST. TSOU AND JP. FRANÇOISE (eds), *Encyclopedia of Mathematical Physics*. Vol. 1. Oxford: Elsevier. <http://trautman.fuw.edu.pl>.