

Research Paper

Cite this article: Bruno A, Fedynich AM, Rollins D, Wester DB (2019). Helminth community and host dynamics in northern bobwhites from the Rolling Plains Ecoregion, U.S.A. *Journal of Helminthology* **93**, 567–573. <https://doi.org/10.1017/S0022149X18000494>

Received: 8 March 2018
Accepted: 23 May 2018
First published online: 29 June 2018

Key words:

Colinus virginianus; helminths; northern bobwhite; parasites; Rolling Plains Ecoregion; Texas

Author for correspondence:

A. Bruno, E-mail: andrea.bruno@students.tamuk.edu

Helminth community and host dynamics in northern bobwhites from the Rolling Plains Ecoregion, U.S.A.

A. Bruno¹, A.M. Fedynich¹, D. Rollins² and D.B. Wester¹

¹Caesar Kleberg Wildlife Research Institute, Texas A&M University-Kingsville, 700 University Boulevard, MSC 218, Kingsville, Texas 78363, USA and ²Rolling Plains Quail Research Foundation, P.O. Box 61517, San Angelo, Texas, 76906, USA

Abstract

One hundred and sixty-one northern bobwhites (*Colinus virginianus*; hereafter ‘bobwhite’) were examined from the Rolling Plains ecoregion of Texas and western Oklahoma from 2011 to 2013. Complete necropsies yielded 13 species, of which two are new host (*Gongylonema phasianella*) and region (*Eucoleus contortus*) records and three (*Dispharynx nasuta*, *Tetrameres pattersoni* and *Oxyspirura petrowi*) are known to cause morbidity and mortality. Of the species found, *Aulonocephalus pennula* commonly occurred, *Oxyspirura petrowi* was intermediate in prevalence, and the remaining species were rare. Species richness was similar compared to studies from the southeastern U.S., but higher than studies from the same region. In addition, 12 of the 13 species were heteroxenous helminths, supporting the theory that heteroxenous helminths in semi-arid regions are more successful than monoxenous helminths. Prevalence and abundance of *A. pennula* and *O. petrowi* were higher in adult bobwhites than in juveniles. Abundance of *A. pennula* and *O. petrowi* was higher at southern locations compared to northern locations in the study area. Our study is the first to provide a current assessment of the bobwhite helminth community across the Rolling Plains ecoregion of the U.S.

Introduction

The northern bobwhite (*Colinus virginianus*, hereafter bobwhite) is an upland game bird species with populations persisting in many ecologically and climatically diverse regions in the eastern half of North America. A slow, but apparent, range-wide decline in bobwhite populations has been occurring over the past 50 years (Hernández *et al.*, 2013; Sauer *et al.*, 2013; Texas Parks and Wildlife Department, 2013). Bobwhite populations fluctuate in a boom and bust pattern, where years with increased productivity are shown to be positively associated with breeding season (April–August) precipitation (Kiel, 1976; Tri *et al.*, 2013). When periods of above-average precipitation in 2010 preceded below-average bobwhite population estimates (Texas Parks and Wildlife Department, 2013), the relationship between helminths and bobwhite populations was revisited as a potential factor in the decline (Bruno, 2014; Dunham *et al.*, 2016; Villarreal *et al.*, 2016).

Much of the early research on bobwhite helminths in the Rolling Plains lacked analyses at the infra and component community levels, reporting results from incomplete necropsies focusing on particular helminth species (Webster & Addis, 1945; Jackson & Greene, 1965; Kocan *et al.*, 1979; Rollins, 1980; Boggs *et al.*, 1990), or based on studies with small geographical scale (two counties; Villarreal *et al.*, 2016), whereas research using a helminth community approach based on complete necropsies has been conducted in the Southern Coastal Plains in Florida and Georgia (Moore *et al.*, 1986; Moore & Simberloff, 1990; Davidson *et al.*, 1991). Because the Southern Coastal Plains differs ecologically and climatically (which influence helminth populations and communities) from the Rolling Plains ecoregion, host–helminth dynamics cannot be extrapolated between regions.

To learn more about diseases and potential disease agents within the Rolling Plains of Texas and Oklahoma, a large-scale bobwhite disease study, called Operation Idiopathic Decline (OID), was initiated in August 2011 in partnership with the Rolling Plains Quail Research Foundation, Texas Tech University, Texas A&M University-Kingsville, Texas A&M University, Texas A&M AgriLife Extension, and the Oklahoma Department of Wildlife and Conservation. We initiated a large-scale helminth study as part of the OID project. Our goals were to assess helminth prevalence, intensity, abundance and community structure (infra and component) in bobwhites and determine how infections are related to intrinsic (host age and host sex) and extrinsic variables (month, year of collection, and northing), thereby providing a current description of this host–helminth system at a regional scale.

Materials and methods

Study area

Bobwhite collections predominantly occurred in the Rolling Plains ecoregion of Texas and western Oklahoma (i.e. Central Great Plains; *fig. 1*). The Rolling Plains is characterized by rolling rangelands dominated by honey mesquite (*Prosopis glandulosa*), Pinchot's juniper (*Juniperus pinchotii*), and prickly pear (*Opuntia* spp.) (Rollins, 2007). Annual rainfall varies from 400 mm along the westward edge to 750 mm along the east; average annual temperatures are 15–17°C across the region (Rollins, 2007; Texas Parks and Wildlife Department, 2013).

Host collection

Bobwhites were live trapped using wire mesh funnel traps baited with grain sorghum (*Sorghum bicolor*) and covered with vegetation to mitigate thermal stress and aid in concealment from predators (for more detail see Bruno, 2014; Dunham *et al.*, 2016). We trapped bobwhites for two weeks in August and October in 2011, 2012 and 2013. This sampling interval was selected to avoid the peak breeding and nesting season during spring and summer (March–July) and the hunting season in autumn and winter (November–February). We processed bobwhites within 12 h of capture. We determined age and sex according to plumage characteristics. As part of a landowner agreement under the OID project protocol, a maximum of 30 bobwhites could be captured per study site over a 3-day sample period. Computer-generated random selection was used to determine which birds went to the various projects being conducted under OID. Bobwhites designated for necropsy were euthanized using cervical dislocation, and flash frozen immediately following death. Carcasses were wrapped in aluminium foil and placed within a small Styrofoam cooler with dry ice and ethanol to rapidly cool samples and prevent autolysis of helminths (Pence *et al.*, 1988). Bobwhite carcasses remained frozen until necropsy.

Helminth collection, processing and identification

Frozen carcasses were thawed in a refrigerator the night before necropsy to reduce damage to parasites from repeated freezing and thawing. Complete necropsies were performed on all sample hosts (see Bruno, 2014 and Bruno *et al.*, 2015 for a detailed description of procedures and specific helminth microhabitats examined). Nematodes were fixed in glacial acetic acid and preserved in 70% alcohol and 8% glycerin in individually marked vials corresponding to each bird and organ. Acanthocephalans and cestodes were fixed in acid–formalin–ethyl alcohol (AFA) solution and preserved in 70% alcohol in individual vials. Specimens were examined in ethyl alcohol wet mounts, identified and counted. Acanthocephalans and cestodes were stained using Mayer's Carmine Alum and mounted in Canada balsam on microscope slides. Cestodes were counted based on the number of scolices found within each cestode-infected host. Voucher specimens were deposited at the Sam Houston State University Parasite Museum, Sam Houston State University, Huntsville, Texas (accession numbers SHSUP000336–381 and SHSUP000474–483). Parasitological terms follow Bush *et al.* (1997). Rare species were defined as those with <25% prevalence, intermediate species with ≥25 and <75% prevalence, and common species with ≥75% prevalence across the collective host sample (Landgrebe *et al.*, 2007).

Statistical analyses

Statistical analyses were conducted using the intermediate and common helminth species. A chi-square analysis using JMP® 10.0 software (SAS Institute Inc., Cary, North Carolina, USA) was used to determine if prevalence for the intermediate and common species differed by host age [adult, juvenile], host sex [male, female], month [August, October], and year of collection [2011, 2012, 2013]. Abundance data for the intermediate and common species were aggregated towards zero. To account for non-normality, data were fitted with a negative binomial distribution and regressed using a generalized linear mixed model procedure (PROC GLIMMIX) with SAS 9.2 software (SAS Institute Inc.). Models were developed to explain variation in helminth abundance of the intermediate and common species. The initial model included the main effects variables (host age, host sex, month, year of collection, and northing [Universal Trans Mercator Northing coordinate; distance north in km]), and their relevant interactions (host age*host sex, host age*year, and host age*month). Backwards variable elimination was used to estimate the best-fitting model at $P \leq 0.05$ (for each included independent variable). This method allowed estimation of the relationship between the number of helminths and each variable, while accounting for effects of multicollinearity among parameters. A pseudo- r^2 (Nakagawa & Schielzeth, 2013) was calculated to estimate the amount of variation in helminth abundance explained by the model. Least squares means were estimated using the inverse of a log link function to compare effects among levels of categorical variables. These estimates are different from the arithmetic means reported in the descriptive analyses. Descriptive statistics are presented as mean ± SE. Statistical tests were considered significant at $P \leq 0.05$.

Two types of community analyses were used on the helminth data. The Percent Similarity Index (Ps_j ; Krebs, 1989) compared proportions of species among host component communities. Numerical similarity of helminth species was measured using Jaccard's Coefficient of Similarity Index (C_j ; Magurran, 1988).

Results

From August and October 2011–2013, we collected 161 bobwhites. Of these, 13 birds were collected from three adjacent ecoregions (Edwards Plateau, High Plains, and Southwestern Tablelands) within 60 km of the Rolling Plains ecoregion boundary (*fig. 1*). The sample comprised 31 adult males, 21 adult females, 53 juvenile males and 56 juvenile females. By collection period, sample size was as follows: August and October 2011, 29 adults, 12 juveniles; August and October 2012, 10 adults, 46 juveniles; August and October 2013, 13 adults, 51 juveniles.

Helminth fauna

We identified 13 species of helminths (9 nematodes, 3 acanthocephalans (1 species being a cystacanth) and 1 cestode) from 12 microhabitats representing 14,428 helminth individuals (*table 1*). *Aulonocephalus pennula* (= *A. lindquisti*) was the most prevalent (73%) and numerically dominant species (95% of all helminth individuals), followed by *Oxyspirura petrowi* (40% prevalence; 3% of all helminth individuals); the remaining species rarely




Fig. 1. Helminths were collected from northern bobwhites (*Colinus virginianus*) during August and October 2011–2013 from collection sites in the Rolling Plains ecoregion of Texas and western Oklahoma, and three adjacent ecoregions. Note: Oklahoma does not categorize its ecoregions similarly to Texas; Rolling Plains ecoregion of Texas = Central Great Plains of Oklahoma.

occurred (table 1). Intensity and abundance followed the same pattern of species numerical dominance as prevalence.

Prevalence by age, sex, month and year

Prevalence and abundance were examined statistically for two species (*A. pennula* and *O. petrowi*), based on their overall prevalence of $\geq 25\%$. Prevalence of *A. pennula* in adult bobwhites (90%) was significantly higher ($P = 0.0005$) than in juveniles (64%), but prevalence did not differ ($P = 0.712$) between males (73%) and females (73%). Prevalence of *O. petrowi* was significantly greater ($P = 0.0001$) in adult bobwhites (63%) compared to juveniles (28%), but prevalence did not differ ($P = 0.400$) between males (43%) and females (36%). Prevalence of *A. pennula* in October (83%) was significantly higher ($P = 0.001$) than in August (59%). Prevalence of *O. petrowi* in August (35%) was similar ($P = 0.265$) to October (43%). Prevalence of *A. pennula* was similar ($P = 0.107$) across 2011 (85%), 2012 (68%) and 2013 (69%). Prevalence of *O. petrowi* in 2012 (21%) was significantly lower ($P = 0.002$) than in 2011

(51%) and 2013 (48%). Prevalence of *O. petrowi* was similar ($P = 0.785$) between 2011 and 2013.

Trend analyses of parasite abundance

The best model for *A. pennula* was the following: abundance = age + northing (associated F and P values presented in table 2). The model explained 12.5% of the variation in *A. pennula* abundance (pseudo- $r^2 = 0.125$). Estimated mean of *A. pennula* in adults (161.2 ± 40.9) was significantly higher than in juveniles (43.8 ± 7.7). For any given age, an estimated slope of 0.00139 is interpreted as follows: estimated mean number of *A. pennula* decreases by 0.139% for each 1 km increase in northing.

The best model for *O. petrowi* was the following: abundance = age + sex + age*sex + year + northing (associated F and P values presented in table 2). The model explained 31% of the variation in *O. petrowi* abundance (pseudo- $r^2 = 0.31$). The main effect of host sex was not significant on its own, but it was included in the model because of the age*sex interaction. The interaction effect between age and sex showed the estimated mean of *O. petrowi*

Table 1. Prevalence (number of infected, % infected, and 95% confidence intervals [CI]), intensity (mean \pm SE; range), and abundance (mean \pm SE) of helminths from 161 northern bobwhites (*Colinus virginianus*) collected during August and October 2011–2013 within the Rolling Plains ecoregion of Texas and western Oklahoma.

Helminth species ^a	Prevalence		Intensity		Abundance	
	No.	% (95% CI)	Mean \pm SE	Range	Mean \pm SE	Total
<i>Aulonocephalus pennula</i> (SI, L, C)	117	73% (65.8–79.5)	116.9 \pm 11.6	1–617	84.9 \pm 9.1	13,675
<i>Oxyspirura petrowi</i> (E)	64	40% (32.2–47.3)	7.7 \pm 1.5	1–61	3.1 \pm 0.6	495
Acanthocephalan larvae (E, N, C, SI)	15	9% (4.8–13.8)	5.3 \pm 2.1	1–32	0.5 \pm 0.2	80
<i>Tetrameres pattersoni</i> (P)	23	14% (8.8–19.7)	3.0 \pm 0.5	1–10	0.4 \pm 0.1	69
<i>Physaloptera</i> sp. (BM)	11	7% (2.9–10.7)	4.1 \pm 1.1	1–12	0.3 \pm 0.1	45
<i>Rhabdometra odiosa</i> (SI)	12	7% (4.8–13.8)	3.7 \pm 1.0	1–12	0.3 \pm 0.1	44
<i>Gongylonema phasaniella</i> (C)	3	2% (<0.1–3.9)	3.3 \pm 0.3	1–4	0.1 \pm <0.1	10
<i>Cheliospirura spinosa</i> (G)	2	1% (<0.1–2.9)	1.5 \pm 0.5	1–2	<0.1 \pm <0.1	3
<i>Dispharynx nasuta</i> (P)	2	1% (<0.1–3.9)	1.0 \pm 0.0	1	<0.1 \pm <0.1	2
<i>Mediorhynchus papillosus</i> (SI)	2	1% (<0.1–2.9)	1.0 \pm 0.0	1	<0.1 \pm <0.1	2
<i>Eucoleus contortus</i> (C)	1	1% (N/A)	1.0 \pm N/A	1	<0.1 \pm <0.1	1
<i>Mediorhynchus</i> sp. (SI)	1	1% (N/A)	1.0 \pm N/A	1	<0.1 \pm <0.1	1
<i>Subulura brumpti</i> (C)	1	1% (N/A)	1.0 \pm N/A	1	<0.1 \pm <0.1	1

^a Microhabitats: BM, breast muscle; C, ceca; E, eye surface, nictitating membrane, associated glands, and ducts; G, gizzard; L, large intestine; N, neck muscle; P, proventriculus; SI, small intestine

Table 2. *F* and *P* values generated from the final main and interaction effects variables of the negative binomial generalized linear regression model for *Aulonocephalus pennula* and *Oxyspirura petrowi* from 161 northern bobwhites (*Colinus virginianus*) collected during August and October 2011–2013 within the Rolling Plains ecoregion of Texas and western Oklahoma.

<i>Aulonocephalus pennula</i>			<i>Oxyspirura petrowi</i>		
Effect	F	P	Effect	F	P
Age	17.75	<0.0001	Age	32.41	<0.0001
Northing	4.61	0.0334	Year	6.18	0.0026
Age * Sex ^a	1.54	0.2165	Northing	21.04	<0.0001
Age * Year ^a	0.73	0.4852	Age * Sex	5.51	0.0049
			Age * Year ^a	0.23	0.7954

^aNon-significant interaction terms retained in final model when it included a significant main effects variable.

in adult males (11.0 \pm 3.7) was significantly higher than in adult females (1.8 \pm 0.8), juvenile males (0.4 \pm 0.2), and juvenile females (0.6 \pm 0.2). The estimated mean of *O. petrowi* by year was similar between 2011 (1.7 \pm 0.5) and 2013 (3.2 \pm 0.9), and similar between 2011 and 2012 (0.7 \pm 0.2), but was significantly higher in 2013 compared to 2012. For any given age*sex combination in each year, an estimated slope of -0.00376 is interpreted as follows: estimated mean number of *O. petrowi* decreases by 0.376% for each 1 km increase in northing. Although there is a large amount of variation in abundance not explained by the variables included in regression models (low r^2 values), the significant variables in these models can still be used to draw important conclusions about helminth abundance in this sample (see Discussion).

Helminth community dynamics

Infracommunity species richness ranged from 1 to 5, with 1 to 2 species occurring in a majority (64%) of the hosts. Species distribution by year showed a similar pattern among hosts, with 1 to 2 occurring in a majority of the hosts for each year (fig. 2). There

was only one year (2012) in which host individuals ($n = 9$) had more than four helminth species.

Based on PS_i values, component communities were almost completely similar among host age, host sex, month and year (table 3). Component communities within each comparison were numerically dominated by *A. pennula*, accounting for 90 to 98% of all helminth individuals. *Oxyspirura petrowi* accounted for 2 to 7% of the total individuals for each comparison, and all other species accounted for < 1%.

Jaccard's Coefficient of Similarity (C_j) found seven (54%) shared species between adults (10 helminth species) and juveniles (10 species), and seven (54%) shared species between male (11 species) and female (9 species) host helminth communities (table 3). The host helminth community in August (11 species) and October (11 species) shared nine (70%) of the species that occurred in each month. Component communities between 2011 (7 species) and 2012 (10 species) shared six (50%) of the species found in each year. Component communities between 2011 and 2013 (9 species) were slightly less similar, sharing five (45%) of the species found in each year. Component communities




Fig. 2. Distribution of helminth species by year in 161 northern bobwhites (*Colinus virginianus*) collected during August and October 2011 (black bars), 2012 (grey bars) and 2013 (white bars) within the Rolling Plains ecoregion of Texas and western Oklahoma.

Table 3. Comparisons of Percent Similarity (PS_i) and Jaccard's Coefficient of Similarity (C_j) indices for helminth communities by host age, host sex, month and year from 161 northern bobwhites (*Colinus virginianus*) collected during August and October 2011–2013 within the Rolling Plains ecoregion of Texas and western Oklahoma.

Comparison	PS_i^a	C_j^b
Adults ($n = 52$) vs Juveniles ($n = 109$)	96.8	0.54
Males ($n = 84$) vs Females ($n = 77$)	96.4	0.54
August ($n = 84$) vs October ($n = 77$)	98.3	0.70
2011 ($n = 41$) vs 2012 ($n = 56$)	98.0	0.50
2011 ($n = 41$) vs 2013 ($n = 64$)	98.2	0.45
2012 ($n = 56$) vs 2013 ($n = 64$)	99.1	0.67

^a Values for PS_i range from 0 to 100, where 0 = completely dissimilar communities, and 100 = completely similar communities.

^b Values for C_j range from 0 to 1, where 0 = completely dissimilar communities, and 1 = completely similar communities.

between 2012 and 2013 were more similar than comparisons between other years, sharing eight (67%) species found in each year (table 3). In each comparison, the helminths not shared by both component communities represent rare species and occurred in <2% of the population.

Discussion

Helminth community dynamics

The helminth component community was composed largely of heteroxenous species (12 of 13 species), but was numerically

dominated by a single species (*A. pennula*). Helminth species richness values in the present study are similar to those found in the southeastern U.S., but higher than in other bobwhite studies from Texas (Parmalee, 1952; Purvis *et al.*, 1998; Villarreal *et al.*, 2016). Cram *et al.* (1931) reported 12 species of helminths in wild bobwhites throughout the eastern U.S., and several surveys from Florida reported 12–15 species of helminths (Davidson *et al.*, 1980, 1991; Moore *et al.*, 1986; Moore & Simberloff, 1990). These component communities consisted of a similar distribution of intermediate and rare species to our study, but they had higher occurrences of common species ($n = 3$). Species composition also differed between the component communities in the southeastern U.S. and our study. *Aulonocephalus pennula* was the numerically dominant species in our study and is a common (>75% prevalence) and abundant species found in quail species from Texas (Webster & Addis, 1945; Lehmann, 1984; Landgrebe *et al.*, 2007; Bedford, 2015; Olsen & Fedynich, 2016; Villarreal *et al.*, 2016) and New Mexico (Campbell & Lee, 1953), but is not found in bobwhites from Florida, Georgia (Kellogg & Prestwood, 1968), North Carolina (Cram *et al.*, 1931; Blakeney & Dimmick, 1971) or Kansas (Williams *et al.*, 2004). Component communities in the southeastern U.S. also consist of more monoxenous nematode species (range: 2–4), which occurred in high intensities (>30; Moore *et al.*, 1986; Moore & Simberloff, 1990; Davidson *et al.*, 1991). The difference in species composition and variation in life cycles of helminths across the bobwhite's geographical range is probably the result of mesic conditions found in the southeastern U.S., compared to the more semi-arid Rolling Plains.

It has been suggested that the dominance of heteroxenous species within a helminth community in semi-arid to arid regions

may be a survival strategy (Fedynich *et al.*, 2001; Landgrebe *et al.*, 2007). Helminths can persist longer in harsher environments if they maintain infective stages in intermediate hosts (Mackiewicz, 1988). This may be why direct life cycle helminths such as *Heterakis* spp., *Trichostrongylus tenuis*, *Strongyloides* spp. and *Capillaria* spp. are more common in bobwhites from mesic areas in the southeastern U.S. (Cram *et al.*, 1931; Kellogg & Prestwood, 1968; Davidson *et al.*, 1980, 1991; Moore *et al.*, 1987; Moore & Simberloff, 1990). This is supported by studies that have found a high prevalence of *T. tenuis* in bobwhites from Texas (Demarais *et al.*, 1987; Purvis *et al.*, 1998), where study areas were dominated by poorly drained clay soils which received precipitation that exceeded the long-term average for the Rolling Plains. Similarly, high intensities of *Heterakis gallinae* were found in bobwhites from north-eastern Texas, a forested region where precipitation and humidity are relatively higher than other areas of Texas (Parmalee, 1952).

Factors influencing helminth population dynamics

Host age was an important variable for both *A. pennula* and *O. petrowi* populations, with more adults infected and having a higher intensity of infection than juveniles. An age effect has been found for various helminth species in bobwhites (Davidson *et al.*, 1980; Forrester *et al.*, 1984; Moore *et al.*, 1986; Davidson *et al.*, 1991; Dunham *et al.*, 2014, 2016; Villarreal *et al.*, 2016). Host age-related differences are attributed to time-related accumulation in adults resulting from longer exposures to infective stages of helminths (Davidson *et al.*, 1980).

Host sex alone was not an important variable in *A. pennula* and *O. petrowi* prevalence or abundance. Host sex is usually not a significant variable in parasite infections of bobwhites (Blakeney & Dimmick, 1971; Moore *et al.*, 1987). Similarities in male and female social behaviour and diet probably create equal exposure probabilities to helminth infective stages. Higher numbers of *O. petrowi* were found in adult males compared to any other age*sex combination. The biological significance of this interaction is unknown. We would expect this result in adult females because of their higher protein (mainly sourced from arthropods) requirements during reproduction (Hernández & Peterson, 2007), as found by Dunham *et al.* (2014).

Concerns have been raised regarding negative impacts of *A. pennula* on bobwhites (Dunham *et al.*, 2017). Similar to Olsen & Fedynich (2016), we observed distension in the terminal portions of the ceca when *A. pennula* infections were high (>300 worms). Nematodes may take advantage of regulatory molecules to evade the host immune system (Maizels *et al.*, 2001), creating tolerance and allowing infections to persist. This phenomenon has not been investigated for the helminths occurring in bobwhites, but it may be possible that the success of *A. pennula* in bobwhites is because of immunomodulators. Consistent with previous descriptions (Chandler, 1935), we did not find *A. pennula* attached to the cecal wall, and Dunham *et al.* (2017) observed no inflammation or lesions in bobwhites with high infections. It is possible that intensities of *A. pennula* in bobwhites are high because an antibody response is not occurring as a result of non-invasive contact with host tissues.

The only significant difference detected by month was prevalence of *A. pennula*, increasing from August to October. Villarreal *et al.* (2016) found a similar seasonal effect, where the prevalence of *A. pennula* increased from the summer to early winter period. *Oxyspirura petrowi* prevalence and abundance in our

study was lowest in 2012, but similar between 2011 and 2013. One explanation is that 82% of the bobwhites sampled in 2012 were juvenile birds and only six individuals were infected with *O. petrowi*.

Counts of both *A. pennula* and *O. petrowi* were found to increase in the study area from north to south. Jackson & Green (1965) similarly noted that *O. petrowi* prevalence was higher in their southern collection sites in the Rolling Plains compared to their northern sites. This spatial pattern could be affected by the variation of bobwhite density, distribution of the insect intermediate hosts (Keymar & Anderson, 1979), precipitation and habitat across the study area.

Our study documented bobwhite helminth community composition at a regional scale in a landscape where bobwhites are highly valued both ecologically and economically. The prevalence and spatial patterns of *O. petrowi* in bobwhites found in earlier studies (Jackson & Green, 1965) were similar to our study, reflecting relatively average stable trends for this species when sampled across the entire ecoregion. We also confirm recent documentations of high prevalence, intensity and abundance of *A. pennula* (Villarreal *et al.*, 2016; Dunham *et al.*, 2017). Current research focusing on understanding the life cycles of *A. pennula* and *O. petrowi* (Kistler *et al.*, 2016; Henry *et al.*, in press) may help clarify their patterns of numerical dominance in the region. Additionally, continued study of the bobwhite-helminth system will provide insight into the infection dynamics of helminths currently reported within the Rolling Plains ecoregion and aid in monitoring current pathogenic helminths (i.e. *O. petrowi*, *Dispharynx nasuta* and *Tetrameres pattersoni*) and the possible establishment of additional helminth species within the region.

Acknowledgements. We thank Texas Tech University, Texas A&M University at College Station, and the Oklahoma Department of Wildlife and Conservation for their assistance in trapping, field processing, and logistics. We thank the landowners for their hospitality and allowing us to sample on their properties. Additionally, we thank Dr Mike Kinsella for his assistance in the identification of helminths. This is manuscript No. 17-124 of the Caesar Kleberg Wildlife Research Institute.

Financial support. This research was funded by the Rolling Plains Quail Research Foundation and Park Cities Quail, and supported by the Caesar Kleberg Wildlife Research Foundation.

Conflict of interest. None.

Ethical standards. Bobwhites were collected in accordance with established permits and protocols approved by the TAMUK Institutional Animal Care and Use Committee (2009-09-21A), TAMU (AUP 2011-193), TAMUK Institutional Biosafety Committee (IBC-ID#009-2011), and Texas Parks and Wildlife Department Scientific Research permit (SPR-0690-152).

References

- Bedford KA (2015) *Parasitological Survey of Scaled Quail from West Texas* (MS thesis). Texas A&M University-Kingsville, Kingsville, Texas.
- Blakeney WC Jr and Dimmick RW (1971) Gizzard and intestinal helminths of bobwhite quail in Tennessee. *Journal of Wildlife Management* **35**, 559–562.
- Boggs JF *et al.* (1990) Occurrence and pathology of physalopterid larvae infections in bobwhite quail from western Oklahoma. *Proceedings of the Oklahoma Academy of Science* **70**, 29–31.
- Bruno A (2014) *Survey for Trichomonas gallinae and Assessment of Helminth Parasites in Northern Bobwhites from the Rolling Plains Ecoregion* (MS thesis). Texas A&M University-Kingsville, Kingsville, Texas.

- Bruno A et al.** (2015) Pathological response of northern bobwhites to *Oxyspirura petrowi* infections. *Journal of Parasitology* **101**, 364–368.
- Bush AO et al.** (1997) Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* **83**, 575–583.
- Campbell H and Lee L** (1953) *Studies on Quail Malaria in New Mexico and Notes on Other Aspects of Quail Populations*. New Mexico Department of Game and Fish report.
- Chandler AC** (1935) A new genus and species of Subuluriae (Nematodes). *Transactions of the American Microscopical Society* **54**, 33–35.
- Cram EB, Jones MF and Allen EA** (1931) Internal parasites and parasitic diseases of the bobwhite. In Stoddard HL (ed.), *The Bobwhite Quail: Its Habits, Preservation, and Increase*. New York, NY: Charles Scribner's Sons, pp. 229–313.
- Davidson WR, Kellogg FE and Doster GL** (1980) Seasonal trends of helminth parasites of bobwhite quail. *Journal of Wildlife Diseases* **16**, 367–374.
- Davidson WR et al.** (1991) Ecology of helminth parasitism in bobwhites from northern Florida. *Journal of Wildlife Diseases* **27**, 185–205.
- Demarais S, Everett DD and Pons ML** (1987) Seasonal comparison of endoparasites of northern bobwhites from two types of habitat in southern Texas. *Journal of Wildlife Diseases* **23**, 256–260.
- Dunham NR et al.** (2014) Evidence of an *Oxyspirura petrowi* epizootic in northern bobwhites (*Colinus virginianus*). *Journal of Wildlife Diseases* **50**, 552–558.
- Dunham NR et al.** (2016) Eyeworms (*Oxyspirura petrowi*) in northern bobwhites (*Colinus virginianus*) from the Rolling Plains ecoregion of Texas and Oklahoma, 2011–2013. *Journal of Wildlife Diseases* **52**, 562–567.
- Dunham NR et al.** (2017) Caecal worm, *Aulonocephalus pennula*, infection in the northern bobwhite quail, *Colinus virginianus*. *International Journal for Parasitology: Parasites and Wildlife* **6**, 35–38.
- Fedynich AM, Monasmith T and Demarais** (2001) Helminth community structure and pattern in Merriam's kangaroo rats, *Dipodomys merriami* Mearns, from the Chihuahuan Desert of New Mexico, U. S. A. *Comparative Parasitology* **68**, 116–121.
- Forrester DJ et al.** (1984) Ecology of helminth parasitism of bobwhites in Florida. *Proceedings of the Helminthological Society of Washington* **51**, 255–260.
- Henry C et al.** (in press) Molecular identification of potential intermediate hosts of *Aulonocephalus pennula* from the order Orthoptera. *Journal of Helminthology*. doi: 10.1017/S0022149X18000111.
- Hernández F and Peterson MJ** (2007) Northern bobwhite ecology and life history. In Brennan LA (ed.), *Texas Quails: Ecology and Management*. College Station, TX: Texas A&M University Press, pp. 40–64.
- Hernández F et al.** (2013) On reversing the northern bobwhite population decline: 20 years later. *Wildlife Society Bulletin* **37**, 177–188.
- Jackson AS and Green H** (1965) *Dynamics of Bobwhite Quail in the West Texas Rolling Plains: Parasitism in Bobwhite Quail* (Federal Aid Project No. W-88-R-4). Austin, TX: Texas Parks and Wildlife Department.
- Kellogg FE and Prestwood AK** (1968) Gastrointestinal helminths from wild and pen-raised bobwhites. *Journal of Wildlife Management* **32**, 468–475.
- Keymar AE and Anderson RM** (1979) The dynamics of infection of *Tribolium confusum* by *Hymenolepis diminuta*: the influence of infective-stage density and spatial distribution. *Parasitology* **79**, 195–207.
- Kiel WH Jr** (1976) Bobwhite quail population characteristics and management implications in South Texas. *Transactions of the North American Wildlife and Natural Resources Conference* **41**, 407–420.
- Kistler WM et al.** (2016) Plains lubber grasshopper (*Brachystola magna*) as a potential intermediate host for *Oxyspirura petrowi* in northern bobwhites (*Colinus virginianus*). *Parasitology Open* **2**, 1–8.
- Kocan AA, Hannon L and Eve JH** (1979) Some parasitic and infectious diseases of bobwhite quail from Oklahoma. *Proceedings of the Oklahoma Academy of Science* **59**, 20–22.
- Krebs CJ** (1989) Other similarity measures. In *Ecological Methodology*. New York, NY: Harper Collins Publishers, pp. 304–307.
- Landgrebe JN et al.** (2007) Helminth community of scaled quail (*Callipepla squamata*) from western Texas. *Journal of Parasitology* **93**, 204–208.
- Lehmann VW** (1984) *Bobwhites in the Rio Grande Plain of Texas*. College Station, TX: Texas A&M University Press, 371 pp.
- Mackiewicz JS** (1988) Cestode transmission patterns. *Journal of Parasitology* **74**, 60–70.
- Magurran AE** (1988) *Ecological Diversity and its Measurement*. Princeton, NJ: Princeton University Press, 179 pp.
- Maizels RM, Blaxter ML and Scott AL** (2001) Immunological genomics of *Brugia malayi*: filarial genes implicated in immune evasion and protective immunity. *Parasite Immunology* **23**, 327–344.
- Moore J and Simberloff D** (1990) Gastrointestinal helminth communities of bobwhite quail. *Ecology* **71**, 344–359.
- Moore J, Freehling M and Simberloff D** (1986) Gastrointestinal helminths of the northern bobwhite in Florida: 1968 and 1983. *Journal of Wildlife Diseases* **22**, 497–501.
- Moore J et al.** (1987) Host age and sex in relation to intestinal helminths of bobwhite quail. *Journal of Parasitology* **73**, 230–233.
- Nakagawa S and Schielzeth H** (2013) A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution* **4**, 133–142.
- Olsen A and Fedynich AM** (2016) Helminth infections in northern bobwhites (*Colinus virginianus*) from a legacy landscape in Texas, USA. *Journal of Wildlife Diseases* **53**, 562–567.
- Parmalee PW** (1952) Ecto- and endoparasites of the bobwhite: their numbers, species, and possible importance in the health and vigor of quail. *The North American Wildlife Conference* **17**, 176–188.
- Pence DB et al.** (1988) Letters to the editors: Critical comments on a recent letter to the editors regarding the use of frozen carcasses in parasite surveys. *Journal of Parasitology* **74**, 197–198.
- Purvis JR et al.** (1998) Northern bobwhites as disease indicators for the endangered Attwater's prairie chicken. *Journal of Wildlife Diseases* **34**, 348–354.
- Rollins D** (1980) *Comparative Ecology of Bobwhite and Scaled Quail in Mesquite Grassland Habitats* (MS thesis). Stillwater, OK: Oklahoma State University.
- Rollins D** (2007) Quails on the Rolling Plains. In Brennan LA (ed.), *Texas Quails: Ecology and Management*. College Station, TX: Texas A&M University Press, pp. 117–141.
- Sauer JR et al.** (2013) The North American breeding bird survey, results and analysis 1966–2011, Version 07.03.2013. Laurel, Maryland, USGS Patuxent Wildlife Research Center.
- Texas Parks and Wildlife Department** (2013) Bobwhite and scaled quail in the Rolling Plains. Available at https://tpwd.texas.gov/huntwild/hunt/planning/quail_forecast/forecast/rolling_plains/ (accessed 1 March 2013).
- Tri AN et al.** (2013) Impacts on northern bobwhite sex ratios, body mass, and annual production in South Texas. *Journal of Wildlife Management* **77**, 579–586.
- Villarreal SM et al.** (2016) Helminth infections across a northern bobwhite (*Colinus virginianus*) annual cycle in Fisher County, Texas. *Western North American Naturalist* **76**, 275–280.
- Webster JD and Addis CJ** (1945) Helminths from the bobwhite quail in Texas. *Journal of Parasitology* **31**, 286–287.
- Williams CK et al.** (2004) Health status of northern bobwhite quail (*Colinus virginianus*) in eastern Kansas. *Avian Diseases* **44**, 953–956.