

Pine chemical volatiles promote dauer recovery of a pine parasitic nematode, *Bursaphelenchus xylophilus*

Research Article

Cite this article: Zhang W, Li Y, Pan L, Wang X, Feng Y, Zhang X (2020). Pine chemical volatiles promote dauer recovery of a pine parasitic nematode, *Bursaphelenchus xylophilus*. *Parasitology* **147**, 50–57. <https://doi.org/10.1017/S0031182019001264>


Received: 16 May 2019
Revised: 18 August 2019
Accepted: 21 August 2019
First published online: 17 September 2019

Key words:

Chemical volatiles; dauer recovery; pine wood nematode; transformation

Author for correspondence:

Yongxia Li, E-mail: liyongxiaxjs@163.com

Wei Zhang^{1,2} , Yongxia Li^{1,2}, Long Pan^{1,2}, Xuan Wang^{1,2}, Yuqian Feng^{1,2} and Xingyao Zhang^{1,2}

¹Lab. of Forest Pathogen Integrated Biology, Research Institute of Forestry New Technology, Chinese Academy of Forestry, Beijing 100091, China and ²Co-Innovation Center for Sustainable Forestry in Southern China, Nanjing Forestry University, Nanjing 210037, China

Abstract

Pinewood nematode, *Bursaphelenchus xylophilus*, a pine parasitic nematode, poses a serious threat to its host pine forests globally. When dispersal-stage larvae 4 (dauer, DL4) of *B. xylophilus* enters the host pine, it moults into propagative adult (dauer recovery) and reproduces quickly to kill the host pine. Here, we found pine chemical volatiles, rather than the common dauer recovery factors of nematodes (e.g. suitable temperatures, nutrient availability or density), promote *B. xylophilus* dauer recovery. The results showed that volatilization of chemicals in host pines could attract DL4 and promote DL4 recovery. To identify which chemicals promote this process, we determined the stimulated activity of the main volatiles of pines including six monoterpenes and two sesquiterpenes. Results showed that all the six monoterpenes promoted dauer recovery, especially β -pinene and β -myrcene, but the two sesquiterpenes have no effect on the transformation. Furthermore, β -pinene performed gradient effects on dauer recovery. We hypothesized that when DL4 infect pine trees, the pine volatiles released from the feeding wounds are used as chemical signals for DL4 transformation to adult to reproduce and rapidly kill the pines. Our study identified the *B. xylophilus* dauer recovery chemical signal and may contribute to preventing pine wilt disease.

Introduction

The pine wood nematode (PWN), *Bursaphelenchus xylophilus*, is a highly invasive species that causes pine wilt disease (PWD). It has devastating effects to pine forests of Japan, China, Korea and Europe (Nickle *et al.*, 1981; Cheng *et al.*, 1983; Yi *et al.*, 1989; Mota *et al.*, 1999; Zamora *et al.*, 2015), and poses a serious threat to pine forests globally. The life cycle of *B. xylophilus* involves two forms, propagative and dispersal (Mamiya, 1975, 1983b). Under suitable environmental conditions, the nematodes will reproduce and grow very rapidly from egg to adult. However, when the environment becomes unsuitable, the dispersal-stage larvae 3 (DL3) and dispersal-stage larvae 4 (dauer, DL4) appear (Mamiya, 1983a; Fukushima, 1991; Maehara and Futai, 2000). The DL3 nematodes moult to DL4 to enter the trachea of enclosing beetles (Zhao *et al.*, 2016). Following dispersal, while the vector beetles are feeding on healthy pines, DL4 nematodes leave the trachea, enter the pines and moult into adults. These propagative nematodes reproduce and kill the pines (Zhao *et al.*, 2013). Thus, the dauer recovery (DL4 transformation to adult) of *B. xylophilus* is the final step leading to the successful spread of PWD to healthy pines.

In response to harsh environmental conditions, most nematode species undergo a diapause stage, dauer arrest, prior to the reproductive stage. In *Caenorhabditis elegans*, a free-living nematode, the dauer developmental decision hinges on the integration of three environmental parameters: population density, nutrient supply and ambient temperature. A high population density initiates the dauer developmental program, while high temperatures and reduced nutrient resources strongly potentiate this decision (Golden and Riddle, 1982, 1984; Ouellet *et al.*, 2008). Thus, dauer recovery, which leads to the propagative stage, requires a low population density, sufficient nutrients and suitable temperatures (Ouellet *et al.*, 2008). As with *C. elegans*, the food supply-related signal could also induce the recovery of entomopathogenic *Heterorhabditis* spp. nematodes (Strauch and Ehlers, 1998; Aumann and Ehlers, 2001; Dolan *et al.*, 2003). However, requirements for dauer recovery are not clearly understood as in many other nematodes.

Terpenes, mainly composed of monoterpenes, sesquiterpenes and diterpenes (Chen *et al.*, 2018), are the main pine volatile components responsible for plant defence (Smith, 2000; Xu *et al.*, 2018) and kairomone attraction of phytophagous pests (Fan *et al.*, 2007; Xu *et al.*, 2016). In *B. xylophilus*, propagative larvae are attracted to terpene volatiles (α -pinene, β -pinene and longifolene) produced by the host pine (Zhao *et al.*, 2007). DL4 are attracted to β -myrcene and are recovered by some monoterpenes like β -myrcene, limonene and α -pinene (Hinode *et al.*, 1987; Stamps and Linit, 1998a). While population density, nutrient supply or ambient




Fig. 1. Transformation rate of *B. xylophilus* DL4 promoted by different nutrients. *G. biloba* is from Ginkgoopsida, as an outgroup. Statistical differences in the means are indicated with different letters, $P < 0.05$. Error bars represent \pm s.e.

temperature are well-known factors promoting nematode dauer recovery, the role that volatile terpenes like sesquiterpenes of pines play in dauer recovery of *B. xylophilus* is unclear.

Here, we investigated the effects of food supply, population density, temperature and volatile terpenes of pines on dauer recovery. The results showed that pine volatiles, but the common dauer recovery stimulators, could promote the dauer recovery of *B. xylophilus* from DL4 to adult. Understanding the mechanisms that trigger dauer recovery may lead to new applications in PWD prevention, for example, if a technique is developed to sustain dauer arrest even after transmission, then it may prevent PWD from developing.

Materials and methods

DL4 of *B. xylophilus*

DL4 nematodes were obtained from vector beetles of *Monochamus alternatus*. To ensure the stability of the DL4, we dissected the beetles collected at peak eclosion time (April to August, 2018) from Zhejiang and Guangdong Provinces, China. The dissected beetle was soaked in ddH₂O in a 60-mm Petri dish for 2 h. The DL4 nematodes swam out of the beetle trachea and were then sequentially transferred three times into new petri dishes using a 10- μ L pipette tip for decontamination. The collected DL4 nematodes were stored at 4 °C for further research.

Stimulation of the *B. xylophilus* dauer recovery process with different nutrients

Cellulose, glucose, *Botrytis cinerea* (the fungal food of *B. xylophilus*) and the twigs of different trees (*Ginkgo biloba*, *Pinus massoniana* and *Pinus thunbergii*) were collected at the vector beetle's peak eclosion time (May) on the campus of the Chinese

Academy of Forestry and used as nutrient sources for the dauer recovery of DL4. In total, 30 DL4 nematodes were soaked in 4 mL ddH₂O supplemented with different nutrients in a 35-mm petri dish. The concentrations of cellulose and glucose were 5×10^{-2} g mL⁻¹. The sizes of *B. cinerea*, *G. biloba*, *P. massoniana* and *P. thunbergii* were 5 mm \times 5 mm. After cultivating for 2 d in a 25 °C incubator, the numbers of adult stage nematodes were counted using an optical microscope equipped with a camera (CZX51 and BX51, Olympus, Japan), and the transformation rates of the DL4 were calculated. These tests were performed with three replicates.

Transformation rates of *B. xylophilus* DL4 stimulated by population density

To test the effects of DL4 population density on dauer recovery, we used one nematode and 100 DL4 nematodes per well as different population densities (Hirao and Ehlers, 2010). In a 96-well plate, every well had one nematode soaked in 100 μ L ddH₂O, with or without a \sim 2 mm \times 2 mm pine chip. Totally, there were 96 nematodes in 96 wells respectively for one nematode in one well tests. Meanwhile, in the well of 96-well plate, every well had 100 nematodes soaked in 100 μ L ddH₂O, with or without a \sim 2 mm \times 2 mm pine chip for one hundred nematodes in one well test. The test methods were as above, and the transformation rates of the DL4 were calculated. These tests were performed with three replicates.

Transformation rates of *B. xylophilus* DL4 stimulated by temperature

To test the effects of temperature on dauer recovery, a temperature gradient was used. In total, 30 DL4 nematodes were cultivated at 4, 10, 15, 20, 25, 30, 35, 40 and 45 °C. The nematodes were soaked in 4 mL ddH₂O, with or without a \sim 5 mm \times 5 mm pine chip. The




Fig. 2. Transformation rates of *B. xylophilus* DL4 promoted by different densities, with or without pine chips. (a) Transformation rates of *B. xylophilus* DL4 of one nematode in one well, with or without pine chips. Statistical differences in the means are indicated with '***', $P < 0.001$. Error bars represent \pm s.e. (b) Transformation rates of *B. xylophilus* DL4 of one hundred nematodes in one well, with or without pine chips. Statistical differences in the means are indicated with '***', $P < 0.001$. Error bars represent \pm s.e.

transformation rates of DL4 were calculated as above. These tests were performed with three replicates.

Transformation rates of *B. xylophilus* DL4 stimulated by pine volatiles

In these tests, petri dishes divided into two parts with a physical barrier in the middle were used (Fig. 4). In total, 50 DL4 nematodes were inoculated into the centre of 2% agarose media on one side of the petri dish. A 20-mm dish, with *G. biloba* or *P. massoniana* chips, and filled with 5 mL ddH₂O was adhered to the opposite side. After cultivating for 2 d in a 25 °C incubator, the numbers of nematodes at different transformational stages in both sides of the petri dish were counted, and the transformation rates of the DL4 were calculated. These tests were performed with three replicates.

Volatile analysis of *G. biloba*, *P. massoniana* and *P. thunbergii*

Volatiles of *G. biloba*, *P. massoniana* and *P. thunbergii* were collected by solid phase micro-extraction (SPME, 57310-U, Supelco, USA) and analysed by Gas chromatography-Mass spectrometry (GC-MS, Agilent, USA, Agilent 6980N GC coupled 5973 mass selective detector) equipped with a DB-WAX capillary column (30 m \times 0.25 mm, Agilent Technologies, USA), and the column temperature was programmed from an initial temperature 50 °C for 1 min, then increased by 5 °C min⁻¹ to 160 °C and held for 2 min, and last increased by 20 °C min⁻¹ to 250 °C and held for 5 min (Zhou *et al.*, 2017). Data files were analysed with the

automated mass spectral and identification system for peak deconvolution, and spectra were matched using the mass spectral library (NIST 2008) and a custom library. To further identify the main volatiles of these trees, the volatile samples were compared with candidate standard chemicals α -pinene (Sigma, USA, 98% purity), camphene (TCI, Japan, >78% purity, containing 20% Tricyclene), D-limonene (Sigma-Aldrich, USA, 98% purity), β -pinene (Sigma-Aldrich, USA, 99% purity), β -myrcene (Sigma-Aldrich, USA, 2000 μ g mL⁻¹ in hexane), β -phellandrene (TRC, Canada, 100% purity), longifolene (Sigma-Aldrich, USA, \geq 75% purity) and *trans*-caryophyllene (Sigma-Aldrich, USA, \geq 98% purity) using above methods. The contents of the main volatiles of the tested trees were measured using GC analysis of the hexane-extracted tree chip samples containing heptyl acetate as an internal standard (Xu *et al.*, 2015).

Transformation rates of *B. xylophilus* DL4 stimulated by the main volatiles

We tested the effects of the pine volatiles including α -pinene, camphene, D-limonene, β -pinene, β -myrcene, β -phellandrene, longifolene and *trans*-caryophyllene on dauer recovery. In total, 30 DL4 nematodes were cultivated in 5% Triton X-100 with or without 10⁻² g mL⁻¹ of each main volatile independently. These tests revealed the most promotive volatile, which was then used to test the gradient effects on dauer recovery. The gradient concentrations of this volatile were 0, 10⁻⁵, 10⁻⁴, 10⁻³, 10⁻² and 10⁻¹ g mL⁻¹. After cultivating for 2 d in a 25 °C incubator, the transformation rates of the DL4 were calculated as above. These tests were performed with three replicates.

Statistical analyses

In all experiments, the normality of data was measured using the Kolmogorov–Smirnov test, and the homogeneity of group variances was screened using Levene's test. The statistical significance of the population density promotive tests was evaluated using the unpaired two-tailed Student's *t* test. Different pine and pine volatile promotive tests were evaluated using one-way ANOVA (analysis of variance) with Tukey's test or Dunnett's T3 depending on normality and homogeneity. A two-way ANOVA was used to evaluate the between-subject effects on transformation rates promoted by *G. biloba* or *P. massoniana* chips. Data were analysed using SPSS 18.0 software (SPSS, Inc., Chicago, USA). All the quantitative data were represented as means \pm s.e. (standard error).

Results

B. xylophilus dauer recovery promoted by different foods, temperatures and densities

Here, we used cellulose, glucose, *B. cinerea*, *G. biloba*, *P. thunbergii* and *P. massoniana* as stimulators to monitor the DL4 transformation process from DL4 to adult. As *B. xylophilus* hosts, *P. thunbergii* and *P. massoniana* promoted high transformation rates of DL4 at \sim 80% significantly higher than the control at \sim 13%. However, The DL4 transformation rates stimulated by cellulose, *B. cinerea* and especially glucose were not significantly different compared to controls. In addition, *G. biloba* from Ginkgoopsida was tested as outgroup. The transformation rate when stimulated by *G. biloba* was similar to controls at \sim 15%. In addition, in the control test without a stimulator, some nematodes also moulted successfully (Fig. 1, one-way ANOVA, $F_{6,14} = 248.90$, $P < 0.001$).




Fig. 3. Transformation rates of *B. xylophilus* DL4 promoted by different temperatures, with or without pine chips.

In addition to nutrient supply, temperature and population density are known effectors of *C. elegans* dauer recovery. Consequently, the effects of these factors on *B. xylophilus* DL4 transformation were tested. For population density, one nematode per well and 100 nematodes per well were used to represent low- and high-population density levels, respectively. Independent of the population level, the transformation rates were low without pine stimulation. However, with pine stimulation, nematodes at both population density levels had significant transformation rates compared with control (Fig. 2a, $df = 4$, $t = 15.27$, $P < 0.001$; Fig. 2b, $df = 4$, $t = 22.01$, $P < 0.001$).

Without pine chips, the transformation rates of DL4 at all the tested temperatures were less than 30%, even after 5 d (Fig. 3). However, with pine chip stimulation, the transformation rates of DL4 at 25 and 30 °C increased rapidly to 85% after 2 d (Fig. 3). At less than 25 °C, the transformation rate decreased as the temperature decreased, until 4 °C, when the nematodes stopped transforming at all over 5 d (Fig. 3). Interestingly, the DL4 transformation was inhibited when temperature was greater

than 35 °C. At 35 °C, DL4 transformed rapidly during the first day, but the transformations stopped in the following days (Fig. 3). At 40 °C, like at 4 °C, the nematodes did not transform. All the DL4 nematodes died after 1 d at 45 °C (Fig. 3).

B. xylophilus DL4 recovery promoted by pine volatiles of *P. massoniana*

P. massoniana was used as the representative pine for investigating transformation-stimulating signals. After cellulose and nutrients, volatiles are the most important components of pines. The promotive capability of volatiles from *P. massoniana* on DL4 transformation was first determined. The percentage of nematodes in the water section of the dish attracted by pine chips was 84% (Fig. 4b), while it was only 23% with *G. biloba* (Fig. 4a). Independent of the section, water or agarose, when nematodes were exposed to pine volatiles, the DL4 transformation rates were ~80%. When nematodes were exposed to ginkgo volatiles from *G. biloba* chips, the transformation rates were only




Fig. 4. Distribution and transformation rates of *B. xylophilus* DL4 promoted by the volatiles from *P. massoniana* or *G. biloba*. (a) DL4 nematode distributions in each part of divided Petri dishes with *G. biloba* chips attraction. In the test diagram the left and right parts contained agarose and water, respectively. An empty 20-mm dish with *G. biloba* chips was adhered to the right side. The distribution of nematodes on agarose and in water are displayed in pie graphs. (b) DL4 nematode distributions in each part of divided Petri dishes with pine chips attraction. A 20-mm dish with pine chips was adhered to the right side. (c) Transformation rate of *B. xylophilus* DL4 promoted by the volatilization of chemicals from *P. massoniana* or *G. biloba*. Transformation rates of DL4 on agarose and in water were calculated with *P. massoniana* or *G. biloba*. Statistical differences in the means are indicated with different letters, $P < 0.05$. Error bars represent \pm s.e.

~20% (Fig. 4c, one-way ANOVA, $F_{3,8} = 123.98$, $P < 0.001$). To eliminate the effect of agarose on DL4 recovery, we evaluated the between-subject effects on the transformation rates promoted by pine chips or agarose (Table S1). The main effect on DL4 recovery was attributed to pine chips ($P < 0.001$), while agarose had no effect ($P > 0.05$). There was no interaction between pine chips and agarose ($P > 0.05$). In conclusion, the volatiles from pine promoted the DL4 transformation into an adult.

Analysis of main volatiles of *P. massoniana*

The main volatiles of *P. massoniana* were α -pinene, camphene, β -pinene, β -myrcene, β -phellandrene, longifolene and *trans*-caryophyllene. Among the main volatiles, the content of α -pinene was highest, followed by β -phellandrene and β -pinene. The contents of other volatiles were no more than 10%. While, the main volatiles of *P. thunbergii* were α -pinene, camphene, β -pinene, β -myrcene, D-limonene and *trans*-caryophyllene. Among the main volatiles, the content of α -pinene was highest, followed by D-limonene and β -pinene. Concentrations of all the volatile components from *P. thunbergii* were higher than those

of *P. massoniana*. In addition, *G. biloba* did not contain any pine volatiles (Table 1).

B. xylophilus DL4 recovery promoted by standard pine volatiles

Authentic standards of seven main volatiles in tested pines were used to investigate their promotive effects on DL4 transformation. After 2 d, the transformation rates of DL4 stimulated by β -myrcene and β -pinene were highest, followed by D-limonene, and the remain chemicals had limited effects on DL4 transformation (Fig. 5a, one-way ANOVA, $F_{12,26} = 127.05$, $P < 0.001$), suggesting pine chips have a stronger promotive ability than that of the standard chemicals (Fig. 1a). After 4 d, β -myrcene and β -pinene were still the most effective promoters for DL4 transformation (Fig. 5a, one-way ANOVA, $F_{12,26} = 158.93$, $P < 0.001$), and no significant effects were found for the longifolene and *trans*-caryophyllene. β -Pinene was chosen for the effect of gradient chemical concentration on DL4 transformation. After 2 d, the transformation rates of DL4 increased correspondingly with the enhanced β -pinene concentrations (Fig. 5b), and all the tested β -pinene concentrations promoted DL4 transformation (Fig. 5b,

Table 1. Quantification of the main volatiles of tested trees

Compounds	Peak time (Min)	Content in different species ($\mu\text{g g}^{-1}$ wet weight)		
		<i>G. biloba</i>	<i>P. massoniana</i>	<i>P. thunbergii</i>
α -Pinene	9.71	0.00	907.54 \pm 52.11	5451.09 \pm 91.83
Camphene	10.14	0.00	16.60 \pm 6.54	132.04 \pm 2.97
β -Pinene	10.91	0.00	171.69 \pm 37.50	441.71 \pm 132.65
β -Myrcene	11.21	0.00	5.81 \pm 3.36	99.69 \pm 9.56
D-limonene	12.38	0.00	0.00	537.16 \pm 18.40
β -Phellandrene	12.4	0.00	148.49 \pm 65.04	0.00
Longifolene	23.16	0.00	43.31 \pm 9.00	0.00
<i>trans</i> -Caryophyllene	23.43	0.00	65.58 \pm 5.62	246.13 \pm 29.95

The value indicates mean \pm s.e.




Fig. 5. Transformation rate of *B. xylophilus* DL4 promoted by the standard volatiles detected in tested pine trees. (a) Transformation rates of *B. xylophilus* DL4 promoted by standard volatiles detected in *P. massoniana* after 2 and 4 d. (b) Transformation rates of *B. xylophilus* DL4 promoted by different concentrations of β -pinene after 2 and 4 d. Statistical differences in the means are indicated with different letters, $P < 0.05$. Error bars represent \pm s.e.

one-way ANOVA, $F_{5,12} = 130.51$, $P < 0.001$) though the chemical had little effect on the transformation when its concentration is less than 10^{-3} g mL $^{-1}$ (Fig. 5b, one-way ANOVA, $F_{5,12} = 77.49$, $P < 0.001$).

Discussion

Dauer recovery is necessary for nematodes to complete their life cycles (Dolan *et al.*, 2003; Murgatroyd and Spengler, 2010; Zhao *et al.*, 2013). For *B. xylophilus*, we found that pine volatiles could promote dauer recovery from DL4 to adult. The

determinant factors of dauer recovery for many other nematodes are population density, food supply and ambient temperature (Dolan *et al.*, 2003; Fielenbach and Antebi, 2008). However, it was not clear whether these three factors promote the DL4 recovery of *B. xylophilus*. In *C. elegans*, its dauer will recover under 27 °C (Fielenbach and Antebi, 2008). However, *B. xylophilus* dauer is not sensitive to temperature. In our experiments, we tested nine temperatures, ranging from 4 to 45 °C to assay their effects on *B. xylophilus* dauer recovery. Some specific temperatures (20, 25 and 30 °C) could promote dauer recovery slightly. The transformation rates were less than 40% in the control group, but the

transformation rates of nematodes rearing at 20, 25 and 30 °C were higher than those in the control, which increased with the rising temperature. Moreover, extreme temperatures of 4, 35 and 40 °C inhibited DL4 transformation even with pine chips. Independent of the population density, the transformation rates of DL4 were very low without pine chips. This is corroborated in the lack of density-dependent effects on the dauer recovery of Entomopathogenic nematodes *Steinernema carpocapsae* and *Steinernema feltiae* (Hirao and Ehlers, 2010). However, with pine chips, the transformation rates of nematodes, no matter at low density or high density, were very high. From the above, we thus deduct that the pine chips play important roles for *B. xylophilus* DL4 recovery, but temperature and density were not.

Although pine chips were necessary for DL4 recovery, DL4 do not feed during the process of leaving the beetle trachea and entering the pines (Van Gundy, 1967; Storey, 1984; Zhao *et al.*, 2013). The main energy reserves of DL4s are neutral lipids (Stamps and Linit, 1995). These lipids are converted into energy or undergo histogenesis into digestive and reproductive organs during dauer recovery to adult. Here, we found that, unlike pine volatiles, nutrition had no effect on DL4 transformation (Figs 2 and 3). Pine volatiles, such as β -myrcene and β -pinene, promoted DL4 transformation (Fig. 5). In addition, they also attracted DL4 (Fig. 4) (Stamps and Linit, 1998b; Linit and Stamps, 2001). Nematodes are very sensitive to chemicals and *B. xylophilus* DL4 appears to respond to a variety of chemical cues to leave the trachea of vector beetles (Futai, 2013).

When the vector beetle feeds on pine trees, the volatile concentrations from the pine increase rapidly (Su *et al.*, 2008; Niu *et al.*, 2012; Zhao *et al.*, 2014; Chen *et al.*, 2018). The accumulation of volatiles in response to herbivore or pathogen attack is an important component of host defence (Lewinsohn *et al.*, 1991; Keeling and Bohlmann, 2006; Hansen *et al.*, 2017). However, *B. xylophilus* might take advantage of these volatile accumulations for its survival (Figs 4 and 5). Pine volatiles could attract DL4 and promote DL4 transformation in a gradient-dependent manner. After a 2-d exposure to β -pinene, the transformation rates increased with the rising concentration of β -pinene. In our experiments, we tested eight volatiles released from *P. massoniana* or *P. thunbergii*. Different terpenes had varied promotive effects. Among them, the monoterpenes β -myrcene and β -pinene were the most effective stimulators of DL4 transformation, following closely by D-limonene, α -pinene, camphene and β -phellandrene. However, the sesquiterpenes longifolene and *trans*-caryophyllene had no effects on DL4 transformation (Fig. 5). Since the results cannot fully explain the different promotive effects of terpenes to *B. xylophilus* DL4 transformation and the sample size is small, more studies are needed to further explore and confirm the molecular mechanisms of the transformation.

Although pine volatiles have been shown to promote *B. xylophilus* DL4 transformation (Fig. 5), a few nematodes were capable of transforming to the propagative stage without any stimulators regardless of the low transformation rate (see controls in all Figures). Thus, other factors may initiate the DL4 transformation. For example, the amounts of neutral storage lipids in DL4 may act as an internal clock that influences the decision to remain in the body of the vector beetle or enter the pine host (Stamps and Linit, 1995; Stamps and Linit, 1998b; Linit and Stamps, 2001). Additionally, different carbon dioxide concentrations from trachea produced by the beetle's breathing may attract or repel DL4 (Maehara and Futai, 2001). These signals might also be the stimulators of *B. xylophilus* DL4 transformation. Our study identified a new dauer recovery signal of nematodes and this may contribute to preventing dauer recovery, which would aid in decreasing the incidence of pine wilt disease.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182019001264>.

Financial support. This study was funded by Fundamental Research Funds of Research Institute of Forest New Technology, CAF (CAFYBB2018SY037), National Key Research and Development Program of China (2017YFD0600100) and the National Natural Science Foundation of China (NSFC 31901315).

Conflict of interest. None.

Ethical standards. Not applicable.

References

- Aumann J and Ehlers R-U (2001) Physico-chemical properties and mode of action of a signal from the symbiotic bacterium *Photorhabdus luminescens* inducing dauer juvenile recovery in the entomopathogenic nematode *Heterorhabditis bacteriophora*. *Nematology* **3**, 849–853.
- Chen R, He X, Chen J, Gu T, Liu P, Xu T, Teale SA and Hao D (2018) Traumatic resin duct development, terpenoid formation, and related synthase gene expression in *Pinus massoniana* under feeding pressure of *Monochamus alternatus*. *Journal of Plant Growth Regulation*, 1–12.
- Cheng H, Lin M, Li W and Fang Z (1983) The occurrence of a pine wilting disease caused by a nematode found in Nanjing. *Forest Pest and Disease* **4**, 1–5.
- Dolan KM, Jones JT and Burnell AM (2003) Detection of changes occurring during recovery from the dauer stage in *Heterorhabditis bacteriophora*. *Parasitology* **125**, 71–81.
- Fan J, Sun J and Shi J (2007) Attraction of the Japanese pine sawyer, *Monochamus alternatus*, to volatiles from stressed host in China. *Annals of Forest Science* **64**, 67–71.
- Fielenbach N and Antebi A (2008) *C. elegans* dauer formation and the molecular basis of plasticity. *Genes & Development* **22**, 2149–2165.
- Fukushige H (1991) Propagation of *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae) on Fungi Growing in Pine-Shoot Segments. *Applied Entomology and Zoology* **26**, 371–376.
- Futai K (2013) Pine wood nematode, *Bursaphelenchus xylophilus*. *Annual Review of Phytopathology* **51**, 61–83.
- Golden J and Riddle D (1982) A pheromone influences larval development in the nematode *Caenorhabditis elegans*. *Science* **218**, 578–580.
- Golden JW and Riddle DL (1984) The *Caenorhabditis elegans* dauer larva – developmental effects of pheromone, food, and temperature. *Developmental Biology* **102**, 368–378.
- Hansen SC, Stolter C, Imholt C and Jacob J (2017) Like or dislike: response of rodents to the odor of plant secondary metabolites. *Integrative Zoology* **12**, 428–436.
- Hinode Y, Shuto Y and Watanabe H (1987) Stimulating effects of β -myrcene on molting and multiplication of the pine wood nematode, *Bursaphelenchus xylophilus*. *Agricultural and Biological Chemistry* **51**, 1393–1396.
- Hirao A and Ehlers RU (2010) Influence of inoculum density on population dynamics and dauer juvenile yields in liquid culture of biocontrol nematodes *Steinernema carpocapsae* and *S. feltiae* (nematoda: Rhabditida). *Applied Microbiology and Biotechnology* **85**, 507–515.
- Keeling CI and Bohlmann J (2006) Genes, enzymes and chemicals of terpenoid diversity in the constitutive and induced defence of conifers against insects and pathogens. *New Phytologist* **170**, 657–675.
- Lewinsohn E, Gijzen M and Croteau R (1991) Defense mechanisms of conifers: differences in constitutive and wound-induced monoterpene biosynthesis among species. *Plant Physiology* **96**, 44–49.
- Linit M and Stamps WT (2001) Interaction of intrinsic and extrinsic chemical cues in the behaviour of *Bursaphelenchus xylophilus* (Aphelenchida: Aphelenchoididae) in relation to its beetle vectors. *Nematology* **3**, 295.
- Maehara N and Futai K (2000) Population changes of the pinewood nematode, *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae), on fungi growing in pine-branch segments. *Applied Entomology and Zoology* **35**, 413–417.
- Maehara N and Futai K (2001) Presence of the cerambycid beetles *Psacotha hilaris* and *Monochamus alternatus* affecting the life cycle strategy of *Bursaphelenchus xylophilus*. *Nematology* **3**, 455.
- Mamiya Y (1975) The life history of the pine wood nematode, *Bursaphelenchus lignicolus*. *Japanese Journal of Nematology* **5**, 16–25.
- Mamiya Y (1983a) The effect of wood tissues on the molting rate of the dauer-larvae of *Bursaphelenchus xylophilus*. *Japanese Journal of Nematology* **13**, 6–13.

- Mamiya Y (1983b) Pathology of the pine wilt disease caused by *Bursaphelenchus xylophilus*. *Annual Review of Phytopathology* **21**, 201–220.
- Mota MM, Braasch H, Bravo MA, Penas AC, Burgermeister W, Metge K and Sousa E (1999) First report of *Bursaphelenchus xylophilus* in Portugal and in Europe. *Nematology* **1**, 727–734.
- Murgatroyd C and Spengler D (2010) Histone tales: echoes from the past, prospects for the future. *Genome Biology* **11**, 105.
- Nickle W, Golden A, Mamiya Y and Wergin W (1981) On the taxonomy and morphology of the pine wood nematode, *Bursaphelenchus xylophilus* (Steiner & Buhner 1934) Nickle 1970. *Journal of Nematology* **13**, 385.
- Niu H, Zhao L, Lu M, Zhang S and Sun J (2012) The ratio and concentration of two monoterpenes mediate fecundity of the pinewood nematode and growth of its associated fungi. *PLoS One* **7**, e31716.
- Ouellet J, Li S and Roy R (2008) Notch signalling is required for both dauer maintenance and recovery in *C. elegans*. *Development* **135**, 2583–2592.
- Smith R (2000) Xylem monoterpenes of pines: distribution, variation, genetics, function. *USDA Forest Service General Technical Report* **177**, 454.
- Stamps WT and Linit MJ (1995) A rapid and simple method for staining lipid in fixed nematodes. *Journal of Nematology* **27**, 244–248.
- Stamps WT and Linit MJ (1998a) Chemotactic response of propagative and dispersal forms of the pinewood nematode *Bursaphelenchus xylophilus* to beetle and pine derived compounds. *Fundamental and Applied Nematology* **21**, 243–250.
- Stamps WT and Linit MJ (1998b) Neutral storage lipid and exit behavior of *Bursaphelenchus xylophilus* fourth-stage dispersal juveniles from their beetle vectors. *Journal of Nematology* **30**, 255–261.
- Storey RMJ (1984) The relationship between neutral lipid reserves and infectivity for hatched and dormant juveniles of Globodera spp. *Annals of Applied Biology* **104**, 511–520.
- Strauch O and Ehlers RU (1998) Food signal production of *Photorhabdus luminescens* inducing the recovery of entomopathogenic nematodes *Heterorhabditis* spp. in liquid culture. *Applied Microbiology and Biotechnology* **50**, 369–374.
- Su J-W, Zeng J-P, Qin X-W and Ge F (2008) Effect of needle damage on the release rate of Masson pine (*Pinus massoniana*) volatiles. *Journal of Plant Research* **122**, 193.
- Van Gundy SD (1967) Aging and starvation in larvae of *Meloidogyne javanica* and *Tylenchulus semipenetrans*. *Phytopathology* **57**, 559–571.
- Xu L, Lou Q, Cheng C, Lu M and Sun J (2015) Gut-associated bacteria of *Dendroctonus valens* and their involvement in verbenone production. *Microbial Ecology* **70**, 1012–1023.
- Xu L, Shi Z, Wang B, Lu M and Sun J (2016) Pine defensive monoterpene alpha-pinene influences the feeding behavior of *Dendroctonus valens* and its gut bacterial community structure. *International journal of molecular sciences* **17**, 1734.
- Xu D, Xu L, Zhou F, Wang B, Wang S, Lu M and Sun J (2018) Gut bacterial communities of *Dendroctonus valens* and monoterpenes and carbohydrates of *Pinus tabulaeformis* at different attack densities to host pines. *Frontiers in Microbiology* **9**, 1251.
- Yi CK, Byun BH, Park JD, Yang S and Chang KH (1989) First finding of the pine wood nematode, *Bursaphelenchus xylophilus* (Steiner et Buhner) Nickle and its insect vector in Korea. *Research Reports of the Forestry Research Institute (Seoul)* **38**, 141–149.
- Zamora P, Rodriguez V, Renedo F, Sanz AV, Dominguez JC, Perez-Escobar G, Miranda J, Alvarez B, Gonzalez-Casas A, Mayor E, Duenas M, Miravalles A, Navas A, Robertson L and Martin AB (2015) First report of *Bursaphelenchus xylophilus* causing pine wilt disease on *Pinus radiata* in Spain. *Plant Disease* **99**, 1449–1449.
- Zhao LL, Wei W, Kang L and Sun JH (2007) Chemotaxis of the pinewood nematode, *Bursaphelenchus xylophilus*, to volatiles associated with host pine, *Pinus massoniana*, and its vector *Monochamus alternatus*. *Journal of Chemical Ecology* **33**, 1207–1216.
- Zhao LL, Zhang S, Wei W, Hao H, Zhang B, Butcher RA and Sun J (2013) Chemical signals synchronize the life cycles of a plant-parasitic nematode and its vector beetle. *Current Biology* **23**, 2038–2043.
- Zhao L, Mota M, Vieira P, Butcher RA and Sun J (2014) Interspecific communication between pinewood nematode, its insect vector, and associated microbes. *Trends in Parasitology* **30**, 299–308.
- Zhao L, Zhang X, Wei Y, Zhou J, Zhang W, Qin P, Chinta S, Kong X, Liu Y, Yu H, Hu S, Zou Z, Butcher RA and Sun J (2016) Ascarosides coordinate the dispersal of a plant-parasitic nematode with the metamorphosis of its vector beetle. *Nature Communications* **7**, 12341.
- Zhou F, Xu L, Wang S, Wang B, Lou Q, Lu M and Sun J (2017) Bacterial volatile ammonia regulates the consumption sequence of d-pinitol and d-glucose in a fungus associated with an invasive bark beetle. *The ISME Journal* **11**, 2809.