

# Factors affecting the component community structure of haemoparasites in bank voles (*Clethrionomys glareolus*) from the Mazury Lake District region of Poland

A. BAJER<sup>1</sup>, A. PAWEŁCZYK<sup>1</sup>, J. M. BEHNKE<sup>2\*</sup>, F. S. GILBERT<sup>2</sup> and E. SINISKI<sup>1</sup>

<sup>1</sup>Department of Parasitology, Institute of Zoology, University of Warszawa, ul. Krakowskie Przedmieście 26/28, 00-927 Warszawa, Poland

<sup>2</sup>School of Biological Sciences, University Park, University of Nottingham, Nottingham NG7 2RD, UK

(Received 22 March 2000; revised 19 July 2000; accepted 30 July 2000)

## SUMMARY

The prevalence and abundance of infections with haemoparasites were studied over a 3 year period in *Clethrionomys glareolus* (bank vole,  $n = 420$ ) sampled from forests in the NE of Poland. Total species richness was 5 (Prevalence = *Haemobartonella* sp. 63.1%, *Bartonella grahamii* 27.4%, *Hepatozoon erhardovae* 31.4%, *Trypanosoma evotomys* 15% and *Babesia microti* 1.0%) with 81.9% of the voles carrying at least 1 species and a mean infracommunity species richness of 1.4. Variation in species richness was determined primarily by season and year, and the interaction of these factors. The observed frequency distribution of infracommunity species richness did not differ from that predicted by a null model, suggesting that there were no marked associations between the species. Analyses of prevalence and abundance of infection with each species in turn, revealed that overall the principal causes of variation were temporal and seasonal and their interaction, intrinsic factors such as age and sex playing only a minor role. However, the relative importance of specific extrinsic, and rarely intrinsic, factors varied and was distinct for each of the species in the study. Prevalence data revealed 4 sets of 2-way associations between species, mostly varyingly dependent on combinations of intrinsic and extrinsic factors. Analysis of quantitative associations suggested 4 sets of positive 2-way interactions, 3 of which remained after controlling for the effect of extrinsic and intrinsic factors on the abundance of each species, but only one could be unequivocally accepted (*Haemobartonella* sp. + *B. grahamii*) after correction for multiple comparisons. These data are discussed in the context of the changing ecological profiles in this region of Eastern Europe and, in a wider context, in relation to current understanding of the factors which shape component community structures of haemoparasites in wild rodents.

Key words: bank vole, *Clethrionomys glareolus*, *Babesia microti*, *Bartonella grahamii*, *Haemobartonella*, *Hepatozoon erhardovae*, *Trypanosoma evotomys*, component community, haemoparasites, co-occurrence.

## INTRODUCTION

Published studies on the component communities of haemoparasites in European wild rodents are mostly based on data collected in western, northern and southern Europe (Healing, 1981; Turner, 1986; Santos-Gomez *et al.* 1993; Wiger, 1979). There is comparatively little information on naturally occurring haemoparasites of wild rodents from eastern Europe (see Baker, Chitty & Phipps, 1963; Sebek, 1978; Sebek, Rosicky & Sixl, 1977; Walter & Liebisch, 1980; Karbowski & Sinski, 1996*a, b*; Karbowski *et al.* 1999), and in a broader context few studies have comprehensively evaluated the contribution of intrinsic and extrinsic factors in structuring these communities.

As part of a long-term study of parasite communities in wild rodents in the northeastern region of Poland, we collected data on blood parasites,

including both protozoan and bacterial infections, over a period of 3 years. The Mazury lake district region of Poland is of interest because many of the collective farms, which comprised intensively cultivated and enriched agricultural land, were abandoned following political changes in the early 1990s. Much of this land is now reverting to nature, and natural succession has given rise to extensive, naturally seeded, young woodland. These ecological changes in the region are being documented (Rogozinska, 1997) and studied from various perspectives, not least regarding disease implications for local communities and visitors from urban centres. There are important consequences for human health from the increasing wild rodent populations because Lyme disease is present and increasing numbers of tourists are visiting the region for recreational purposes (Sinski & Pawelczyk, 1999).

Over a 3-year period, and in specific seasons, we systematically sampled rodent populations in a study site that covered both abandoned farmland and the surrounding forests. In this paper we report on the blood parasite communities of the bank vole (*Clethrionomys glareolus*), which is numerically the

\* Corresponding author: School of Biological Sciences, University Park, University of Nottingham, Nottingham NG7 2RD, UK. Tel: +0115 951 3208. Fax: +0115 951 3252. E-mail: jerzy.behnke@nottingham.ac.uk

dominant rodent species inhabiting the forest part of our study site. We evaluate the role of intrinsic (host age and sex) and extrinsic (year and season) factors in structuring the component community of haemo-parasites and in influencing both prevalence and abundance of each of the species involved. Finally, we determine the extent to which co-occurring species interact with each other.

## MATERIALS AND METHODS

### Study sites

Our study site was located east of the nature reserve surrounding Lake Łuknajno, and north of Lake Śniardwy, near the town of Mikołajki in the Mazury lake district region of NE Poland. Trapping was conducted in mature woodland with Scots pine (*Pinus sylvestris*), oak (*Quercus robur*), Norway spruce (*Picea abies*), silver birch (*Betula verucosa*) and common alder (*Alnus glutinosa*) as the dominant trees, forming a high canopy over the entire site. The dominant undergrowth was hazel (*Corylus avellana*) and ground cover comprised mostly *Oxalis acetosella*, *Convallaria mayalis* and *Stellaria holostea*.

### Collection of mice

Rodents were caught live in locally constructed wooden traps, with a small metal platform internally which, when triggered, released a metal door. These were set out at 15 m intervals in parallel lines, 10 m either side of a track running through the woodland site. Trapping sessions comprised a minimum of 4 days, with traps being inspected in the early morning and just before dusk. Fresh traps replaced any traps containing animals and the animals were brought to the University of Warszawa's field station at Urwitalt. Visits to the study sites varied, but generally comprised at least a 4 day duration, at approximately 4 week intervals from late March until mid-November. For both practical and animal welfare reasons, it was not possible to visit the sites in the period between November and March because the ground was mostly hard frozen and deep snow covered our sites for most of the winter period. The months between March and November were divided into 3 seasons, comprising spring (March to early June), summer (late June to August) and autumn (September to November).

### Sampling of hosts

At the field station in Urwitalt, all animals were inspected, identified, sexed, relevant morphometric data were recorded and they were weighed (to the nearest 0.5 g). Ectoparasites were removed from the ears and limbs (mostly larval and nymph ticks and some mites) and the fur was inspected carefully for

fleas. Thin blood smears were prepared from drops of blood taken from the tail vein. After inspection most animals were marked and released as near as possible to the original site of capture, whilst others were killed for recovery of endoparasites (data to be published elsewhere).

Blood smears were air-dried, fixed in absolute methanol, stained for 45 min in Giemsa's stain (diluted 1:3) in buffer at pH 7.2 and were brought back to the Department of Parasitology at the University of Warszawa for examination. Each smear was examined under oil immersion (using an Olympus AX70 microscope). Initially, sufficient fields of vision were examined to enable up to 50 leukocytes to be inspected for the presence of *H. erhardovae* (approximately 200 fields of vision under  $\times 100$  objective lens). Each field of vision was also examined for the presence of other species, although these were not quantified at this stage. If the blood smear revealed the presence of other species, during this initial phase of examination, an additional 1000 erythrocytes were inspected and the number of cells infected with *B. grahamii*, *B. microti* and *Haemobartonella* sp. were recorded and expressed as no. of infected cells/100 erythrocytes. The concentration of trypanosomes was also expressed per 100 erythrocytes.

Age classes were established on the basis of weight (Morris, 1971) and sexual development as described by Mazurkiewicz (1972). Age class 1 comprised immature, juvenile voles < 15 g in weight and approximately less than 1.5 months old. Age class 2 comprised young mature voles, 15–19.5 g in weight representing animals approximately 1.5–2.5 months old. Age class 3 comprised adult voles, weighing > 19.5 g and 2.5 months and older.

### Statistical analysis

The frequency distribution of infracommunity species richness was tested for goodness of fit to the positive binomial distribution (assumption of the null model is a regular distribution), the Poisson distribution (assumption of the null model is a random distribution) and to the null model of Janovy *et al.* (1995) (assumption of the null model is that, in the absence of associations and interactions between species, the frequency distribution of infracommunity species richness is predicted by prevalence values). Goodness of fit in each case was tested by  $\chi^2$ .

Prevalence (percentage of animals infected) was analysed by maximum likelihood techniques based on log linear analysis of contingency tables, implemented by the software package, Statgraphics Version 7. For each species in turn we entered prevalence of infection as a binary factor (infected = 1, not infected = 0) and then year (3 levels, 1997, 1998 and 1999), season (3 levels, spring, summer and autumn),

host age (3 levels) and host sex (2 levels) as factors. Beginning with the most complex model, involving all possible main effects and interactions, those combinations not contributing significantly to explaining variation in the data were eliminated stepwise, beginning with the highest-level interaction. A minimum sufficient model was then obtained, for which the likelihood ratio of  $\chi^2$  was not significant, indicating that the model was sufficient in explaining the data.

Quantitative data reflecting parasite abundance within hosts were expressed as geometric means because the data were highly overdispersed (Elliott, 1977; Dash, Hall & Barger, 1988). In some cases arithmetic means and standard errors of the means are also provided. These means reflect the abundance of infection as defined by Margolis *et al.* (1982) and include all subjects within the specified group, infected and not infected, for which relevant data were available. The degree of aggregation in quantitative data was calculated by the Index of Dispersion ( $I$  = the variance to mean ratio) and the Index of Discrepancy ( $D$ ) as described by Poulin (1993) (a value of 0 indicates an even distribution of counts across all hosts and a value of 1 indicates all parasites aggregated in a single host). Frequency distributions of individual species were also tested for goodness of fit to negative binomial, positive binomial and Poisson models by  $\chi^2$  as described by Elliott (1977) and the negative binomial exponent  $k$  is given as appropriate.

Parasite abundance was analysed by GLIM (A statistical system for generalized linear interactive modeling; GLIM 4, PC version, Royal Statistical Society 1993) as described previously, using models with normal errors after normalization of the data by  $\log_{10}(x+1)$  transformation (Crawley, 1993; Wilson & Grenfell, 1997; Behnke *et al.* 1999). Year, season, host age and host sex (see above for levels) were entered as factors. For models with normal errors the change in deviance is divided by the scale parameter and the result divided by the change in degrees of freedom D.F. following each deletion, to give a variance ratio,  $F$ . Complex models are given in full ANOVA tables, but simpler models are summarized in the text and only  $F$  values are presented. The residuals from all models were checked for approximately normal distribution.

Quantitative associations between parasites were examined by multiple correlation analysis (Spearman rank-order correlation test) of raw parasite data from animals carrying both of 2 species, in each of the possible 2-way combinations (excluding *B. microti* because of very low prevalence of this species). To avoid the risk of Type 1 errors we adopted the Dunn–Sidak correction, lowering the cut-off value of  $P$ , according to the number of comparisons implemented (Sokal & Rohlf, 1981). We also examined the correlations between the

residuals from minimum sufficient 3, 2 or 1-way ANOVAs, as appropriate for individual species, in order to control for established differences between the years, seasons, host age and sex.

## RESULTS

### *Clethrionomys glareolus*

A total of 420 *C. glareolus* were sampled in the 3 year period. The structure of the sampled host population by year, host sex and age is summarized in Table 1. Host density was estimated to enable comparison of vole populations between the 3 years of the study. Vole numbers were expressed as the number of trapped voles divided by the product of the number of traps set and the duration of trapping hours (Table 2). In agreement with other studies (Alibhai & Gipps, 1985) we observed the same seasonal pattern each year with vole density beginning low in spring (May) and then rising to peak in the autumn (September and November). However, despite some differences between the years and the trapping sessions in which a proportion of voles was culled, overall the population of *C. glareolus* remained relatively stable across the 3 years of our study.

### Total species richness in the component community

Five species of blood parasites were recorded (Table 3) and 344 voles (81.9%) carried at least 1 of these species. *Haemobartonella* sp. was the most common and *B. microti* the rarest.

### Frequency distribution of infracommunity species richness

The observed distribution of infracommunity species richness, illustrated in Fig. 1, showed a good fit to the positive binomial ( $\chi^2 = 0.49$ , 4 D.F.,  $P = 0.972$ ) but not to the Poisson distribution ( $\chi^2 = 21.8$ , 4 D.F.,  $P = 0.0004$ ). It was not significantly different from that predicted by the null model for interactions of parasite species in an assemblage (Janovy *et al.* 1995) and  $\chi^2 = 6.14$  (4 D.F.,  $P > 0.05$ ).

### Mean infracommunity species richness

The overall mean number of species of blood parasites harboured per host was  $1.4 \pm 0.05$  (S.E.M.), with a variance to mean ratio of 0.665 indicating a positive binomial distribution. Analysis of these data (4-way ANOVA in GLIM with normal errors and year, season, host age and sex as factors, and Fig. 2) revealed that only year and season played a significant role in determining infracommunity species richness. The main effect of season was highly significant ( $F_{2,409} = 15.039$ ,  $P < 0.001$ ) and that of year was marginal ( $F_{2,409} = 3.014$ ,  $0.05 > P > 0.025$ )

Table 1. The structure of the sampled host population by year of capture, host sex and age

Year	Sex	Age			Total by sex	Total by year
		1	2	3		
1997	Male	26	47	17	90	165
	Female	20	31	24	75	
	Combined	46	78	41		
1998	Male	18	25	21	64	117
	Female	23	9	21	53	
	Combined	41	34	42		
1999	Male	20	32	12	64	138
	Female	25	32	17	74	
	Combined	45	64	29		
Total by age		132	176	112		

Table 2. Estimation of population density of voles in the study site

Month	Relative population density*		
	1997	1998	1999
March	15.6	N.D.	N.D.
April	8.3	10.2	7.6
May	4.0	1.1	4.1
June	7.2	14.4	N.D.
July	18.0	13.4	52.3
August	31.5	34.1	85.0
September	34.5	44.5	188.3
October	20.0	33.5	N.D.
November	35.4	N.D.	N.D.

\* Population density was calculated as the number of voles trapped divided by trap hour  $\times 10^4$ . N.D., Not done.

Table 3. The overall prevalence of infection with haemoparasites in male and female voles

Species	Males (n = 218)	Females (n = 202)	Combined
<i>Haemobartonella</i> sp.	65.1	60.9	63.1
<i>Bartonella grahamii</i>	23.9	31.2	27.4
<i>Hepatozoon erhardovae</i>	31.2	31.7	31.4
<i>Trypanosoma evotomys</i>	15.1	14.9	15.0
<i>Babesia microti</i>	1.4	0.5	1.0

but there was also a strong significant interaction between these factors ( $F_{4,393} = 12.090$ ,  $P < 0.001$ ). Fig. 2 shows that mean species richness increased between spring and summer 1997 and then fell in the autumn. In 1998 mean species richness again began low and rose to peak in the summer, but this time stayed comparatively high in the autumn. In the final year of the study, mean species richness began




Fig. 1. Frequency distribution of infracommunity species richness. The observed data are in the filled columns and those predicted by the null model of Janovy *et al.* (1995) in the open columns. See text for full explanation and statistical analysis.




Fig. 2. Annual and seasonal changes in mean infracommunity species richness. See text for statistical analysis.

higher in the spring than in the 2 earlier years and there was little further change. Thus there were season-dependent changes in mean species richness, but the pattern of seasonal changes varied from one year to the next.

#### Prevalence of species

We analysed the influence of year, season, host sex and age on prevalence of infection by maximum likelihood methods, testing respectively models for each of the 5 species and the resulting minimum sufficient models are summarized in Table 4. The age  $\times$  sex interaction arose in each case because there were significant differences in the distribution of the

Table 4. Minimum sufficient maximum likelihood statistical models of factors affecting the prevalence of infection

(For each species the interactions listed are significant and therefore form part of the final model.)

Species	Principal interactions and components in explaining variation in data*	Test of individual effects			Goodness of fit of min.suf. model		
		$\chi^2$	D.F.	$P\ddagger$	$\chi^2$	D.F.	$P\ddagger$
<i>Haemobartonella</i> sp.	Year $\times$ season $\times$ age $\times$ infection	20.96	8	0.0073	52.04	51	0.43
	Sex $\times$ age	6.65	2	0.0360			
<i>B. grahamii</i>	Year $\times$ season $\times$ infection	17.90	4	0.0013	69.33	69	0.46
	Year $\times$ season $\times$ age	32.34	8	0.0001			
	Sex $\times$ age	6.65	2	0.0360			
<i>H. erhardovae</i>	Year $\times$ season $\times$ sex $\times$ infection	10.98	4	0.0268	24.07	24	0.46
	Year $\times$ season $\times$ sex $\times$ age	20.58	8	0.0083			
	Sex $\times$ age $\times$ infection	6.13	2	0.0465			
	Season $\times$ age $\times$ infection	9.85	4	0.0431			
	Year $\times$ age $\times$ infection	17.20	4	0.0018			
<i>T. evotomys</i>	Year $\times$ season $\times$ age	32.34	8	0.0001	79.07	75	0.35
	Sex $\times$ age	6.65	2	0.0360			
	Season $\times$ infection	8.19	2	0.0166			
<i>B. microti</i>	Year $\times$ season $\times$ age	32.34	8	0.0001	52.06	77	0.98
	Sex $\times$ age	6.65	2	0.0360			
	Infection	537.05	1	< 0.0001			

\* Year – 3 levels (1997, 1998, 1999); season – 3 levels (spring, summer and autumn); age – 3 levels (age cohorts 1, 2 and 3); sex – 2 levels (male and female), infection – 2 levels (infected or uninfected).

† Probability that excluding the effect will make a significant change to the model.

‡ Probability that the data do not differ significantly from the minimum sufficient model described by the principal interactions and components.

two sexes of voles among age classes. Similarly the year  $\times$  season  $\times$  age interaction indicated significant differences in the distribution of age classes between seasons and years.

For *B. microti* none of the factors entered into the analyses influenced prevalence. *T. evotomys* was only affected by season and these data are summarized in Fig. 3C, where it can be seen that prevalence was at its lowest in the spring in all 3 years. Prevalence then rose to peak in the summer in 1997 and 1998, but continued to rise in 1999 to peak later, in the autumn.

Infections with *B. grahamii* also varied in relation to season but in addition there was a significant year effect. In Fig. 3A the similarity in the seasonal pattern in 1997 and 1998 is clearly apparent with low prevalence in spring, a peak in the summer and a drop by the autumn. In 1999 there was a contrasting pattern with little change in prevalence across all three seasons.

*Haemobartonella* sp. yielded a more complex model with year, season and age all contributing to variation in the data as shown in Fig. 4. There was no consistent pattern of prevalence across the age classes. In some years and seasons the youngest mice showed low prevalence and the older classes higher prevalence (e.g. spring 1999). In others, the pattern was reversed (e.g. autumn 1997) and again in other years and seasons, all the age classes showed similar

prevalence (e.g. summer 1999 and 1997). The only general conclusion to emerge from this analysis is that in 1998, prevalence was generally high across all seasons and age classes in comparison to other years.

*H. erhardovae* generated the most complex model, with all 4 factors contributing to interactions in various combinations. Essentially, prevalence varied among years and seasons, but also in relation to age and sex. We summarize the year  $\times$  seasonal effects only in Fig. 3B. The greatest discrepancy in prevalence, between the years, was in spring and quite clearly overall prevalence was highest in 1999.

#### Abundance of infection among infracommunities

The frequency distribution of abundance of infection with *B. grahamii* was not well described by any of the distributions tested, but was closest to the negative binomial distribution ( $k = 0.146 \pm 0.0175$ ,  $I = 63.0$  and  $D = 0.862$ ). We analysed these data further after normalization (by  $\log_{10}[x+1]$  transformation) by 4-way ANOVA in GLIM. This showed that the main effects of season and year were significant ( $F_{2, 409} = 3.422$ ,  $0.05 > P > 0.025$  and  $F_{2, 409} = 3.931$ ,  $0.025 > P > 0.01$ , respectively) but there was also a significant 2-way interaction between year  $\times$  age ( $F_{4, 399} = 3.019$ ,  $0.025 > P > 0.01$ ). From Fig. 5A it can be seen that in all 3 years mean abundance increased between spring and summer. In 1997 and




Fig. 3. Annual and seasonal changes in the prevalence of *Bartonella grahamii*, *Hepatozoon erhardovae* and *Trypanosoma evotomys*. See text for statistical analysis.

1998, but not in 1999, abundance then fell in the autumn.

*Haemobartonella* sp. also did not conform well to any of the distributions tested but again the discrepancy was least in the case of the negative binomial distribution ( $k = 1.199 \pm 0.169$ ,  $I = 2.7$  and  $D = 0.596$ ). Abundance was analysed further through a 4-way ANOVA in GLIM on  $\log_{10}(x+1)$  transformed data. This revealed no significant main effects but there were some highly significant 2- and 3-way interactions (Table 5). Therefore variation in the abundance of *Haemobartonella* sp. was compounded by all 4 factors in varying ways and no clear picture emerged with respect to seasonal and annual trends.

The frequency distribution of *H. erhardovae* gave a good fit to the negative binomial distribution only ( $\chi^2 = 8.7$ , D.F. = 7,  $P = \text{N.S.}$ ,  $k = 0.158 \pm 0.018$ ,  $I = 240.8$  and  $D = 0.861$ ). Analysis by 4-way ANOVA in




Fig. 4. Annual, seasonal and host age-related changes in the prevalence of *Haemobartonella* sp. (○) age class 1; (△) age class 2; (□) age class 3. See text for statistical analysis.

GLIM on  $\log_{10}(x+1)$  transformed data indicated that abundance varied significantly between years (main effect of year  $F_{2, 409} = 5.024$ ,  $P < 0.001$ ), but this was compounded by a year  $\times$  season interaction ( $F_{4, 393} = 3.428$ ,  $P < 0.01$ ). Fig. 5B shows that seasonal changes in abundance were very similar in 1997 and 1998, but quite different in 1999, when the spring cohort of voles carried an unusually high abundance of infection. Perhaps surprisingly, there was no significant main effect of season but there were 2 significant 2-way interactions (year  $\times$  age  $F_{4, 399}$ ,  $P = 0.05$ ; season  $\times$  sex  $F_{2, 401} = 4.185$ ,  $0.025 > P > 0.01$ ) indicating additional discrepancies in relation to host sex and age.

Abundance of *T. evotomys* infection conformed well to the negative binomial distribution ( $\chi^2 = 3.2$ , D.F. = 4,  $P = \text{N.S.}$ ;  $k = 0.0728 \pm 0.011$ ;  $I = 7.4$  and  $D = 0.9225$ ). These data were analysed by a 4-way ANOVA (GLIM with  $\log_{10}(x+1)$  transformed data) which revealed that only a single 2-way interaction was significant (year  $\times$  season,  $F_{4, 393} = 3.877$ ,  $P < 0.005$ ). Fig. 5C shows that in 1997 abundance peaked in the summer, whereas in 1998 and 1999 peak abundance was recorded in the autumn. Despite the relatively high peak abundance in the autumn of 1999, there was no significant main effect of year.

#### Associations between parasites based on categorical data

We tested the idea that any existing interactions between parasites should be reflected in co-occurrence or exclusion of species. In addition to data on the parasites (presence of infection/absence of infection) we also entered the 4 factors, year, season, host age and sex in a log-linear analysis, since these




Fig. 5. Annual and seasonal changes in the abundance of infection with *Bartonella grahamii*, *Hepatozoon erhardovae* and *Trypanosoma evotomys*. Data are the geometric mean abundance in (A) no. of cells/100 erythrocytes, (B) no. of infected cells/100 leucocytes and (C) no. of parasites/100 erythrocytes. See text for statistical analysis.

had been shown earlier to influence the prevalence of the parasites in various ways. We tested a model with 9 factors, and this is summarized in Table 6. Only 4 terms in this model incorporated 2 sets of parasites each. The association between *B. microti* and *B. grahamii* arose because 3 of the 4 *B. microti* infections were detected in voles that also carried *B. grahamii*, but this was the weakest term in the model. *H. erhardovae* and *T. evotomys* were also associated, (prevalence of *T. evotomys* in voles carrying *H. erhardovae* = 18.9%, compared with 13.1% among voles without *H. erhardovae*). However, this association was year and age dependent and was also a weak term in the model. A stronger association was detected between *H. erhardovae* and *Haemo-*

*bartonella* sp. (prevalence of *Haemobartonella* sp. = 67.4% among voles carrying *H. erhardovae* and 61.1% among those without), but again this was not a simple association and was dependent on season and age of the hosts. Finally, *B. grahamii* associated with *Haemobartonella* sp. (prevalence of *Haemobartonella* sp. = 70.4% in voles with *B. grahamii* and 60.3% in those without) in a host sex and age dependent manner.

#### *Interactions between species based on quantitative data*

Interactions between species should be reflected in quantitative associations in hosts carrying both species. We first carried out a multiple correlation analysis on abundance data of 4 of the species in the study (excluding infection with *B. microti* because only 4 animals carried this species) using data only from animals that carried both species of each pair (Table 7). Because of the possible risk of Type I errors, we implemented the Dunn–Sidak correction and accordingly this allowed acceptance of just one of the 6 possible associations as significant, i.e. that between *Haemobartonella* sp. and *B. grahamii* (Fig. 6). However, it is worth noting that 3 of the associations involving *T. evotomys* also suggest some underlying quantitative interaction between the species involved (with *Haemobartonella* sp., *B. grahamii* and *H. erhardovae*).

Some of these relationships could have arisen as a consequence of differences in parasite burdens attributable to year, season, host age and sex effects. Therefore, to control for the possible bias arising out of the factors known to affect abundance of parasites, we next carried out a multiple correlation analysis on the residuals of each species, following minimum sufficient ANOVAs as described in the preceding section. Three associations had high correlation coefficients, but because of the Dunn–Sidak correction, we could only accept the association between *Haemobartonella* sp. and *B. grahamii* as significant (Table 7 and Fig. 6).

In a further approach we carried out a principal components analysis on the standardized residuals from each of the 5 ANOVAs. This gave approximately similar amounts of variation on each axis (PCA1 = 30.1, PCA2 = 25.2, PCA3 = 23.4, PCA4 = 21.3%). Hence no further reduction of the data was possible.

#### DISCUSSION

This study constitutes the first comprehensive study of haemoparasites in a wild rodent population from north-east Europe. To our knowledge, apart from species identification (Sebek, 1978; Karbowski & Sinski, 1996a, b), no other studies have reported on the component community structure of haemoparasites from this region nor have probed the

Table 5. Statistical analysis of factors affecting abundance of infection ( $\log_{10}[x + 1]$  transformed data) with *Haemobartonella* sp. through a 4-way ANOVA with normal errors

(The full model deviance was 6.7144 with a scale parameter of 0.01825.)

Source of variation	Change in deviance*	Degrees of freedom	Scale parameter	Scaled deviance†	P
Sex	0.0353	1	0.0211	1.6703	N.S.
Age	0.1193	2	0.0213	2.8005	N.S.
Season	0.1069	2	0.1069	2.5129	N.S.
Year	0.0527	2	0.0211	1.2470	N.S.
Age × sex	0.0296	2	0.0211	0.7023	N.S.
Season × age	0.0366	4	0.0211	0.4329	N.S.
Season × sex	0.0259	2	0.0213	0.6079	N.S.
Year × age	0.7128	4	0.0213	8.3662	< 0.001
Year × sex	0.0003	2	0.0197	0.0073	N.S.
Year × season	0.3738	4	0.0198	4.7173	< 0.001
Year × season × age	0.2866	6	0.0191	2.5074	0.025 > P > 0.01
Season × sex × age	0.1225	4	0.0186	1.6465	N.S.
Year × sex × age	0.0752	4	0.0185	1.0168	N.S.
Year × season × sex	0.1885	4	0.0185	2.5514	0.05 > P > 0.025
Year × season × sex × age	0.0243	3	0.0182	0.4464	N.S.

\* Change in deviance following removal of combination specified in 'source of variation' column from the full factorial model. We begin by removing the 4-way interaction from the model. We then progressively remove the combinations in order from the base of the table towards the top. The main effect of year, was removed to assess the change in deviance but then replaced before removing the effect of season, and this was repeated for age and sex respectively.

† Scaled deviance = measure of contribution of factor specified under column labelled 'source of variation' to explaining variation in the data, calculated by fitting ANOVA with normal errors through GLIM. It is distributed as *F*.

n.s., Not significant.

Table 6. Minimum sufficient maximum likelihood statistical model incorporating all factors quantified and all haemoparasitic species detected in the study

(Each line in the upper compartment represents a significant interaction that therefore forms part of the final model.)

Principal interactions in explaining variation in data*			Test of individual effects		
Extrinsic factors	Intrinsic factors	Parasites	$\chi^2$	D.F.	P†
Year, season	Sex, age	<i>H. erhardovae</i>	16.42	8	0.0368
	Sex, age	<i>B. grahamii</i> , <i>Haemobartonella</i> sp.	7.52	2	0.0233
Season	Age	<i>H. erhardovae</i> , <i>Haemobartonella</i> sp.	10.90	4	0.0277
Year	Age	<i>H. erhardovae</i> , <i>T. evotomys</i>	9.97	4	0.0410
Year, season	Age	<i>Haemobartonella</i> sp.	20.06	8	0.0101
Year, season		<i>B. grahamii</i>	17.36	4	0.0016
Year, season		<i>T. evotomys</i>	9.69	4	0.0461
Year, season		<i>B. microti</i> , <i>B. grahamii</i>	3.96	1	0.0467
Goodness-of-fit of the minimum sufficient model specified by the interactions listed above			281.8	1535	1.0‡

\* Year – 3 levels (1997, 1998, 1999); season – 3 levels (spring, summer and autumn); age – 3 levels (age cohorts 1, 2 and 3); sex – 2 levels (male & female), infection – 2 levels (infected or uninfected).

† Probability that excluding the effect will make a significant change to the model.

‡ Probability that the data do not differ significantly from the minimum sufficient model described by the principal interactions and components.

factors that structure such communities. Moreover, even in studies based elsewhere, the dynamics of these infections, and their mutual interactions, have seldom been evaluated comprehensively (Healing, 1981; Turner & Cox, 1985) with quantifiable intrinsic and extrinsic factors taken into account simultaneously.

We recorded 5 species of haemoparasites in *C. glareolus*, all known to infect bank voles in other regions of Europe (Young, 1970; Baker, 1974; Turner, 1986; Walter & Liebisch, 1980). *Haemobartonella* sp. was clearly the dominant (adjudged by prevalence) and *B. microti* an incidental species. Whilst total species richness in the bank voles was 5,



Table 7. Analysis of quantitative associations between species

(Each test was carried out on voles carrying both species in the pair specified.)

Combination	$n^*$	Raw data		Residuals§	
		$r_s \dagger$	$P \ddagger$	$r_s$	$P$
<i>H. erhardovae</i> + <i>T. evotomys</i>	24	0.42	0.044	0.34	0.10
<i>H. erhardovae</i> + <i>B. grahamii</i>	36	-0.12	N.S.	< 0.01	N.S.
<i>H. erhardovae</i> + <i>Haemobartonella</i> sp.	88	-0.02	N.S.	0.11	N.S.
<i>T. evotomys</i> + <i>B. grahamii</i>	20	0.48	0.038	0.47	0.04
<i>T. evotomys</i> + <i>Haemobartonella</i> sp.	43	0.46	0.003	0.40	0.01
<i>B. grahamii</i> + <i>Haemobartonella</i> sp.	81	0.41	0.0002	0.38	0.0007

\* Number of voles carrying both species.

† Spearman's Correlation Coefficient.

‡ Probability (2-tailed). Applying the Dunn-Sidak correction for multiple comparisons, we accept  $P = 0.0085$  as the cut-off for rejecting the null hypothesis (Sokal & Rohlf, 1981).

§ Residuals from minimum sufficient ANOVAs in GLIM as explained in the text.

N.S., Not significant.




Fig. 6. Positive association between the abundance of infection of *Haemobartonella* sp. with *Bartonella grahamii*, and with *Trypanosoma evotomys*. (A and B) Raw abundance data (units as in Fig. 5). (C and D) Residuals from minimum sufficient ANOVAs to control for the compounding effects of year, season, host sex and age. The lines are best-fit linear relationships fitted in Cricket Graph III, Version 1.5. See text for statistical analysis.

mean infracommunity species richness was 1.4, no animals carried all 5 species, and only 7 voles carried various combinations of 4 species. Perhaps surprisingly, neither age nor sex influenced mean species richness significantly but seasonal changes were apparent in 2 of the years. In 1998 there was little seasonal variation generating a statistically significant interaction, which explained more of the variation in the model than the relatively weak between-year difference. Thus, overall mean infra-

community species richness remained relatively stable across the 3 years of our study.

Fluctuations in the prevalence and abundance of individual species of parasites were considerably more marked. Both prevalence and abundance of *T. evotomys* were highly season dependent and varied between the years, dropping low in two of the 3 years in the spring and then climbing to peak either in the summer or autumn. These observations concur with earlier work reporting marked reductions in preva-

lence in the winter among voles in the UK (Young, 1970; Turner, 1986). *T. evotomys* belongs to the *lewisi* group of trypanosomes and is mainly transmitted by fleas (Molyneux, 1969). In experimental primary infections in voles parasitaemia is relatively short compared with other species, but can last for 3 weeks (Molyneux, 1969), sustained by reproducing amastigote stages in the spleen and lymphoid tissues (Molyneux, 1969). As with other related trypanosomes, it is likely that strong acquired immunity is a feature of the host–parasite relationship (Maraghi & Molyneux, 1989; Albright & Albright, 1991). Hence, we expected to see this species mainly in young animals, as found by Healing (1981), but the lack of a clear age effect in our study signifies that voles of all ages were exposed to infection and that many achieved maturity before becoming infected for the first time. Overall prevalence was 15%, intermediate in comparison to prevalence rates reported by Sebek (1979), Healing (1981), Karbowski & Sinski (1996b) and that found by Turner (1986). In this part of Poland winters are severe (the ground may be frozen and covered by deep snow for months at a time) when it is likely that flea populations are less active, surviving only in the close confines of rodent nests. Such close proximity of vectors and hosts during the stressful winter period (and hence possibly with weakened immune systems) may have resulted in extended transmission between vector and host in those nests where infection was present. Consequently, any underlying age dependence in susceptibility to infection may have become occluded resulting in higher prevalence among older hosts than otherwise expected.

The ex-erythrocytic schizont stages of *H. erhardovae* reside in the lungs (Krampitz & Haberkorn, 1988), where they can last for 7–9 months after initial infection (Krampitz, 1964). Since only gametocytes are quantifiable from blood, the real prevalence was probably much higher than reflected in our data. Both prevalence and abundance varied significantly from year to year and there were seasonal differences, but the overall pattern was complex and subject to host age and sex. It showed some similarity to that reported by Healing (1981), whose data suggest between-year differences but show no clear seasonal pattern, and contrasted with Young (1970) and Turner (1986), who found very marked seasonal patterns characterized by a sharp reduction in prevalence in the winter months. Whilst we were not able to sample rodents in the winter, a complex picture was generated in the 3 seasons that we monitored, in which the between-year effect was stronger than any underlying seasonal effect and compounded by sex- and age-based interactions.

Some species of *Bartonella* (*B. baciliformis*, *B. quintana*, *B. henselae*, *B. elizabethae*) are considered to be infective to humans (Birtles *et al.* 1995; Anderson & Neuman, 1997) but *B. grahamii* (=

*Grahamella* sp.; Birtles *et al.* 1995), probably transmitted by fleas, is a parasite of rodents (Krampitz, 1962) in which it causes chronic infections lasting up to 120 days (Tyzzer, 1942). In our study the abundance of *B. grahamii* varied significantly with seasons, increasing from a low in spring to peak in either summer or autumn, as found by Young (1970) and Turner (1986), but in contrast to the latter study the seasonal pattern varied between years.

Giemsa-stained blood smears do not allow confident distinction between some of the Anaplasmataceae. In all likelihood the organisms infecting our animals were *Haemobartonella* sp., but we cannot be totally confident about this. Several species have been described from other rodent hosts (e.g. *H. muris musculi* from *Mus musculus* (Schilling, 1928); *H. microti* from *Microtus p. pennsylvanicus* (Tyzzer & Weinman, 1939); *H. arvicolae* from *M. arvali* (Yakimoff, 1928 after Weinman, 1944)), but to our knowledge the *Haemobartonella* sp. infecting *C. glareolus* has not been characterized. We observed small round cocci, mostly orientated centrally on cells, although in 5% of cases marginal attachment on the edge of cells suggested that at least some were extracellular. Infections were generally low, with the highest recorded intensity being 16 parasitized cells/1000 erythrocytes, but the vast majority (75.1%) were 1–2 parasitized cells/1000 erythrocytes with only 1 organism per cell. Our analysis revealed a complex situation and it was not possible to prioritize the relative importance of the influence of extrinsic and intrinsic factors on variation in *Haemobartonella* sp. because only compound interactions were detected. Arthropod vectors transmit the Anaplasmataceae, and likely candidates in this region are fleas and lice (Weinman, 1944; Gothe & Kreier, 1977) and hence cycles of transmission will inevitably be complex because of the different seasonal occurrence of these vectors.

Overall, perhaps the most surprising outcome of our analysis was the absence of any dominant role for intrinsic factors, particularly age, in structuring the component communities of haemoparasites in voles, albeit both sex and age contributed to some of the 2 and 3-way interactions recorded. It may be relevant that earlier workers also disagreed about age-based patterns of prevalence: young voles (Healing, 1981), old voles (Turner, 1986) and voles of intermediate age (Young, 1970) have all been reported as showing high prevalence of *T. evotomys* in different studies. Similar disagreements are recorded for *H. erhardovae* and *B. grahamii*. However, consistent with our data, but contrasting with Young (1970), neither Turner (1986) nor Healing (1986) found any sex bias in prevalence of the parasites they monitored.

Since the frequency distribution of infra-community species richness did not differ stat-

istically from that predicted by a null model, the frequency of 2, 3 and 4 species co-infections was consistent with those expected from their respective prevalence rates. We conclude that at this level of analysis, our data did not support the existence of strong associations between species. Moreover, at the level of individual parasite species we found only weak support for interactions. *B. grahamii* with *T. evotomys*, reported earlier as co-varying (Turner & Cox, 1985), did not do so in our study although a weak positive quantitative association (which we could not accept as significant without risking a Type I error) was evident. As reported by Young (1970), *T. evotomys* associated with *H. erhardovae*, but weakly and in a year and age-dependent manner. The latter species also showed a positive association with *Haemobartonella* sp., but in a season and age-dependent manner and there was no support for a quantitative association of these species. The only combination which was convincingly supported by both prevalence (albeit in a sex and age-dependent manner) and quantitative data sets was that between *B. grahamii* and *Haemobartonella* sp. Both of these species are Anaplasmatataceae and transmitted by arthropods, although little is known about their transmission dynamics and relationships with their vector hosts.

While our study was not designed specifically to relate host population size, or climatic changes, to parasite prevalence, some relationships are worth pointing out. The over-wintering population of voles was similar in each of the 3 years and fell further in May before peaking at varying levels in the summer. The relatively high density in the autumn of 1999 followed a long period of warm, relatively dry weather in the region. Interestingly, the rise in prevalence and abundance of *B. grahamii*, *H. erhardovae* and *T. evotomys* from winter to summer paralleled the increasing host populations in 1997 and 1998 but it was in 1999 that we observed the greatest contrast with other years (least apparent seasonal variation in mean infracommunity species richness and prevalence of *B. grahamii*). In contrast, *H. erhardovae* showed the highest prevalence in 1999 and a reversed pattern of seasonal occurrence. These contrasting patterns may have been related to fluctuations in host and vector populations and this will be worth exploring in future work.

The forest site in which our study was conducted has been relatively stable for about 70–80 years and now comprises mature mixed woodland, known to be the preferred habitat of bank voles (Alibhai & Gipps, 1985). However, this site borders closely on abandoned fields, which have lain fallow for 8 years and been allowed to revert to nature (since 1991, Rogozinska, 1997), with succession to woodland apparent throughout in the form of young, high and dense silver birch copses. The bank vole population in the woodland now has much more opportunity to

expand, to mix with other vole populations (and to be exposed to infections from neighbouring populations) and the corridors of new woodland are likely to increase with each passing year if no intervention is implemented. The long-term consequences of the changing ecology of the region for parasitic infections are largely unknown, and therefore, it is all the more important to sustain the monitoring activities for the future.

Finally, this study, the first comprehensive analysis of the component community structure of haemoparasites in *C. glareolus* from NE Europe, has established the highly dynamic nature of the community with marked variation between years and seasons. These extrinsic factors, and their interaction, consistently explained the greatest proportion of the variation in our data. It is also pertinent that despite the underlying fluctuation in the prevalence and abundance of individual species, variation of infracommunity species richness was less marked with moderate seasonal fluctuations, and between-year differences, within a relatively narrow range. At this level the component community of haemoparasites in *C. glareolus* was therefore relatively stable, despite the changing environment on the periphery of our study site.

We are grateful to the British Council, UK and the State Committee for Scientific Research (KBN), Poland for financial support for this study (KBN Grants UM 855 and UM 930). We thank Dr R. Halba for making the facilities at the field station in Urwitalt available to us.

#### REFERENCES

- ALBRIGHT, J. W. & ALBRIGHT, J. F. (1991). Rodent trypanosomes: their conflict with the immune system of the host. *Parasitology Today* **7**, 137–140.
- ALIBHAI, S. K. & GIPPS, J. H. W. (1985). The population dynamics of bank voles. In *The Ecology of Woodland Rodents Bank Voles and Wood Mice. Symposia of the Zoological Society of London* **55** (ed. Flowerdew, J. R., Gurnell, J. & Gipps, J. H. W.), pp. 277–305. Clarendon Press, Oxford.
- ANDERSON, B. E. & NEUMAN, M. A. (1997). *Bartonella* spp. as emerging human pathogens. *Clinical Microbiology Reviews* **10**, 203–219.
- BAKER, J. R., CHITTY, D. & PHIPPS, E. (1963). Blood parasites of wild voles, *Microtus agrestis*, in England. *Parasitology* **53**, 297–301.
- BEHNKE, J. M., LEWIS, J. W., MOHD ZAIN, S. N. & GILBERT, F. S. (1999). Helminth infections in *Apodemus sylvaticus* in southern England: interactive effects of host-age, sex and year on prevalence and abundance of infections. *Journal of Helminthology* **73**, 31–44.
- BIRTLES, R. J., HARRISON, T. G., SAUNDERS, N. A. & MOLYNEUX, D. H. (1995). Proposal to unify the genera *Grahamella* and *Bartonella*, with descriptions of *Bartonella talpae* comb. nov., and three new species, *Bartonella grahamii* sp. nov., *Bartonella taylorii* sp. nov., and *Bartonella doshiae* sp. nov.. *International Journal of Systematic Bacteriology* **45**, 1–8.

- CRAWLEY, M. T. (1993). *GLIM for Ecologists*. Blackwell Scientific Press, Oxford.
- DASH, K. M., HALL, E. & BARGER, I. A. (1988). The role of arithmetic and geometric mean worm egg counts in faecal egg count reduction tests and in monitoring strategic drenching programs in sheep. *Australian Veterinary Journal* **65**, 66–68.
- ELLIOTT, J. M. (1977). *Some Methods for the Statistical Analysis of Samples of Benthic Invertebrates*. Freshwater Biological Association, Cumbria, UK.
- GOTHE, R. & KREIER, J. P. (1977). *Aegyptianella, Eperythrozoon and Haemobartonella*. In *Parasitic Protozoa*, Vol. IV (ed. Kreier, J. P.), pp. 251–294. Academic Press, New York.
- HEALING, T. D. (1981). Infections with blood parasites in small British rodents *Apodemus sylvaticus*, *Clethrionomys glareolus* and *Microtus agrestis*. *Parasitology* **83**, 179–189.
- JANOVY, J. JR., CLOPTON, R. E., CLOPTON, D. A., SNYDER, S. D., EFTING, A. & KREBS, L. (1995). Species density distributions as null models for ecologically significant interactions of parasite species in an assemblage. *Ecological Modelling* **77**, 189–196.
- KAROWIAK, G. & SINSKI, E. (1996a). The finding of *Babesia microti* in bank vole *Clethrionomys glareolus* in the district of Mazury Lakes (Poland). *Acta Parasitologica* **43**, 50–51.
- KAROWIAK, G. & SINSKI, E. (1996b). Occurrence and morphological characteristics of a *Trypanosoma evotomys* strain from North Poland. *Acta Parasitologica* **41**, 105–107.
- KAROWIAK, G., STANKO, M., RYCHLIK, L., NOWAKOWSKI, W. & SIUDA, K. (1999). The new data about zoonotic reservoir of *Babesia microti* in small mammals in Poland. *Acta Parasitologica* **44**, 142–144.
- KRAMPIITZ, H. E. (1962). Weitere Untersuchungen an *Grahamella*, Brumpt, 1911. *Zeitschrift fuer Tropenmedizin und Parasitologie* **13**, 34–53.
- KRAMPIITZ, H. E. (1964). Über das Vorkommen und Verhalten von Haemococcidien der Gattung *Hepatozoon*, Müller 1908 (Protozoa, Adeleida) in mittel- und südeuropäischen Säugern. *Acta Tropica* **21**, 114–154.
- KRAMPIITZ, H. E. & HABERKORN, A. (1988). Experimental treatment of hepatozoon infections with anticoccidial agent toltrazuril. *Journal of Veterinary Medicine, B* **35**, 131–137.
- MARAGHI, S. & MOLYNEUX, D. H. (1989). Studies on cross-immunity in *Herpetosoma* trypanosomes of *Microtus*, *Clethrionomys* and *Apodemus*. *Parasitology Research* **75**, 175–177.
- MARGOLIS, L., ESCH, G. W., HOLMES, J. C., KURIS, A. M. & SCHAD, G. A. (1982). The use of ecological terms in parasitology (report of an *ad hoc* committee of The American Society of Parasitologists). *Journal of Parasitology* **68**, 131–133.
- MAZURKIEWICZ, M. (1972). Density and weight structure of populations of the bank vole in open and enclosed areas. *Acta Theriologica* **34**, 455–465.
- MOLYNEUX, D. H. (1969). The morphology and biology of *Trypanosoma* (*Herpetosoma*) *evotomys* of the bank-vole, *Clethrionomys glareolus*. *Parasitology* **59**, 843–857.
- MORRIS, P. (1972). A review of mammalian age determination methods. *Mammal Review* **2**, 69–104.
- POULIN, R. (1993). The disparity between observed and uniform distributions: a new look at parasite aggregation. *International Journal for Parasitology* **23**, 937–944.
- ROGOZINSKA, M. (1997). Sukcesja w zgrupowaniu drobnych ssaków ugorowanego pola uprawnego. Stan w 4 late po rozpoczęciu ugorowania. (Changes in the population structures of small mammals on fallow fields, previously cultivated. Appraisal 4 years after the cessation of cultivation). M.Sc. thesis, Department of Ecology, University of Warszawa.
- SANTOS-GOMEZ, G. M., ABRANCHES, P., MARAGHIS, S., DIRIE, M. F., SILVA-PEREIRA, M. C., VALVERDE, D. & MOLYNEUX, D. H. (1993). Laboratory and field studies on *Herpetosoma trypanosomes* from Portugal. *Annales de Parasitologie Humaine et Comparée* **68**, 163–168.
- SCHILLING, V. (1928). *Eperythrozoon coccoides*, eine neue durch Splenectomie aktivierbare Dauerinfektion der weissen Maus. *Klinische Wochenschrift* **7**, 1854–1855.
- SEBEK, Z. (1978). Blood parasites of small mammals in Western Hungary. *Parasitologia Hungarica* **11**, 17–21.
- SEBEK, Z., ROSICKY, B. & SIXL, W. (1977). The occurrence of babesiosis affecting small terrestrial mammals and the importance of this zoonosis in Europe. *Folia Parasitologica* **24**, 221–228.
- SINSKI, E. & PAWELCZYK, A. (1999). Detection of reservoirs for Lyme borreliosis in the Mazury Lake district, Poland. *Zentralblatt für Bakteriologie* **289**, 689–703.
- SOKAL, R. R. & ROHLF, F. J. (1981). *Biometry*, 2nd Edn. Freeman and Company, New York.
- TURNER, C. M. R. (1986). Seasonal and age distributions of *Babesia*, *Hepatozoon*, *Trypanosoma* and *Grahamella* species in *Clethrionomys glareolus* and *Apodemus sylvaticus* populations. *Parasitology* **93**, 279–289.
- TURNER, C. M. R. & COX, F. E. G. (1985). Interspecific interactions between blood parasites in a wild rodent community. *Annals of Tropical Medicine and Parasitology* **79**, 463–465.
- TYZZER, E. E. (1942). A comparative study of *Grahamella*, *Haemobartonella* and *Eperythrozoon* in small mammals. *Proceedings of the American Philosophical Society* **85**, 359–398.
- TYZZER, E. E. & WEINMAN, D. (1939). *Haemobartonella* n.g. (*Bartonella* olim. pro parte). *H. microti* n. sp. of the field vole. *American Journal of Hygiene* **30**, 141–157.
- WALTER, G. & LIEBISCH, A. (1980). Studies of the ecology of some blood protozoa of mammals in north Germany. *Acta Tropica* **37**, 31–40.
- WIGER, R. (1979). Seasonal and annual variations in the prevalence of blood parasites in cyclic species of small rodents in Norway with special reference to *Clethrionomys glareolus*. *Holarctic Ecology* **2**, 169–175.
- WILSON, K. & GRENFELL, B. T. (1997). Generalized linear modelling for parasitologists. *Parasitology Today* **13**, 33–38.
- WEINMAN, D. (1944). Infectious anaemia due to *Bartonella* and related red cell parasites. *Transactions of the Philosophical Society* **33**, 243–350.
- YOUNG, A. S. (1970). Studies on the blood parasites of small mammals with special reference to piroplasms. Ph.D. thesis, University of London, UK.