

Seasonal dynamics of tuberculosis epidemics and implications for multidrug-resistant infection risk assessment

Y.-J. LIN AND C.-M. LIAO*

Department of Bioenvironmental Systems Engineering, National Taiwan University, Taipei, Taiwan, ROC

*Received 27 November 2012; Final revision 31 March 2013; Accepted 15 April 2013;
first published online 16 May 2013*

SUMMARY

Understanding how seasonality shapes the dynamics of tuberculosis (TB) is essential in determining risks of transmission and drug resistance in (sub)tropical regions. We developed a relative fitness-based multidrug-resistant (MDR) TB model incorporated with seasonality and a probabilistic assessment model to assess infection risk in Taiwan regions. The model accurately captures the seasonal transmission and population dynamics of TB incidence during 2006–2008 and MDR TB in high TB burden areas during 2006–2010 in Taiwan. There is ~3% probability of having exceeded 50% of the population infected attributed to MDR TB. Our model not only provides insight into the understanding of the interactions between seasonal dynamics of TB and environmental factors but is also capable of predicting the seasonal patterns of TB incidence associated with MDR TB infection risk. A better understanding of the mechanisms of TB seasonality will be critical in predicting the impact of public control programmes.

Key words: Infection risk, modelling, multidrug-resistant, transmission, seasonality, tuberculosis.

INTRODUCTION

Recently, the World Health Organization (WHO) documented the diagnosis of nearly 8·7 million new cases of tuberculosis (TB) in 2011 with an estimated 1·4 million deaths from TB in the same year [1]. Although the absolute number of TB cases has been falling since 2006, TB remains a leading cause of global morbidity and mortality in that one-third of the human population is infected. The bacterium *Mycobacterium tuberculosis* that causes TB is generally spread in airborne droplets when people with active disease cough or sneeze.

Multidrug-resistant (MDR) TB has been documented in 114 countries and regions worldwide and has emerged as a global public health issue [2]. MDR TB is caused by strains resistant to at least isoniazid and rifampicin, the two principal first-line drugs used in combination chemotherapy [3]. Treatment for MDR TB patients requires use of second-line drugs for at least 24 months [4]. Therefore, MDR TB is increasingly becoming a serious threat to TB control, and the recognition of extensively drug-resistant TB has further highlighted this threat [5].

In Taiwan, TB has always had the highest incidence rate in all communicable diseases. The incidence rate of TB in Taiwan was 62·0–74·6/100 000 population for the period 2002–2008, respectively [6]. About 149–164 new MDR TB cases were reported in Taiwan during 2007–2010 [7]. Although MDR TB represents only 1·2% of total new TB cases in

* Author for correspondence: Dr C.-M. Liao, Department of Bioenvironmental Systems Engineering, National Taiwan University, Taipei, Taiwan 10617, ROC.
(Email: cmliao@ntu.edu.tw)

Taiwan, controlling MDR TB is challenging because it is becoming increasingly hard to diagnose and treat.

Previous studies have reported a strong association between seasonality and TB in temperate, tropical, and subtropical regions [8–12]. Their findings indicated that seasonal peaks of TB cases generally occurred at the end of winter and at the beginning of the spring [8–10] or summer seasons [11, 12]. They also implied that indoor activity in winter, diagnostic delays, seasonal change in human immunity, and an association with sunlight and vitamin D levels might play the crucial factors in affecting TB seasonality [10, 12]. Thus, there are strong interactions between the effects of seasonality and the effects of weather because seasonality might influence the season of emergence of TB, making both sensitive to additional stress such as climate change.

The simplest mathematical model for modelling drug resistant (DR) and MDR TB epidemics came from Blower & Gerberding [13]. However, their model did not consider the contribution of exogenous re-infection to the overall disease incidence. Over the past two decades many expanded and sophisticated models have been used to predict the future burden of DR and MDR TB. Dye & Williams [14] and Cohen & Murray [15] developed the MDR TB model, incorporating re-infection at a reduced rate by partial immunity applying to latent or recovered individuals. In view of these models, it is recognized that the assumptions about the relative fitness (RF) of MDR strains play a crucial role in describing MDR dynamics [15, 16].

The seasonality and meteorological factors might influence the dynamics of TB; however, little is known about the time-series dynamics of TB in the (sub)tropical Taiwan region taking into account seasonal patterns and weather effects. The purpose of this study was to examine seasonal population dynamics of TB and to estimate implicitly the MDR TB infection risk in the Taiwan region. We sought to develop a RF-based MDR TB model built on previous models, seasonal transmission to predict the potential impact of seasonality on TB and MDR TB transmission. A probabilistic model was also developed to estimate the site-specific infection risks of TB and MDR TB in Taiwan regions. We anticipated that our study could provide insights into the dynamics of TB/MDR TB transmission, which would inform the uncertainty over the different regions for populations at high-burden risk.

MATERIALS AND METHODS

Study data

Monthly-based disease burden of TB data in Taiwan were obtained from the Centers for Disease Control of Taiwan (Taiwan CDC) (<http://www.cdc.gov.tw/>) for 2005–2008. We geographically divided Taiwan into four regions (northern, central, southern, eastern) in order to estimate TB incidence rates. The annual disease burden of MDR TB for each county was adopted from the Taiwan tuberculosis control report [17] and Taiwan CDC national notified disease surveillance system [7] for 2006–2010 to calculate MDR TB incidence rates. The monthly-based mean temperature, relative humidity, and rainfall intensity were adopted from the Taiwan Central Weather Bureau (<http://www.cwb.gov.tw/>) for 2005–2008.

We found that the average incidence rates during 2005–2008 in northern, central, southern, and eastern regions were $57.55 \pm 3.88/100\,000$, $67.32 \pm 6.27/100\,000$, $81.52 \pm 5.18/100\,000$ and $111.55 \pm 19.45/100\,000$ population, respectively. Moreover, we also found that the higher TB epidemic areas appeared in Hualien and Taitung counties in east Taiwan and Pingtung county in southern Taiwan.

The annual incidence rates of TB and MDR TB in Hualien county were $119.72 \pm 13.84/100\,000$ and $4.91 \pm 2.37/100\,000$ respectively, whereas the annual TB and MDR TB incidence rates were $103.39 \pm 9.82/100\,000$ and $2.82 \pm 1.79/100\,000$ respectively, in Taitung county. The annual incidence rates of TB and MDR TB in Pingtung county in south Taiwan were $105.98 \pm 10.36/100\,000$ and $0.97 \pm 0.34/100\,000$ respectively, whereas Taipei city in north Taiwan experienced the lowest incidences of TB at $48.46 \pm 3.39/100\,000$ and MDR TB at $0.43 \pm 0.11/100\,000$. We thus used TB epidemic data of Hualien, Taitung, Pingtung counties, and Taipei City to investigate the seasonal transmission dynamics of TB and to estimate implicitly the MDR TB infection risk.

To model drug resistance dynamics, the data of RF of DR strains have to be determined. One of the methods to measure RF of resistant strains is based on the results of genotype clustering studies in that a cluster can be defined as two or more cases having an identical DNA fingerprint [3]. Based on the genotype clustering method, RF can be estimated by calculating the odds ratio as $RF = (C_R/G_R)/(C_S/G_S)$, where C_R , C_S , G_R , and G_S are the numbers of resistant (R) and sensitive (S) cases that appear singly (G) or in clusters (C) [3].

García-García *et al.* [18] have recently provided a valuable genotype clustering study that can be used to estimate RF of MDR TB. They found that the overall rate of resistance was 28.4%, in that 10.8% had MDR TB. DNA fingerprinting was performed on 188 (81%) of 232 culture-confirmed cases for whom bacterial DNA was available. Of the 188 patients with a DNA fingerprint, 120 (64%) were infected with unique genotypes and 68 (36%) with identical DNA fingerprints were grouped into a total of 20 clusters. Based on the genotype clustering analysis, Garcia-Garcia *et al.* [18] estimated the odds ratio of MDR TB compared to drug-sensitive (DS) TB was 0.16 [95% confidence interval (CI) 0.04–0.6]. Therefore, we adopted this odds ratio and used the Monte Carlo (MC) technique to obtain the best-fitted distribution of RF to capture the uncertainty.

Seasonal transmission dynamic model

The previous well-developed DR TB transmission models [13–16, 19, 20] were adopted and modified to describe parsimoniously the population dynamics of MDR TB in Taiwan. Thus a two-strain TB model was used (see Supplementary Fig. S1). Briefly, (i) susceptible individuals may be infected with either DS or MDR strains, (ii) two certain types of primary progressive TB (i.e. fast TB) and latently infected TB caused by endogenous reactivation or exogenous re-infection (i.e. slow TB) were included, (iii) a case may be spontaneously cured at a cure rate and move into the latent non-infectious state, and (iv) the MDR TB epidemic can be composed of two main subepidemics in which individuals are primarily infected/re-infected with MDR TB (i.e. primary resistance) and have DS TB treatment failure resulting in resistance (i.e. acquired resistance). The system equations of the present two-strain TB model incorporated with seasonal transmission are listed in Table 1.

This study used a regression model incorporating seasonality, temperature, relative humidity, and rainfall as potential predictors of TB to assess the characteristics of TB epidemics in Taiwan during 2005–2008. The model was fitted to data to estimate TB trends as

$$Y_{RM}(t) = \beta_0 + \sum_{n=1}^5 \left(\beta_{1,n} \sin\left(\frac{2n\pi t}{12}\right) + \beta_{2,n} \cos\left(\frac{2n\pi t}{12}\right) \right) + \beta_3 \text{Temp} + \beta_4 \text{RH} + \beta_5 \text{Rain}, \tag{1a}$$

where $Y_{RM}(t)$ is the expected monthly number of TB new cases at time t estimated based on the regression model, β_0 is the intercept, $\beta_{1,n}$, $\beta_{2,n}$, and β_3 to β_5 represent the fitted coefficients, $\sin(2n\pi t/12)$ and $\cos(2n\pi t/12)$ represent seasonality in which n is the number of seasonal cycles per year, and Temp, RH, and Rain are the mean temperature (°C), relative humidity (%), and rainfall (mm), respectively.

However, the expected monthly number of new TB cases at time t can also be calculated based on the two-strain TB model by incorporating equations (T4) and (T5) and is designated as $Y_{TM}(t)$:

$$Y_{TM}(t) = p\beta_S(t)T_S S + (v + p\sigma\beta_S(t)T_S)L_S + p\beta_R(t)T_R S + p\sigma\beta_R(t)T_R L_S + (v + p\sigma\beta_S(t)T_S + p\sigma\beta_R(t)T_R)L_R + c_{Fr}T_S. \tag{1b}$$

$\beta_S(t)$ can then be written as

$$\beta_S(t) = \frac{Y_{TM}(t) - vL_S - vL_R - c_{Fr}T_S}{pT_S S + p\sigma T_S L_S + pRFT_R S + p\sigma RFT_R L_S + p\sigma T_S L_R + p\sigma RFT_R L_R}. \tag{2}$$

To estimate $\beta_S(t)$, we linked the two-strain TB model and regression models based on the relationship between $\beta_S(t)$ and expected monthly number of new TB cases. We combined equations (1) and (2) to solve $\beta_S(t)$.

The expressions for basic reproduction number (R_0) [20, 21] quantifying the transmission potential of *M. tuberculosis* due to the subepidemics driven by DS TB (R_{0S}) and MDR TB (R_{0R}) are summarized in Table 1. R_0 is defined as the average number of successful secondary infectious cases generated by a typical primary infected case in an entirely susceptible population [22]. When $R_0 > 1$ it implies that the epidemic is spreading within a population and incidence is increasing, whereas $R_0 < 1$ means that the disease is dying out. An average R_0 of 1 means the disease is in endemic equilibrium within the population.

Probabilistic TB risk model

To develop a probabilistic DS/MDR TB risk model, a dose–response model describing the relationships between transmission potential of DS/MDR *M. tuberculosis* quantified by R_0 and the total proportion of infected population must be constructed. Generally, the probability of infection for each susceptible person each day is based on the transmission probabilities for each potentially infected contact. According to Anderson & May [22], in a homogeneous and

Table 1. Equations for the present proposed two-strain tuberculosis (TB) model

Equation	Meaning
Two-strain TB model*	
$\dot{S}(t) = \pi - (\beta_S(t)T_S + \beta_R(t)T_R + \mu) S$	(T1) Susceptible individuals
$\dot{L}_S(t) = (1 - p)\beta_S(t)T_S S + c_S T_S - (v + p\sigma\beta_S(t)T_S + \sigma\beta_R(t)T_R + \mu)L_S$	(T2) Latently infected individuals with drug-sensitive (DS) TB
$\dot{L}_R(t) = (1 - p)\beta_R(t)T_R S + (1 - p)\sigma\beta_R(t)T_R L_S + c_R T_R - (v + p\sigma\beta_S(t)T_S + p\sigma\beta_R(t)T_R + \mu)L_R$	(T3) Latently infected individuals with multidrug-resistant (MDR) TB
$\dot{T}_S(t) = p\beta_S(t)T_S S + (v + p\sigma\beta_S(t)T_S)L_S - (c_S + \mu + \mu_S + c_F r) T_S$	(T4) DS infectious TB
$\dot{T}_R(t) = p\beta_R(t)T_R S + p\sigma\beta_R(t)T_R L_S + (v + p\sigma\beta_S(t)T_S + p\sigma\beta_R(t)T_R)L_R + c_F r T_S - (c_R + \mu + \mu_R) T_R$	(T5) MDR infectious TB
Basic reproduction number	
$R_{0S} = \frac{\beta_S N_T p(\mu + v)}{(\mu + v)(\mu + \mu_S + c_S + (1 - c_S)c_F) - c_S v}$	(T6) Basic reproduction number of DS TB
$R_{0R} = \frac{\beta_R N_T p(\mu + v)}{(\mu + v)(\mu + \mu_R + c_R) - c_R v}$	(T7) Basic reproduction number of MDR TB

π , Recruitment rate (person yr⁻¹); $\beta_S(t)$, seasonal transmission rate for DS TB at time t (person⁻¹ yr⁻¹); $\beta_R(t) = RF \times \beta_S(t)$, seasonal transmission rate for MDR TB at time t (person⁻¹ yr⁻¹), where RF is the relative fitness of MDR strains; μ , background mortality rate (yr⁻¹); p , probability of new infections that develop progressive primary active TB within 1 year; c_S , cure rate of active DS TB (yr⁻¹); v , progression rate from latency to active TB (yr⁻¹); σ , partial immunity that decreases probability of fast progression after re-infection; c_R , cure rate of active MDR TB (yr⁻¹); μ_S , DS TB-caused mortality rate (yr⁻¹); c_F , rate of treatment failure (yr⁻¹); r , proportion of DS TB treatment failure acquiring resistant; μ_R , MDR TB-caused mortality rate (yr⁻¹); N_T , total population size (person).

* See Supplementary Figure S1.

unstructured population, the total proportion of infected population during the epidemic (I) depends only on R_0 , and can be theoretically expressed as

$$I = 1 - \exp(-R_0 I). \tag{3}$$

Equation (3) cannot be solved analytically. Thus, we solved equation (3) numerically by using a nonlinear regression model to best fit the profile describing the relationship between I and R_0 [equation (3)] for R_0 , ranging from 1 to 10. We found that I can be expressed as a function of R_0 only,

$$I(R_0) = 1 - \exp(1.63 - 1.66 R_0), \quad 1 < R_0 < 10, \tag{4}$$

$$(r^2 = 0.99)$$

Equation (4) can be seen as a conditional response distribution describing the dose–response relationship between I and R_0 and can be expressed as: $P(I|R_0)$. Thus, following Bayesian inference, DS/MDR TB infection risk (the posterior probability) can be calculated as the product of the probability distribution of R_0 (the prior probability) and the conditional response probability of proportion of the population expected to be infected, given R_0 [the likelihood $P(I|R_0)$]. This

results in a joint probability distribution or a risk profile as

$$R(I) = P(R_0) \times P(I|R_0), \tag{5}$$

where $R(I)$ is the cumulative distribution function describing the probabilistic infection risk of a TB epidemic in a susceptible population at specific R_0 , and $P(R_0)$ is the probability density function of R_0 . The exceedance risk profile can be obtained by $1 - R(I)$.

Model parameterization and validation

The parameter values used in the two-strain TB model [Table 1, equations (T1)–(T5)] can be estimated based on site-specific TB data obtained from Taiwan CDC, Department of Statistics, Ministry of the Interior, ROC (Taiwan) [23], and published data adopted from the literature [18, 19, 24–26]. We used the two-strain TB model to predict the seasonal incidence dynamics of TB in our four study sites for 2006–2016 with 95% credible intervals.

We validated the model performance by using the root mean squared error (RMSE) to compare the MDR TB incidence rates between predictions and

Table 2. Fitting of regression models for Hualien, Taitung, Pingtung counties, and Taipei city, respectively, in the period 2005–2008

Regression model								
Parameter	Hualien county		Taitung county		Pingtung county		Taipei city	
Intercept	✓	✓	✓	✓	✓	✓	✓	✓
sin(2nπt/12)								
n = 1	✓	✓	✓	✓	✓	✓	✓	✓
n = 2	✓	✓	✓	✓	✓	✓	✓	✓
n = 3	✓	✓	✓	✓	✓	✓	✓	✓
n = 4	✓	✓	✓	✓	✓	✓	✓	✓
n = 5	✓	✓	✓	✓	✓	✓	✓	✓
cos(2nπt/12)								
n = 1	✓	✓	✓	✓	✓	✓	✓	✓
n = 2	✓	✓	✓	✓	✓	✓	✓	✓
n = 3	✓	✓	✓	✓	✓	✓	✓	✓
n = 4	✓	✓	✓	✓	✓	✓	✓	✓
n = 5	✓	✓	✓	✓	✓	✓	✓	✓
Environmental factors								
Temperature	✓		✓		✓		✓	
Relative humidity	✓		✓		✓		✓	
Rainfall	✓		✓		✓		✓	
AIC	210·11	206·28	183·91	182·43	280·68	274·76	272·79	267·69
ΔAIC	Ref.*	3·83	Ref.	1·48	Ref.	5·92	Ref.	5·10
r	0·65	0·63	0·52	0·46	0·36	0·36	0·53	0·52

AIC, Akaike’s Information Criterion.

* Ref. = reference model [equation (1a)].

The best fitting models are highlighted in bold font.

observed data obtained from Taiwan CDC for 2006–2010. The RMSE is a measure of the differences between the predictive values by the two-strain TB model and observation data obtained from Taiwan CDC. The RMSE is calculated as

$$RMSE = \sqrt{\sum_{k=1}^K (I_{o,n} - I_{s,n})^2 / K},$$

where K denotes the number of observations, $I_{o,n}$ is the observed incidence rate, and $I_{s,n}$ is the simulation result corresponding to data point k [27].

Uncertainty analyses and simulation scheme

TableCurve 2D package (AISN Software Inc., USA) and Statistica (version 9, Statsoft Inc., USA) were used to perform model-fitting techniques and statistical analyses. A MC technique was implemented to quantify the uncertainty and its impact on the estimation of expected risk. A MC simulation was also performed with 10000 iterations to generate 2·5 and 97·5 percentiles as the 95% CI for all fitted models. Crystal Ball software (version 2000·2, Decisioneering Inc., USA) was employed to implement MC simulation. Model simulations were performed using

Berkeley Madonna 8·0·1 (Berkeley Madonna was developed by Robert Macey and George Oster, University of California at Berkeley).

Akaike’s Information Criterion (AIC) was used to measure the relative goodness of fit for the regression model. AIC is an assessment for regression model selection and can be expressed as $AIC = 2k - 2\ln(L)$, where k is the number of predictors in the regression model and L is the maximized value of the likelihood function for the estimated model [28]. In the regression model selection procedure, we used equation (1a) as reference model to compare another model and then calculated the differences in AIC (ΔAIC). A model was considered more likely when $\Delta AIC \geq 4$. When two models had similar values with $\Delta AIC < 4$, the model with the lowest AIC value was selected [29]. A P value $< 0·05$ was taken as significant.

RESULTS

Optimal regression model determination

We first used a sine wave function to optimally fit the meteorological data to obtain likelihood




Fig. 1. Comparison of monthly number of new tuberculosis (TB) cases between regression model-fitting outcomes with 95% confidence intervals (CI) and observed data for July (2005–2008) in (a) Hualien county, (b) Taitung county, (c) Pingtung county, (d) Taipei city.

distribution of mean temperature ($r^2=0.91-0.96$), relative humidity ($r^2=0.18-0.62$), and rainfall ($r^2=0.32-0.45$) in four study sites for 2005–2008 (see Supplementary Table S1 and Supplementary Fig. S2). We then incorporated sine wave functions of mean temperature, relative humidity, and rainfall into the regression model, $Y_{RM}(t)$ [equation (1a)] to capture the TB trends, allowing both the magnitude (quantitative) (Table 2 and Supplementary Fig. S2) and shape (qualitative) (Fig. 1) of trends to be estimated. We found that meteorological factors showed no significant effects on TB incidence (Table 2). The best-fitting models with $\Delta AIC \geq 4$ were in Pingtung county ($\Delta AIC=5.92$, $r=0.36$) and Taipei city ($\Delta AIC=5.10$, $r=0.52$), whereas the smallest AIC values were in Hualien (AIC=206.28, $r=0.63$) and Taitung (AIC=182.43, $r=0.46$) counties (Table 2).

Seasonal population dynamics of TB/MDR TB

The results of model parameterization were listed in Table 3. We optimally fitted the published data [18] to obtain the likelihood distribution of RF by MC simulation. This resulted in a normal (N) distribution

of RF with a mean of 0.32 and a standard deviation (S.D.) of 0.14 (Table 3). The fitted coefficients in $Y_{RM}(t)$ [equation (1a)] for Hualien, Taitung, Pingtung counties, and Taipei city during 2005–2008 are listed in Supplementary Table S2 and were used in equation (2) to estimate $\beta_S(t)$. $\beta_R(t)$ can then be estimated by $\beta_S(t) \times \text{RF}$. We incorporated $\beta_S(t)$ [equation (2)], estimated probability distributions of the model parameters, and site-specific initial population size in 2006 (Table 3) into the two-strain TB model [Table 1, equations (T1)–(T5)] to project future site-specific population dynamics of TB and MDR TB incidence for 2006–2016 (Figs 2 and 3).

Figure 2 demonstrates the comparison of the TB incidence data with our model simulation output with 95% credible intervals, indicating that the predictions are in apparent agreement with the observed data for 2006–2008. Figure 3 shows the comparison of MDR TB incidence rates between predictions varying with different percentile estimates of β_R by the two-strain TB model for 2006–2016, indicating that the predictions were consistent with observed data for 2007–2010.

The model was also extended to project the TB and MDR TB incidence rate for periods 2009–2016 and

Table 3. Probability distributions (N =normal, LN =lognormal) of parameter values and initial population sizes used in the two-strain TB model and basic reproduction number (R_0) estimations^a

	Probability distribution			
	Hualien county	Taitung county	Pingtung county	Taipei city
Model parameter				
p^b			N(0.08, 0.03)	
v (yr^{-1}) ^c			N(0.0003, 0.00004)	
μ (yr^{-1}) ^d	LN(0.031, 2.05)	LN(0.031, 2.05)	LN(0.030, 2.11)	LN(0.027, 2.00)
μ_S (yr^{-1}) ^e	N(0.037, 0.015)	N(0.040, 0.019)	N(0.052, 0.021)	N(0.033, 0.013)
μ_R (yr^{-1}) ^f			N(0.30, 0.05)	
c_S (yr^{-1}) ^e	N(0.64, 0.07)	N(0.61, 0.08)	N(0.68, 0.09)	N(0.72, 0.01)
c_R (yr^{-1}) ^f	LN(0.10, 2.68)	LN(0.08, 2.83)	LN(0.18, 1.89)	LN(0.28, 1.41)
c_F (yr^{-1}) ^e	N(0.37, 0.06)	N(0.41, 0.08)	N(0.39, 0.09)	N(0.31, 0.06)
r^e	N(0.034, 0.018)	N(0.024, 0.022)	N(0.017, 0.007)	N(0.016, 0.005)
σ^c			N(0.65, 0.05)	
N_T (person) ^d	N(345297, 2748)	N(236156, 3174)	N(893289, 5625)	N(2625962, 8435)
π (person yr^{-1})	2960	2228	7284	21217
β_S^g	N(2.6×10^{-5} , 5.7×10^{-6})	N(3.6×10^{-5} , 6.4×10^{-6})	N(1.1×10^{-5} , 1.5×10^{-6})	N(3.7×10^{-6} , 3.7×10^{-7})
β_R^h	N(8.6×10^{-6} , 5.5×10^{-6})	N(1.2×10^{-5} , 7.6×10^{-6})	N(3.4×10^{-6} , 2.2×10^{-6})	N(1.2×10^{-6} , 7.3×10^{-7})
RF ⁱ			N(0.32, 0.14)	
Initial population size ^j				
N_T	348223	239658	899249	2619424
S	346469	238536	895303	2609262
L_S	1268	873	3276	9543
L_R	13	9	33	96
T_S	819	225	623	1818
T_R	19	15	14	9

^a See Table 1 for explanation of symbols.

^b Estimated by 0.04 (0.015–0.14) for <15-year-olds and 0.14 (0.08–0.25) for >15-year-olds [24].

^c Estimated from Lin *et al.* [25].

^d Estimated based on data from Department of Statistics, Ministry of the Interior, Taiwan [23].

^e Estimated based on Taiwan CDC data.

^f Estimated based on Dye & Espinal [19].

^g β_S were estimated based on equations (1) and (2).

^h $\beta_R = \beta_S \times RF$.

ⁱ RF is the relative fitness that is estimated based on Garcia-Garcia *et al.* [18].

^j The initial population size in 2006 of N_T , T_S , and T_R are adopted from Taiwan Tuberculosis Control Report (2007) [17]. $S = N_T - L_S - L_R - T_S - T_R$. $L_S = 0.004 \times 0.92 \times 0.99 \times N_T$ and $L_R = 0.004 \times 0.92 \times 0.01 \times N_T$, where 0.004 is adopted from Yeh *et al.* [26], 0.92=(1–0.08) adopted from Dye *et al.* [24], and the proportions of infections that develop L_S (0.99) and L_R (0.01) are assumed.

2011–2016, respectively. The RMSE of MDR TB simulations (Fig. 3) ranging from 0.38 to 3.85 were comparable to the data’s average standard deviation (s.d.) of 0.11–2.37. However, our model had the lowest RMSE values for the predictions with 50 (RMSE = 2.29), 50 (RMSE = 1.65), 2.5 (RMSE = 0.28), and 25 (RMSE = 0.10) percentiles in Hualien, Taitung, Pingtung counties, and Taipei city, respectively, indicating that all RMSE values were less than the s.d. of observed data. Overall, the model captures the seasonal transmission and population dynamics of TB incidence for 2006–2008 and MDR TB in high TB incidence areas in Taiwan during 2006–2010.

DS/MDR TB infection risk estimates

To estimate the probabilities of DS and MDR TB infection risk, the transmission potential quantified by R_{0S} and R_{0R} had to be determined. The site-specific seasonal R_{0S} (Fig. 4a, c, e, g) and R_{0R} (Fig. 4b, d, f, h) were calculated based on equations (3) and (4) with input parameter values listed in Table 3. The MC simulation results showed that the optimized log-normal distribution was the most suitable fitted model for seasonal values of R_{0S} and R_{0R} (Fig. 4i). We found that, for instance, in the highest TB epidemic area of Hualien county, the R_{0S} and R_{0R}




Fig. 2. Modelling seasonal tuberculosis (TB) incidence rates (per 100 000 population) with 95% credible intervals from 2006 to 2016 (for July) based on the two-strain TB model and the comparison of incidence data with model simulation outcomes for 2006–2008 in (a) Hualien county, (b) Taitung county, (c) Pingtung county, (d) Taipei city.

estimates were 0.97 (95% CI 0.26–2.03) and 0.44 (95% CI 0.06–1.45), respectively, whereas R_{0S} and R_{0R} values were 0.96 (95% CI 0.26–1.97) and 0.44 (95% CI 0.06–1.38), respectively, in Taitung county. The R_{0S} and R_{0R} estimates in Pingtung county were 0.97 (95% CI 0.26–1.92) and 0.41 (95% CI 0.05–1.25), respectively, whereas in Taipei city, the R_{0S} value was 0.98 (95% CI 0.27–1.77) with a lowest R_{0R} value of 0.37 (95% CI 0.05–1.02).

Figure 5a presents the conditional dose–response profile showing the estimate for the total proportion of TB-infected population (I) that depended only on R_0 based on equation (4). Given the site-specific R_{0S} and R_{0R} distributions (Fig. 4i) and conditional dose–response relationship $P(I|R_0)$ (Fig. 5a), the site-specific exceedance risk probability of DS and MDR TB infection can then be estimated by using equation (5) (Fig. 5b, c). We found that DS TB in Hualien, Taitung, Pingtung counties, and Taipei city, respectively, had nearly 17.6%, 15.5%, 14.5%, and 12.8% probabilities of the total proportion of infected population exceeding 50%, whereas there was a 29–32% probability of having exceeded 20% of the total proportion of infected population (Fig. 5b). Our results also indicated that the selected four regions had only

~3% probability of having exceeded 50% of the population infected attributed to MDR TB (Fig. 5c).

DISCUSSION

Seasonal dynamics of TB/MDR TB transmission

This study linked the two-strain TB and the statistical regression models to describe the seasonal transmission and population dynamics of TB and MDR TB in Taiwan. Our study showed that the seasonal peak of new TB cases generally occurred during late spring to early summer seasons in Taiwan. Thus dynamic consequences of seasonal variation in TB appeared in Taiwan. Several studies have found variable periods of peak seasonality of TB at the end of winter to late spring in Spain [8], in spring to late autumn in the USA [10], during summer in Hong Kong [11], and during spring and summer in Japan [12].

Our results revealed that meteorological factors showed weak effects on new TB cases based on the regression model analysis. In previous analysis, we also found that there was weak relationship between (lagged) temperature and TB incidence in a Poisson




Fig. 3. Annual incidence rates (per 100 000 population) of multidrug-resistant tuberculosis (MDR TB) estimated by the two-strain TB model varying with different percentile estimates of β_R during 2006–2016 and the comparison of incidence rates between predictions and observed data for 2006–2010 in (a) Hwalien county, (b) Taitung county, (c) Pingtung county, (d) Taipei city.

regression model; however, it is interesting to note that seasonality, aborigines, gender, and age showed stronger association with TB trends in Taiwan [28].

Greenman *et al.* [30] indicated that external forcing can cause some cycles, oscillations, and even chaotic phenomenon in disease dynamics of population. This study applied five cycles of seasonality to examine periodic trends of monthly new TB cases, which was similar to those of in China [9] and USA [10] and the seasonality was significant in the regression model, except in Pingtung county. Therefore, we conclude that seasonality is a major factor in shaping the TB dynamics and can also be seen as a predictor to better understanding the TB incidence patterns in Taiwan.

The causes of TB seasonality are not well understood. Some studies indicated that the transmission of TB may be intensified by increased time spent in overcrowded, poorly ventilated places, and by an increased frequency of viral infections like flu in winter [9]. Marais *et al.* [31] also indicated that viral

respiratory co-infection may increase susceptibility to TB infection and progression to disease. Many works suggest that vitamin D supplementation induces antimycobacterial immunity which can decrease the risk for TB infection [32, 33]. The seasonality of TB may reflect the seasonality of human immunity to TB [10, 12].

Furthermore, it is also important to consider that the seasonal peak of TB in spring may be due in part to delay in diagnosis of wintertime disease [12]. Missed early diagnosis of TB may occur more often in winter as coughs and fever are common seasonal symptoms, reflecting the viral respiratory infection [10]. The interval between observable immunological response and infection may be in excess of 7 weeks [33]. This could lead to a longer infectious period and increased transmission during winter, as well as more new TB cases will be detected in spring.

We used RMSE to validate the model fitness and test model predictive power in this study. The RMSE is a measure of the differences between




Fig. 4. Site-specific seasonal basic reproduction numbers of (a, c, e, g) drug-sensitive tuberculosis (TB) (R_{0S}) and (b, d, f, h) multidrug-resistant (MDR) TB (R_{0R}) in Hwaiien, Taitung, and Pingtung counties, and Taipei city. (i) The box-and-whisker plot illustrates the overall R_{0S} and R_{0R} .

the observed data and predictive values by optimal model. The RMSE values were comparable to the s.d. of the data. Our simulation results by the two-strain TB model showed that all RMSE values were less than the s.d. of observed data, demonstrating that the model-simulated values agreed with the observed data. Overall, our present model captures

the seasonal transmission and population dynamics of total TB incidence and MDR TB in high TB incidence areas in Taiwan for the periods 2006–2008 and 2006–2010, respectively.

Given the high frequency of MDR TB in eastern Taiwan, our simulation showed that the incidence of MDR TB seems to be falling by 2013–2016, which




Fig. 5. (a) The conditional dose–response profile representing the relationship between total proportion of tuberculosis (TB)-infected population (I) and R_0 . The site-specific exceedance risks of total proportions of TB infections estimated for (b) drug-sensitive (DS) TB and (c) multidrug-resistant (MDR) TB in Hwalien, Taitung, and Pingtung counties, and Taipei city.

is consistent with the downward trend of observed MDR TB incidence rate due to the effective TB control programmes. In particular, the implementation of the directly observed treatment short-course (DOTS) strategy since 2006 to prevent the occurrence of new MDR TB resulted from treatment failure and the MDR TB programme (DOTS-plus) since 2007 in providing MDR TB patients with complete and non-stop care [34]. Our finding also implicitly provides information that the ongoing control programmes implemented in Taiwan may succeed in curing patients with TB and MDR TB and reduce TB incidence nationwide.

Infection risk estimates of DS/MDR TB

The median values of the R_{OR} estimate were higher in Hwalien and Taitung counties in east Taiwan, whereas Taipei city had the lowest R_{OR} value. Our results also show that all R_{OS} were larger than R_{OR} in the four study sites. The persistence of both DS and DR TB (i.e. co-existence) occurs if $R_{OS} > 1$ and $R_{OS} > R_{OR}$. Under these conditions, co-existence can even occur when $R_{OR} < 1$ [3, 13].

The results for infection risk assessment show that the incidences of DS TB in Hwalien, Taitung, Pingtung counties, and Taipei city had nearly 17.6%, 15.5%, 14.5%, and 12.8% probabilities, respectively, of the total proportion of infected population exceeding 50%, whereas there was only ~3% probability of having exceeded 50% of the population infected attributable to MDR TB. Although MDR TB seems unlikely to result in an emergence, yet the case reproduction numbers of DS TB are alarming from a conservative point of view. As long as patients are carrying sensitive strains, there will always be some MDR TB cases, because MDR TB occurs from treatment failure, mutation at some constant frequency, and is transmitted occasionally for MDR strains.

Limitations and implications

The quality of the results depend on the input data. In our study, the limited datasets used in the two-strain TB model and regression model may pose the greatest sources of uncertainty. In the present study, the most unknown important parameter is RF of MDR strains. The MDR TB incidence rate of Mexico in 2008 was estimated to be 0.6/100 000 population (range 0.3–0.9/100 000) [35]. The average MDR TB incidence rate of Taiwan during 2007–2010 (0.7/100 000 population) was similar to Mexico. Thus, we estimated RF for MDR strains based on Garcia-Garcia *et al.* [18].

Several researchers used low values of RF for mathematical modelling of MDR TB such as values ranging from 0.7–1.0 [14] and 0.04–0.6 [19]. Cohen & Murray [15] assumed that the unfit MDR strains have a low fitness (0.3), whereas the fit MDR strain has RF ranging from 0.8–1.2. Burgos *et al.* [36] investigated the relative secondary-case ratio of MDR TB to DS TB based on genotype clustering method, indicating that there were no secondary cases associated with MDR strains. Therefore, MDR strains may

have lower transmissibility than DS strains, resulting in a low RF value.

Moreover, TB epidemiology values in the two-strain TB model are not easy to parameterize in Taiwan due to data limitation. To compensate for these shortcomings, we adopted published TB data [24, 25] together with our study results to improve model predictability. Although the data sources were adopted from England and Wales and The Netherlands, these parameter values could be used to examine the population dynamics of TB in different countries, such as Cambodia, China, India, Russia, South African, and Brazil [15, 37].

In addition to seasonality and meteorological factors, we did not consider age, gender, socioeconomic factors, and subpopulation effects at the local level, where data remain scarce, and has limited MDR TB burden and time-series data. In Taiwan, the incidence rate of TB rises with age; of all new patients, 53% were aged ≥ 65 years [17]. The subpopulation effects such as number of aborigines, alcoholics, smokers, and diabetics that have proven critical to TB trend variability [38]. Aborigines in east Taiwan are a subpopulation with high alcohol and cigarette consumption. Notably, the immune system is affected by age, co-infections, and past therapeutic history [39]. We thus anticipated that those factors may be incorporated into our models to improve their predictability and to reflect important dynamic consequences that are worth exploring in the future.

Our integrated-level model provided an opportunity to understand the complex interactions between seasonal dynamics of TB and environmental factors and could predict the seasonal patterns of TB incidence and estimate implicitly the MDR TB infection risk based on previous data information. Our Taiwan-based analysis could be applied to assess the DS/MDR TB infection potential in high-burden countries, where *M. tuberculosis* remains a substantial cause of morbidity and mortality.

A better understanding of the mechanisms of TB seasonality will help to develop more effective public control programmes. The quality of the local data allows us a rare opportunity to estimate the site-specific R_0 values that could be used to establish an early warning system. We could then evaluate the site-specific probabilistic infection risk that depends only on R_0 values. The practical implications of infection risk might be initiated for risk management. We suggest that more preventive implementation in TB control should be initiated during the period of higher

risk infection such as actively seeking TB cases in high-risk populations, accurately diagnosing TB and MDR TB, and preventing transmission in clinics and hospitals [40].

SUPPLEMENTARY MATERIAL

For supplementary material accompanying this paper visit <http://dx.doi.org/10.1017/S0950268813001040>.

ACKNOWLEDGEMENTS

The authors acknowledge the financial support of the National Science Council of the Republic of China under Grant NSC 100-2313-B-002-012-MY3.

DECLARATION OF INTEREST

None.

REFERENCES

1. **World Health Organization.** Global tuberculosis control 2012 (http://apps.who.int/iris/bitstream/10665/75938/1/9789241564502_eng.pdf). Accessed 22 October 2012.
2. **World Health Organization.** Multidrug and extensively drug-resistant TB (M/XDR-TB): 2010 global report on surveillance and response (http://whqlibdoc.who.int/publications/2010/9789241599191_eng.pdf). Accessed 20 April 2011.
3. **Dye C, et al.** Erasing the world's slow strain: strategies to beat multidrug-resistant tuberculosis. *Science* 2002; **295**: 2042–2046.
4. **Iseman MD.** Treatment of multidrug-resistant tuberculosis. *New England Journal of Medicine* 1993; **329**: 784–791.
5. **Shah NS, et al.** Worldwide emergence of extensively drug-resistant tuberculosis. *Emerging Infectious Diseases* 2007; **13**: 380–387.
6. **Centers for Disease Control, Department of Health, ROC (Taiwan).** Taiwan tuberculosis incidence and mortality rate, 2002–2008 (<http://www2.cdc.gov.tw/public/Data/9123117163971.pdf>). Accessed 6 August 2010.
7. **Centers for Disease Control, Department of Health, ROC. (Taiwan).** National notifiable disease surveillance system (<http://www.cdc.gov.tw/>). Accessed 19 August 2011.
8. **Luquero FJ, et al.** Trend and seasonality of tuberculosis in Spain, 1996–2004. *International Journal of Tuberculosis and Lung Disease* 2008; **12**: 221–224.
9. **Liu L, Zhao XQ, Zhou Y.** A tuberculosis model with seasonality. *Bulletin of Mathematical Biology* 2010; **72**: 931–952.
10. **Willis MD, et al.** Seasonality of tuberculosis in the United States, 1993–2008. *Clinical Infectious Diseases* 2012; **54**: 1553–1560.

11. **Leung CC, et al.** Seasonal pattern of tuberculosis in Hong Kong. *International Journal of Epidemiology* 2005; **34**: 924–930.
12. **Nagayama N, Ohmori M.** Seasonality in various forms of tuberculosis. *International Journal of Tuberculosis and Lung Disease* 2006; **10**: 1117–1122.
13. **Blower SM, Gerberding JL.** Understanding, predicting and controlling the emergence of drug-resistant tuberculosis: a theoretical framework. *Journal of Molecular Medicine* 1998; **76**: 624–636.
14. **Dye C, Williams BC.** Criteria for the control of drug-resistant tuberculosis. *Proceedings of the National Academy of Sciences USA* 2000; **97**: 8180–8185.
15. **Cohen T, Murray M.** Modeling epidemics of multidrug-resistant *M. tuberculosis* of heterogeneous fitness. *Nature Medicine* 2004; **10**: 1117–1121.
16. **Luciani F, et al.** The epidemiological fitness cost of drug resistance in *Mycobacterium tuberculosis*. *Proceedings of the National Academy of Sciences USA* 2009; **106**: 14711–14715.
17. **Centers for Disease Control, Department of Health, ROC. (Taiwan).** Taiwan Tuberculosis Control Report 2007, 2008, 2009, and 2010 (<http://www.cdc.gov.tw/>). Accessed 13 August 2010.
18. **García-García ML, et al.** Clinical consequences and transmissibility of drug-resistant tuberculosis in Southern Mexico. *Archives of Internal Medicine* 2000; **160**: 630–636.
19. **Dye C, Espinal MA.** Will tuberculosis become resistant to all antibiotics? *Proceedings of the Royal Society of London, Series B: Biological Sciences* 2001; **268**: 45–52.
20. **Rodrigues P, Gomes MG, Rebelo C.** Drug resistance in tuberculosis – a reinfection model. *Theoretical Population Biology* 2007; **71**: 196–212.
21. **van den Driessche P, Watmough J.** Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical Biosciences* 2002; **180**: 29–48.
22. **Anderson RM, May RM.** *Infectious Diseases of Humans: Dynamics and Control*. Oxford: Oxford University Press, 1991.
23. **Ministry of the Interior, ROC. (Taiwan).** Department of Statistics (<http://www.moi.gov.tw/stat/index.aspx>). Accessed 14 January 2010.
24. **Dye C, et al.** Prospects for worldwide tuberculosis control under the WHO DOTS strategy: directly observed short-course therapy. *Lancet* 1998; **352**: 1886–1891.
25. **Lin HH, et al.** Effects of smoking and solid-fuel use on COPD, lung cancer, and tuberculosis in China: a time-based, multiple risk factor, modelling study. *Lancet* 2008; **372**: 1473–1783.
26. **Yeh YP, et al.** Tuberculin reactivity in adults after 50 years of universal bacille Calmette-Guérin vaccination in Taiwan. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2005; **99**: 509–516.
27. **Hyndman RJ, Koehler AB.** Another look at measures of forecast accuracy. *International Journal of Forecasting* 2006; **22**: 679–688.
28. **Liao CM, et al.** Assessing trends and predictors of tuberculosis in Taiwan. *BMC Public Health* 2012; **12**: 29.
29. **Burnham KP, Anderson DR.** *Model Selection and Multi-model Inference: A Practical Information-Theoretic Approach*. New York: Springer, 2002.
30. **Greenman J, Kamo M, Boots M.** External forcing of ecological and epidemiological systems: a resonance approach. *Physica D: Nonlinear Phenomena* 2004; **190**: 136–151.
31. **Marais BJ, et al.** The clinical epidemiology of childhood pulmonary tuberculosis: a critical review of literature from the pre-chemotherapy era. *International Journal of Tuberculosis and Lung Disease* 2004; **8**: 278–285.
32. **Martineau AR, et al.** Vitamin D in the treatment of pulmonary tuberculosis. *Journal of Steroid Biochemistry and Molecular Biology* 2007; **103**: 793–798.
33. **Fares A.** Seasonality of tuberculosis. *Journal of Global Infectious Diseases* 2011; **3**: 46–55.
34. **Huang SH, et al.** Evolution of MDR-TB Control Strategy in Taiwan. *Taiwan Epidemiology Bulletin* 2012; **28**: 269–285.
35. **World Health Organization.** Towards universal access to diagnosis and treatment of multidrug-resistant and extensively drug-resistant tuberculosis by 2015. WHO progress report 2011 (http://whqlibdoc.who.int/publications/2011/9789241501330_eng.pdf). Accessed 2 April 2012.
36. **Burgos M, et al.** Effect of drug resistance on the generation of secondary cases of tuberculosis. *Journal of Infectious Diseases* 2003; **188**: 1878–1884.
37. **Legrand J, et al.** Modeling the impact of tuberculosis control strategies in highly endemic overcrowded prisons. *PLoS One* 2008; **7**: e2100.
38. **Hsueh PR, et al.** *Mycobacterium tuberculosis* in Taiwan. *Journal of Infection* 2006; **52**: 77–85.
39. **Keeler E, et al.** Reducing the global burden of tuberculosis: the contribution of improved diagnostics. *Nature* 2006; **444**: 49–57.
40. **Dye C, Williams BG.** The population dynamics and control of tuberculosis. *Science* 2010; **328**: 856–861.