



Unified bounds for the independence number of graphs

Jiang Zhou

Abstract. The Hoffman ratio bound, Lovász theta function, and Schrijver theta function are classical upper bounds for the independence number of graphs, which are useful in graph theory, extremal combinatorics, and information theory. By using generalized inverses and eigenvalues of graph matrices, we give bounds for independence sets and the independence number of graphs. Our bounds unify the Lovász theta function, Schrijver theta function, and Hoffman-type bounds, and we obtain the necessary and sufficient conditions of graphs attaining these bounds. Our work leads to some simple structural and spectral conditions for determining a maximum independent set, the independence number, the Shannon capacity, and the Lovász theta function of a graph.

1 Introduction

The *independence number* $\alpha(G)$ of graph G is the maximum size of independent sets in G , which is an important graph parameter in graph theory. The following is a celebrated upper bound given by Hoffman.

Theorem 1.1 [5, Theorem 3.5.2] *Let G be a k -regular ($k \neq 0$) graph with n vertices, and let τ be the minimum adjacency eigenvalue of G . Then*

$$\alpha(G) \leq n \frac{|\tau|}{k - \tau}.$$

If an independent set C meets this bound, then every vertex not in C is adjacent to exactly $|\tau|$ vertices of C .

The upper bound in Theorem 1.1 is often called Hoffman (ratio) bound for $\alpha(G)$. Haemers generalized the Hoffman bound to the general case.

Theorem 1.2 [17, p. 17] *Let G be a graph with minimum degree δ , and let λ and τ be the maximum and the minimum adjacency eigenvalue of G , respectively. Then*

$$\alpha(G) \leq n \frac{\lambda |\tau|}{\delta^2 - \lambda \tau}.$$

Received by the editors August 1, 2022; revised May 19, 2023; accepted December 4, 2023.

Published online on Cambridge Core December 11, 2023.

This work is supported by the National Natural Science Foundation of China (Grant No. 12071097) and the Natural Science Foundation for The Excellent Youth Scholars of the Heilongjiang Province (Grant No. YQ2022A002).

AMS subject classification: 05C69, 05C50, 94A15, 15A09.

Keywords: Independence number, graph matrix, Lovász theta function, Shannon capacity, generalized inverse, eigenvalue.



There are many Hoffman-type ratio bounds, which are useful in graph theory, coding theory, and extremal combinatorics [8–10, 12, 13]. Some generalized Hoffman ratio bounds involving adjacency eigenvalues, Laplacian eigenvalues, and normalized Laplacian eigenvalues of graphs can be found in [14, 17, 18, 20].

For a graph $G = (V, E)$, let G^k denote the graph whose vertex set is V^k , in which two vertices $u_1 \dots u_k$ and $v_1 \dots v_k$ are adjacent if and only if for each $i \in \{1, \dots, k\}$ either $u_i = v_i$ or $\{u_i, v_i\} \in E$ (see [1]). The *Shannon capacity* [5, 23] of G is defined as

$$\Theta(G) = \lim_{k \rightarrow \infty} \alpha(G^k)^{1/k} = \sup_{k \rightarrow \infty} \alpha(G^k)^{1/k},$$

and represents the number of distinct messages per use the channel can communicate with no error while used many times [1]. It is difficult to compute $\Theta(G)$ for a general graph G , we do not even know the Shannon capacity of the cycle C_7 (see [4, 21]).

The Lovász theta function $\vartheta(G)$ is an upper bound for $\alpha(G)$ introduced in [19], which is a powerful tool for studying the Shannon capacity of graphs. Lovász [19] proved that $\alpha(G) \leq \Theta(G) \leq \vartheta(G)$, and posed the problem of finding graphs with $\Theta(G) = \vartheta(G)$ (see Problem 1 in [19]). The Schrijver theta function $\vartheta'(G)$ is a smaller upper bound for $\alpha(G)$. Schrijver [22] proved that $\alpha(G) \leq \vartheta'(G) \leq \vartheta(G)$. By considering the relation between $\alpha(G)$ and the rank of matrices associated with G , Haemers [16] proved that $\Theta(G)$ is at most the minimum rank of a class of graph matrices.

Generalized inverses of graph matrices have important applications in random walks on graphs [6] and the sparsification of graphs [24]. This paper uses generalized inverses methods to study $\alpha(G)$, $\Theta(G)$, $\vartheta(G)$, and $\vartheta'(G)$, and gives bounds which unify the Lovász theta function, Schrijver theta function, and Hoffman-type bounds. Based on the generalized inverses method, we obtain simple structural and spectral conditions to determine $\alpha(G)$, $\Theta(G)$, $\vartheta(G)$, and a maximum independent set in G . Our conditions can also be used to find graphs with $\Theta(G) = \vartheta(G)$, which is a partial answer to the problem posed by Lovász.

The paper is organized as follows: In Section 2, we introduce some auxiliary lemmas. In Section 3, we give bounds for the sizes of independent sets and the independence number of graphs by using generalized inverses and eigenvalues of graph matrices. In Sections 4–6, we show that our bounds unify the Lovász theta function, Schrijver theta function, and Hoffman-type bounds in [5, 14, 18, 20], and obtain the necessary and sufficient conditions of graphs attaining the bounds. In Section 7, we give some simple structural and spectral conditions for determining a maximum independent set, the independence number, the Shannon capacity, and the Lovász theta function of a graph. Some concluding remarks are given in Section 8.

2 Preliminaries

For a real square matrix M , the *group inverse* of M , denoted by $M^\#$, is the matrix X such that $MXM = M$, $XXM = X$, and $MX = XM$. It is known [2, p. 156] that $M^\#$ exists if and only if $\text{rank}(M) = \text{rank}(M^2)$. If $M^\#$ exists, then $M^\#$ is unique.

For a positive semidefinite real matrix M , there exists an orthogonal matrix U such that

$$M = U \text{diag}(\lambda_1, \dots, \lambda_n) U^\top,$$

where $\text{diag}(\lambda_1, \dots, \lambda_n)$ denotes a diagonal matrix with nonnegative diagonal entries $\lambda_1, \dots, \lambda_n$. We set $M^{\frac{1}{2}} = U\text{diag}(\lambda_1^{\frac{1}{2}}, \dots, \lambda_n^{\frac{1}{2}})U^T$ and $M^{-\frac{1}{2}} = U\text{diag}((\lambda_1^+)^{\frac{1}{2}}, \dots, (\lambda_n^+)^{\frac{1}{2}})U^T$, where $\lambda_i^+ = \lambda_i^{-1}$ if $\lambda_i > 0$, and $\lambda_i^+ = 0$ if $\lambda_i = 0$. Then

$$M^\# = U\text{diag}(\lambda_1^+, \dots, \lambda_n^+)U^T = (M^{-\frac{1}{2}})^2.$$

Lemma 2.1 [2, pp. 162–163] *Let M be a square matrix such that $M^\#$ exists, and let $\lambda \neq 0$ be an eigenvalue of M with an eigenvector x . Then*

$$M^\#x = \lambda^{-1}x.$$

For a real matrix A , the *Moore–Penrose inverse* of A is the real matrix X such that $AXA = A$, $XAX = X$, $(AX)^T = AX$, and $(XA)^T = XA$. Let A^+ denote the Moore–Penrose inverse of A . It is well known that A^+ exists and is unique.

Let $R(M) = \{x : x = My, y \in \mathbb{R}^n\}$ denote the range of an $m \times n$ real matrix M .

Lemma 2.2 [2, pp. 43, 49] *Let A be a real matrix. Then*

$$\begin{aligned} R(AA^T) &= R(A) = R(AA^+), \\ (A^T A)^\# A^T &= A^+. \end{aligned}$$

Lemma 2.3 *Let A be an $n \times m$ real matrix, and let x be a unit real vector of dimension n . Then*

$$x^T AA^+ x \leq 1,$$

with equality if and only if $x \in R(A)$.

Proof It is known [2] that AA^+ is a real symmetric idempotent matrix. So each eigenvalue of AA^+ is 0 or 1. For a unit real vector x , we have $x^T AA^+ x \leq 1$, with equality if and only if $x \in R(AA^+)$. By Lemma 2.2, $x \in R(AA^+)$ is equivalent to $x \in R(A)$. ■

We now introduce the Lovász theta function defined in [19]. An *orthonormal representation* of an n -vertex graph G is a set $\{u_1, \dots, u_n\}$ of unit real vectors such that $u_i^T u_j = 0$ if i and j are two nonadjacent vertices in G . The *value* of an orthonormal representation $\{u_1, \dots, u_n\}$ is defined to be

$$\min_c \max_{1 \leq i \leq n} \frac{1}{(c^T u_i)^2},$$

where c ranges over all unit real vectors. The *Lovász theta function* $\vartheta(G)$ is the minimum value of all orthonormal representations of G .

Let $(M)_{ij}$ denote the (i, j) -entry of a matrix M , and let $\lambda_1(A)$ denote the largest eigenvalue of a real symmetric matrix A . The independence number $\alpha(G)$, the Shannon capacity $\Theta(G)$, and the Lovász theta function $\vartheta(G)$ have the following relations.

Lemma 2.4 [19] *For any graph G , we have*

$$\alpha(G) \leq \Theta(G) \leq \vartheta(G) = \min_A \lambda_1(A),$$

where A ranges over all real symmetric matrices indexed by vertices of G such that $(A)_{ij} = 1$ when $i = j$ or i, j are two nonadjacent vertices.

For a graph G , its *Schrijver theta function* $\vartheta'(G)$ is defined as [22]

$$\vartheta'(G) = \min_A \lambda_1(A),$$

where A ranges over all real symmetric matrices indexed by vertices of G such that $(A)_{ij} \geq 1$ when $i = j$ or i, j are two nonadjacent vertices.

Lemma 2.5 [22] *For any graph G , we have*

$$\alpha(G) \leq \vartheta'(G).$$

For an n -vertex graph G , the *adjacency matrix* A of G is an $n \times n$ symmetric matrix with entries

$$(A)_{ij} = \begin{cases} 1, & \text{if } \{i, j\} \in E(G), \\ 0, & \text{if } \{i, j\} \notin E(G), \end{cases}$$

where $E(G)$ denotes the edge set of G . Eigenvalues of the adjacency matrix of G are called *adjacency eigenvalues* of G . The largest adjacency eigenvalue of a graph has the following upper bound.

The following lemma will be used later in the proof of Example 4.3.

Lemma 2.6 [3] *Let G be a graph with adjacency matrix A . Then*

$$\lambda_1(A) \leq \max_{\{u,v\} \in E(G)} \sqrt{d_u d_v},$$

where d_u denotes the degree of a vertex u .

3 Bounds for independent sets and independence number

Let $A[S]$ denote the principal submatrix of a square matrix A determined by the rows and columns whose index set is S . A real vector $x = (x_1, \dots, x_n)^\top$ is called *total nonzero* if $x_i \neq 0$ for $i = 1, \dots, n$.

For an n -vertex graph G , let $V(G)$ denote the vertex set of G , and let $\mathcal{P}(G)$ denote the set of real matrix-vector pairs (M, x) such that:

- (a) M is a positive semidefinite $n \times n$ matrix indexed by vertices of G .
- (b) $x = (x_1, \dots, x_n)^\top \in R(M)$ is total nonzero and $(M)_{ij}x_i x_j \leq 0$ if i, j are two nonadjacent distinct vertices.

For $(M, x) \in \mathcal{P}(G)$ and vertex subset $T \subseteq V(G)$, let $F(M, x)$ and $F_T(M, x)$ denote the following functions:

$$F(M, x) = x^\top M^\# x \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2},$$

$$F_T(M, x) = \frac{x^\top M^\# x}{|T|} \sum_{u \in T} \frac{(M)_{uu}}{x_u^2}.$$

In [14], Godsil and Newman gave bounds on the size of an independent set by using positive semidefinite graph matrices. Motivated by the ideas of Godsil and Newman, we give the following bounds in terms of generalized inverses of positive semidefinite graph matrices.

Theorem 3.1 Let $(M, x) \in \mathcal{P}(G)$ be a matrix-vector pair associated with an n -vertex graph G . Then the following statements hold:

(1) For any independent set S of G , we have

$$|S| \leq F_S(M, x),$$

with equality if and only if $M[S]$ is diagonal and there exists a constant c such that

$$\begin{aligned} \frac{(M)_{vv}}{x_v^2} &= c \quad (v \in S), \\ cx_v &= \sum_{u \in S} (M)_{vu} x_u^{-1} \quad (v \notin S). \end{aligned}$$

(2) The independence number $\alpha(G)$ satisfies

$$\alpha(G) \leq F(M, x),$$

with equality if and only if G has an independent set C such that $M[C]$ is diagonal and

$$\begin{aligned} \frac{(M)_{vv}}{x_v^2} &= \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2} = c \quad (v \in C), \\ cx_v &= \sum_{u \in C} (M)_{vu} x_u^{-1} \quad (v \notin C). \end{aligned}$$

Moreover, if an independent set C satisfies the above conditions, then

$$\alpha(G) = |C| = F(M, x).$$

(3) Suppose that $\frac{(M)_{11}}{x_1^2} \geq \dots \geq \frac{(M)_{nn}}{x_n^2}$ are arranged in a decreasing order. If $\alpha(G) \geq t$ for some positive integer t , then

$$\alpha(G) \leq F_T(M, x),$$

where $T = \{1, \dots, t\}$.

Proof Since $M^{\frac{1}{2}} M^{-\frac{1}{2}} M = M$ and $x \in R(M)$, we have

$$M^{\frac{1}{2}} M^{-\frac{1}{2}} x = x.$$

For any independent set S of G , let $y = (y_1, \dots, y_n)^T$ be the vector such that

$$y_u = \begin{cases} x_u^{-1}, & \text{if } u \in S, \\ 0, & \text{if } u \notin S. \end{cases}$$

By the Cauchy–Schwarz inequality, we have

$$|S|^2 = (y^T x)^2 = (y^T M^{\frac{1}{2}} M^{-\frac{1}{2}} x)^2 \leq (y^T M y)(x^T M^{\#} x).$$

Since $(M)_{ij} x_i x_j \leq 0$ when i, j are two nonadjacent vertices, we get

$$\begin{aligned} |S|^2 &\leq (y^T M y)(x^T M^{\#} x) \leq x^T M^{\#} x \sum_{u \in S} \frac{(M)_{uu}}{x_u^2}, \\ |S| &\leq F_S(M, x), \end{aligned}$$

with equality if and only if $M[S]$ is diagonal and there exists a constant c such that $M^{\frac{1}{2}}y = cM^{-\frac{1}{2}}x$, that is, $My = cx$ (because $M^{\frac{1}{2}}M^{-\frac{1}{2}}x = x$). When $M[S]$ is diagonal, $My = cx$ is equivalent to

$$\begin{aligned} \frac{(M)_{vv}}{x_v^2} &= c \quad (v \in S), \\ cx_v &= \sum_{u \in S} (M)_{vu}x_u^{-1} \quad (v \notin S). \end{aligned}$$

So part (1) holds.

Notice that $|S| \leq F_S(M, x) \leq F(M, x)$ for any independent set S of G . Hence

$$\alpha(G) \leq F(M, x).$$

If the equality holds, then G has an independent set C such that $M[C]$ is diagonal and

$$\begin{aligned} \frac{(M)_{vv}}{x_v^2} &= \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2} = c \quad (v \in C), \\ cx_v &= \sum_{u \in C} (M)_{vu}x_u^{-1} \quad (v \notin C). \end{aligned}$$

If an independent set C satisfies the above conditions, then

$$\begin{aligned} |C|^2 &= (z^\top x)^2 = (z^\top M^{\frac{1}{2}}M^{-\frac{1}{2}}x)^2 = (z^\top Mz)(x^\top M^\#x) = |C|F(M, x), \\ |C| &= F(M, x), \end{aligned}$$

where $z = (z_1, \dots, z_n)^\top$ is the vector such that

$$z_u = \begin{cases} x_u^{-1}, & \text{if } u \in C, \\ 0, & \text{if } u \notin C. \end{cases}$$

Since $\alpha(G) \leq F(M, x)$, we have

$$|C| = \alpha(G) = F(M, x).$$

So part (2) holds.

Suppose that $\frac{(M)_{11}}{x_1^2} \geq \dots \geq \frac{(M)_{nn}}{x_n^2}$ are arranged in a decreasing order. Let C be an independent set such that $|C| = \alpha(G)$. If $\alpha(G) \geq t$, then

$$\alpha(G) \leq F_C(M, x) \leq F_T(M, x) \quad (T = \{1, \dots, t\}).$$

So part (3) holds. ■

Remark 3.2 For a graph G , let M be a positive definite matrix indexed by vertices of G such that $(M)_{ij} = 0$ if i, j are two nonadjacent distinct vertices. Then $M^\# = M^{-1}$ and $(M, x) \in \mathcal{P}(G)$ for any real vector x . By part (2) of Theorem 3.1, we have

$$\alpha(G) \leq F(M, x) = x^\top M^{-1}x \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2}$$

for any total nonzero vector x .

The $\{1\}$ -inverse of M is a matrix X such that $MXM = M$. If M is singular, then it has infinite $\{1\}$ -inverses (see [2, page 41]).

Remark 3.3 Let M be a real symmetric matrix. Since $MM^\#M = MM^{(1)}M = M$ for any $\{1\}$ -inverse $M^{(1)}$ of M , we have $x^\top M^\#x = x^\top M^{(1)}x$ for any $x \in R(M)$. Hence, the upper bounds given in Theorem 3.1 can be obtained from any $\{1\}$ -inverse of M .

The *signless Laplacian matrix* of a graph G is defined as $D + A$, where D is the diagonal matrix of vertex degrees of G , A is the adjacency matrix of G . The *subdivision graph* of G , denoted by $S(G)$, is a graph obtained from G by replacing each edge of G by a path of length 2.

We can find concrete examples such that an upper bound derived from Theorem 3.1 is smaller than the Hoffman bound given in Theorem 1.2.

Example 3.4 Let G be a nonregular graph with n vertices, m edges, and minimum degree $\delta \geq 2$. Then there exists $(M, x) \in \mathcal{P}(S(G))$ such that

$$\alpha(S(G)) = x^\top M^\#x \max_{u \in V(S(G))} \frac{(M)_{uu}}{x_u^2} < \frac{(m+n)\lambda^2}{4+\lambda^2},$$

where λ is the maximum adjacency eigenvalue of $S(G)$.

Proof The signless Laplacian matrix of $S(G)$ can be written as $M = \begin{pmatrix} 2I & B^\top \\ B & D \end{pmatrix}$, where B is the vertex-edge incidence matrix of G , D is the diagonal matrix of vertex degrees of G . Let $x = \begin{pmatrix} 2I & B^\top \\ B & D \end{pmatrix} \begin{pmatrix} e \\ 0 \end{pmatrix} = \begin{pmatrix} 2e \\ y \end{pmatrix}$, where e is the all-ones vector, y is the vector of vertex degrees of G . Let C be the independent set in $S(G)$ corresponding to the edge set of G , then $(M, x) \in \mathcal{P}(S(G))$ and C satisfy the conditions given in part (2) of Theorem 3.1. By part (2) of Theorem 3.1, we get

$$\alpha(S(G)) = x^\top M^\#x \max_{u \in V(S(G))} \frac{(M)_{uu}}{x_u^2} = |C| = m.$$

Let λ be the maximum adjacency eigenvalue of $S(G)$, then $-\lambda$ is the minimum adjacency eigenvalue of $S(G)$ (because $S(G)$ is bipartite). By Theorem 1.2, we have

$$\alpha(S(G)) = x^\top M^\#x \max_{u \in V(S(G))} \frac{(M)_{uu}}{x_u^2} = m \leq \frac{(m+n)\lambda^2}{4+\lambda^2}.$$

We next show that the inequality is strict. It is known that λ^2 is the maximum eigenvalue of the signless Laplacian matrix of G , and $\lambda^2 \geq \frac{4m}{n}$, with equality if and only if G is regular (see [7, Theorems 2.4.4 and 7.8.6]). Since G is nonregular, we have $\lambda^2 > \frac{4m}{n}$, that is $m < \frac{(m+n)\lambda^2}{4+\lambda^2}$. ■

For a connected graph G , if M be a nonnegative matrix indexed by $V(G)$ such that $(M)_{ij} > 0$ if and only if i, j are two adjacent vertices, then M is irreducible. By the Perron–Frobenius Theorem, the spectral radius ρ of M is a positive eigenvalue of M , and there exists unique positive unit eigenvector associated with ρ . The following eigenvalue bound follows from Theorem 3.1.

Corollary 3.5 *Let M be a positive semidefinite nonnegative matrix associated with a connected graph G , and $(M)_{ij} > 0$ if and only if i, j are two adjacent vertices. Let ρ be the spectral radius of M , and let x be the corresponding positive unit eigenvector. Then*

$$\alpha(G) \leq \rho^{-1} \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2}.$$

Proof Since $Mx = \rho x$, we have $(M, x) \in \mathcal{P}(G)$. By Lemma 2.1, we get

$$M^\# x = \rho^{-1} x.$$

By part (2) of Theorem 3.1, we get

$$\alpha(G) \leq F(M, x) = x^\top M^\# x \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2} = \rho^{-1} \max_{u \in V(G)} \frac{(M)_{uu}}{x_u^2}. \quad \blacksquare$$

The Laplacian matrix of a graph G is defined as $L = D - A$, where D is the diagonal matrix of vertex degrees of G , A is the adjacency matrix of G . It is well known that L is positive semidefinite. The following result follows from Theorem 3.1.

Corollary 3.6 *Let G be a connected graph with Laplacian matrix L . For any total nonzero vector x satisfying $\sum_{u \in V(G)} x_u = 0$, we have*

$$\alpha(G) \leq F(L, x).$$

Proof Since G is connected, the null space of L has dimension one. Since $Le = 0$ for the all-ones vector e , we have $x \in R(L)$ for any vector x satisfying $e^\top x = \sum_{u \in V(G)} x_u = 0$. By part (2) of Theorem 3.1, we get

$$\alpha(G) \leq F(L, x). \quad \blacksquare$$

4 Lovász theta function and Shannon capacity

For a graph G , let $\mathcal{M}(G)$ denote the set of real matrix-vector pairs (M, x) such that:

(a) M is a positive semidefinite matrix indexed by vertices of G such that $(M)_{ij} = 0$ if i, j are two nonadjacent distinct vertices.

(b) $x \in R(M)$ is a total nonzero vector.

Since $\mathcal{M}(G)$ is a subset of $\mathcal{P}(G)$, the notations $F(M, x)$ and $F_T(M, x)$ defined in Section 3 can be used for $(M, x) \in \mathcal{M}(G)$ directly.

In the following theorem, we show that the Lovász theta function $\vartheta(G)$ is a special upper bound given in Theorem 3.1, and obtain the necessary and sufficient conditions of graphs satisfying $\alpha(G) = \vartheta(G)$.

Theorem 4.1 *For any graph G , we have*

$$\alpha(G) \leq \min_{(M, x) \in \mathcal{M}(G)} F(M, x) = \vartheta(G),$$

with equality if and only if there exist $(M, x) \in \mathcal{M}(G)$ and an independent set C satisfying the conditions given in part (2) of Theorem 3.1.

Proof By Theorem 3.1, we have

$$\alpha(G) \leq \min_{(M,x) \in \mathcal{M}(G)} F(M, x),$$

with equality if and only if there exist $(M, x) \in \mathcal{M}(G)$ and an independent set C satisfying the conditions in part (2) of Theorem 3.1. We need to prove that the minimum value equals to $\vartheta(G)$.

We first prove that the minimum value does not exceed $\vartheta(G)$. Let c, u_1, \dots, u_n be unit real vectors such that $\{u_1, \dots, u_n\}$ is an orthonormal representation of G and

$$\vartheta(G) = \max_{1 \leq i \leq n} \frac{1}{(c^\top u_i)^2}.$$

Let U be the matrix whose i th column is u_i , then each diagonal entry of $U^\top U$ is 1. Take $z = U^\top c$, then $(U^\top U, z) \in \mathcal{M}(G)$ and

$$z_i = c^\top u_i \neq 0, \quad i = 1, \dots, n.$$

By Lemmas 2.2 and 2.3, we have

$$F(U^\top U, z) = z^\top (U^\top U)^\# z \max_{1 \leq i \leq n} \frac{1}{z_i^2} = \vartheta(G) c^\top U U^+ c \leq \vartheta(G).$$

So we have

$$\min_{(M,x) \in \mathcal{M}(G)} F(M, x) \leq F(U^\top U, z) \leq \vartheta(G).$$

We turn to prove that $\min_{(M,x) \in \mathcal{M}(G)} F(M, x)$ is at least $\vartheta(G)$. There exists $(M_0, y) \in \mathcal{M}(G)$ such that

$$F(M_0, y) = y^\top (M_0)^\# y \max_{1 \leq i \leq n} \frac{(M_0)_{ii}}{y_i^2} = \min_{(M,x) \in \mathcal{M}(G)} F(M, x).$$

Since M_0 is positive semidefinite, there exists real matrix B such that $M_0 = B^\top B$. Since $y \in R(M_0)$, we can choose y such that $y = B^\top c$ for a unit real vector $c \in R(B)$. Let b_i be the i th column of B . By Lemmas 2.2 and 2.3, we have

$$F(M_0, y) = y^\top (M_0)^\# y \max_{1 \leq i \leq n} \frac{(M_0)_{ii}}{y_i^2} = c^\top B B^+ c \max_{1 \leq i \leq n} \frac{b_i^\top b_i}{(c^\top b_i)^2} = \max_{1 \leq i \leq n} \frac{b_i^\top b_i}{(c^\top b_i)^2}.$$

We next prove that $F(M_0, y) \geq \vartheta(G)$ by using the method similar with proof of [19, Theorem 3]. Let $A = (a_{ij})_{n \times n}$ be the matrix with entries

$$\begin{aligned} a_{ii} &= 1, \quad i = 1, \dots, n, \\ a_{ij} &= 1 - \frac{b_i^\top b_j}{(c^\top b_i)(c^\top b_j)} = 1 - \frac{(M_0)_{ij}}{y_i y_j}, \quad i \neq j. \end{aligned}$$

By Lemma 2.4, we get

$$\lambda_1(A) \geq \vartheta(G).$$

We set

$$\beta = F(M_0, y) = \max_{1 \leq i \leq n} \frac{b_i^\top b_i}{(c^\top b_i)^2},$$

then

$$-a_{ij} = \left(c - \frac{b_i}{(c^\top b_i)} \right)^\top \left(c - \frac{b_j}{(c^\top b_j)} \right), \quad i \neq j,$$

$$\beta - a_{ii} = \left(c - \frac{b_i}{(c^\top b_i)} \right)^\top \left(c - \frac{b_i}{(c^\top b_i)} \right) + \beta - \frac{b_i^\top b_i}{(c^\top b_i)^2}.$$

Hence, $\beta I - A$ (I is the identity matrix) is positive semidefinite, that is,

$$\min_{(M,x) \in \mathcal{M}(G)} F(M, x) = \beta \geq \lambda_1(A) \geq \vartheta(G). \quad \blacksquare$$

The following conclusion comes from Theorems 3.1 and 4.1 and Lemma 2.4.

Theorem 4.2 *Let $(M, x) \in \mathcal{M}(G)$ be a matrix-vector pair for graph G . If G has an independent set C such that (M, x) and C satisfy the conditions given in part (2) of Theorem 3.1, then*

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = F(M, x).$$

A bipartite graph G is called *semiregular* with parameters (n_1, n_2, r_1, r_2) if $V(G)$ has a bipartition $V(G) = V_1 \cup V_2$ such that $|V_1| = n_1, |V_2| = n_2$ and vertices in the same color class have the same degree (n_i vertices of degree $r_i, i = 1, 2$).

Let I denote the identity matrix. The following are some examples for Theorem 4.2.

Example 4.3 *Let G be a graph with a spanning subgraph H , and let H be a semiregular bipartite graph with parameters (n_1, n_2, r_1, r_2) ($n_1 r_1 = n_2 r_2 > 0, n_1 \leq n_2$). If the independent set with n_2 vertices in H is also an independent set in G , then*

$$\alpha(G) = \Theta(G) = \vartheta(G) = n_2.$$

Proof The adjacency matrix of H can be written as $A = \begin{pmatrix} 0 & B^\top \\ B & 0 \end{pmatrix}$, and each row (column) sum of $B \in \mathbb{R}^{n_1 \times n_2}$ is r_1 (r_2). By Lemma 2.6, we have $\lambda_1(A) \leq \sqrt{r_1 r_2}$. Then

$$\lambda_1(BB^\top) = \lambda_1(A^2) \leq r_1 r_2.$$

Let $M = \begin{pmatrix} r_1 I & B^\top \\ B & r_2 I \end{pmatrix}$, then the all-ones vector $e \in R(M)$ because $Me = (r_1 + r_2)e$. Since the Schur complement

$$r_2 I - r_1^{-1} B B^\top = r_1^{-1} (r_1 r_2 I - B B^\top)$$

is positive semidefinite, M is positive semidefinite. Hence $(M, e) \in \mathcal{M}(G)$. Let C be the independent set with n_2 vertices in H , which is also an independent set in G . Since (M, e) and C satisfy the conditions given in part (2) of Theorem 3.1, then by Theorem 4.2, we have

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = \alpha(H). \quad \blacksquare$$

For two disjoint graphs G_1 and G_2 , let $G_1 \vee G_2$ denote the graph obtained from the union of G_1 and G_2 by joining each vertex of G_1 to each vertex of G_2 . Let \overline{G} denote the complement of a graph G .

Example 4.4 Let G be an n -vertex graph with minimum degree δ . For any integer $s \geq n - \delta$, we have

$$\alpha(G \vee \overline{K_s}) = \Theta(G \vee \overline{K_s}) = \vartheta(G \vee \overline{K_s}) = s,$$

where K_s is the complete graph with s vertices.

Proof The adjacency matrix of $G \vee \overline{K_s}$ can be written as $\begin{pmatrix} 0 & J_{s \times n} \\ J_{s \times n}^\top & A \end{pmatrix}$, where A is the adjacency matrix of G , $J_{s \times n}$ is the $s \times n$ all-ones matrix. Let L be the Laplacian matrix of G , and let $M = \begin{pmatrix} sI & J_{s \times n} \\ J_{s \times n}^\top & nI - L \end{pmatrix}$. Then the all-ones vector $e \in R(M)$ because $Me = (s + n)e$. Since the Schur complement

$$nI - L - s^{-1}J_{s \times n}^\top J_{s \times n}$$

equals to the Laplacian matrix of \overline{G} (which is positive semidefinite), M is positive semidefinite. Hence $(M, e) \in \mathcal{M}(G)$. Let C be the independent set of size s in $G \vee \overline{K_s}$. Since (M, e) and C satisfy the conditions given in part (2) of Theorem 3.1, then by Theorem 4.2, we have

$$\alpha(G \vee \overline{K_s}) = \Theta(G \vee \overline{K_s}) = \vartheta(G \vee \overline{K_s}) = |C| = s. \quad \blacksquare$$

Let $e = (1, \dots, 1)^\top$ denote the all-ones vector.

Corollary 4.5 Let G be a graph with adjacency matrix A and $A + \lambda I$ is positive definite for some $\lambda > 0$. Then

$$\alpha(G) \leq \lambda e^\top (A + \lambda I)^{-1} e,$$

with equality if and only if there exists an independent set C such that every vertex not in C is adjacent to exactly λ vertices of C . Moreover, if an independent set C satisfies the above condition, then

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = \lambda e^\top (A + \lambda I)^{-1} e.$$

Proof Since $A + \lambda I$ is positive definite, we have $(A + \lambda I, e) \in \mathcal{M}(G)$ and $F(A + \lambda I, e) = \lambda e^\top (A + \lambda I)^{-1} e$. The conclusions follow from Theorems 3.1 and 4.2. \blacksquare

The following is an example for Corollary 4.5.

Example 4.6 For the path P_{2k+1} with $2k + 1$ vertices, let C be the independent set in P_{2k+1} with $k + 1$ vertices. Then every vertex not in C is adjacent to exactly two vertices of C . By Corollary 4.5, we have

$$\alpha(P_{2k+1}) = \Theta(P_{2k+1}) = \vartheta(P_{2k+1}) = |C| = 2e^\top (A + 2I)^{-1} e = k + 1,$$

where A is the adjacency matrix of P_{2k+1} .

A *clique covering* of G is a set $\varepsilon = \{Q_1, \dots, Q_r\}$ of cliques in G that cover all edges of G , that is, each edge of G belongs to at least one Q_i . For a clique covering ε and a vertex u of G , the ε -*degree* of u is the number of cliques in ε containing u , denoted by d_u^ε . The *clique covering matrix* A_ε is the $n \times n$ symmetric matrix with entries

$$(A_\varepsilon)_{ij} = \begin{cases} d_i^\varepsilon, & \text{if } i = j, \\ w_{ij}, & \text{if } \{i, j\} \in E(G), \\ 0, & \text{if } \{i, j\} \notin E(G), \end{cases}$$

where w_{ij} is the number of cliques in ε containing the edge $\{i, j\}$.

Theorem 4.7 *Let G be a graph without isolated vertices. For any clique covering ε of G , we have*

$$\alpha(G) \leq F(A_\varepsilon, x),$$

where x is a vector satisfying $x_u = d_u^\varepsilon$. The equality holds if and only if there exists an independent set C such that $d_v^\varepsilon = \min_{u \in V(G)} d_u^\varepsilon$ for each $v \in C$, and $|C \cap Q| = 1$ for each $Q \in \varepsilon$. Moreover, if an independent set C satisfies the above conditions, then

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = F(A_\varepsilon, x).$$

Proof For a clique covering $\varepsilon = \{Q_1, \dots, Q_r\}$ of G , the corresponding vertex-clique incidence matrix B is a $|V(G)| \times r$ matrix with entries

$$(B)_{ij} = \begin{cases} 1, & \text{if } i \in V(G), i \in Q_j, \\ 0, & \text{if } i \in V(G), i \notin Q_j. \end{cases}$$

Then $BB^T = A_\varepsilon$. Notice that $x = Be \in R(B) = R(A_\varepsilon)$, so $(A_\varepsilon, x) \in \mathcal{M}(G)$. The conclusions follow from Theorems 3.1 and 4.2. ■

The following is an example for Theorem 4.7.

Example 4.8 The graph G in Figure 1 has a clique covering $\varepsilon = \{Q_1, Q_2, Q_3\}$, where $Q_1 = \{1, 2, 3\}$, $Q_2 = \{2, 4, 5\}$, $Q_3 = \{3, 5, 6\}$. Then G has ε -degrees $d_1^\varepsilon = 1$, $d_2^\varepsilon = 2$, $d_3^\varepsilon = 2$, $d_4^\varepsilon = 1$, $d_5^\varepsilon = 2$, $d_6^\varepsilon = 1$ (see [25]). The independent set $C = \{1, 4, 6\}$ satisfies the conditions in Theorem 4.7. So we have

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = 3.$$

Let $m(H)$ denote the matching number of a hypergraph H . The *intersection graph* $\Omega(H)$ of H has vertex set $E(H)$, and two vertices f_1, f_2 of $\Omega(H)$ are adjacent if and only if $f_1 \cap f_2 \neq \emptyset$. Clearly, we have $m(H) = \alpha(\Omega(H))$. The vertex-edge incidence matrix B of H is a $|V(H)| \times |E(H)|$ matrix with entries

$$(B)_{uf} = \begin{cases} 1, & \text{if } u \in V(H), u \in f \in E(H), \\ 0, & \text{if } u \in V(H), u \notin f \in E(H). \end{cases}$$

We say that H is k -uniform if each edge of H has exactly k vertices.




Figure 1: Graph G.

Theorem 4.9 Let H be a k -uniform hypergraph without isolated vertices, and let B be the vertex-edge incidence matrix of H . Then

$$m(H) \leq ke^T (B^T B)^{\#} e,$$

with equality if and only if H has a perfect matching. If H has a perfect matching, then

$$m(H) = \Theta(\Omega(H)) = \vartheta(\Omega(H)) = ke^T (B^T B)^{\#} e.$$

Proof Notice that $e = k^{-1}B^T e \in R(B^T) = R(B^T B)$, so $(B^T B, e) \in \mathcal{M}(\Omega(H))$ and $F(B^T B, e) = ke^T (B^T B)^{\#} e$. The conclusions follow from Theorems 3.1 and 4.2. ■

5 Schrijver theta function

In the following theorem, we show that the Schrijver theta function $\vartheta(G)$ is the minimum upper bound given in part (2) of Theorem 3.1, and obtain the necessary and sufficient conditions of graphs satisfying $\alpha(G) = \vartheta'(G)$.

Theorem 5.1 For any graph G , we have

$$\alpha(G) \leq \min_{(M,x) \in \mathcal{P}(G)} F(M, x) = \vartheta'(G),$$

with equality if and only if there exist $(M, x) \in \mathcal{P}(G)$ and an independent set C satisfying the conditions given in part (2) of Theorem 3.1.

Proof By Theorem 3.1, we have

$$\alpha(G) \leq \min_{(M,x) \in \mathcal{P}(G)} F(M, x),$$

with equality if and only if there exist $(M, x) \in \mathcal{P}(G)$ and an independent set C satisfying the conditions given in part (2) of Theorem 3.1. We will prove that the minimum value equals to $\vartheta'(G)$ by using the method similar with proof of [19, Theorem 3].

We first prove that the minimum value does not exceed $\vartheta'(G)$. Let $A = (a_{ij})_{n \times n}$ be the real symmetric matrix indexed by vertices of G such that $\lambda_1(A) = \vartheta'(G)$, and $a_{ij} \geq 1$ when $i = j$ or i, j are two nonadjacent vertices. Then $\vartheta'(G)I - A$ is positive semidefinite. So there exists real matrix T such that

$$\vartheta'(G)I - A = T^T T$$

and the rank of T is smaller than the number of rows of T . Let c be a unit real vector such that $c^T T = 0$. Suppose that t_i is the i th column of T , then

$$t_i^T t_j = \vartheta'(G)\delta_{ij} - a_{ij},$$

where $\delta_{ij} = 1$ if $i = j$, and $\delta_{ij} = 0$ if $i \neq j$. We set

$$u_i = \vartheta'(G)^{-\frac{1}{2}}(c + t_i), \quad i = 1, \dots, n,$$

and let U be the matrix whose i th column is u_i . Then

$$\begin{aligned} (U^T U)_{ii} &= \vartheta'(G)^{-1}(1 + \vartheta'(G) - a_{ii}) \leq 1, \quad i \in V(G), \\ (U^T U)_{ij} &= \vartheta'(G)^{-1}(1 - a_{ij}) \leq 0, \quad \{i, j\} \notin E(G). \end{aligned}$$

Take $z = U^T c$, then $z_i = c^T u_i = \vartheta'(G)^{-\frac{1}{2}}$ ($i = 1, \dots, n$) and $(U^T U, z) \in \mathcal{P}(G)$. By Lemmas 2.2 and 2.3, we have

$$F(U^T U, z) = z^T (U^T U)^\# z \max_{1 \leq i \leq n} \frac{(U^T U)_{ii}}{z_i^2} \leq \vartheta'(G)c^T U U^+ c \leq \vartheta'(G).$$

So we have

$$\min_{(M,x) \in \mathcal{P}(G)} F(M, x) \leq F(U^T U, z) \leq \vartheta'(G).$$

We next prove that $\min_{(M,x) \in \mathcal{P}(G)} F(M, x)$ is at least $\vartheta'(G)$. There exists $(M_0, y) \in \mathcal{P}(G)$ such that

$$F(M_0, y) = y^T (M_0)^\# y \max_{1 \leq i \leq n} \frac{(M_0)_{ii}}{y_i^2} = \min_{(M,x) \in \mathcal{P}(G)} F(M, x).$$

Since M_0 is positive semidefinite, there exists matrix B such that $M_0 = B^T B$. Since $y \in R(M_0)$, we can choose y such that $y = B^T c$ for a unit real vector $c \in R(B)$. Let b_i be the i th column of B . By Lemmas 2.2 and 2.3, we have

$$F(M_0, y) = y^T (M_0)^\# y \max_{1 \leq i \leq n} \frac{(M_0)_{ii}}{y_i^2} = c^T B B^+ c \max_{1 \leq i \leq n} \frac{b_i^T b_i}{(c^T b_i)^2} = \max_{1 \leq i \leq n} \frac{b_i^T b_i}{(c^T b_i)^2}.$$

Let $A = (a_{ij})_{n \times n}$ be the matrix with entries

$$\begin{aligned} a_{ii} &= 1, \quad i = 1, \dots, n, \\ a_{ij} &= 1 - \frac{b_i^T b_j}{(c^T b_i)(c^T b_j)} = 1 - \frac{(M_0)_{ij}}{y_i y_j}, \quad i \neq j. \end{aligned}$$

Since $(M_0)_{ij}y_iy_j \leq 0$ for nonadjacent vertices i, j , we have $a_{ij} \geq 1$ when $i = j$ or i, j are two nonadjacent vertices. By the definition of $\vartheta'(G)$, we have

$$\lambda_1(A) \geq \vartheta'(G).$$

We set

$$\beta = F(M_0, y) = \max_{1 \leq i \leq n} \frac{b_i^\top b_i}{(c^\top b_i)^2},$$

then

$$-a_{ij} = \left(c - \frac{b_i}{(c^\top b_i)} \right)^\top \left(c - \frac{b_j}{(c^\top b_j)} \right), \quad i \neq j,$$

$$\beta - a_{ii} = \left(c - \frac{b_i}{(c^\top b_i)} \right)^\top \left(c - \frac{b_i}{(c^\top b_i)} \right) + \beta - \frac{b_i^\top b_i}{(c^\top b_i)^2}.$$

Hence, $\beta I - A$ is positive semidefinite, that is,

$$\min_{(M,x) \in \mathcal{P}(G)} F(M, x) = \beta \geq \lambda_1(A) \geq \vartheta'(G).$$

■

The following is an example for a graph G satisfying $\alpha(G) = \vartheta'(G) < \vartheta(G)$ (see [22]).

Example 5.2 [22] Let G be a graph with vertex set $\{0,1\}^6$, and two vertices are adjacent if and only if their Hamming distance is at most 3. Then $\{000000, 001111, 110011, 111100\}$ is an independent set of G . It is known that $4 = \vartheta'(G) < \vartheta(G) = 16/3$.

By Theorems 3.1 and 5.1 and Lemma 2.5, we can get the following conclusion.

Theorem 5.3 Let $(M, x) \in \mathcal{P}(G)$ be a matrix-vector pair for graph G . If G has an independent set C such that (M, x) and C satisfy the conditions given in part (2) of Theorem 3.1, then

$$\alpha(G) = \vartheta'(G) = |C| = F(M, x).$$

6 Hoffman-type bounds

In this section, we show that the Hoffman-type bounds in [5, 14, 18, 20] are special cases of Theorem 3.1.

By Theorem 3.1, we can get the following classical Hoffman bound, and obtain the necessary and sufficient conditions of graphs attaining the Hoffman bound. The necessary condition of the equality case given in Theorem 1.1 is indeed a sufficient condition.

Corollary 6.1 Let G be a k -regular ($k \neq 0$) graph with n vertices, and let τ be the minimum adjacency eigenvalue of G . Then

$$\alpha(G) \leq n \frac{|\tau|}{k - \tau},$$

with equality if and only if there exists an independent set C such that every vertex not in C is adjacent to exactly $|\tau|$ vertices of C . Moreover, if an independent set C satisfies the above condition, then

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = n \frac{|\tau|}{k - \tau}.$$

Proof Since G is k -regular, we have $(A - \tau I, e) \in \mathcal{M}(G)$, where A is the adjacency matrix of G . By Lemma 2.1, we get

$$(A - \tau I)^\# e = (k - \tau)^{-1} e,$$

$$F(A - \tau I, e) = |\tau| e^\top (A - \tau I)^\# e = n \frac{|\tau|}{k - \tau}.$$

The conclusions follow from Theorems 3.1 and 4.2. ■

Let d_u denote the degree of a vertex u , and let $\bar{d}_S = |S|^{-1} \sum_{u \in S} d_u$ denote the average degree in a vertex set S . The largest Laplacian eigenvalue of a graph G is the largest eigenvalue of the Laplacian matrix of G .

The following bound for independent sets was given in Theorem 4.3 of [14] without sufficient condition of the equality case.

Corollary 6.2 [14] *Let G be an n -vertex graph with the largest Laplacian eigenvalue $\mu > 0$. For any independent set S of G , we have*

$$|S| \leq \frac{n(\mu - \bar{d}_S)}{\mu},$$

with equality if and only if there exists a constant d such that $d_u = d$ for each $u \in S$, and every vertex not in S is adjacent to exactly $\mu - d$ vertices of S .

Proof Let L be the Laplacian matrix of G . Since $(\mu I - L)e = \mu e$, we have $(\mu I - L, e) \in \mathcal{M}(G)$. By Lemma 2.1, we get

$$(\mu I - L)^\# e = \mu^{-1} e,$$

$$F_S(\mu I - L, e) = e^\top (\mu I - L)^\# e \sum_{u \in S} \frac{\mu - d_u}{|S|} = \frac{n}{\mu} (\mu - \bar{d}_S).$$

The conclusion follows from part (1) of Theorem 3.1. ■

The following bound was given independently in [20] and [14], and the necessary condition of the equality case was given in Theorem 3.2 of [20].

Corollary 6.3 [14, 20] *Let G be an n -vertex graph with the largest Laplacian eigenvalue $\mu > 0$ and the minimum degree δ . Then*

$$\alpha(G) \leq \frac{n(\mu - \delta)}{\mu},$$

with equality if and only if there exists an independent set C such that $d_u = \delta$ for each $u \in C$, and every vertex not in C is adjacent to exactly $\mu - \delta$ vertices of C . Moreover, if

an independent set C satisfies the above conditions, then

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = \frac{n(\mu - \delta)}{\mu}.$$

Proof Let L be the Laplacian matrix of G . Since $(\mu I - L)e = \mu e$, we have $(\mu I - L, e) \in \mathcal{M}(G)$. By Lemma 2.1, we get

$$\begin{aligned} (\mu I - L)^\# e &= \mu^{-1} e, \\ F(\mu I - L, e) &= (\mu - \delta) e^\top (\mu I - L)^\# e = \frac{n(\mu - \delta)}{\mu}. \end{aligned}$$

The conclusions follow from Theorems 3.1 and 4.2. ■

For a graph G without isolated vertices, the *normalized Laplacian matrix* \mathcal{L} of G is the $n \times n$ symmetric matrix with entries

$$(\mathcal{L})_{ij} = \begin{cases} 1, & \text{if } i = j, \\ -(d_i d_j)^{-\frac{1}{2}}, & \text{if } \{i, j\} \in E(G), \\ 0, & \text{if } \{i, j\} \notin E(G). \end{cases}$$

The largest normalized Laplacian eigenvalue of G is the largest eigenvalue of \mathcal{L} .

The following bound was given in [18] without necessary and sufficient conditions of the equality case.

Corollary 6.4 [18] *Let G be a graph with m edges, the largest normalized Laplacian eigenvalue μ and the minimum degree $\delta > 0$. Then*

$$\alpha(G) \leq \frac{2m(\mu - 1)}{\mu\delta},$$

with equality if and only if there exists an independent set C such that $d_u = \delta$ for each $u \in C$, and every vertex v not in C is adjacent to exactly $(\mu - 1)d_v$ vertices of C . Moreover, if an independent set C satisfies the above conditions, then

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C| = \frac{2m(\mu - 1)}{\mu\delta}.$$

Proof Let \mathcal{L} be the normalized Laplacian matrix of G , and let $x = (d_1^{\frac{1}{2}}, \dots, d_n^{\frac{1}{2}})^\top$. Since $(\mu I - \mathcal{L})x = \mu x$, we have $(\mu I - \mathcal{L}, x) \in \mathcal{M}(G)$. By Lemma 2.1, we get

$$\begin{aligned} (\mu I - \mathcal{L})^\# x &= \mu^{-1} x, \\ F(\mu I - \mathcal{L}, x) &= \frac{\mu - 1}{\delta} x^\top (\mu I - \mathcal{L})^\# x = \frac{2m(\mu - 1)}{\mu\delta}. \end{aligned}$$

The conclusions follow from Theorems 3.1 and 4.2. ■

The following bound was given in Theorem 6.1 of [14] without necessary and sufficient conditions of the equality case.

Corollary 6.5 [14] *Let G be an n -vertex graph, and M is a positive semidefinite matrix indexed by vertices of G such that:*

- (a) $(M)_{ij} \leq 0$ if i, j are two nonadjacent vertices.
 - (b) M has constant row sum $r > 0$.
 - (c) Every diagonal entry of M is t .
- Then

$$\alpha(G) \leq \frac{nt}{r},$$

with equality if and only if G has an independent set C such that $M[C]$ is diagonal and $\sum_{u \in C} (M)_{vu} = t$ ($v \notin C$). Moreover, if an independent set C satisfies the above conditions, then

$$\alpha(G) = \vartheta'(G) = |C| = \frac{nt}{r}.$$

Proof Since $Me = re$, we have $(M, e) \in \mathcal{P}(G)$. By Lemma 2.1, we get

$$M^\# e = r^{-1}e,$$

$$F(M, e) = te^\top M^\# e = \frac{nt}{r}.$$

The conclusions follow from Theorems 3.1 and 5.3. ■

7 Structural and spectral conditions

Let $\delta(G)$ denote the minimum degree of a graph G . In the following theorem, we give some simple structural and spectral conditions for determining the independence number $\alpha(G)$, the Shannon capacity $\Theta(G)$, and the Lovász theta function $\vartheta(G)$. These conditions mean that if a given independent set C satisfies some properties, then C must be a maximum independent set in G , and $\Theta(G) = \vartheta(G) = |C|$.

Theorem 7.1 *Let G be a graph with an independent set C . Then*

$$\alpha(G) = \Theta(G) = \vartheta(G) = |C|$$

if C satisfies one of the following conditions:

- (1) Every vertex not in C is adjacent to exactly $\lambda > -\tau$ vertices of C , where τ is the minimum adjacency eigenvalue of G .
- (2) Every vertex not in C is adjacent to exactly $-\tau$ vertices of C and G is regular, where $\tau < 0$ is the minimum adjacency eigenvalue of G .
- (3) $d_u = \delta(G)$ for each $u \in C$, and every vertex not in C is adjacent to exactly $\mu - \delta(G)$ vertices of C , where $\mu > 0$ is the largest Laplacian eigenvalue of G .
- (4) $d_u = \delta(G) > 0$ for each $u \in C$, and every vertex v not in C is adjacent to exactly $(\mu - 1)d_v$ vertices of C , where μ is the largest normalized Laplacian eigenvalue of G .
- (5) Every vertex not in C is adjacent to all vertices of C and $2|C| \geq |V(G)| - \delta(G - C)$.
- (6) G has a spanning subgraph H ($\delta(H) > 0$) which is semiregular bipartite with parameters (n_1, n_2, r_1, r_2) ($n_1 \leq n_2$), and C is the independent set with n_2 vertices in H .
- (7) There exists a clique covering ε such that $d_v^\varepsilon = \min_{u \in V(G)} d_u^\varepsilon > 0$ for each $v \in C$, and $|C \cap Q| = 1$ for each $Q \in \varepsilon$.
- (8) There exists a uniform hypergraph H such that $G = \Omega(H)$ and C corresponds to a perfect matching in H .

Proof Parts (1) and (2) follow from Corollaries 4.5 and 6.1, respectively. Parts (3) and (4) follow from Corollaries 6.3 and 6.4, respectively. Parts (5) and (6) follow from Examples 4.4 and 4.3, respectively. Parts (7) and (8) follow from Theorems 4.7 and 4.9, respectively. ■

A family \mathcal{F} of k -subsets of $\{1, \dots, n\}$ is called *intersecting* if $F_1 \cap F_2 \neq \emptyset$ for any $F_1, F_2 \in \mathcal{F}$. For $n \geq 2k$, Erdős, Ko, and Rado [11] proved that the maximum size of intersecting families of $\{1, \dots, n\}$ is $\binom{n-1}{k-1}$.

The *Kneser graph* $K(n, k)$ is a graph whose vertices consist of all k -subsets of $\{1, \dots, n\}$, and two vertices $U_1, U_2 \subseteq \{1, \dots, n\}$ are adjacent if and only if $U_1 \cap U_2 = \emptyset$. The Erdős–Ko–Rado Theorem is equivalent to say that $\alpha(K(n, k)) = \binom{n-1}{k-1}$ for $n \geq 2k$. It is known [15, Theorem 9.4.3] that the minimum adjacency eigenvalue of $K(n, k)$ is $-\binom{n-k-1}{k-1}$. By computation, the Hoffman bound for $K(n, k)$ is $\binom{n-1}{k-1}$. So we have $\alpha(K(n, k)) = \binom{n-1}{k-1}$ because $K(n, k)$ has an independent set of size $\binom{n-1}{k-1}$.

We can also obtain the EKR Theorem without using the Hoffman bound. All k -subsets containing a fixed vertex 1 forms an independent set C of size $\binom{n-1}{k-1}$, and every vertex not in C is adjacent to exactly $\binom{n-k-1}{k-1}$ vertices of C . From part (2) of Theorem 7.1, we have

$$\alpha(K(n, k)) = \Theta(K(n, k)) = \vartheta(K(n, k)) = |C| = \binom{n-1}{k-1}.$$

8 Concluding remarks

Let $M^{\otimes k}$ denote the Kronecker product of k copies of matrix M . If there is some k such that $\alpha(G^k) = \vartheta(G)^k$, then $\Theta(G) = \vartheta(G)$. We can derive the following characterization of graphs satisfying $\alpha(G^k) = \vartheta(G)^k$.

Theorem 8.1 *Let $(M, x) \in \mathcal{M}(G)$ be a matrix-vector pair for graph G such that $\vartheta(G) = F(M, x)$. There is some k such that $\alpha(G^k) = \vartheta(G)^k$ if and only if G^k has an independent set C such that $(M^{\otimes k}, x^{\otimes k})$ and C satisfy the conditions given in part (2) of Theorem 3.1. Moreover, if G^k has an independent set C satisfies the above conditions, then*

$$\alpha(G^k) = \vartheta(G)^k = \Theta(G)^k = |C|.$$

Proof Since $(M, x) \in \mathcal{M}(G)$, we have $(M^{\otimes k}, x^{\otimes k}) \in \mathcal{M}(G^k)$. By Theorem 3.1, we have

$$\alpha(G^k) \leq F(M^{\otimes k}, x^{\otimes k}) = F(M, x)^k = \vartheta(G)^k,$$

with equality if and only if G^k has an independent set C such that $(M^{\otimes k}, x^{\otimes k})$ and C satisfy the conditions given in part (2) of Theorem 3.1. Moreover, if G^k has an independent set C satisfies the above conditions, then by Theorem 4.2, we have

$$\alpha(G^k) = \vartheta(G^k) = \Theta(G^k) = |C| = F(M, x)^k = \vartheta(G)^k.$$

Since $\alpha(G^k)^{\frac{1}{k}} \leq \Theta(G) \leq \vartheta(G)$, we have $\Theta(G) = \vartheta(G)$. ■

In [19], Lovász proved that $\Theta(C_5) = \sqrt{5}$. The cycle C_5 is also an example for Theorem 8.1.

Example 8.2 Let A be the adjacency matrix of the cycle C_5 , then $(A + 2I, e) \in \mathcal{M}(C_5)$ and $((A + 2I)^{\otimes 2}, e^{\otimes 2}) \in \mathcal{M}(C_5^2)$. Suppose that $\{0, 1, 2, 3, 4\}$ is the vertex set of C_5 and $E(C_5) = \{\{0, 1\}, \{1, 2\}, \{2, 3\}, \{3, 4\}, \{4, 0\}\}$. Then $S = \{00, 12, 24, 31, 43\}$ is an independent set of C_5^2 . By computation, we know that $((A + 2I)^{\otimes 2}, e^{\otimes 2})$ and S satisfy the conditions given in part (2) of Theorem 3.1. By Theorem 8.1, we have $\alpha(C_5^2) = \vartheta(C_5)^2 = \Theta(C_5)^2 = 5$.

Acknowledgment The authors would like to thank the reviewer and Editors for giving valuable comments and suggestions.

References

- [1] N. Alon, *The Shannon capacity of a union*. *Combinatorica* 18(1998), 301–310.
- [2] A. Ben-Israel and T. N. E. Greville, *Generalized inverses: Theory and applications*, 2nd ed., Springer, New York, 2003.
- [3] A. Berman and X. D. Zhang, *On the spectral radius of graphs with cut vertices*. *J. Combin. Theory Ser. B* 83(2001), 233–240.
- [4] T. Bohman, *A limit theorem for the Shannon capacities of odd cycles I*. *Proc. Amer. Math. Soc.* 131(2003), 3559–3569.
- [5] A. E. Brouwer and W. H. Haemers, *Spectra of graphs*, Springer, New York, 2012.
- [6] F. R. K. Chung and S. T. Yau, *Discrete Green's functions*. *J. Combin. Theory Ser. A* 91(2000), 191–214.
- [7] D. Cvetković, P. Rowlinson, and S. Simić, *An introduction to the theory of graph spectra*, Cambridge University Press, Cambridge, 2010.
- [8] P. Delsarte, *An algebraic approach to the association schemes of coding theory*. *Philips Res. Rep. Suppl.* 10(1973), vi+97.
- [9] D. Ellis, Y. Filmus, and E. Friedgut, *Triangle-intersecting families of graphs*. *J. Eur. Math. Soc.* 14(2012), 841–885.
- [10] D. Ellis, E. Friedgut, and H. Pilpel, *Intersecting families of permutations*. *J. Amer. Math. Soc.* 24(2011), 649–682.
- [11] P. Erdős, C. Ko, and R. Rado, *Intersection theorems for systems of finite sets*. *Quart. J. Math. Oxford Ser. 12*(1961), 313–320.
- [12] P. Frankl and R. M. Wilson, *The Erdős–Ko–Rado theorem for vector spaces*. *J. Combin. Theory Ser. A* 43(1986), 228–236.
- [13] E. Friedgut, *On the measure of intersecting families, uniqueness and stability*. *Combinatorica* 28(2008), 503–528.
- [14] C. D. Godsil and M. W. Newman, *Eigenvalue bounds for independent sets*. *J. Combin. Theory Ser. B* 98(2008), 721–734.
- [15] C. D. Godsil and G. Royle, *Algebraic graph theory*, Springer, New York, 2001.
- [16] W. H. Haemers, *On some problems of Lovász concerning the Shannon capacity of a graph*. *IEEE Trans. Inform. Theory* 25(1979), 231–232.
- [17] W. H. Haemers, *Eigenvalue techniques in design and graph theory*. Ph.D. thesis, Technical University Eindhoven, 1979, Mathematical Centre Tracts, 121, Amsterdam, 1980.
- [18] J. Harant and S. Richter, *A new eigenvalue bound for independent sets*. *Discrete Math.* 338(2015), 1763–1765.
- [19] L. Lovász, *On the Shannon capacity of a graph*. *IEEE Trans. Inform. Theory* 25(1979), 1–7.
- [20] M. Lu, H. Liu, and F. Tian, *Laplacian spectral bounds for clique and independence numbers of graphs*. *J. Combin. Theory Ser. B* 97(2007), 726–732.
- [21] S. C. Polak and A. Schrijver, *New lower bound on the Shannon capacity of C_7 from circular graphs*. *Inform. Process. Lett.* 143(2019), 37–40.
- [22] A. Schrijver, *A comparison of the Delsarte and Lovász bounds*. *IEEE Trans. Inform. Theory* 25(1979), 425–429.
- [23] C. E. Shannon, *The zero-error capacity of a noisy channel*. *IRE Trans. Inform. Theory* 2(1956), 8–19.

- [24] D. A. Spielman, *Graphs, vectors, and matrices*. Bull. Amer. Math. Soc. 54(2017), 45–61.
- [25] J. Zhou and C. Bu, *Eigenvalues and clique partitions of graphs*. Adv. Appl. Math. 129(2021), 102220.

College of Mathematical Sciences, Harbin Engineering University, Harbin, PR China
e-mail: zhoujiang@hrbeu.edu.cn