

Testicular structure and spermatogenesis of Amazonian freshwater cururu stingray *Potamotrygon cf. histrix*

Sérgio Fonseca Zaiden¹, Richard Phillip Brinn², Jaydione Luiz Marcon³ and Elisabeth Criscuolo Urbinati⁴

Department of Biology, University of Rio Verde – FESURV, Campus Universitário, s/n. Rio Verde, Goiás, Brazil; Department of Biology, Florida International University, Florida USA; Department of Physiological Sciences, Federal University of Amazonas (UFAM), Manaus, Amazonas, Brazil; and Department of Animal Morphology and Physiology, Faculty of Agriculture and Veterinarian Sciences/Aquaculture Center, São Paulo State University (UNESP), São Paulo, Brazil

Date submitted: 24.02.10. Date accepted: 13.05.10

Summary

The cururu stingray *Potamotrygon cf. histrix*, a new and endemic Amazonian freshwater species, presents appropriate characteristics for fish keeping and is exploited from its natural environment. The present study identified the testicular structure and spermatogenesis of this species. Gonads from adult male specimens were dissected, fixed and processed for histological analysis. The testes were of testicular/epigonial type. The presence of germinal papillae was observed in the upper portion of organ with primordial germ cells and Sertoli cell precursors. The testis was lobular with zonal organization and cystic gametogenesis, with the occurrence of spermatoblasts. The Sertoli cells underwent morphological modifications over the course of gamete formation. The spermatozooids had long heads and were spiraled on their own axis. Information on the reproductive biology will serve as basis for studies on the reproduction and phylogeny of this peculiar group of cartilaginous fish.

Keywords: Epigonial testes, Germinal papilla, Intratesticular duct, Sertoli cell, Spermatoblast

Introduction

The family Potamotrygonidae (Elasmobranchii: Myliobatiformes) is the only living chondrichthyan family restricted to freshwater habitats (Lovejoy, 1996) and they are distributed throughout most of the major river systems of South America. The Amazon Basin contains the greatest number of valid species of freshwater stingrays and Brazil is the country

with the highest diversity of species (approximately 18 species) (Rosa, 1985; Compagno & Cook, 1995; Charvet-Almeida *et al.*, 2002; Carvalho *et al.*, 2003). The cururu stingray (*Potamotrygon cf. histrix*) is endemic to the Negro river basin (Carvalho *et al.*, 2003) and as a new species still deserves an appropriate taxonomic description. This high endemism has led researchers to express concern that some species may be endangered (Rosa, 1985; Compagno & Cook, 1995).

The reproductive strategy of these stingrays (Wourms, 1977), associated with the position they occupy in food chains, signifies that they are present in low densities in their habitats (Camhi *et al.*, 1998). Because of exploitation, environmental changes and degradation of the aquatic environment, these species might have some difficulty in maintaining their stock levels (Hoenig & Gruber, 1990; Compagno & Cook, 1995; Araújo *et al.*, 2004; Martin, 2005). Temperature appears to play an important role in plasma testosterone regulation, as levels in wild caught male round stingrays (*Urobatis halleri*) were negatively correlated with water temperature (Mull

¹All correspondence to: Sérgio Fonseca Zaiden. Department of Biology, University of Rio Verde – FESURV, Campus Universitário, s/n. Rio Verde, Goiás, CEP: 75901-970. Brazil. Tel: +55 64 3621-2200. Fax: +55 64 3621-2210. e-mail: sfzaiden@hotmail.com

²Department of Biology, Florida International University, Florida USA.

³Department of Physiological Sciences, Federal University of Amazonas (UFAM), Manaus, Amazonas, Brazil.

⁴Department of Animal Morphology and Physiology, Faculty of Agriculture and Veterinarian Sciences/Aquaculture Center, São Paulo State University (UNESP), Via de Acesso Prof. Paulo Donato Castelane, s/n. Jaboticabal, São Paulo, Brazil.

et al., 2008). Otherwise, freshwater stingrays exhibit a very small range of tolerance to both natural and anthropogenic impacts (Compagno & Cook, 1995).

The reproductive cycle of stingrays of the genus *Potamotrygon* is directly related to the hydrological cycle, taking the seasonality (dry and rainy seasons) into account, with defined reproductive periods and differentiated periodicity for each species, although all of them present periods of gonad maturation, copulation, pregnancy, birth and resting (Charvet-Almeida *et al.*, 2005).

The age of sexual maturation among freshwater stingrays is highly variable. Regarding the cururu stingray, it has been estimated to be 2 years of age, with gonad maturation lasting 2 months, copulation during the rainy period, a gestational period of 3 months and parturition in the dry season. However, data on the reproductive biology of freshwater stingrays are still limited, especially concerning information on microscopic morphology (Charvet-Almeida *et al.*, 2005).

Freshwater stingrays have regularly been captured for ornamental purposes for decades. Currently, at least six species from the genus *Potamotrygon* are legally permitted to be exported as ornamental fish in Brazil (IN N. 204/08, IBAMA, 2008). Their capture represents an income source for fishermen in the northern region of Brazil (Araújo *et al.*, 2004; Charvet-Almeida, 2005), as well as in other regions of South America like the Peruvian Amazonian region (Moreau & Coomes, 2007). These cartilaginous fish are generally not consumed as food. However, populations of *Paratrygon aiereba* are receiving increasing capture pressure from the commercial fishery in Pará State (Brazil), due to its larger size and appreciable muscle mass (Charvet-Almeida & Almeida, 2007). Fishery management and conservation of freshwater stingrays is a delicate topic because of their restricted distribution and the lack of official fisheries records. *Potamotrygon cf. histrix* represents 50% of all freshwater stingray legally exported from Brazil (Araújo *et al.*, 2004). Further studies on their reproductive biology are, therefore, very much needed to help ensure sustainable management and the protection and conservation of South American stingrays. Studies on the reproduction of stingrays in captive environment are scarce and they are developed in public aquariums, universities and zoos (Oldfield, 2005a, b; Henningsen, 2000).

The present study describes the testicular structure and spermatogenesis of the freshwater Amazonian cururu stingray (*Potamotrygon cf. histrix*), in order to provide better understanding of its reproductive biology. Data from structural analysis might contribute towards clarifying the reproductive mechanisms of this group of fish, thus providing tools for future

studies on the biology and management of neotropical freshwater stingrays.

Material and methods

Ten mature male specimens of the cururu stingray were collected from the Puxurituba Igarapé, in the municipality of Barcelos, State of Amazonas, Brazil, between November and December 2005. Following capture, the body size and weight of the specimens were measured and recorded (mean total weight = 266.6 g; mean disc width = 20.6 cm and mean total length = 31.8 cm). They were then euthanized using high doses of benzocaine (1.0 g in 15 litres). The abdominal cavity was opened and the gonads were exposed for gross examination. The testes on both sides were removed and fixed in buffered 10% paraformaldehyde. Following 24 h of fixing, the samples were dehydrated in a graded series of alcohol (70–100%), cleared in xylol and infiltrated into molten paraffin. Sections of 5 µm in thickness were stained with hematoxylin–eosin and Masson's trichrome for light microscopy. Photomicrographs were produced using a Leica DM 2500 microscope and morphological measurements were made using the Leica QWin V3 software. Data were subjected to analysis of variance (ANOVA) and the means were compared using Tukey's test (5%), by means of the SAS 8.12 software.

Results

Gross structure of testes

The testes were paired, oval-shaped, reddish and elongated organs located in the abdominal cavity, ventrally to the vertebral column and suspended by a mesorchium. The dorsal surface was flattened and totally covered with the epigonial organ. Both organs were developed but the left testis was larger than the right one.

Microscopic structure of testes

The testes were enclosed in a capsule of connective tissue, the tunica albuginea. Between the simple squamous epithelium and the collagenous fibres, germinal papillae or testicular appendages appeared as projections of the dorsal wall of the testes (Fig. 1A). The germinal papilla zone presented undifferentiated mesenchymal-like cells and primary spermatogonia surrounded by Sertoli cell precursors, thus characterizing their endodermal origin.

Proliferation of cells from the germinative lineage caused projection of the germinal papillae ventrally,




Figure 1 (A) Germinal papilla (GP), epithelium (E), collagen (C), epigonal organ (EO). (B) Germinal papilla projection (GPp). (C) Primary lobule (PL).

with consequent formation of spermatogenic lobules, thus indicating that this was the site of origin and development of the spermatocysts (Fig. 1B).

The testicular parenchyma was defined and was filled with lobules at different phases of development. The primary lobules were covered by a thin layer of connective tissue and the parenchyma was delimited by the epigonal organ (with content resembling lymphoid tissue), with the distribution of varied cell types, according to the development stage of the seminiferous lobules (Fig. 1C). In each lobule, there were spermatocysts at different phases of spermatogenic development. As they developed, they migrated radially to the periphery of the lobule. Thus, gonad development was characterized by cell differentiation associated with changes in cell type and diameter within the spermatocysts.

The testicular lobules were rounded, presenting zonal organization. Each zone was characterized by the presence of spermatocysts at similar phases of spermatogenesis. These zones had dorsal–ventral orientation and the upper zones were at earlier phases of spermatogenic development while the lower zones were at more advanced phases of spermatogenesis. The organization of the secondary spermatogenic lobules was characterized by the presence of germinal papillae in its upper dorsal position, followed by the germinal, spermatogonial, spermatocyte, spermatid, spermatozoid and degenerative zones (Fig. 2A).

The germinal zone was characterized by the presence of primary spermatogonia that were always accompanied by precursor of the Sertoli cells, encrusted in the tissue forming the spermatogenic cysts (Fig. 2A).

The spermatogonial zone was formed by cysts containing primary spermatogonia undergoing division, secondary spermatogonia, Sertoli cells and lumen. Proliferation of the spermatogonia and Sertoli cells led

to an increase in the size of the spermatogenic cysts and, possibly, to disappearance of the lumen and to the process of Sertoli cell nucleus migration to the periphery of the spermatocysts (Fig. 2A).

The spermatocyte zone was totally filled by spermatocysts containing primary spermatocytes at different phases of the first meiotic division. Groups of spermatocytes surrounded by the same Sertoli cell (called spermatoblasts), gave rise to a subdivision within the spermatogenic cyst and it seemed that all the cells within each spermatoblast were at the same phase of meiotic division. There were also spermatocysts containing secondary spermatocytes with characteristics similar to the primary ones, despite being half their size. The Sertoli cells were already seen in the periphery of the spermatocysts (Fig. 2A).

The spermatid zone consisted of spermatocysts containing spermatids at different phases of spermiogenesis, in which the shape changed from oval to elongated and finally spiraled. The nuclei were turned towards the peripheral region of the spermatocysts, pointing towards the nuclei of the Sertoli cells, since their cytoplasm surrounded the spermatids (Fig. 2A).

The spermatozoid zone presented these cells in groups, initially with their heads turned towards the Sertoli cell nucleus and subsequently free within the lumen of the spermatocysts. The spermatocysts opened up in the intratesticular ducts, with release inside them. The ducts were formed by a simple ciliated columnar epithelium and were also associated with the epigonal organ, with or without spermatozoids inside it (Fig. 2A).

The degeneration zone was dispersed, including the spermatozoid zone, intratesticular ducts, spermatocysts (in the degeneration and reabsorption phase) and epigonal organ tissue (through direct contact) (Fig. 2C).




Figure 2 (A) Secondary lobule (SL), germinal zone (GZ), spermatogonial zone (SZ), spermatocyte (CZ), spermatid zone (TZ), spermatozoa zone (ZZ). (B) Duct (D), lumen (L), spermatozoa (Z). (C) Degenerative zone (DZ).

Germinative lineage

The following cells from the male germinative lineage were identified:

- a. Primordial germinative cells were present in small number only inside the testicular papillae. They stood out from the other cell types that were also present there because of their size and rounded shape (Table 1). They presented defined regular outlines, acidophilic cytoplasm and spherical basophilic nuclei. During differentiation into spermatogonia, changes to the nuclei, such as chromatin condensation to form chromosomes, were visible. This cell type was always associated with precursors of Sertoli cells (Fig. 3A).
- b. Primary spermatogonia were located within the papillae and the germinal zone and were the biggest cell type of the germinative lineage (Table 1). These cells were large, rounded and well defined, with abundant cytoplasm and low affinity for staining. The spherical nuclei contained uniform euchromatin and heterochromatin, scattered irregularly. The nucleoli were eccentric. These cells were always associated with the Sertoli cells, which had comma-shaped nuclei that were completely occupied by heterochromatin (Fig. 3B).
- c. Secondary spermatogonia had characteristics similar to the primary spermatogonia. However, they were half their size and were within well defined cysts (Table 1). The cysts presented increasing sizes because of the process of spermatogonium multiplication (Fig. 3B).
- d. Primary spermatocytes were present within the spermatogenic cysts and presented spherical and well-defined nuclei, with chromatin at different phases of condensation, depending on which sub-phase of prophase I they were in or, in addition, whether they were concentrated in the periphery of the nucleus, which gave them a ring appearance. These spermatocytes were also found in distinct phases of the first meiotic division, in subunits within the same spermatocyst, called spermatoblasts (Fig. 3C). Within single cysts, there were spermatoblasts that contained primary spermatocytes and others that contained secondary ones (Table 1; Fig. 3D).
- e. Secondary spermatocytes were also found within the spermatogenic cysts, similar to the primary spermatocytes, but half their size (Table 1). Within single cysts, secondary spermatocytes at different phases of the second meiotic division and spermatoblasts containing round spermatids were found (Fig. 3E).
- f. Spermatids were also observed within the cysts, with half the size of the preceding cell type and with spherical nuclei containing very condensed chromatin, which gave it a dark appearance with high staining affinity. As spermiogenesis progressed, the spermatids went through a process of elongation, tail formation and spiraling of the




Figure 3 Germinative cell lineage. (A) Germinative primordial cells (\Rightarrow), precursor Sertoli precursor cell (\blacktriangleright). (B) Primary spermatogonia (G1), secondary spermatogonia (G2). (C) Spermatogenic cyst with spermatoblast (Sb). (D) Primary spermatocyte (S1), mitosis 1 (M1). (E) Secondary spermatocyte (S2), mitosis 2 (M2). (F) Round spermatid (T1). (G) Drop-like spermatids (T2). (H) Elongating spermatid I (T3). (I) Elongating spermatid II (T4). (J) Elongating spermatid III (T5). (K) Elongated spermatid (T6). (L) Spiraling spermatid (T7). (M, N) Spermatozoa (Z). (O) Degenerating cyst (DC). *, Sertoli cell.

anterior part where the nucleus is located. The phases identified are described below:

- f1. Round spermatids were found within the cysts, with half the size of the secondary spermatocytes (Table 1) and with spherical nuclei containing very condensed chromatin, which gave it a dark appearance with high staining affinity (Fig. 3F).
- f2. Oval or drop-shaped spermatids presented a strongly colored nucleus, with a round posterior pole and a triangular anterior one, giving the

appearance of a drop. The nuclei were turned towards the Sertoli cells (Table 2; Fig. 3G).

- f3. Spermatids going through the first stage of elongation had rod-shaped strongly basophilic nuclei, with a funneled anterior extremity that was turned towards the Sertoli cell nuclei and a rounded posterior region. There was initial formation of an acidophilic tail (Table 2; Fig. 3H).
- f4. Spermatids going through the second stage of elongation had rod-shaped basophilic nuclei, with the anterior third funneled and a

Table 1 Nuclear area of the germ cells.

Cell type	Area (μm^2)*
Germinative cell	43.35 \pm 0.89 ^e
Primary spermatogonia	111.22 \pm 2.20 ^a
Secondary spermatogonia	67.39 \pm 1.25 ^c
Primary spermatocyte	74.79 \pm 0.99 ^b
Secondary spermatocyte	51.83 \pm 0.65 ^d
Round spermatid	27.16 \pm 0.65 ^f

^{a-f}Means followed by the same letters in the column do not differ (5%). *Mean \pm standard error.

Table 2 Length of the cell nuclei during spermiogenesis.

Cell type	Length (μm)*
Round spermatid	4.73 \pm 0.12 ^f
Oval spermatid (drop-like)	5.05 \pm 0.08 ^f
Elongating spermatid (I)	7.40 \pm 0.14 ^e
Elongating spermatid (II)	11.63 \pm 0.25 ^d
Elongating spermatid (III)	18.20 \pm 0.35 ^c
Elongated spermatid	36.71 \pm 0.39 ^a
Spiraled spermatid	36.60 \pm 0.38 ^a
Spermatozoa	29.85 \pm 0.45 ^b

^{a-f}Means followed by the same letters in the column do not differ (5%). *Mean \pm standard error.

point-shaped extremity that was turned towards the Sertoli cell nuclei. The posterior extremity was rounded, with an acidophilic tail that was half the size of the head, undergoing formation (Table 2; Fig 3I).

- f5. Spermatids undergoing the third stage of elongation had long basophilic nuclei that were two-thirds funnelled, with a sharply pointed anterior extremity that was turned towards the Sertoli cell nuclei. There was an acidophilic tail that was undergoing formation and was smaller than the head (Table 2; Fig 3J).
- f6. Long spermatids also had basophilic, long, thin and straight nuclei throughout their whole length that were grouped and turned towards the Sertoli cell nuclei. An acidophilic tail was present (Table 2; Fig. 3K).

f7. Spiraling spermatids had basophilic nuclei that were long, thin and undergoing the spiraling process around their own axis, beginning at the posterior region and heading towards the anterior region. These were in groups and were turned towards the Sertoli cell nuclei (Table 2; Fig. 3L).

- g. Spermatozooids were elongated cells with an acidophilic tail and a long basophilic head, with a spiral shape around their own axis. These were found within the cysts with their anterior extremity turned towards the Sertoli cell nuclei (Fig. 3M). They were free in the lumen of the cysts (Fig. 3N), in the degeneration zone and within the intratesticular ducts (Fig. 2B).

Table 1 presents the length of the cell nuclei during spermiogenesis and Table 2 presents the nuclear area of the germ cells.

Sertoli cells

These cells accompanied the germinative lineage throughout spermatogenesis and underwent drastic modifications, with migration and morphological modification of their nuclei and increased cytoplasm (Table 3). The nuclei changed from comma-shaped (typical of precursor cells) to a triangular shape and then to an oval shape and these were larger by the end of spermatogenesis (Fig. 3B,K–M,O). After spermatozoid release, the degeneration process began. While the precursor cells were rich in heterochromatin, the Sertoli cells presented euchromatin, thus characterizing the stage of cell functionality.

Intratesticular ducts

The intratesticular ducts developed simultaneously with the seminiferous lobules, interspersed with them. They were covered with a thin connective septum and were formed by simple columnar epithelium with cells presenting basophilic elongated nuclei and acidophilic cytoplasm. These ducts, which were present throughout the degeneration zone, showed elongations that connected them directly to the efferent

Table 3 Length, width and area of the Sertoli cell nuclei associated with different cell types during spermatogenic development.

Cell types	Length (μm)*	Width (μm)	Area (μm^2)
Germinative cell	7.83 \pm 0.13 ^c	3.15 \pm 0.69 ^d	23.27 \pm 1.02 ^e
Primary spermatogonia	9.96 \pm 0.15 ^b	5.28 \pm 0.12 ^b	49.11 \pm 1.06 ^c
Secondary spermatogonia	8.02 \pm 0.12 ^c	4.31 \pm 0.07 ^c	34.72 \pm 0.71 ^d
Primary and secondary spermatocytes	11.02 \pm 0.23 ^a	4.38 \pm 0.09 ^c	48.32 \pm 1.51 ^c
Spermatids/spermatozoa	10.33 \pm 0.22 ^{a,b}	6.47 \pm 0.19 ^a	68.11 \pm 2.46 ^a
Degenerative cells	10.14 \pm 0.17 ^b	6.12 \pm 0.12 ^a	57.23 \pm 1.31 ^b

^{a-e}Means followed by the same letters in the column do not differ (5%). *Mean \pm standard error.

Table 4 Length, area and height of the duct epithelial cell nuclei, with and without presence of spermatozoa inside the duct.

Spermatozoa	Length* (μm)	Area (μm^2)	Height (μm)
Absent	10.42 ± 0.83^a	40.66 ± 6.43^a	17.47 ± 1.81^a
Present	06.47 ± 0.88^b	29.63 ± 6.04^b	9.05 ± 1.09^b

^{a,b}Means followed by the same letters in the column do not differ (5%). *Mean \pm standard error.

channel, which went through the whole length of the testicle (Fig. 2B). The length and area of the nuclei and height of the duct epithelium were greater when there were no spermatozoids inside the ducts (Table 4).

Discussion

The cururu stingray is considered to be a small species in relation to the other species of the same genus and is known as the 'dwarf freshwater stingray', with a mean disc size of 160 mm for sexually mature males (Chavert-Almeida *et al.*, 2005). The stingrays of the present study presented a mean disc size of 206 mm, which is within the expected range for this species at the reproductive stage and they presented functional testicles.

Testicular structure resembling that of cururu stingray, i.e. associated with the epigonal organ, with formation of lobules proceeding radially and presence of ducts, has previously been described to *Dasyatis sabina* (Maruska *et al.*, 1996), *Himantura signifier* (Chatchavalvanich *et al.*, 2005) and also to rajids *Atlantoraja cyclophora* (Oddone & Vooren, 2005), *Leucoraja ocellata* (Sulikowski *et al.*, 2005) and other elasmobranchs (Pratt, 1988). This latter author defined the seminiferous lobes as seminiferous follicles or ampullae, whereas Teshima & Takeshita (1992), for *Potamotrygon magdalenae*, used the term seminiferous follicles.

A difference in size between the right and left testicle has also been described in relation to the Atlantic stingray (*Dasyatis sabina*) (Maruska *et al.*, 1996) that was attributed to the size and position of the stomach and spiral intestine, which occupied the greater part of the abdominal cavity.

The differences in the cell types in the epigonal organ observed in cururu stingrays, with regard to testicles with lobules at different phases of spermatogenesis, were also described by Lutton & Callard (2008) in relation to *Leucoraja erinacea*. These authors found greater proliferation of epigonal leukocytes and greater release into blood vessels when the gonads were at the stage of reproductive activity.

This situation characterizes interaction between the immune and reproductive systems (Lutton & Callard, 2006). In a study on *Squalus acanthias*, focusing on the immunological action of the epigonal organ in the gonads, the presence of a growth inhibition factor (EGIF) was considered responsible for inhibiting spermatogenesis in that species (Piferrer & Callard, 1995). In another study, changes to the immune and reproductive systems resulting in blockage of spermatogenesis was described among Atlantic stingrays that were exposed to environmental contamination due to organochlorates (Gelsleichter *et al.*, 2002).

Genital papillae and immature or maturing genital appendices, forming primary lobules, were identified in the testicular structure of the cururu stingrays. In addition, there were papillae that were completely developed, characterized by the presence of both primary and secondary lobules. However, among *Urolophus halleri* stingrays, Babel (1967) identified the presence of primary lobules throughout the annual reproductive cycle and secondary ones only during the reproduction period, thus characterizing the latter as temporary. Germinal papillae have also been found among other elasmobranchs, such as *Squalus acanthias* (McClusky, 2005).

The organization of the spermatogenic lobules, which was histologically characterized in the cururu stingrays by their defined development zones, followed cell differentiation and changes in spermatocyst composition and diameter that were distributed in stages equivalent to the zones described for *Himantura signifier* (Chatchavalvanich *et al.*, 2004) and *Dasyatis sabina* (Maruska *et al.*, 1996; Tricas *et al.*, 2000); and for other elasmobranchs such as *Mustelus canis* (Conrath & Musick, 2002), *Centroscyrmnus coelolepis* and *Centrophorus squamosus* (Girard *et al.*, 2000) and *Squalus acanthias* (McClusky, 2005, 2006).

The germinative lineage found in cururu stingray was similar to that described for *Dasyatis sabina* (Maruska *et al.*, 1996; Tricas *et al.*, 2000), *Himantura signifier* (Chatchavalvanich *et al.*, 2004) and *Raja asterias* (Barone *et al.*, 2007). At the time of cell division, it was observed that within the same spermatocyst, there were spermatoblasts containing cells at different phases of cell division, thus resulting in the presence of different cell types within the same spermatocyst, with the presence of secondary spermatocytes and some spermatoblasts and spermatids within others. Prisco *et al.* (2003) also identified different stages of cell differentiation in the same spermatogenic cyst in *Torpedo marmorata* and also among other elasmobranchs, such as *Centroscyrmnus coelolepis* and *Centrophorus squamosus* (Girard *et al.*, 2000).

Length measurements on the spermatids during the spermiogenesis were also made by Maruska *et al.* (1996) on *Dasyatis sabina* stingray. The results

showed that a significant difference appeared as the differentiation phenomena evolved, thus indicating that the increase in length of the reproductive cells undergoing formation is considerable. The spiraling process observed in the final events of spermiogenesis in cururu stingray is characteristic of elasmobranchs and, among these, freshwater stingrays (Maruska *et al.*, 1996; Tricas *et al.*, 2000; Chatchavalvanich *et al.*, 2004, 2005).

Helical spermatozooids with rotation on their own axis, as identified in the present study, is characteristic of elasmobranchs in general (Wourms, 1977) and has also been identified in *Potamotrygon magdalenae* (Teshima & Takeshita, 1992) and *Himantura signifier* freshwater stingrays (Chatchavalvanich *et al.*, 2005).

The structural modifications identified in the Sertoli cells, at different times during spermatogenesis, indicate that this cell type is directly related to the phenomena of gamete production. This is due to the greater metabolic need for their functions to be performed with the germinative lineage. The cells present in cysts in which spermatozoid release had already occurred began to degenerate (Maruska *et al.*, 1996; Tricas *et al.*, 2000; Prisco *et al.*, 2002; Chatchavalvanich *et al.*, 2004). This process occurs at the end of their function, when mature gametes are released and spermatocysts disappear (Prisco *et al.*, 2003; Liguoro *et al.*, 2004).

The intratesticular ducts found in cururu stingray are similar to those described in other stingrays, such as *Urolophus halleri* (Babel, 1967) and also in *Raja erinacea* (Pratt, 1988). The differences in the histological structure of the intratesticular ducts that were observed in cururu stingray show the functionality of the organ, in which the cells undergo losses of cytoplasm in order to form the seminal fluid. These changes accompany the phenomena of spermatogenesis and the release of spermatozooids. Piercy *et al.* (2003) analysed the genital ducts of Atlantic stingrays and identified extensive histological changes in their architecture, in relation to both size and secretion activity, during the reproductive period. The information obtained from the present study will serve as a tool for gaining knowledge of the reproductive mechanism of this group of freshwater chondrichthyes and as support for further studies on reproduction in captivity or in degraded and polluted natural environments.

Acknowledgements

This work was supported by a grant from National Geographic Conservation Fund (No. C47-04). The scientific expedition was authorized by the Brazilian

Ministry of Science and Technology (MCT/CNPq, process No. EXC 023/05 to J.L.M.) and by the Federal Environmental Agency (process No. 098/2005 DIFAP/IBAMA-DF).

References

- Araújo, M.L.G., Charvet-Almeida, P. & Almeida, M.P. (2004). Conservation status of freshwater stingrays (Chondrichthyes: Potamotrygonidae) in the Brazilian Amazon. In *Biology and Conservation of Elasmobranchs: Symposium Proceedings. VI International Congress on the Biology of Fish, American Fisheries Society (Physiology Section)*, Manaus, Hotel Tropical, pp. 49–59.
- Babel, J.S. (1967). Reproduction, life history and ecology of the round stingray, *Urolophus halleri* Cooper. *Calif. Fish Game Bull.* **137**, 1–104.
- Barone, M., De Ranieri, S., Fabiani, O., Pirone, A. & Serena, F. (2007). Gametogenesis and maturity stages scale of *Raja asterias* Delaroché, 1809 (Chondrichthyes, Rajidae) from the South Ligurian Sea. *Hydrobiologia* **580**, 245–54.
- Camhi, M., Fowler, S., Musick, J., Brauntingam, A. & Fordham, S. (1998). *Sharks and their relatives—ecology and conservation*. IUCN/SSC Shark Specialist Group. IUCN. Switzerland and Cambridge, UK: Gland. 43 pp.
- Carvalho, M.R., Lovejoy, N.R. & Rosa, R.S. (2003). *Potamotrygonidae* (river stingrays). In *Checklist of the Freshwater Fishes of South and Central America*. (eds. R.E. Reis, & S.O. Kullander Ferraris, Jr), pp. 22–8. C.J. Porto Alegre: EDIPUCRS, Brasil.
- Charvet-Almeida, P. & Almeida, M.P. Contribuição ao conhecimento, distribuição e aos desafios para a conservação dos elasmobrânquios (raias e tubarões) no Sistema Solimões-Amazonas. (2007). In *Conservação da várzea: identificação e caracterização de regiões biogeográficas* (Albernaz, A.L.K.M., Org.). pp. 199–235. ProVárzea Ibama, [Technical report in Portuguese].
- Charvet-Almeida, P., Araújo, M.L.G., Rosa, R.S. & Rincon, G. (2002). Neotropical freshwater stingrays: diversity and conservation status. *IUCN Shark News*, **14**, 47–51.
- Charvet-Almeida, P., Araújo, M.L.G. & Almeida, M.P. (2005). Reproductive aspects of freshwater stingrays (Chondrichthyes: Potamotrygonidae) in the Brazilian Amazon basin. *J. Northw. Atl. Fish. Sci.* **35**, 165–71.
- Chatchavalvanich, K., Thogpan, A. & Nakai, M. (2004). Structure of the testis and genital duct of freshwater stingray, *Himantura signifier* (Elasmobranchii: Myliobatiformes: Dasyatidae). *Ichthyol. Res.* **52**, 123–31.
- Chatchavalvanich, K., Thogpan, A. & Nakai, M. (2005). Ultrastructure of spermiogenesis in a freshwater stingray, *Himantura signifier*. *Ichthyol. Res.* **52**, 379–85.
- Conrath, C.L. & Musick, J.A. (2002). Reproductive biology of the smooth dogfish, *Mustelus canis*, in the northwest Atlantic Ocean. *Environ. Biol. Fishes* **64**, 367–77.
- Compagno, L.J.V. & Cook, S.F. (1995). The exploitation and conservation of freshwater elasmobranchs: status of taxa and prospects for the future. *J. Aquaricult. Aquatic Sci.* **7**, 62–90.
- Gelsleichter, J., Rasmussen, L.E.L., Marine, C.A., Tyminki, J., Chang, B. & Lombardi-Carlson, L. (2002). Serum steroid concentrations and development of reproductive organs

- during puberty in male bonnethead sharks, *Sphyrna tiburo*. *Fish Physiol. Biochem.* **26**, 389–401.
- Girard, M., Rivalan, P. & Siquin, G. (2000). Testis and sperm morphology in two deep-water squaloid sharks, *Centroscyrmus coelolepis* and *Centrophorus squamosus*. *J. Fish Biol.* **57**, 1575–89.
- Henningsen, A.D. (2000). Notes on reproduction in the southern stingray, *Dasyatis americana* (Chondrichthyes: Dasyatidae), in a captive environment. *Copeia* **3**, 826–8.
- Hoenig, J.M. & Gruber, S.H. (1990). Life-history patterns in the elasmobranchs: implications for fisheries management. In: *Elasmobranchs as living resources: advances in the biology, ecology, systematics, and the status of fisheries*, vol. 90 (eds H.L. Pratt, Jr., S.H. Gruber & T. Taniuchi), pp. 1–16. U.S. Department of Commerce, NOAA Technical Report NMFS. IBAMA (2008). IN 204/2008. Accessible at http://www.ibama.gov.br/recursos-pesqueiros/wp-content/files/in_ibama_204_2008_raiasdoce_retificada.pdf. Captured on 12 February 2010.
- Liguoro, A., Prisco, M., Mennella, C., Ricchiari, L., Angelini, F. & Andreuccetti, P. (2004). Distribution of terminal sugar residues in the testis of the spotted ray *Torpedo marmorata*. *Mol. Reprod. Dev.* **68**, 524–30.
- Lovejoy, N.R. (1996). Systematics of myliobatoid elasmobranchs: with emphasis on the phylogeny and historical biogeography of neotropical freshwater stingrays (Potamotrygonidae: Rajiformes). *Zool. J. Linn. Soc.* **117**, 207–57.
- Lutton, B.V. & Callard, I.P. (2006). Evolution of reproductive-immune interactions. *Integr. Comp. Biol.* **46**, 1060–9.
- Lutton, B.V. & Callard, I.P. (2008). Influence of reproductive activity, sex steroids and seasonality on epigonal organ cellular proliferation in the skate (*Leucoraja erinacea*). *Gen. Comp. Endocrinol.* **155**, 116–25.
- Martin, R.A. (2005). Conservation of freshwater and euryhaline elasmobranchs: a review. *J. Mar. Biol. Ass. U.K.* **85**, 1049–73.
- Maruska, K.P., Cowie, E.G. & Tricas, T.C. (1996). Periodic gonadal activity and protacted mating in elasmobranch fishes. *J. Exp. Zool.* **276**, 219–32.
- McClusky, L.M.A. (2005). Stage and season effects on cell cycle and apoptotic activities of germ cells and Sertoli cells during spermatogenesis in the spiny dogfish (*Squalus acanthias*). *Reproduction* **129**, 89–102.
- McClusky, L.M.A. (2006). Stage-dependency of apoptosis and the blood-testis barrier in the dogfish shark (*Squalus acanthias*): cadmium-induced changes as assessed by vital fluorescence techniques. *Cell Tissue Res.* **325**, 541–53.
- Moreau, M-A. & Coomes, O.T. (2007). Aquarium fish exploitation in Western Amazonia: conservation issues in Peru. *Environ. Cons.* **34**, 12–22.
- Mull, C.G., Lowe, C.G. & Young, K.A. (2008). Photoperiod and water temperature regulation of seasonal reproduction in male round stingray (*Urobatis halleri*). *Comp. Biochem. Physiol., Part A.* **151**, 717–25.
- Oddone, M.C. & Vooren, C.M. (2005). Reproductive biology of *Atlantoraja cyclophora* (Regan 1903) (Elasmobranchii: Rajidae) of southern Brazil. *ICES J. Mar. Sci.* **62**, 1095–1103.
- Oldfield, R.G. (2005a). Biology, husband and reproduction of freshwater stingrays I. *Tropical Fish Hobbyist* **53**(12), 114–6.
- Oldfield, R.G. (2005b). Biology, husband and reproduction of freshwater stingrays II. *Tropical Fish Hobbyist* **54**(1), 110–2.
- Piercy, A., Gelsleichter, J. & Snelson Jr., F.F. (2003). Morphological and histological changes in the genital ducts of the male atlantic stingray, *Dasyatis sabina* during the seasonal reproductive cycle. *Fish Physiol. Biochem.* **29**, 23–35.
- Piferrer, F.C. & Callard, G.V. (1995). Inhibition of deoxyribonucleic acid synthesis during premeiotic stages of spermatogenesis by a factor from testis-associated lymphomyeloid tissue in the dogfish shark (*Squalus acanthias*). *Biol. Reprod.* **53**, 390–8.
- Pratt, Jr, H.L. (1988). Elasmobranch gonad structure: a description and survey. *Copeia* **3**, 719–29.
- Prisco, M., Liguoro, A., D'Onghia, B., Ricchiari, L., Andreuccetti, P. & Angelini, F. (2002). Fine structure of Leydig and Sertoli cells in the testis of immature and mature spotted ray *Torpedo marmorata*. *Mol. Reprod. Dev.* **63**, 192–201.
- Prisco, M., Liguoro, A., Comitato, R., Cardone, A., D'Onghia, B., Ricchiari, L., Angelini, F. & Andreuccetti, P. (2003). Apoptosis during spermatogenesis in the spotted ray *Torpedo marmorata*. *Mol. Rep. Dev.* **64**, 341–8.
- Rosa, R.S. 1985. *A systematic revision of the south american freshwater stingrays (Chondrichthyes: Potamotrygonidae)*. Doctorate Thesis, Williamsburg, College of William and Mary, 523 pp.
- Sulikowski, J.A., Tsang, P.C.W. & Howell, W.H. (2005). Age and size at sexual maturity for the winter skate, *Leucoraja ocellata*, in the Western Gulf of Maine based on morphological, histological and steroid hormone analyses. *Environ. Biol. Fish.* **72**, 429–41.
- Teshima, K. & Takeshita, K. (1992). Reproduction of the freshwater stingray, *Potamotrygon magdalenae* taken from the Magdalena River system in Colombia, South America. *Bull. Seikai Nat. Fish. Res. Inst.* **70**, 11–27.
- Tricas, T.C., Maruska, K.P. & Rasmussen, L.E.L. (2000). Annual cycles of steroid hormone production, gonad development and reproductive behavior in the Atlantic stingray. *Gen. Comp. Endocrinol.* **118**, 209–25.
- Wourms, J.P. (1977). Reproduction and development in Chondrichthyan fishes. *Am. Zool.* **17**, 379–410.