

## SEPARATING THE FAN THEOREM AND ITS WEAKENINGS

ROBERT S. LUBARSKY AND HANNES DIENER

**Abstract.** Varieties of the Fan Theorem have recently been developed in reverse constructive mathematics, corresponding to different continuity principles. They form a natural implicational hierarchy. Some of the implications have been shown to be strict, others strict in a weak context, and yet others not at all, using disparate techniques. Here we present a family of related Kripke models which separates all of the as yet identified fan theorems.

**§1. Introduction.** To be able to talk about fans, Cantor space, and similar objects properly, we will start by introducing some notation. The space of all infinite binary sequences, endowed with the standard topology (wherein a basic open set is given by a finite binary sequence), will be denoted by  $2^{\mathbb{N}}$ ; the set of all finite binary sequences will be denoted by  $2^*$ . The concatenation of  $u, v \in 2^*$  will be denoted by  $u * v$ . For  $\alpha \in 2^{\mathbb{N}}$  and  $n \in \mathbb{N}$ , the first  $n$  elements of  $\alpha$  form a finite sequence denoted by  $\bar{\alpha}n$ . A subset  $B \subseteq 2^*$  is called a **bar** if

$$\forall \alpha \in 2^{\mathbb{N}} \exists n \in \mathbb{N} (\bar{\alpha}n \in B),$$

and a bar is called **uniform** if

$$\exists n \in \mathbb{N} \forall \alpha \in 2^{\mathbb{N}} \exists m \leq n (\bar{\alpha}m \in B).$$

Notice that if a bar  $B$  is closed under extensions, that is if

$$\forall u \in 2^* (u \in B \implies \forall v \in 2^* u * v \in B),$$

then it is uniform if and only if

$$\exists n \in \mathbb{N} \forall \alpha \in 2^{\mathbb{N}} (\bar{\alpha}n \in B).$$

Not all of the bars we consider will be closed under extensions.

There are currently four versions of Brouwer's Fan Theorem in common use. All of them enable one to conclude that a given bar is uniform. The differences among them lie in the definitional complexity demanded (as an upper bound) of the bar in order for the theorem to apply to it, which ranges from the very strongest

---

Received October 17, 2012.

2010 *Mathematics Subject Classification.* 03C90, 03F50, 03F60, 03H05.

*Key words and phrases.* fan theorems, Kripke models, forcing, non-standard models, Heyting-valued models.

© 2014, Association for Symbolic Logic  
0022-4812/14/7903-0008/\$3.40  
DOI:10.1017/jsl.2014.9

requirement to no restriction on the bar at all. A bar  $C \subset 2^*$  is **decidable** if it is decidable as a set:

$$\forall u \in 2^* u \in C \vee u \notin C.$$

A bar  $C \subset 2^*$  is called a  **$c$ -bar** if there exists a decidable set  $C' \subset 2^*$  such that

$$u \in C \iff \forall v \in 2^* (u * v \in C').$$

A bar  $B \subset 2^*$  is called a  **$\Pi_1^0$ -bar** if there exist a decidable set  $S \subset 2^* \times \mathbb{N}$  such that

$$u \in B \iff \forall n(u, n) \in S.$$

(The  $\Pi_n^0$ -nomenclature alludes to the arithmetical hierarchy in computability theory.) We can now state four commonly used versions of the Fan Theorem.

- FAN $_{\Delta}$ : Every decidable bar is uniform.
- FAN $_c$ : Every  $c$ -bar is uniform.
- FAN $_{\Pi_1^0}$ : Every  $\Pi_1^0$ -bar is uniform.
- FAN $_{full}$ : Every bar is uniform.

Notice that every decidable bar can be taken to be closed under extensions; that is, the closure of a decidable bar under extension is still decidable. If there is no restriction on the definability of a bar, then every bar can be taken to be so closed, by working with the closure of any given bar. Every  $c$ -bar is already closed under extension. In contrast,  $\Pi_1^0$ -bars seemingly cannot be replaced by their closures while remaining  $\Pi_1^0$ .

By way of motivation, these principles were developed within reverse constructive mathematics, because they are equivalent with certain continuity principles. In particular, over a weak base theory, FAN $_{\Delta}$  is equivalent with the assertion that every uniformly continuous, positively valued function from  $[0,1]$  to  $\mathbb{R}$  has a positive infimum [8], FAN $_c$  with the uniform continuity of every continuous  $f : 2^{\mathbb{N}} \rightarrow \mathbb{N}$  [2], and FAN $_{\Pi_1^0}$  with the uniform equicontinuity of every equicontinuous sequence of functions from  $[0,1]$  to  $\mathbb{R}$  [6]. (For background on analysis in a constructive context, see [4].)

The following implications hold trivially [2, 5] and over a weak base theory:

$$FAN_{full} \implies FAN_{\Pi_1^0} \implies FAN_c \implies FAN_{\Delta}.$$

One naturally wonders whether any of the implications can be reversed, including whether FAN $_{\Delta}$  is outright provable in constructive set theory. Some such non-implications have already been determined.

- It is well-known (see [1] for instance) that FAN $_{\Delta}$  is not provable, via recursive realizability. That is, there is an infinite (Turing) computable sub-tree of  $2^*$  with no infinite computable branch, which fact translates to a failure of FAN $_{\Delta}$  under IZF (Intuitionistic ZF, the constructive correlate to classical ZF) via recursive realizability, and also to the independence of WKL (Weak König’s Lemma) over RCA $_0$  in reverse mathematics [10].
- Berger [3] shows that FAN $_{\Delta}$  does not imply FAN $_c$  over a very weak base system. His argument is in its essence a translation of the reverse mathematics proof that WKL $_0$  does not imply ACA [10], by coding the Turing jump into a  $c$ -bar. In order for this argument to work, he must be in a context in which the existence of the Turing jump is not outright provable, hence the use of a weak base system.

- Fourman and Hyland [7] present a Heyting-valued, almost topological, model in which  $\text{FAN}_{\text{full}}$  fails; they do not address which fragments of the Fan Theorem might hold, since these distinctions were not available at the time. We show below that  $\text{FAN}_{\Pi_1^0}$  holds in their model, separating the left-most pair of principles in the diagram above.

We are not aware of any prior proofs separating  $\text{FAN}_c$  and  $\text{FAN}_{\Pi_1^0}$ .

The goal of this paper is to separate all of these principles via a uniform technique. This has several benefits. For one, it separates  $\text{FAN}_c$  and  $\text{FAN}_{\Pi_1^0}$ . For another, it separates  $\text{FAN}_{\Delta}$  and  $\text{FAN}_c$  over full IZF. That is new because Berger's argument still leaves open the possibility that IZF would allow that implication to go through; independence of  $\text{FAN}_c$  over  $\text{IZF} + \text{FAN}_{\Delta}$  means that it does not. In addition, since the arguments employed rather handily provide four separation results, they seem to provide a flexible tool that might be useful elsewhere. This seems not to be the case for the other techniques that have been used. It could well be the case, for instance, that realizability could produce all of the results discussed here. But no one has been able to do this yet. As for the Fourman-Hyland argument, they also show in the same work that all topological models satisfy  $\text{FAN}_{\text{full}}$ . So for the separations of interest here, topological models are just out. To be sure, variants of topological models, along the lines used by Fourman and Hyland, might still do the trick. But before coming up with the arguments below that's exactly what we tried, and got nowhere. In short, we cannot say that the techniques used here get you anything that could not be gotten by other means, but at least it seems to be easier to use. Beyond that, it could be the case that the proofs here really are in some sense the right ones for these results. In the face of the perfectly nice realizability and Heyting-valued models that provide some of these separations, we are not at this point making that claim. While the constructions below are natural enough, they are not so compelling as to seem canonical. Nonetheless, since they seem to work so well, it might be that with further reflection and development, it turns out that proofs along these lines are the way to go for a large class of problems.

As for what the techniques employed actually are, we would like to provide some motivation for how we happened upon them. Since it seemed that realizability and Heyting algebras weren't working, we turned to the only other kind of model we know of, Kripke models. To build a tree we could control, along with its paths, over set theory with full Separation and Collection, we turned to forcing. In order to have the trees be decidable, yet not completely pinned down, as required by the theories in question, we were forced to use non-standard integers, to provide non-standard levels on the trees.

Since this is a paper about constructive mathematics, a word about the meta-theory used is in order. It is classical through and through. We work in ZFC. Presumably most if not all of the arguments are fully constructive, as in so many mathematical papers in all fields. We did not check, and so have no idea.

In the next section we discuss the Fan Theorem in topological and related models, including giving a proof that the Fourman-Hyland model satisfies  $\text{FAN}_{\Pi_1^0}$ . The following sections provide the advertised separation results, going right-to-left in the diagram above. We then close with some questions.

**§2. The Fan Theorem in Heyting-Valued Models.** To make this paper somewhat self-contained, we repeat the proof that explains why the construction afterwards is more complicated than just a topological model.

PROPOSITION 2.1 (Fourman-Hyland [7]). *In any topological model  $FAN_{full}$  holds.*

PROOF. Let  $T$  be a topological space, and suppose

$$T \Vdash \text{“} B \subseteq 2^* \text{ is a bar closed under extension.”}$$

Then, in particular, for any external sequence  $\alpha \in 2^{\mathbb{N}}$  (that is, one from the ground model)

$$T = \llbracket \exists n \bar{\alpha}n \in B \rrbracket = \bigcup_{n \in \mathbb{N}} \llbracket \bar{\alpha}n \in B \rrbracket. \tag{1}$$

Let  $A_u$  denote the open set

$$\llbracket \text{the bar } \{w \mid u * w \in B\} \text{ is uniform} \rrbracket.$$

If  $T \not\Vdash \text{“} B \text{ is uniform,”}$  then choose some  $p \notin A_{()}.$  Define a set  $Tr = \{u \in 2^* \mid p \notin A_u\}.$  Since  $A_u = A_{u*0} \cap A_{u*1}$  for any  $u \in 2^*,$   $Tr$  is a tree (i.e., closed under restriction) with no terminal nodes. Since in addition  $() \in Tr$  (that is,  $Tr$  is non-empty),  $Tr$  is infinite. Thus, by Weak König’s Lemma, there exists an infinite path  $\beta$  in  $Tr.$  By the definition of  $Tr,$   $p \notin A_{\bar{\beta}n}$  for all  $n \in \mathbb{N}.$  Now Equation 1 yields the existence of  $n \in \mathbb{N}$  such that

$$p \in \llbracket \bar{\beta}n \in B \rrbracket;$$

but this contradicts  $\llbracket \bar{\beta}n \in B \rrbracket \subset A_{\bar{\beta}n}.$  ⊥

This suggests that if we are looking for models in which some form of the Fan Theorem fails we need to “delete points”. This was done in [7], Section 4, where they consider  $K(T),$  the coperfect open sets of a topological space  $T.$  This can be viewed as the equivalence classes of the open sets of  $T,$  under which an open set is identified with its smallest coperfect superset. In this setting, removing a point from an open set does not change the set.

DEFINITION 2.2. A Heyting algebra is connected if  $A \vee B = \top$  and  $A \wedge B = \perp$  implies that either  $A = \top$  or  $A = \perp.$

Let  $\Omega$  be  $K([0, 1] \times [0, 1]).$  It is easy to see that  $\Omega$  is connected.

PROPOSITION 2.3. *If  $H$  is a connected Heyting algebra, then  $H \Vdash FAN_{\Pi_1^0}.$*

PROOF. Suppose  $H \Vdash \text{“} B \text{ is a } \Pi_1^0\text{-bar, given say by } S : u \in B \text{ iff } \forall n \in \mathbb{N} (u, n) \in S.”$  Since  $H \Vdash \text{“} S \text{ is decidable,”}$  for any  $u \in 2^*$  and  $n \in \mathbb{N},$

$$H \Vdash \text{“}(u, n) \in S \vee (u, n) \notin S.”$$

By the connectedness of  $H$  either  $H \Vdash \text{“}(u, n) \in S”}$  or  $H \Vdash \text{“}(u, n) \notin S.”}$  So define a set  $\tilde{B} \subset 2^*$  in the metatheory by

$$u \in \tilde{B} \iff \forall n \in \mathbb{N} H \Vdash \text{“}(u, n) \in S.”$$

$\tilde{B}$  is itself a bar, as follows. Let  $\alpha \in 2^{\mathbb{N}}$  be arbitrary. If  $\bar{\alpha}n \notin \tilde{B}$  for all  $n \in \mathbb{N}$  then for all  $n$  there exist  $i_n$  such that  $\llbracket (\bar{\alpha}n, i_n) \in B \rrbracket = \perp.$  Thus

$$\llbracket \forall m \in \mathbb{N} (\bar{\alpha}n, m) \in B \rrbracket = \perp$$

for any  $n \in \mathbb{N}$ , and therefore

$$\llbracket \exists n \in \mathbb{N} \forall m \in \mathbb{N} (\bar{\alpha}n, m) \in B \rrbracket = \perp ;$$

a contradiction to  $B$  being a bar internally. Hence  $\tilde{B}$  is a bar externally, and therefore, working with a classical metatheory (or simply the Fan Theorem), it is uniform. So there exists  $N$  such that for all  $u$  of length  $N$  some initial segment of  $u$  is in  $\tilde{B}$ . Then it is easy to see, that this same  $N$  witnesses the uniformity of  $B$  internally.  $\dashv$

**COROLLARY 2.4.** *FAN<sub>□</sub> does not imply FAN<sub>full</sub> (over IZF).*

**PROOF.** In [7] it is shown that  $\Omega \not\vdash$  FAN<sub>full</sub>.  $\dashv$

**§3. FAN<sub>Δ</sub> is not Provable.** As discussed in the introduction, recursive realizability shows that IZF does not prove FAN<sub>Δ</sub>. However, we do not see how to adapt that, or the Heyting-valued model from the previous section, to the other desired separation results. Hence we are hoping not merely to provide here a different model falsifying FAN<sub>Δ</sub> as a technical exercise, but rather to provide a technique more flexible than those referenced, to produce the other separation results. Of course, if this really is a flexible technique, it should work for the known separations too.

We will build a Kripke model, working within ZFC. To construct a bar, it will be crucial to control what paths exist. This is most easily done with a generic set, in the sense of forcing.

**DEFINITION 3.1.** Let the forcing partial order  $P$  be the set of appropriate labelings of finitely many nodes from  $2^*$ . A labeling of nodes assigns to each one either IN, OUT, or  $\infty$ , with the following restrictions. Any node labeled IN has no descendant, the idea being that once a node gets into the eventual bar so are all of its descendants automatically, so nothing more need be said. Any descendant of a node labeled OUT must be labeled IN or OUT. Finally, for any node labeled  $\infty$ , if both children are labeled, then at least one of them must be labeled  $\infty$ .

Let  $G$  be a generic through the condition that labels  $\langle \rangle$  with  $\infty$ . By straightforward density arguments, any node labeled OUT by  $G$  has a uniform bar above it (or below it, depending on how you draw your trees) all labeled IN, and every node labeled  $\infty$  has a path through it always labeled  $\infty$ , in fact a perfect set of such.

Let  $B = \{\alpha \in 2^* \mid \text{for some } n \ G(\alpha \upharpoonright n) = \text{IN}\}$ .  $B \in M[G]$  is the interpretation  $\sigma_B^G$  of the term  $\sigma_B = \{\langle p, \hat{\alpha} \rangle \mid \text{for some } n \ p(\alpha \upharpoonright n) = \text{IN}\}$ . (As usual, the function  $\hat{\cdot}$  is the canonical injection of the ground model into the terms:  $\hat{x} = \{\langle \emptyset, \hat{y} \rangle \mid y \in x\}$ .) Because of these latter  $\infty$ -paths,  $B$  is not a bar. However, we might reasonably think that if we no longer had access to the distinction between the OUT and the  $\infty$  nodes, we might no longer be able to build a path avoiding  $B$ . This intuition is confirmed by the next proposition.

**DEFINITION 3.2.** The shadow forcing  $Q$  is the set of functions from finite subtrees of  $2^*$  to  $\{\text{IN}, \text{OUT}\}$  such that any node labeled IN has no descendant. Equivalently,  $Q$  is the sub-partial order of  $P$  beneath the condition labeling  $\langle \rangle$  with OUT (together with the condition which labels  $\langle \rangle$  IN, which has no extension). The canonical projection  $\text{proj}_Q$  of  $P$  onto  $Q$  replaces all occurrences of  $\infty$  with OUT. The canonical projection of the terms of  $P$ 's forcing language to those of  $Q$ 's, ambiguously also

called  $\text{proj}_Q$ , acts by applying  $\text{proj}_Q$  to the conditions that appear in the terms, hereditarily. (Notice that  $Q$  term are also  $P$  terms.)

Notice that a  $P$ -filter projects to a  $Q$ -filter. If  $G$  is a generic  $P$ -filter, then  $\text{proj}_Q(G)$  will not be  $Q$ -generic, because in  $Q$  the terminal conditions are dense. Still,  $\text{proj}_Q(G)$  induces an interpretation  $\sigma^{\text{proj}_Q(G)}$  of the terms  $\sigma$  of  $Q$ . These interpretations are in  $M[G]$ , as they are easily definable from  $\sigma$  and  $G$ ; alternatively,  $\sigma^{\text{proj}_Q(G)} = (\text{proj}_Q^{-1''}\sigma)^G$ .

For any  $P$ -filter  $G$ ,  $\text{proj}_Q(\sigma_B)^{\text{proj}_Q(G)} = B$ : the induced interpretation of the projection of  $B$  is just  $B$  itself. Effectively,  $B$  as a  $P$ -term is already a  $Q$ -term.

**PROPOSITION 3.3.** *If  $\sigma$  is a  $Q$ -term and  $p \Vdash_P$  “ $\text{proj}_Q^{-1''}\sigma$  is an infinite branch through  $2^*$ ,” then  $p \Vdash_P$  “ $\text{proj}_Q^{-1''}\sigma$  goes through  $\sigma_B$ .”*

**PROOF.** By standard forcing technology, it suffices to extend  $p$  to some condition forcing “ $\text{proj}_Q^{-1''}\sigma$  goes through  $\sigma_B$ ,” as then it will be dense beneath  $p$  to force as much, and so will happen generically.

First extend  $p$  so that every sequence in  $2^*$  of length  $2^{n-1}$  for some  $n$  either is labeled OUT or  $\infty$  or has a proper initial segment labeled IN. Then extend again by adjoining both children to all nodes of length  $2^{n-1}$ , and labeling them  $\infty$  whenever possible (otherwise IN or OUT). For a technical reason soon to become clear, we must extend yet again. This time have the domain include all length  $k$  descendants of the length  $n$  nodes not labeled IN, and label them so that every length  $n$  node labeled  $\infty$  has a unique descendant of length  $k$  labeled  $\infty$ , and, most importantly, for each pair of nodes  $\alpha$  and  $\beta$  of length  $k$  labeled  $\infty$ , there is some  $i$  with  $\alpha(i) = 1$  and  $\beta(i) = 0$ . One way of doing this is to let  $s$  be the number of nodes of length  $n$  labeled  $\infty$ , to let  $k$  be  $n + s$ , and to build the  $\infty$ -labeled descendant of the  $j^{\text{th}}$  such node by adjoining to it  $j - 1$  0's, a 1, and then  $s - j$  0's, all other descendants of length  $k$  being labeled OUT.

Extend one last time to  $q \Vdash \text{proj}_Q^{-1''}\sigma(\hat{k}) = \hat{\alpha}$  for some fixed  $\alpha$ , where as usual  $\hat{x}$  is the standard term for the internalization of the set  $x$ . Moreover,  $q$  should force the equality in the strong sense that for each  $j < k$  there is a term  $\tau$  and a condition  $r \geq q$  with  $\langle r, \tau \rangle \in \text{proj}_Q^{-1''}\sigma$  and  $q \Vdash \tau = \langle \hat{j}, \hat{\alpha}(\hat{j}) \rangle$ ; even further, if  $\alpha(j) = 1$  then  $q$  forces a particular element to be in  $\tau$ 's second component.

If  $q$  labels some initial segment of  $\alpha$  IN then we're done.

If  $q$  labels  $\alpha$  OUT then it is dense beneath  $q$  that all descendants of  $\alpha$  of some fixed length are labeled IN, and again we're done.

If  $q$  labels  $\alpha$   $\infty$  then let  $q_{alt}$  be identical to  $q$  except that all descendants of  $\alpha \upharpoonright n$  labeled  $\infty$  by  $q$  are labeled OUT by  $q_{alt}$ . Observe first that  $q_{alt}$  extends  $p$ . Then note that, because  $\text{proj}_Q(q_{alt}) = \text{proj}_Q(q)$ , the strong forcing facts posited of  $q$  hold for  $q_{alt}$  as well: for the same  $\tau$  and  $j$  as above,  $q_{alt} \Vdash \tau \in \text{proj}_Q^{-1''}\sigma$  and  $q_{alt} \Vdash$  “ $\tau$  is an ordered pair with first component  $\hat{j}$ ,” and if  $q$  forced  $\tau$ 's second component to be non-empty,  $q_{alt}$  also forces it to be non-empty, containing the same term as for  $q$ . The difference between  $q$  and  $q_{alt}$ , from  $\sigma$ 's point of view, is that  $q_{alt}$  has more extensions than  $q$ : there are conditions extending  $q_{alt}$  which bar the tree beneath  $\alpha$ , which is not so for  $q$ . That means that it is possible for extensions of  $q_{alt}$  to force sets into  $Q$ -terms that no extension of  $q$  could. In the case of  $\text{proj}_Q^{-1''}\sigma(\hat{k})$ , though, such opportunities are limited. That term is already forced by  $p$  to be a function

with domain  $k$ ; for each  $j < k$  there is already a fixed term forced to stand for  $\langle j, (proj_Q^{-1''}\sigma(\hat{k}))(j) \rangle$ ; if that function value at  $j$  was forced by  $q$  to be 1 then it must retain a member and so is also forced by  $q_{alt}$  to be 1. The only change possible is that something formerly forced to be empty (i.e., be 0) could now be forced by some extension to have an element (i.e., be 1). Recall, though, the construction of  $q$  on level  $k$ : if  $proj_Q^{-1''}\sigma(\hat{k})$  is ever forced by some  $r \leq q_{alt}$  to be some  $\beta \neq \alpha$  by flipping some 0's to 1's, by  $\alpha$ 's distinguished 1  $r$  cannot label  $\beta \infty$ . So  $r$  can be extended so that all extensions of  $\beta$  of a certain length are labeled IN, forcing  $proj_Q^{-1''}\sigma$  to hit  $\sigma_B$ . Of course, any extension of  $q_{alt}$  forcing  $proj_Q^{-1''}\sigma(\hat{k})$  to be  $\alpha$  works the same way as such an  $r$  does, since  $q_{alt}$  already labels  $\alpha$  OUT. In either case we have an extension of  $p$  forcing  $proj_Q^{-1''}\sigma$  go through  $\sigma_B$ .  $\dashv$

Even though we have just seen that  $B$  is a bar relative to the  $Q$ -paths, we will perhaps surprisingly have occasion to consider weaker situations, where  $B$  is larger and hence even easier to hit. The case of interest is if we were to change some  $\infty$ 's in  $G$  to OUTs, thereby allowing uniform bars above those nodes. Notice that if  $\alpha$ 's sibling is not labeled  $\infty$ , then  $\alpha$ 's label could not consistently be changed from  $\infty$ , as then  $\alpha$ 's parent, labeled  $\infty$ , would then have both children not labeled  $\infty$ . Such considerations do not apply when  $\alpha = \langle \rangle$ .

**DEFINITION 3.4.**  $H$  is a legal weakening of  $G$  if  $H$  can be constructed by choosing finitely many nodes labeled  $\infty$  by  $G$ , changing those labels (to either IN or OUT), also changing the labeling of finitely many descendants of those nodes from  $\infty$  or OUT to OUT or IN in such a way that each node labeled OUT has a uniform bar above it labeled IN, and then eliminating all descendants of nodes labeled IN. Furthermore, this must be done in such a manner that  $H$  is a filter through  $P$  (avoiding, for instance, the problem posed just before this definition).

Notice that the difference between  $H$  and  $G$  can be summarized in one condition  $p$ , which contains the new bars, all labeled IN, and all of their ancestors. Hence we use the notation  $G_p$  to stand for this  $H$ : to build  $G_p$ , make the minimal change to each condition in  $G$  in order to be consistent with  $p$ .

**LEMMA 3.5.** *If  $G_p$  is a legal weakening of  $G$  then  $G_p$  is generic through  $p$ .*

**REMARK 3.6.** Notice that if  $p$  labels the empty sequence IN or OUT then  $p = G_p$  is a terminal condition in  $P$ , trivially satisfying the lemma.

**PROOF.** Let  $D$  be dense beneath  $p$ . Notice that  $G \upharpoonright dom(p)$  is a condition in  $P$  contained in  $G$ . It is not hard to define the notion of projection beneath  $p$ ,  $proj_p$ , by making the minimal changes in a condition necessary to be compatible with  $p$ . We claim that  $proj_p^{-1''}D$  is dense beneath  $G \upharpoonright dom(p)$ . To see this, let  $q \leq G \upharpoonright dom(p)$ . Extend  $proj_p(q)$  to  $r \in D$ . The only way  $r$  can extend  $proj_p(q)$  is by labeling extensions  $\alpha$  of nodes which are unchanged by  $proj_p$ : if  $\alpha \in dom(r) \setminus dom(proj_p(q))$  then, for  $\alpha \upharpoonright n \in dom(q)$ ,  $q(\alpha \upharpoonright n) = proj_p(q)(\alpha \upharpoonright n)$ . Extend  $q$  to  $q_r$  by labeling those same extensions the same way: for  $\alpha \in dom(r) \setminus dom(proj_p(q))$   $q_r(\alpha) = r(\alpha)$ . We have that  $proj_p(q_r) = r$ , hence  $q_r \in proj_p^{-1''}D$ . So  $proj_p^{-1''}D$  is dense beneath  $G \upharpoonright dom(p)$ , hence contains a member of  $G$ , say  $q$ . Then  $proj_p(q)$  is in both  $D$  and  $G_p$ .  $\dashv$

We can now describe the ultimate Kripke model. Recall that  $G$  is generic for  $P$  over  $M$  and labels the empty sequence with  $\infty$ . The bottom node  $\perp$  of the Kripke model consists of the  $Q$ -terms, with membership (not equality!) as interpreted by  $proj_Q(G)$ . Let  $N$  be an ultrapower of  $M[G]$  using any non-principal ultrafilter on  $\omega$ , with elementary embedding  $f : M[G] \rightarrow N$ . This necessarily produces non-standard integers. Let  $\mathcal{H}$  be the set of legal weakenings of  $f(G)$ , as defined in  $N$ , which induce the same  $B$  on the standard levels of  $2^*$ , which restriction is definable only in  $M[G]$ . That is, any standard node labeled  $\infty$  by  $G$  can only be changed to OUT by the legal weakening.  $\mathcal{H}$  will index the successors of  $\perp$ . At the node indexed by  $f(G)_p$ , the universe will be the  $Q$ -terms of  $N$  as interpreted by  $proj_Q(f(G)_p)$ . Regarding the embeddings from  $\perp$ , for a  $Q$ -term  $\sigma \in M$ ,  $f(\sigma)$  is an  $f(Q)$ -term in  $N$ , so send  $\sigma$  to  $f(\sigma)$ . If  $f(G)_p$  is a terminal condition in  $P$ , then the node indexed by  $f(G)_p$  is terminal in the Kripke ordering. Else iterate. That is, suppose  $f(G)_p$  is non-terminal. The structure at its node can be built in  $N$ . As an ultrapower of  $M[G]$ ,  $N$  internally looks like  $f(M)[f(G)]$ ; internally,  $f(G)$  is  $f(P)$ -generic over the ground model  $f(M)$ . The structure at node  $f(G)_p$  could be built in  $f(M)[f(G)_p]$ , where, by the previous lemma,  $f(G)_p$  is generic through  $f(P)$ , and also non-terminal. Hence the construction just described, using an ultrapower and legal weakenings to get additional nodes, can be performed in  $f(M)[f(G)_p]$  just as above. Continue through  $\omega$ -many levels. We will ambiguously use  $f$  to stand for any of the elementary embeddings, including compositions of such (making  $f$  a sort-of polymorphic transition function). Notice that the construction relativizes: the Kripke structure from node  $f(G)_p$  onwards is definable in  $f(M)[f(G)_p]$  just as the entire structure is definable in  $M[G]$ .

This defines a Kripke structure interpreting membership. Equality at any node can now be defined as extensional equality beyond that node in this structure, inductively on the ranks of the terms, even though the model is not well-founded, thanks to the elementarity present. That is, working at  $\perp$ , suppose  $\sigma$  and  $\tau$  are terms of rank at most  $\alpha$ , and we have defined equality at  $\perp$  for all terms of rank less than  $\alpha$ . Moreover, suppose (strengthening the inductive assumption here) that this definability was forced in  $M$  by the empty condition  $\emptyset$ . At node  $f(G)_p$  the structure is definable over  $f(M)[f(G)_p]$ , and, by elementarity, in  $f(M)$ ,  $\emptyset \Vdash$  "Equality in the Kripke model is unambiguously definable for all terms of rank less than  $f(\alpha)$ ." So at that node we can see whether there is a witness to  $f(\sigma)$  and  $f(\tau)$  being unequal. If there is such a witness at any node  $f(G)_p$ , then  $\sigma$  and  $\tau$  are unequal at  $\perp$ , else they are equal at  $\perp$ . This extends the definability of equality to all terms of rank  $\alpha$ . Hence inductively equality is definable for all terms.

PROPOSITION 3.7.  $\perp \not\Vdash FAN_\Delta$ .

PROOF. It is immediate that  $B$  is a bar: any node is internally of the form  $f(M)[f(G)_p]$ ; by the lemma,  $f(G)_p$  is always  $f(P)$ -generic; by the proposition, no path given by a  $Q$ -term can avoid the interpretation of the term for  $B$  as given by an  $f(P)$ -generic. Moreover,  $B$  is decidable, as  $f(G)_p$  agrees with  $G$  on the standard part of  $2^*$ , the only part that exists at  $\perp$ , and that argument relativizes to all nodes. However,  $B$  is not uniform at any non-terminal node, since  $f(G)_p$ , when non-terminal, has labels of  $\infty$  at every level.  $\dashv$



What remains to show is that our model satisfies IZF. In order to do this, we will need to get a handle on internal truth in the model. This is actually unnecessary for most of the IZF axioms, but for Separation in particular we will have to deal with truth in the model. When forcing, this is done via the forcing and truth lemmas:  $M[G] \models \phi$  iff  $p \Vdash \phi$  for some  $p \in G$ , where  $\Vdash$  is definable in  $M$ . Since our Kripke model is built in  $M[G]$ , statements about it are statements within  $M[G]$ , and so are forced by conditions in  $G$ . The problem is that the Kripke model internally does not have access to  $G$ , but only to  $B$ . In detail, Separation for  $M[G]$  is proven as follows: given  $\phi$  and  $\sigma$ , it suffices to consider  $\{\langle q, \tau \rangle \mid \text{for some } \langle p, \tau \rangle \in \sigma, q \leq p \text{ and } q \Vdash \phi(\tau)\}$ . The problem we face is that that set seems not to be in the Kripke model, even if  $\sigma$  is. What we need to show is that if  $\sigma$  and  $\phi$ 's parameters are  $Q$ -terms then that separating set is given by a  $Q$ -term.

Recall that  $\text{proj}_Q$  operates by replacing all occurrences of  $\infty$  by  $\text{OUT}$ .

DEFINITION 3.8.  $p \sim p'$  if  $\text{proj}_Q(p) = \text{proj}_Q(p')$ .

DEFINITION 3.9.  $p \Vdash^* \phi$ , for  $\phi$  in the language of the Kripke model, i.e., when  $\phi$ 's parameters are  $Q$ -terms, inductively on  $\phi$  :

- $p \Vdash^* \sigma \in \tau$  if, for some  $\langle q, \rho \rangle \in \tau$ ,  $q \geq_Q \text{proj}_Q(p)$  and  $p \Vdash^* \sigma = \rho$ .
- $p \Vdash^* \sigma = \tau$  if for all  $p' \sim p$ ,  $p'' \leq_P p'$ , and  $\langle q, \rho \rangle \in \sigma$ , if  $\text{proj}_Q(p'') \leq_Q q$  then there is a  $p''' \leq_P p''$  such that  $p''' \Vdash^* \rho \in \tau$ , and symmetrically.
- $p \Vdash^* \phi \wedge \theta$  if  $p \Vdash^* \phi$  and  $p \Vdash^* \theta$ .
- $p \Vdash^* \phi \vee \theta$  if  $p \Vdash^* \phi$  or  $p \Vdash^* \theta$ .
- $p \Vdash^* \phi \rightarrow \theta$  if for all for all  $p' \sim p$  and  $p'' \leq_P p'$ , if  $p'' \Vdash^* \phi$  then there is a  $p''' \leq_P p''$  such that  $p''' \Vdash^* \theta$ .
- $p \Vdash^* \exists x \phi(x)$  if, for some  $Q$ -term  $\sigma$ ,  $p \Vdash^* \phi(\sigma)$ .
- $p \Vdash^* \forall x \phi(x)$  if for all for all  $p' \sim p$ ,  $p'' \leq_P p'$ , and  $Q$ -term  $\sigma$ , there is a  $p''' \leq_P p''$  such that  $p''' \Vdash^* \phi(\sigma)$ .

LEMMA 3.10. If  $p \sim p'$  then  $p \Vdash^* \phi$  iff  $p' \Vdash^* \phi$ .

PROOF. For the cases  $\in, =, \rightarrow$ , and  $\forall$ , that is built right into the definition of  $\Vdash^*$ . The other cases are a trivial induction. ⊢

LEMMA 3.11. If  $q \leq_P p$  and  $p \Vdash^* \phi$  then  $q \Vdash^* \phi$ .

PROOF. Inductively on  $\phi$ . For  $\in$ , use that  $\text{proj}_Q$  is monotone. The cases  $\wedge, \vee$ , and  $\exists$  are trivial inductions. For the remaining cases, suppose  $q'' \leq_P q', q' \sim q$ , and  $q \leq_P p$ . Then  $q'' \leq_P q' \upharpoonright \text{dom}(p) \sim p$ , and use that  $p \Vdash^* \phi$ . ⊢

PROPOSITION 3.12.  $\perp \models \phi$  iff  $p \Vdash^* \phi$  for some  $p \in G$ .

PROOF. Inductively on  $\phi$ .

$\sigma \in \tau$ :  $\perp \models \sigma \in \tau$  iff there are  $p \in G$  and  $\langle q, \rho \rangle \in \tau$  such that  $\text{proj}_Q(p) \leq_Q q$  and  $\perp \models \sigma = \rho$ . Inductively  $\perp \models \sigma = \rho$  iff there is an  $r \in G$  \*-forcing the same. In one direction, using lemma 3.11,  $p \cup r$  suffices, in the other we have  $p = r$ .

$\sigma = \tau$ : Suppose  $p \in G$  and  $p \Vdash^* \sigma = \tau$ . By taking  $p'$  equal to  $p$  in the definition of  $\Vdash^*$ , for every member  $\rho$  of either  $\sigma$  or  $\tau$ , it is dense to \*-force  $\rho$  to be in the other set. By the genericity of  $G$  some such  $p'''$  will be in  $G$ , and so inductively  $\rho$  will end up in the other set. This shows that  $\sigma$  and  $\tau$  have the same members at  $\perp$ . Regarding a future node  $f(G)_{p''}$ , because  $f(G)_{p''}$  is a legal weakening of  $f(G)$ ,

$p'' \upharpoonright \text{dom}(p) \sim p$ , so again it is dense for any member of  $\sigma$  or  $\tau$  to be forced into the other, so they have the same members at node  $f(G)_{p''}$ . Hence  $\perp \Vdash^* \sigma = \tau$ .

Conversely, suppose for all  $p \in G$   $p \not\Vdash^* \sigma = \tau$ . That means there are  $p' \sim p, p'' \leq_P p'$ , and  $\rho$  forced by  $p''$  into  $\sigma$  (without loss of generality), but  $p''$  has no extension  $*$ -forcing  $\rho$  into  $\tau$ . For every natural number  $n$  the set  $D_n = \{q \mid \text{for some } k > n, \text{dom}(q) \subseteq 2^k, \text{ and all binary sequences of length } k \text{ either are labeled } \infty \text{ by } q \text{ or some initial segment is labeled IN by } q\}$  is dense. Hence cofinally many levels of  $G$  are in  $D_0$ . Observe that if  $q$  is in  $D_0 \cap G$  and  $q' \sim q$  then any extension of  $q'$  can be extended again to induce a legal weakening of  $G$ . In  $N$ , by overspill choose  $p \in f(G)$  to be in  $f(D_0)$ . Choose  $p'' \leq_P p' \sim p$  and  $\rho$  as given by the case hypothesis. Extend  $p''$  to  $p'''$  so that  $f(G)_{p'''}$  is a legal weakening of  $f(G)$ . Since  $p'''$  has no extension  $*$ -forcing  $\rho$  into  $\tau$ , inductively at node  $f(G)_{p'''}$   $\rho$  is not a member of  $\tau$ . Hence  $\perp \not\Vdash \sigma = \tau$ .

$\phi \wedge \theta$ : Trivial.

$\phi \vee \theta$ : Trivial.

$\phi \rightarrow \theta$ : Suppose  $p \in G$  and  $p \Vdash^* \phi \rightarrow \theta$ . At any node  $f(G)_{p'}$ , if  $f(G)_{p'} \models \phi$  then inductively choose  $p'' \in f(G)_{p'}$  such that  $p'' \Vdash^* \phi$ . Without loss of generality  $p''$  can be taken to extend  $p'$ . Since  $f(G)_{p'}$  indexes a node in the model,  $p' \upharpoonright \text{dom}(p) \sim p$ , so  $p'' \leq_P p' \upharpoonright \text{dom}(p) \sim p$ . By the case assumption there is a  $p'''$  extending  $p''$  with  $p''' \Vdash^* \theta$ . By the genericity of  $f(G)_{p'}$  there is such a  $p'''$  in  $f(G)_{p'}$ . So inductively  $f(G)_{p'} \models \theta$ . At node  $\perp$  the argument is even simpler, as  $p'$  can be chosen to be  $p$ . So  $\perp \models \phi \rightarrow \theta$ .

Conversely, suppose for all  $p \in G$  that  $p \not\Vdash^* \phi \rightarrow \theta$ . That means there are  $p' \sim p$  and  $p'' \leq_P p'$  with  $p'' \Vdash^* \phi$  but no extension of  $p''$   $*$ -forces  $\theta$ . As in the = case above, in  $N$ , by overspill choose  $p \in f(G)$  to be in  $f(D_0)$ . Choose  $p'' \leq_P p' \sim p$  as given by the case hypothesis. Extend  $p''$  to  $p'''$  so that  $f(G)_{p'''}$  is a legal weakening of  $f(G)$ . Inductively  $f(G)_{p'''}$   $\models \phi$ , but since  $p'''$  has no extension  $*$ -forcing  $\theta$ , inductively  $f(G)_{p'''}$   $\not\models \theta$ . Hence  $\perp \not\models \phi \rightarrow \theta$ .

$\exists x \phi(x)$ : Trivial.

$\forall x \phi(x)$ : Suppose  $p \in G$  and  $p \Vdash^* \forall x \phi(x)$ . For any node  $f(G)_{p'}$  and any  $\sigma$  in the universe there,  $p' \leq_P p' \upharpoonright \text{dom}(p) \sim p$ , so there is a  $p'' \leq_P p'$  such that  $p'' \Vdash^* \phi(\sigma)$ . By genericity there is such a  $p''$  in  $f(G)_{p'}$ . Inductively  $f(G)_{p'} \models \phi(\sigma)$ . So every element at node  $f(G)_{p'}$  satisfies  $\phi$  there. At node  $\perp$  the argument is even easier, since  $p'$  can be chosen to be  $p$ . Hence  $\perp \models \forall x \phi(x)$ .

Conversely, suppose for all  $p \in G$  that  $p \not\Vdash^* \forall x \phi(x)$ . That means there are  $p' \sim p, p'' \leq_P p'$ , and  $Q$ -term  $\sigma$  such that  $p''$  has no extension  $*$ -forcing  $\phi(\sigma)$ . As in the cases of = and  $\rightarrow$  above, in  $N$ , by overspill choose  $p \in f(G)$  to be in  $f(D_0)$ . Choose  $p'' \leq_P p' \sim p$  and  $\sigma$  as given by the case hypothesis. Extend  $p''$  to  $p'''$  so that  $f(G)_{p'''}$  is a legal weakening of  $f(G)$ . Since  $p'''$  has no extension  $*$ -forcing  $\phi(\sigma)$ , inductively  $f(G)_{p'''}$   $\not\models \phi(\sigma)$ . Hence  $\perp \not\models \forall x \phi(x)$ . -1

**THEOREM 3.13.**  $\perp \models IZF$ .

**PROOF.** Empty Set and Infinity are witnessed by  $\emptyset$  and  $\hat{\omega}$  respectively. Pairing is witnessed by  $\{\langle \emptyset, \sigma \rangle, \langle \emptyset, \tau \rangle\}$ , and Union by  $\{\langle q \cup r, \rho \rangle \mid \text{for some } \tau \langle q, \tau \rangle \in \sigma \text{ and } \langle r, \rho \rangle \in \tau\}$ . Extensionality holds because that's how = was defined.

For  $\varepsilon$ -Induction, suppose  $\perp \models “(\forall y \in x \phi(y)) \rightarrow \phi(x).”$  If it were not the case that  $\perp \models “\forall x \phi(x)”$ , then at some later node  $G_p$  there would be a term  $\sigma$  with

$f(G)_p \not\models \phi(\sigma)$ . The restricted Kripke model of node  $f(G)_p$  and its extensions is definable in a model of ZF, say  $N$ , which is a finite iteration of the ultrapower construction, and so is itself a model of ZF. Hence, in  $N$ ,  $\sigma$  can be chosen to be such a term of least  $V$ -rank, say  $\kappa$ . Then at all nodes after  $f(G)_p$ , by elementarity, it holds that  $f(\kappa)$  is the least rank of any term not satisfying  $\phi$ . So all members of  $\sigma$ , being of lower rank, satisfy  $\phi$  at whatever node they appear. By the induction hypothesis,  $\sigma$  must also satisfy  $\phi$ , contradicting the assumption that some term does not satisfy  $\phi$ .

For the powerset of  $\sigma$  take all sets with members of the form  $\langle q, \tau \rangle$ , where  $\langle p, \tau \rangle \in \sigma$  and  $q \leq_Q p$ .

It is easy to give a coarse proof of Bounding. The Kripke model can be built in  $M[G]$ . Given a  $\sigma$  at  $\perp$ , Bounding in  $M[G]$  can be used to bound the range of  $\phi$  on  $\sigma$  at  $\perp$ . Also, the set of nodes is set-sized, so there are only set-many interpretations of  $f(\sigma)$  at the other nodes, so the range of  $\phi$  on them can also be bounded. Since the standard ordinals are cofinal through the ordinals in all of the iterated ultrapowers, by picking  $\kappa$  large enough,  $\hat{V}_\kappa$  suffices for bounding the range of  $\phi$  on  $\sigma$ .

For Separation, given  $\phi$  and  $\sigma$ , let  $\text{Sep}_{\phi,\sigma}$  be  $\{\langle \text{proj}_Q(p), \tau \rangle \mid \text{for some } \langle q, \tau \rangle \in \sigma \text{ with } p \leq q \text{ we have } p \Vdash^* \phi(\sigma)\}$ . By lemmas 3.10 and 3.12, this works.  $\dashv$

Although this model does not satisfy  $\text{FAN}_\Delta$ , it does satisfy  $\neg\neg\text{FAN}_\Delta$ , as the terminal nodes are dense. Admittedly this is a rather weak failure of  $\text{FAN}_\Delta$ . In the final section, we will address the issue of getting stronger failures of  $\text{FAN}_\Delta$ .

**§4.  $\text{FAN}_\Delta$  does not imply  $\text{FAN}_c$ .** We will need a tree similar to that of the last proof. In fact, we will need two trees: the  $c$ -bar  $C$ , and the decidable set  $C'$  from which  $C$  is defined. (Both can be viewed either as  $2^*$  with labels or as subtrees of  $2^*$ .) Mostly we will focus on  $C$ . Because  $\text{FAN}_c$  refers to eventual membership in a tree, the difference between IN and OUT nodes is no longer relevant: the bar is uniform beneath any OUT node. So we can describe the forcing in terms similar to those before, and with some simplifications introduced. The forcing partial order  $P$  will be the set of appropriate labelings of finitely many nodes from  $2^*$ . A labeling of nodes assigns to each one either IN or  $\infty$ , with the following restrictions. Any node labeled IN has no descendant, the idea being that once a node gets into the eventual bar so are all of its descendants automatically, so nothing more need be said. For any node labeled  $\infty$ , if both children are labeled, then at least one of them must be labeled  $\infty$ . Let  $G$  be  $P$ -generic through the condition labeling the empty sequence with  $\infty$ .

As before, we will need to look at weaker trees, ones with bigger bars.

**DEFINITION 4.1.**  $H$  is a legal weakening of  $G$  if  $H$  can be constructed by choosing finitely many nodes labeled  $\infty$  by  $G$ , whose siblings are also labeled  $\infty$  by  $G$ , and changing those labels to IN and eliminating all descendants.

As before, each legal weakening  $H$  can be summarized by one forcing condition  $p$ , which consists of those nodes changed by  $H$  and their ancestors, labeled as in  $G$ .  $H$  is then the set of conditions in  $G$  each minimally changed to be consistent with  $p$ . Hence we refer to  $H$  as  $G_p$ .

LEMMA 4.2. *If  $G_p$  is a legal weakening of  $G$  then  $G_p$  is generic through  $p$ .*

PROOF. As in the corresponding lemma in the previous section. ◻

DEFINITION 4.3. Terms are defined inductively (through the ordinals) as sets of the form  $\{\langle B_i, \sigma_i \rangle \mid i \in I\}$ , where  $I$  is any index set,  $\sigma_i$  a term, and  $B_i$  a finite set of truth values. A truth value is a symbol of the form  $b^+$  or  $b'$  or  $\neg b'$ , for  $b \in 2^*$  a finite binary sequence.

DEFINITION 4.4. Let  $C$  be the term  $\{\{\langle b^+, \hat{b} \rangle \mid b \in 2^*\}$ , and  $C'$  be  $\{\{\langle b', \hat{b} \rangle \mid b \in 2^*\}$ .

In our final model, (the interpretation of)  $C$  will be the  $c$ -bar induced by (the interpretation of)  $C'$ , and  $C$  will not be uniform, thereby falsifying  $FAN_c$ . Furthermore, we will show that  $FAN_\Delta$  holds in this model.

We can now describe the ultimate Kripke model. Recall that  $G$  is generic for  $P$  over  $M$  and labels the empty sequence with  $\infty$ . The bottom node  $\perp$  of the Kripke model consists of the terms. At  $\perp$ ,  $b^+$  counts as true iff  $G(b) = \text{IN}$ ,  $b'$  always counts as true, and  $\neg b'$  never counts as true. Later nodes will have different ways of counting the various literals as true. At any node, for  $\sigma = \{\langle B_i, \sigma_i \rangle \mid i \in I\}$ , if each member of some  $B_i$  counts as true, then at that node  $\sigma_i \in \sigma$ . This induces a notion of extensional equality among the terms. One way of viewing this is at any node to remove from a term  $\sigma$  any pair  $\langle B_i, \sigma_i \rangle$  if some member of  $B_i$  is not true at that node. Then each remaining  $\langle B_i, \sigma_i \rangle$  can be replaced by  $\sigma_i$ . Equality is then as given by the Axiom of Extensionality as interpreted in the model.

As for what the other nodes in the model are, there are two different kinds. As in the last section, let  $N$  be an ultrapower of  $M[G]$  using any non-principal ultrafilter on  $\omega$ , with elementary embedding  $f : M[G] \rightarrow N$ . This necessarily produces non-standard integers. In  $N$ , any forcing condition  $p$  which induces a legal weakening of  $f(G)$  will index a successor node to  $\perp$ . At the node indexed by  $p$ , the universe will be the terms of  $N$  as interpreted by  $f(G)_p$ . That is,  $b^+$  is true if  $f(G)_p(b) = \text{IN}$ ,  $b'$  is always true, and  $\neg b'$  never. Regarding the embeddings from  $\perp$ , for a term  $\sigma \in M$ ,  $f(\sigma)$  is a term in  $N$ , so send  $\sigma$  to  $f(\sigma)$ . In addition, definably over  $M[G]$ , any non-standard  $c \in 2^*$  with  $f(G)(c) = \infty$  also indexes a node. At such a node  $c$ ,  $b'$  counts as true iff  $b \neq c$ ,  $\neg b'$  counts as true iff  $b = c$ , and  $b^+$  counts as true iff  $b \not\subseteq c$  ( $b$  is not an initial segment of  $c$ ). Note that at  $\perp$  any  $b'$  refers only to a standard  $b$ ; for some  $b'$  to be declared false at a later node  $c$ ,  $b$  would have to equal  $c$ , and  $c$  indexes a node only if  $c$  is non-standard. Hence there is no conflict with the Kripke structure: once  $b'$  is deemed true, it remains true. Similarly with  $b^+$ :  $G_p$  is a fattening of  $G$ . Hence membership, being based on finitely many truth values, is monotone.

Any node indexed by such a  $c \in 2^*$  is terminal in the Kripke ordering. Also, among nodes of the other kind, there is one trivial condition  $p$ , the one with  $p(\langle \rangle) = \text{IN}$ . This is also a terminal node, where each  $b^+$  and each  $b'$  is true. At any other node, iterate. That is, suppose  $p$  is not the preceding condition. The model at  $p$  can be built in  $N$ . As an ultrapower of  $M[G]$ ,  $N$  internally looks like  $f(M)[f(G)]$ . The structure at node  $p$  could be built in  $f(M)[f(G)_p]$ , where  $f(G)_p$  is generic through  $f(P)$  (and non-trivial). Hence the construction just described, using an ultrapower and legal weakenings and non-standard binary strings to get additional

nodes, can be performed in  $f(M)[f(G)_p]$  just as above. This provides immediate successors to nodes indexed by (non-trivial)  $p$ 's. Iterate  $\omega$ -many times.

The picture is that, at  $\perp$ ,  $C$  looks like  $G$ , that is, those nodes  $G$  assigns to be IN. This tree gets fatter at later nodes that are legal weakenings. At terminal nodes  $c$ ,  $C$  is everything but the branch up to  $c$ . At most nodes  $C'$  looks like everything; at node  $c$ , where  $c$  is non-standard relative to its predecessor, we find the one thing not in  $C'$ , namely  $c$ .

What we need to show is that this model satisfies IZF and  $FAN_\Delta$ , and falsifies  $FAN_c$ .

LEMMA 4.5.  $\perp \not\models FAN_c$ .

PROOF. It is easy to see that  $C$  is the  $c$ -set induced by  $C'$ : once  $b$  is forced into  $C$ , none of its descendants index terminal nodes, so no descendant is forced out of  $C'$ ; similarly, if  $b$  is not forced into  $C$ , say at node  $p$ , then  $G_p(b) = \infty$ , and in  $N$  some non-standard extension  $c$  of  $b$  will also be labeled  $\infty$  by  $f(G)_p$ , and that  $c$  will index a node at which  $c$  is not in  $C'$ . Clearly,  $C$  is not uniform, and  $C'$  is decidable. So it remains only to show that  $\perp \models C$  is a bar.

Suppose  $\sigma$  is forced to be an infinite binary path at some node. If that node is a terminal node,  $C$  contains cofinitely many members of  $2^*$ , and so certainly intersects  $\sigma$ . Else without loss of generality we can assume the node is  $\perp$ . Then, for some  $p \in G$ ,  $p \Vdash \text{“}\perp \models \sigma \text{ is an infinite binary path.”}$  If it is not dense beneath  $p$  to force the standard part of  $\sigma$  (that is,  $\sigma$  applied to the standard integers) to be in the ground model, then extensions  $q$  and  $r$  of  $p$  force incompatible facts about  $\sigma$ . The only incompatible facts about  $\sigma$  are of the form  $b \frown 0 \in \sigma$  and  $b \frown 1 \in \sigma$ . The positive parts of  $q$  and  $r$  (that is,  $q^{-1}(IN)$  and  $r^{-1}(IN)$ ) induce a legal weakening of  $G$ . That is, there is a canonical condition  $\text{inpart}(q, r)$ , with domain  $\text{dom}(q) \cup \text{dom}(r)$ , that returns IN on any node that either  $q$  or  $r$  returns IN on, as well as on any node if  $\text{inpart}(q, r)$  returns IN on both children, else OUT. Because terms use only positive (i.e., IN) information, at the node  $f(G)_{\text{inpart}(q,r)}$ , both  $b \frown 0$  and  $b \frown 1$  are in  $\sigma$ . (More coarsely and perhaps more simply, at the node induced by the trivial condition sending the empty sequence to IN, the same conclusion holds for the same reason.) Hence  $\perp$  could not have forced  $\sigma$  to be a path in the first place. Therefore  $p$  forces  $\sigma$  on the standard binary tree to be in the ground model. It is easy to see that generically  $G$  labels some node in  $\sigma$  IN. ⊣

LEMMA 4.6.  $\perp \models FAN_\Delta$ .

PROOF. By arguments similar to the above. If a set of nodes  $B$  is forced by  $p$  to be decidable, then no extensions of  $p$  can force incompatible facts about  $B$ . Hence  $B$  is in the ground model. If  $B$  were not a bar in the ground model, there would be a ground model path missing  $B$ . This path would also be in the Kripke model. Hence  $B$  is a bar in the ground model. Since the ground model is taken to be classical,  $B$  is uniform. ⊣

Regarding getting IZF to be true, just as in the previous section, the problem is that truth in the Kripke model is on the surface determined by forcing conditions in the ground model, to which the Kripke model has no access. The essence is to capture truth at a node using those truth values that are allowed in the building of terms.

DEFINITION 4.7. For a forcing condition  $p$ ,  $B_p = \{b^+ \mid \text{for some initial segment } c \text{ of } b, p(c) = \text{IN}\}$ .

For a set of truth values  $B$ ,  $B^+ = B \cap \{b^+ \mid b \in 2^*\}$ . Also,  $B$  is positive if  $B$  contains no truth value of the form  $\neg b'$ .

- DEFINITION 4.8.
1.  $\neg b' \Vdash^* B$  iff  $c^+ \in B \rightarrow c^+ \not\subseteq b'$ ,  $c' \in B \rightarrow c \neq b$ , and  $\neg c' \in B \rightarrow c = b$ .
  2.  $\sigma^{-b'} = \{\sigma_i^{-b'} \mid \text{for some } \langle B_i, \sigma_i \rangle \in \sigma, \neg b' \Vdash^* B_i\}$ .
  3. For  $\phi(\sigma_1, \dots, \sigma_n)$  in the language of the Kripke model, that is, with parameters (displayed) terms,  $\phi^{-b'} = \phi(\sigma_1^{-b'}, \dots, \sigma_n^{-b'})$ .
  4.  $\neg b' \Vdash^* \phi$ , for  $\phi$  in the language of the Kripke model, if  $\phi^{-b'}$  is true (i.e., in  $V$ ). Note that  $\phi^{-b'}$  is a formula with set parameters.

DEFINITION 4.9.  $q \leq_W p$  ( $q$  is a weakening of  $p$  as conditions) if for  $b \in \text{dom}(p)$  either  $p(b) = \infty$  or for some initial segment  $c$  of  $b$   $q(c) = \text{IN}$ .

The idea behind this definition is the  $q$  may change some  $\infty$ 's to  $\text{IN}$ 's, as well as extend the domain of  $p$ . Notice that  $\leq_W$  is a partial order, and  $\text{inpart}(p, q)$ , from lemma 14, is the greatest lower bound of  $p$  and  $q$ .

DEFINITION 4.10.  $p \Vdash^* \phi$ , for  $\phi$  in the language of the Kripke model, i.e., when  $\phi$ 's parameters are terms, inductively on  $\phi$  :

- $p \Vdash^* \sigma \in \tau$  if for some  $\langle B_i, \tau_i \rangle \in \tau$  with  $B_i$  positive,  $B_i^+ \subseteq B_p$  and  $p \Vdash^* \sigma = \tau_i$ .
- $p \Vdash^* \sigma = \tau$  if
  - i) for all  $\langle B_i, \sigma_i \rangle \in \sigma$  and  $q \leq_W p$ , if  $B_i$  is positive and  $B_i^+ \subseteq B_q$  then there is an  $r \leq q$  such that  $r \Vdash^* \sigma_i \in \tau$ , and symmetrically between  $\sigma$  and  $\tau$ , and
  - ii) for all  $b \notin \text{dom}(p)$ , if for no initial segment  $c$  of  $b$  is  $c^+$  in  $B_p$ , then  $\neg b' \Vdash^* \sigma = \tau$ .
- $p \Vdash^* \phi \wedge \theta$  if  $p \Vdash^* \phi$  and  $p \Vdash^* \theta$ .
- $p \Vdash^* \phi \vee \theta$  if  $p \Vdash^* \phi$  or  $p \Vdash^* \theta$ .
- $p \Vdash^* \phi \rightarrow \theta$  if
  - i) for all  $q \leq_W p$  if  $q \Vdash^* \phi$  then there is an  $r \leq q$  such that  $r \Vdash^* \theta$ , and
  - ii) for all  $b \notin \text{dom}(p)$ , if for no initial segment  $c$  of  $b$  is  $c^+$  in  $B_p$ , then  $\neg b' \Vdash^* \phi \rightarrow \theta$ .
- $p \Vdash^* \exists x \phi(x)$  if for some term  $\sigma$   $p \Vdash^* \phi(\sigma)$ .
- $p \Vdash^* \forall x \phi(x)$  if
  - i) for all terms  $\sigma$  and  $q \leq_W p$ , there is an  $r \leq q$  such that  $r \Vdash^* \phi(\sigma)$ , and
  - ii) for all  $b \notin \text{dom}(p)$ , if for no initial segment  $c$  of  $b$  is  $c^+$  in  $B_p$ , then  $\neg b' \Vdash^* \forall x \phi(x)$ .

PROPOSITION 4.11. If  $p \Vdash^* \phi$  and  $q \leq_W p$  then  $q \Vdash^* \phi$ .

PROOF. Trivial induction on  $\phi$ . ⊖

LEMMA 4.12.  $\perp \models \phi$  iff  $p \Vdash^* \phi$  for some  $p \in G$ .

PROOF. Inductively on  $\phi$ .

$\sigma \in \tau$ :  $\perp \models \sigma \in \tau$  iff for some  $\langle B_i, \tau_i \rangle \in \tau$  every member of  $B_i$  is true at  $\perp$  and  $\perp \models \sigma = \tau_i$ . The former clause holds iff  $B_i$  is positive and, for some  $p \in G$ ,  $B_i \subseteq B_p$ . Inductively, the latter clause holds iff, for some  $q \in G$ ,  $B_q \Vdash^* \sigma = \tau_i$ . Given such  $p$  and  $q$ ,  $p \cup q$  suffices. The converse direction is immediate.

$\sigma = \tau$ : Suppose  $p \in G$  and  $p \Vdash^* \sigma = \tau$ . If  $q \in f(P)$  indexes a node then  $q \leq_W p$ . If  $q \Vdash \rho \in \sigma$  then inductively there is a  $q' \in f(G)_q$ ,  $q' \leq q$ , such that  $q' \Vdash^* \rho = \sigma_i \wedge \sigma_i \in f(\sigma)$  for some  $\langle B_i, \sigma_i \rangle \in f(\sigma)$ . By *i*) of the case hypothesis, there is an  $r \leq q'$  with  $r \Vdash^* \rho \in \tau$ . Generically, there is such an  $r$  in  $f(G)$ , so inductively  $q \Vdash \rho \in \tau$ . If  $c$  indexes a node, then by *ii*) of the case hypothesis  $c \Vdash \sigma = \tau$ . Hence  $\perp \Vdash \sigma = \tau$ .

Conversely, suppose there is no such  $p \in G$ . If  $p \not\Vdash^* \sigma = \tau$  because clause *i*) fails, then there is a witness  $q \leq_W p$  to that failure. We say that such a  $q$  is close to  $p$  if  $\text{dom}(q) \subseteq \text{dom}(p)$ . That means that  $q$  comes from  $p$  by changing some  $\infty$ 's to IN's and not adding anything else. Observe that if *i*) fails for  $p$ , then  $p$  can be extended to  $p'$  so that *i*) fails for  $p'$  via a witness  $q$  close to  $p'$ . That's because  $\text{dom}(p')$  can be taken to be  $\text{dom}(p) \cup \text{dom}(q)$ , for  $b \in \text{dom}(p)$   $p'(b)$  can be taken to be  $p(b)$ , and for  $b \in \text{dom}(q) \setminus \text{dom}(p)$   $p'(b)$  can be taken to be  $q(b)$ . Therefore  $D = \{p \mid p \Vdash^* \sigma = \tau, \text{ or } p \text{ violates } i) \text{ with a witness } q \text{ close to } p, \text{ or } p \text{ violates } ii)\}$  is dense.

Suppose there were a  $p \in G$  violating *i*) with a witness  $q$  close to  $p$ . Then  $q$  induces a legal weakening  $f(G)_q$  of  $f(G)$ , and so indexes a node. By the choice of  $q$ ,  $q \Vdash \sigma_i \in \sigma$ . If  $q \Vdash \sigma_i \in \tau$  then inductively that would be  $*$ -forced by some  $r \leq q$ . But by the choice of  $q$  there is no such  $r$ . Hence we would have  $q \not\Vdash \sigma = \tau$ .

If there is no such  $p$  then every  $p \in G$  violates *ii*). Let  $p \in f(G)$  be such that  $p \supseteq G$ . Since *ii*) fails for that  $p$ , then, with  $b$  from that failure,  $b$  indexes a node,  $b \not\Vdash \sigma = \tau$ . In either case,  $\perp \not\Vdash \sigma = \tau$ .

$\phi \wedge \theta$ : Trivial.

$\phi \vee \theta$ : Trivial.

$\phi \rightarrow \theta$ : Suppose  $p \in G$  and  $p \Vdash^* \phi \rightarrow \theta$ . If  $q \Vdash \phi$  then inductively, for some  $q' \in f(G)_q$ ,  $q' \Vdash^* \phi$ . Since we can take  $q' \leq q \leq_W p$ , by *i*) of the hypothesis there is an  $r \leq q$  such that  $r \Vdash^* \theta$ . By genericity, there is such an  $r$  in  $f(G)_q$ . Hence  $q \Vdash \theta$ . If  $c \Vdash \phi$  then use *ii*) of the hypothesis.

Conversely, suppose there is no such  $p \in G$ . If  $p \not\Vdash^* \phi \rightarrow \theta$  because clause *i*) fails, then there is a witness  $q \leq_W p$  to that failure, in which case  $p$  can be extended to  $p'$  so that *i*) fails for  $p'$  via a witness  $q$  close to  $p'$ , where closeness is as defined above in the case for =, for the same reason as above. Therefore  $D = \{p \mid p \Vdash^* \phi \rightarrow \theta, \text{ or } p \text{ violates } i) \text{ with a witness } q \text{ close to } p, \text{ or } p \text{ violates } ii)\}$  is dense.

Suppose there were a  $p \in G$  violating *i*) with a witness  $q$  close to  $p$ . Then  $q$  induces a legal weakening  $f(G)_q$  of  $f(G)$ , and so indexes a node. By the choice of  $q$ ,  $q \Vdash \phi$ . If  $q \Vdash \theta$  then inductively that would be  $*$ -forced by some  $r \leq q$ . But by the choice of  $q$  there is no such  $r$ . Hence we would have  $q \not\Vdash \phi \rightarrow \theta$ .

If there is no such  $p$  then every  $p \in G$  violates *ii*). Let  $p \in f(G)$  be such that  $p \supseteq G$ . Since *ii*) fails for that  $p$ , then, with  $b$  from that failure,  $b$  indexes a node and  $b \not\Vdash \phi \rightarrow \theta$ . In either case,  $\perp \not\Vdash \phi \rightarrow \theta$ .

$\exists x \phi(x)$ : Trivial.

$\forall x \phi(x)$ : As in the cases for = and  $\rightarrow$ . ⊣

LEMMA 4.13.  $\perp \Vdash IZF$ .

PROOF. Just as in the last section, most of the axioms have soft proofs in this model. The only issue is with Separation. Given  $\phi$  and  $\sigma$ , let  $\text{Sep}_{\phi, \sigma}$  be  $\{\langle B, \tau \rangle \mid$

for some  $\langle B', \tau \rangle \in \sigma$  with  $B \supseteq B'$  either *i*)  $B = B_p \Vdash \phi(\sigma)$ , or *ii*)  $\neg b' \in B$  and  $\neg b' \Vdash^* \phi$ . By the previous lemma, this works.  $\dashv$

As in the previous section, this model does not satisfy  $FAN_c$ , but does satisfy  $\neg FAN_c$ . For further discussion, see the questions at the end.

**§5.  $FAN_c$  does not imply  $FAN_{\Pi^0_1}$ .** Let  $G$  be  $P$ -generic exactly as in the last section. By convention, we say that if  $G(\alpha) = IN$  then  $G$  applied to any extension of  $\alpha$  is also  $IN$ . Our goal is to hide  $G$  a bit better than before, so  $FAN_c$  remains true, but not too well, so that  $FAN_{\Pi^0_1}$  is false.

Let  $N$  be an ultrapower of  $M[G]$  using a non-principal ultrafilter on  $\omega$ . The Kripke model has a bottom node  $\perp$ , and the successors of  $\perp$  are indexed by the labels  $\langle n, \alpha \rangle$ , where  $n$  is a non-standard integer, and  $\alpha \in 2^*$  either has non-standard length or  $G(\alpha) = \infty$ .

**DEFINITION 5.1.** A truth value is a symbol of the form  $\langle n, \alpha \rangle$ ,  $\neg \langle n, \alpha \rangle$ , or  $\langle \forall n, \alpha \rangle$ , for  $n$  a natural number (in the first two cases) and  $\alpha \in 2^*$ . Admittedly truth values of the first kind are also used to index nodes; whether truth values or nodes are intended in any particular case should be clear from the context. Terms are defined inductively (through the ordinals) as sets of the form  $\{\langle B_i, \sigma_i \rangle \mid i \in I\}$ , where  $I$  is any index set,  $\sigma_i$  a term, and  $B_i$  a finite set of truth values.

The sets at  $\perp$  will be the terms in  $M$ . The sets at any other node will be analogous, that is, the terms in what  $N$  thinks is the ground model, i.e.,  $\bigcup_{\kappa \in ORD} f(M_\kappa)$ . At  $\perp$ ,  $\langle n, \alpha \rangle$  will always be true,  $\neg \langle n, \alpha \rangle$  always false, and  $\langle \forall n, \alpha \rangle$  true exactly when  $G(\alpha) = IN$ . At node  $\langle m, \beta \rangle$ ,  $\langle n, \alpha \rangle$  is true exactly when  $\langle n, \alpha \rangle \neq \langle m, \beta \rangle$ ,  $\neg \langle n, \alpha \rangle$  is true exactly when  $\langle n, \alpha \rangle = \langle m, \beta \rangle$ , and  $\langle \forall n, \alpha \rangle$  true exactly when  $\alpha \neq \beta$ . (Note that, perhaps perversely, the node  $\langle n, \alpha \rangle$  is exactly the node at which the truth value  $\langle n, \alpha \rangle$  is false. The reason behind this choice is that the node  $\langle n, \alpha \rangle$  is where something special happens to the corresponding truth value. If preferred, the reader can call that node  $\neg \langle n, \alpha \rangle$  instead.) This interpretation of the truth values induces an interpretation of the terms at all nodes.

Let  $T_n$  be the term  $\{\langle \langle n, \alpha \rangle \rangle, \hat{\alpha} \mid \alpha \in 2^*\}$ . Let  $C$  be a term naming the function that on input  $n$  returns  $T_n$ .  $T_n$  at  $\perp$  looks like the full tree  $2^*$ ;  $T_n$  at  $\langle n, \alpha \rangle$  looks like everything except  $\alpha$ ; and  $T_n$  at  $\langle m, \alpha \rangle$ ,  $m \neq n$ , again looks like  $2^*$ . The term for  $\bigcap_n C(n)$  is given by  $\{\langle \langle \forall n, \alpha \rangle \rangle, \hat{\alpha} \mid \alpha \in 2^*\}$ , and is interpreted as  $\{\alpha \mid G(\alpha) = IN\}$  at  $\perp$  and  $2^* \setminus \{\alpha\}$  at  $\langle n, \alpha \rangle$ . Notice that  $\bigcap_n C(n)$  is not closed under extensions.

The proof will be finished once we show that, at  $\perp$ ,  $FAN_c$  holds,  $IZF$  holds, and  $\bigcap_n C(n)$  is a counter-example to  $FAN_{\Pi^0_1}$ .

**LEMMA 5.2.**  $\perp \not\models FAN_{\Pi^0_1}$ .

**PROOF.** It is clear that  $T_n$  is decidable, and so  $\bigcap_n C(n)$  is on the face of it  $\Pi^0_1$ . It is also clear that  $\bigcap_n C(n)$  is not a uniform bar. So it suffices to show that  $\perp \Vdash$  “ $\bigcap_n C(n)$  is a bar.”

Let  $\perp \models$  “ $Br$  is a branch through  $2^*$ .” (Without loss of generality, it suffices to start at  $\perp$  instead of at an arbitrary node.) Work beneath a condition forcing that, so we can assume  $Br$  consists of sets of the form  $\langle B_i, \hat{\alpha} \rangle$ , for various  $\alpha \in 2^*$ . If the standard part of  $Br$ , the part visible at  $\perp$ , is in the ground model  $M$ , then, by the



genericity of  $G$ ,  $Br$  will hit  $G$  (i.e., for some  $\alpha \in Br$ ,  $G(\alpha) = \text{IN}$ ), which is how  $\perp$  interprets  $\bigcap_n C(n)$ . If the standard part of  $Br$  were not in  $M$ , then contradictory facts about  $Br$  would be forced by different forcing conditions. In particular, we would have  $p, q$ , and  $\alpha$  with  $p \Vdash \perp \models \alpha \frown 0 \in Br$  and  $q \Vdash \perp \models \alpha \frown 1 \in Br$ . That means there are  $\langle B_p, \widehat{\alpha \frown 0} \rangle \in Br$  and  $\langle B_q, \widehat{\alpha \frown 1} \rangle \in Br$ , with  $B_p$  and  $B_q$  consisting only of truth values automatically true at  $\perp$  save for some of the form  $\langle \forall n, \alpha \rangle$ . But at some node  $\langle n, \alpha \rangle$  with  $\alpha$  non-standard, all of those latter truth values will be true. Hence  $\langle n, \alpha \rangle \models \widehat{\alpha \frown 0}, \widehat{\alpha \frown 1} \in Br$ , so  $\perp$  could not force  $Br$  to be a path.  $\dashv$

In order to prove the other facts, we will need to deal with truth at  $\perp$ .

**DEFINITION 5.3.** For a forcing condition  $p$ , let  $|p|$ , the length of  $p$ , be the length of the longest  $\alpha \in \text{dom}(p)$ . Let  $B_p$  be  $\{\langle n, \alpha \rangle \mid n, \text{length}(\alpha) \leq |p|\} \cup \{\langle \forall n, \alpha \rangle \mid \text{length}(\alpha) \leq |p|\}$  and, for some initial segment  $\beta$  of  $\alpha$ ,  $p(\beta) = \text{IN}$ .

- DEFINITION 5.4.**
1. For  $B$  a finite set of truth values,  $\neg \langle n, \alpha \rangle \Vdash^* B$  iff  $\langle n, \alpha \rangle \notin B$ ,  $\langle \forall n, \alpha \rangle \notin B$ , and the only truth value of the form  $\neg \langle m, \beta \rangle$  in  $B$  is  $\neg \langle n, \alpha \rangle$  itself.
  2.  $\sigma \neg \langle n, \alpha \rangle = \{\sigma_i \neg \langle n, \alpha \rangle \mid \text{for some } \langle B_i, \sigma_i \rangle \in \sigma, \neg \langle n, \alpha \rangle \Vdash^* B_i\}$ .
  3. For  $\phi(\sigma_1, \dots, \sigma_n)$  in the language of the Kripke model, that is, with parameters (displayed) terms,  $\phi \neg \langle n, \alpha \rangle = \phi(\sigma_1 \neg \langle n, \alpha \rangle, \dots, \sigma_n \neg \langle n, \alpha \rangle)$ .
  4.  $\neg \langle n, \alpha \rangle \Vdash^* \phi$ , for  $\phi$  in the language of the Kripke model, if  $\phi \neg \langle n, \alpha \rangle$  is true (i.e., in  $V$ ). Note that  $\phi \neg \langle n, \alpha \rangle$  is a formula with set parameters.

**DEFINITION 5.5.**  $p \Vdash^* \phi$ , for  $\phi$  in the language of the Kripke model, i.e., when  $\phi$ 's parameters are terms, inductively on  $\phi$ :

- $p \Vdash^* \sigma \in \tau$  if, for some  $\langle B_i, \tau_i \rangle \in \tau$ ,  $B_i \subseteq B_p$  and  $p \Vdash^* \sigma = \tau_i$ .
- $p \Vdash^* \sigma = \tau$  if
  - i) for all  $\langle B_i, \sigma_i \rangle \in \sigma$  and  $q \leq p$ , if  $B_i \subseteq B_q$  then there is an  $r \leq q$  such that  $r \Vdash^* \sigma_i \in \tau$ , and symmetrically between  $\sigma$  and  $\tau$ , and
  - ii) if  $n > |p|$ , and if either  $\text{length}(\alpha) > |p|$  or for no initial segment  $\beta$  of  $\alpha$  do we have  $p(\beta) = \text{IN}$ , then  $\neg \langle n, \alpha \rangle \Vdash^* \sigma = \tau$ .
- $p \Vdash^* \phi \wedge \theta$  if  $p \Vdash^* \phi$  and  $p \Vdash^* \theta$ .
- $p \Vdash^* \phi \vee \theta$  if  $p \Vdash^* \phi$  or  $p \Vdash^* \theta$ .
- $p \Vdash^* \phi \rightarrow \theta$  if
  - i) for all  $q \leq p$ , if  $q \Vdash^* \phi$  then there is an  $r \leq q$  such that  $r \Vdash^* \theta$ , and
  - ii) if  $n > |p|$ , and if either  $\text{length}(\alpha) > |p|$  or for no initial segment  $\beta$  of  $\alpha$  do we have  $p(\beta) = \text{IN}$ , then  $\neg \langle n, \alpha \rangle \Vdash^* \phi \rightarrow \theta$ .
- $p \Vdash^* \exists x \phi(x)$  if for some term  $\sigma$   $p \Vdash^* \phi(\sigma)$ .
- $p \Vdash^* \forall x \phi(x)$  if
  - i) for all terms  $\sigma$  and  $q \leq p$ , there is an  $r \leq q$  such that  $r \Vdash^* \phi(\sigma)$ , and
  - ii) if  $n > |p|$ , and if either  $\text{length}(\alpha) > |p|$  or for no initial segment  $\beta$  of  $\alpha$  do we have  $p(\beta) = \text{IN}$ , then  $\neg \langle n, \alpha \rangle \Vdash^* \forall x \phi(x)$ .

**PROPOSITION 5.6.** If  $p \Vdash^* \phi$  and  $q \leq p$  then  $q \Vdash^* \phi$ .

**PROOF.** Trivial induction on  $\phi$ .  $\dashv$

**LEMMA 5.7.**  $\perp \models \phi$  iff  $p \Vdash^* \phi$  for some  $p \in G$ .

**PROOF.** Inductively on  $\phi$ .

$\sigma \in \tau: \perp \models \sigma \in \tau$  iff for some  $\langle B_i, \tau_i \rangle \in \tau$  every member of  $B_i$  is true at  $\perp$  and  $\perp \models \sigma = \tau_i$ . The former clause holds iff  $B_i$  contains nothing of the form  $\neg \langle n, \alpha \rangle$ , and if  $(\forall n, \alpha) \in B_i$  then  $G(\alpha) = \text{IN}$ . Given such a  $B_i$ , let  $p$  be a sufficiently long initial segment of  $G$  forcing “ $\sigma = \tau_i$ .” Such a  $p$  suffices. The converse direction is immediate.

$\sigma = \tau$ : Suppose  $p \in G$  and  $p \Vdash^* \sigma = \tau$ . Then any member of  $\sigma$  at  $\perp$  is equal at  $\perp$  to some  $\sigma_i$ , where  $\langle B_i, \sigma_i \rangle \in \sigma$  and  $B_i \subseteq B_q$  for some  $q \in G$ . Then by the hypothesis and genericity there will be an extension  $r$  of  $q$  in  $G$  forcing  $\sigma_i$  to be in  $\tau$ . At any other node  $\langle n, \alpha \rangle$ , working in  $N$ ,  $n$  is non-standard and so greater than  $|p|$ , and  $\alpha$  also satisfies the conditions in *ii*) (of the definition of  $*$ -forcing equality), so “ $\sigma = \tau$ ” is true at these other nodes too.

Conversely, suppose there is no such  $p \in G$ . With reference to the definition of  $*$ -forcing equality, observe that  $\{p \mid p \text{ satisfies clause } i)\} \cup \{p \mid \text{for some } \langle B_i, \sigma_i \rangle \in \sigma, B_i \subseteq B_p, \text{ yet } p \text{ has no extension } * \text{-forcing } \sigma_i \text{ into } \tau\}$  is dense. If  $G$  contains a member of the second set of that union, then the induced  $\sigma_i$  witnesses that  $\perp \not\models \sigma = \tau$ . If not, then  $G$  contains  $p$  satisfying *i*). So no  $p \in G$  satisfies *ii*). This also holds in  $N$ . In  $N$ , take  $p$  to be an initial segment of  $G$  of non-standard length. The failure of *ii*) for that  $p$  produces an  $n$  and  $\alpha$  which index a node at which  $\sigma \neq \tau$ , showing  $\perp \not\models \sigma = \tau$ .

$\phi \wedge \theta$ : Trivial.

$\phi \vee \theta$ : Trivial.

$\phi \rightarrow \theta$ : Suppose  $p \in G$  and  $p \Vdash^* \phi \rightarrow \theta$ . Then it is direct that  $\perp \models \phi \rightarrow \theta$ .

Conversely, suppose there is no such  $p \in G$ . With reference to the definition of  $*$ -forcing implication, observe that  $\{p \mid p \text{ satisfies clause } i)\} \cup \{p \mid p \Vdash^* \phi \text{ yet } p \text{ has no extension } * \text{-forcing } \theta\}$  is dense. If  $G$  contains a member of the second set of that union, then inductively  $\perp \models \phi$  and  $\perp \not\models \theta$ , hence  $\perp \not\models \phi \rightarrow \theta$ . If not, then  $G$  contains  $p$  satisfying *i*). So no  $p \in G$  satisfies *ii*). This also holds in  $N$ . In  $N$ , take  $p$  to be an initial segment of  $G$  of non-standard length. The failure of *ii*) for that  $p$  produces an  $n$  and  $\alpha$  which index a node at which  $\phi \rightarrow \theta$  is false, showing  $\perp \not\models \phi \rightarrow \theta$ .

$\exists x \phi(x)$ : Trivial.

$\forall x \phi(x)$ : As in the cases for  $=$  and  $\rightarrow$ . ¬

LEMMA 5.8.  $\perp \models \text{FAN}_c$ .

PROOF. Suppose that at  $\perp$  we have a decidable set  $C' \subseteq 2^*$  inducing a  $c$ -bar  $C$ . We would like to show that at  $\perp$  the  $c$ -bar  $C$  is uniform, which means that, for some  $k$ ,  $C$  contains every sequence of length at least  $k$ ; in notation,  $C \supseteq 2^{\geq k}$ . This is equivalent with  $C'$  containing  $2^{\geq k}$ , which is what we will prove.

Say that  $\alpha \in 2^*$  is *good* if there is a natural number  $k$  such that, whenever  $n \geq k$  and  $\beta \supseteq \alpha$  has length at least  $k$ ,  $\neg \langle n, \beta \rangle \Vdash^* C' \supseteq 2^{\geq k}$ . Observe that if  $\alpha \frown 0$  and  $\alpha \frown 1$  are good then so is  $\alpha$  (by taking  $k$  sufficiently large). So if the empty sequence  $\langle \rangle$  is bad (i.e., not good) then there is a branch  $Br_0$  of bad nodes. For each  $\alpha \in Br_0$ , by the definition of badness, taking  $k$  to be the length  $|\alpha|$  of  $\alpha$ , we have some  $\beta \supseteq \alpha$  and  $n \geq |\alpha|$  such that  $\neg \langle n, \beta \rangle \not\Vdash^* C' \supseteq 2^{\geq k}$ . Because  $\neg \langle n, \beta \rangle \Vdash^* \phi$  is defined as the truth of  $\phi^{\neg \langle n, \beta \rangle}$  in the classical universe  $V$ , we can reason classically and conclude that there is a  $\gamma \in 2^{\geq k}$  such that  $\neg \langle n, \beta \rangle \Vdash^* \gamma \notin C'$ . By choosing  $\alpha$ 's of increasing length, we can get infinitely many  $\gamma$ 's of increasing length, in particular

infinitely many distinct  $\gamma$ 's. Hence there is a branch  $Br_1$  such that each node in  $Br_1$  has infinitely many such  $\gamma$ 's as extensions.

That was all in  $M$ . Now with reference to  $N$ , if  $\alpha \in Br_0^N$  has standard length, then the corresponding choice of  $\gamma$  is also standard, since it's the same  $\gamma$  as in  $M$ . So if we choose a non-standard  $\gamma$  coming from the procedure above, that  $\gamma$  came from a non-standard  $\alpha$ . Since  $N \models$  " $Br_1$  is infinite," there is a non-standard node on  $Br_1^N$ , with some such  $\gamma$  as an extension; since the node chosen from  $Br_1^N$  was non-standard, so is  $\gamma$ , and hence so is the  $\alpha$  that  $\gamma$  came from. From  $\alpha$ , we also have  $\beta \supseteq \alpha$  and  $n \geq |\alpha|$  with  $\neg \langle n, \beta \rangle \Vdash^* \gamma \notin C'$ . In particular,  $\langle n, \beta \rangle$  indexes a node in the model. But at  $\perp$ ,  $C'$  induces a  $c$ -bar, so  $\perp \models$  "there is a node  $\delta \in Br_1$  such that  $\delta \in C$ ; that is, every extension of  $\delta$  is in  $C'$ ." This contradicts the choice of  $\gamma$ .

We conclude from this that  $\langle \rangle$  is good. Fix  $k$  witnessing this goodness. We will show  $\perp \models C' \supseteq 2^{\geq k}$ .

First, if  $\delta \in 2^{\geq k}$  is standard, then, for any  $n$  and  $\beta$  non-standard,  $\neg \langle n, \beta \rangle \Vdash^* \delta \in C'$ , so, with reference to the Kripke node  $\langle n, \beta \rangle$ ,  $\langle n, \beta \rangle \models \delta \in C'$ . Since  $C'$  is decidable,  $\perp \models \delta \in C'$ .

To finish the argument, we need only consider nodes  $\langle n, \beta \rangle$ , and show  $\langle n, \beta \rangle \models C' \supseteq 2^{\geq k}$ . If  $\beta$  has length at least  $k$ , this follows from the goodness of  $\langle \rangle$ . The only other case is  $\beta$  of length less than  $k$  such that  $G(\beta) = \infty$ . It suffices to show that, for any such fixed  $\beta$ , in  $M$  there is a finite  $n$  such that, for all  $m \geq n$ ,  $\neg \langle m, \beta \rangle \Vdash^* C' \supseteq 2^{\geq k}$ .

Toward that end, suppose not. Then for infinitely many  $m$  there is a  $\gamma$  of length at least  $k$  such that  $\neg \langle m, \beta \rangle \Vdash^* \gamma \notin C'$ . If those  $\gamma$ 's are of bounded length then one occurs infinitely often. For that fixed  $\gamma$ , by overspill there is a non-standard  $m$  such that  $\neg \langle m, \beta \rangle \Vdash^* \gamma \notin C'$ . But  $\langle m, \beta \rangle$  is a Kripke node, and we already saw that, for  $\delta \in 2^{\geq k}$ ,  $\perp \models \delta \in C'$ , which is a contradiction. Hence there are infinitely many different  $\gamma$ 's. That means there is a branch  $Br_2$  such that every node on  $Br_2$  has infinitely many different  $\gamma$ 's as extensions. Pick a non-standard  $m$  such that the corresponding  $\gamma$  extends a non-standard node of  $Br_2$ . But again,  $\perp \models$  " $C$  is a  $c$ -bar," so  $\perp \models$  "there is a node  $\delta \in Br_2$  such that  $\delta \in C$ ; i.e., every extension of  $\delta$  is in  $C'$ ." This contradicts the choice of  $\gamma$ .  $\dashv$

LEMMA 5.9.  $\perp \models IZF$ .

PROOF. As before, all of the axioms have soft proofs, save for Separation. Given  $\phi$  and  $\sigma$ , let  $\text{Sep}_{\phi, \sigma}$  be  $\{ \langle B_i \cup B_p, \tau \rangle \mid \langle B_i, \tau \rangle \in \sigma \text{ and } p \Vdash^* \phi(\tau) \} \cup \{ \langle B, \tau \rangle \mid \text{for some } \neg \langle n, \alpha \rangle \in B \text{ and some } B_i, \langle B_i, \tau \rangle \in \sigma, \neg \langle n, \alpha \rangle \Vdash^* B_i, \text{ and } \neg \langle n, \alpha \rangle \Vdash^* \phi(\tau) \}$ . By lemma 5.7, this works.  $\dashv$

**§6.  $\text{FAN}_{\Pi_1^0}$  does not imply  $\text{FAN}_{\text{full}}$ .** As usual, let  $G$  be generic as above. In  $M[G]$ , the Kripke model will have bottom node  $\perp$ , and successor nodes labeled by those  $\alpha \in 2^*$  such that  $G(\alpha) = \infty$ . As is standard, terms are defined inductively, and always subject to the usual restrictions so as to have a Kripke model. That is, to define the full model [9] over any partial order  $\langle P, < \rangle$ , at node  $p \in P$  a term  $\sigma$  is any function with domain  $P^{\geq p}$  such that  $\sigma(q)$  is a set of terms at node  $q$ ; furthermore, with transition function  $f_{qr}$  for  $q < r$ , if  $\tau \in \sigma(q)$  then  $f_{qr}(\tau) \in \sigma(r)$ ; finally,  $f_{pq}$  is extended to  $\sigma$  by restriction:  $f_{pq}(\sigma) = \sigma \upharpoonright P^{\geq q}$ . That is called the full model, because everything possible is being thrown in. For the current construction, we will

take a sub-model of the full model by imposing one additional restriction: a term at any node  $\alpha$  other than  $\perp$  must be in the ground model  $M$ .

Let  $C$  be the term such that  $\perp \models \hat{\beta} \in C$  ( $\beta \in 2^*$ ) iff, for some initial segment  $\beta \upharpoonright n$  of  $\beta$ ,  $G(\beta \upharpoonright n) = \text{IN}$ , and, at node  $\alpha \neq \perp$ ,  $\alpha \models \hat{\beta} \in C$  iff  $\beta$  is not an initial segment of  $\alpha$ .

LEMMA 6.1.  $\perp \models \text{FAN}_{\Pi^0_1}$ .

PROOF. If  $\perp \models \text{“}B \subseteq 2^* \text{ is decidable”}$  then, for any  $\beta \in 2^*$ ,  $\perp \models \hat{\beta} \in B$  iff, for some node  $\alpha \neq \perp$ ,  $\alpha \models \hat{\beta} \in B$  iff the same holds for all  $\alpha \neq \perp$ . Hence  $\perp \models \text{“}B = \hat{B}_M \text{”}$  for some set  $B_M \in M$ . So if  $\perp \models \text{“}B_n \text{ is a sequence of decidable trees,”}$  then that sequence is the image of a sequence of sets from  $M$ . Hence their intersection internally is the image of a set from  $M$ . So if  $\bigcap_n B_n$  is internally a bar, it is the image of a bar, and by the Fan Theorem in  $M$  is uniform.  $\dashv$

LEMMA 6.2.  $\perp \not\models \text{FAN}_{\text{full}}$ .

PROOF. At  $\perp$ ,  $C$  is not uniform, so it suffices to show  $\perp \not\models \text{“}C \text{ is a bar”}$ . If  $\perp \models \text{“}P \text{ is a path through } 2^* \text{”}$  then  $\perp \models \text{“}P \text{ is decidable”}$ , and as above  $P$  is then the image of a ground model path. Generically, for some  $\beta$  along that path,  $G(\beta) = \text{IN}$ . For that  $\beta$ ,  $\perp \models \text{“}P \text{ goes through } \hat{\beta} \text{ and } \hat{\beta} \in C \text{”}$ .  $\dashv$

LEMMA 6.3.  $\perp \models \text{IZF}$ .

PROOF. Not only are most of the axioms trivial to verify, in this case even Separation is too. Given a formula  $\phi$ , term  $\sigma$ , and node  $nd$ , let  $\text{Sep}_{\phi,\sigma}(nd) = \{\tau \mid \tau \in \sigma \text{ and } nd \models \phi(\tau)\}$ . The reason that at node  $\alpha$  this is in  $M$  is that, at  $\alpha$ ,  $\phi$ 's parameters can also be interpreted in  $M$ , and so truth at  $\alpha$  is definable in  $M$ .  $\dashv$

§7. Questions.

- We have seen that  $\text{FAN}_{\text{full}}$  holds in every topological model, and that  $\text{FAN}_{\Pi^0_1}$  holds in the model over any connected Heyting algebra. Are there any other sufficient or necessary properties for any of the various fan theorems we have been considering to hold or fail in a Heyting-valued model?
- As a particular instance of the previous question, if a Heyting algebra satisfies  $\text{FAN}_{\Delta}$  (respectively  $\text{FAN}_c$ ), does it automatically satisfy  $\text{FAN}_c$  (respectively  $\text{FAN}_{\Pi^0_1}$ )?
- Although we were not able to make use of any Heyting algebras other than  $\Omega$ , some seem worthwhile to investigate, as possibly separating some of these fan theorems, or perhaps having some other interesting properties. We would include among these  $K(T)$  for various natural spaces  $T$ , such as  $2^{\mathbb{N}}$ . We would also include other ways of killing points, such as over a measure space  $\tau$  with measure  $\lambda$  modding out by sets of measure 0:

$$U \sim V \iff \lambda(U) = \lambda(V) = \lambda(V \cap U)$$

(two opens are equivalent if their symmetric difference is of measure zero). The space  $\tau / \sim$  should be a Heyting algebra, which we will denote by analogy with  $K$  as  $L(\tau)$ . Of particular interest seem to be  $L(I)$  and  $L(I \times I)$ .

- In the models presented here, the principles in question were not true. That's different from their being false (meaning their negations being true). We expect this could be done by iterating the constructions presented here. That is, to each

terminal node of the model append another model of the same kind, starting with the ambient universe of that terminal node as the new ground model. By iterating this procedure infinitely often, one is left with a Kripke model with no terminal nodes. In order still to have a model of IZF, to get the Power Set Axiom for instance, terms for all of these bars from the iteration might have to be present at  $\perp$ , or perhaps some other fix would work. So this suggestion would at least take some work to implement, and might even demand some new ideas.

It would be even better, or at least different, if we had a model with one fixed counter-example. Maybe the models presented here could be so tweaked. For instance, for  $\text{FAN}_\Delta$ , could we just throw away the terminal nodes? For  $\text{FAN}_c$ , it might work not to stop a node just because  $\neg b'$  is true, but rather to continue extending the node to allow finitely many  $\neg b'$ 's to be true. Or maybe a more radical idea is needed.

- One of the referees asked about the role of Choice here. It is not that hard to see that Dependent Choice fails in most (or all) of these models. Are there some nice choice principles that are true here? Are there other models in which DC or other choice principles of interest hold? Are there significant fragments of Choice that are incompatible with these separations?
- Within reverse classical mathematics, many weakenings of Weak König's Lemma (classically equivalent to the Fan Theorem) have been identified. Of interest to us here is Weak Weak König's Lemma. Whereas WKL states that any bar (closed under extension, for simplicity) contains an entire level of  $2^*$ , WWKL states that any bar contains half of a level. (WWKL has been shown to be connected to the development of measure theory.) In our context, any of the principles we have been considering could be so weakened, yielding Weak  $\text{FAN}_\Delta$ , Weak  $\text{FAN}_c$ , Weak  $\text{FAN}_{\Pi_1^0}$ , and Weak  $\text{FAN}_{\text{full}}$ . Clearly any principle implies its weak correlate (e.g.,  $\text{FAN}_{\Pi_1^0}$  implies Weak  $\text{FAN}_{\Pi_1^0}$ ), and any weak principle implies the weak principles lower down (e.g., Weak  $\text{FAN}_{\Pi_1^0}$  implies Weak  $\text{FAN}_c$ ), forming a bit of a square. Are there any implications along the diagonal (e.g., between  $\text{FAN}_c$  and Weak  $\text{FAN}_{\Pi_1^0}$ )? Are these weak principles even natural or interesting, by being equivalent with interesting theorems?
- Are there any other interesting principles to be found here, for instance  $\Pi_n^0$ -FAN for  $n > 1$ , or adaptations of reverse math principles beneath WKL other than WWKL?

#### REFERENCES

- [1] MICHAEL BEESON, *Foundations of Constructive Mathematics*, Springer-Verlag / Berlin, Germany, 1985.
- [2] JOSEF BERGER, *The logical strength of the uniform continuity theorem*, Logical Approaches to Computational Barriers, Proceedings of CiE 2006, LNCS 3988, 2006, pp. 35–39.
- [3] ———, *A separation result for varieties of Brouwer's Fan Theorem*, Proceedings of the 10th Asian Logic Conference (ALC 10), Kobe University in Kobe, Hyogo, Japan, September 1–6, 2008, 2010, pp. 85–92.
- [4] ERRETT BISHOP and DOUGLAS BRIDGES, *Constructive Analysis*, Springer-Verlag / Berlin, Germany, 1985.
- [5] HANNES DIENER, *Compactness under constructive scrutiny*, Ph.D. thesis, 2008.

[6] HANNES DIENER and IRIS LOEB, *Sequences of real functions on  $[0, 1]$  in constructive reverse mathematics*. *Annals of Pure and Applied Logic*, vol. 157 (2009), no. 1, pp. 50–61.

[7] M. FOURMAN and J. HYLAND, *Sheaf models for analysis*, *Applications of Sheaves* (Michael Fourman, Christopher Mulvey, and Dana Scott, editors.), Lecture Notes in Mathematics, vol. 753, Springer Berlin / Heidelberg, 1979, pp. 280–301.

[8] WILLIAM JULIAN and FRED RICHMAN, *A uniformly continuous function on  $[0,1]$  that is everywhere different from its infimum*. *Pacific Journal of Mathematics*, vol. 111 (1984), no. 2, pp. 333–340.

[9] ROBERT S. LUBARSKY, *Independence Results around Constructive ZF*. *Annals of Pure and Applied Logic*, vol. 132 (2005), no. 2–3, pp. 209–225.

[10] STEPHEN SIMPSON, *Subsystems of Second Order Arithmetic*, ASL/Cambridge University Press / Cambridge, UK, 2009.

DEPT. OF MATHEMATICAL SCIENCES  
FLORIDA ATLANTIC UNIVERSITY  
BOCA RATON, FL 33431, USA  
*E-mail*: robert.lubarsky@alum.mit.edu

DEPARTMENT MATHEMATIK, FAK. IV  
EMMY-NOETHER-CAMPUS, WALTER-FLEX-STR. 3  
UNIVERSITY OF SIEGEN  
57068 SIEGEN, GERMANY  
*E-mail*: diener@math.uni-siegen.de