


TAIL ASYMPTOTICS FOR THE AREA UNDER THE EXCURSION OF A RANDOM WALK WITH HEAVY-TAILED INCREMENTS

DENIS DENISOV ^{*}, *University of Manchester*
ELENA PERFILEV,^{** ****} AND
VITALI WACHTEL,^{** ****} *Universität Augsburg*

Abstract

We study the tail behaviour of the distribution of the area under the positive excursion of a random walk which has negative drift and heavy-tailed increments. We determine the asymptotics for tail probabilities for the area.

Keywords: Random walk; subexponential distribution; heavy-tailed distribution; integrated random walk

2010 Mathematics Subject Classification: Primary 60G50
Secondary 60G40; 60F17

1. Introduction and statement of results

Let $\{S_n; n \geq 1\}$ be a random walk with independent and identically distributed increments $\{X_k; k \geq 1\}$. We shall assume that the increments have negative expected value, $\mathbb{E}X_1 = -a$. Let $\bar{F}(x) = \mathbb{P}(X_1 > x)$ be the tail distribution function of X_1 . Let $\tau := \min\{n \geq 1 : S_n \leq 0\}$ be the first time the random walk exits the positive half-line. We consider the area under the random walks excursion $\{S_1, S_2, \dots, S_{\tau-1}\}$:

$$A_\tau := \sum_{k=0}^{\tau-1} S_k.$$

Since τ is finite almost surely, the area A_τ is finite as well. In this note we will study asymptotics for $\mathbb{P}(A_\tau > x)$, as $x \rightarrow \infty$, in the case when the distribution of increments is heavy-tailed. This paper continues the research of [14], where the light-tailed case was considered.

The area under the random walk excursion appears in a number of combinatorial problems, for example in investigations of the asymptotic number of random trees, see [16, 17, 18]; some further references may be found in [6]. Another application area is statistical physics, see, e.g., [8] or [3] and references therein. Applications to queuing theory for the analysis of the load in Transmission Control Protocol networks and to risk theory are discussed in [2].

In the light-tailed case logarithmic asymptotics for $\mathbb{P}(A_\tau > x)$ was obtained in [10], and exact local asymptotics in [14]. Heavy-tailed asymptotics for $\mathbb{P}(A_\tau > x)$ was previously studied in [2], which considered the case when the increments of the random walk have a distribution

Received 2 July 2019; revision received 27 July 2020.

^{*} Postal address: Department of Mathematics, University of Manchester, Oxford Road, Manchester M13 9PL, UK.
Email address: denis.denisov@manchester.ac.uk

^{**} Postal address: Institut für Mathematik, Universität Augsburg, 86135 Augsburg, Germany.

^{***} Email address: Elena.Perfilev@math.uni-augsburg.de

^{****} Email address: vitali.wachtel@mathematik.uni-augsburg.de

© The Author(s), 2021. Published by Cambridge University Press on behalf of Applied Probability Trust

with regularly varying tail, that is $\bar{F}(x) = x^{-\alpha}L(x)$, where $L(x)$ is a slowly varying function. For $\alpha > 1$ it was shown that

$$\mathbb{P}(A_\tau > x) \sim \mathbb{E}\tau \bar{F}(\sqrt{2ax}), \quad x \rightarrow \infty. \tag{1}$$

Here, note that $\mathbb{E}[\tau] < \infty$ follows from the assumption $\mathbb{E}[X_1] = -a < 0$; see, e.g., [11, Chapter XII.2, Theorem 2]. The asymptotics can be explained by traditional heavy-tailed one-big-jump heuristics. In order to have a huge area, the random walk should have a large jump, say y , at the very beginning of the excursion. After this jump the random walk goes down along the line $y - an$ according to the law of large numbers. Thus, the duration of the excursion should be approximately y/a . As a result, the area will be of order $y^2/2a$. Now, from the equality $x = y^2/2a$ we infer that a jump of order $\sqrt{2ax}$ is needed. Since the same strategy is valid for the maximum $M_\tau := \max_{n < \tau} S_n$ of the first excursion, one can rewrite (1) in the following way:

$$\mathbb{P}(A_\tau > x) \sim \mathbb{P}(M_\tau > \sqrt{2ax}), \quad x \rightarrow \infty.$$

However, the class of regularly varying distributions does not include all subexponential distributions, excluding, in particular, the log-normal distribution and Weibull distribution with parameter $\beta < 1$. The asymptotics for these remaining cases have been raised as an open problem in [13, Conjecture 2.2] for a strongly related workload process. We will reformulate this conjecture as

$$\mathbb{P}(A_\tau > x) \sim \mathbb{P}\left(\tau > \sqrt{\frac{2x}{a}}\right), \quad x \rightarrow \infty, \tag{2}$$

when $F \in \mathcal{S}$ and \mathcal{S} is a subclass of subexponential distributions. Note that using the asymptotics for

$$\mathbb{P}(\tau > x) \sim \mathbb{E}\tau \bar{F}(ax) \tag{3}$$

from [7] for Weibull distributions with parameter $\beta < 1/2$, we can see that in this case the asymptotics in (2) is equivalent to (1). In this note we partially settle (2). It is not difficult to show that the same arguments hold for the workload process and to prove the same asymptotics for the area of the workload process, thus settling the original [13, Conjecture 2.2]. In passing, we note that it is doubtful that (2) holds in full. The reason is that for both τ and A_τ the asymptotics (3) and (2) are no longer valid for Weibull distributions with parameter $\beta > 1/2$. The analysis for $\beta > 1/2$ involves a more complicated optimisation procedure leading to a Cramér series, and it is unlikely that the answers will be the same for the area and for the exit time.

1.1. Main results

We will now present the results. We will start with the regularly varying case. In this case the connection between the tails of A_τ and M_τ is strong and we will be able to use the asymptotics for $\mathbb{P}(M_\tau > x)$ found in [12] (see also a short proof in [4]) to find the asymptotics for $\mathbb{P}(A_\tau > x)$.

Proposition 1. *The following two statements hold.*

- (a) *If $\bar{F}(x) := \mathbb{P}(X_1 > x) = x^{-\alpha}L(x)$ with some $\alpha \geq 1$ and $\mathbb{E}|X_1| < \infty$ then, uniformly in $y \in [\varepsilon\sqrt{x}, \sqrt{2ax}]$, $\varepsilon \in (0, 1)$,*

$$\mathbb{P}(A_\tau > x, M_\tau > y) \sim \mathbb{E}\tau \bar{F}(\sqrt{2ax}). \tag{4}$$

(b) If $\bar{F}(x) \sim x^{-\varkappa} e^{-g(x)}$, where $g(x)$ is a monotone continuously differentiable function satisfying $\frac{g(x)}{x^\beta} \downarrow$ for some $\beta \in (0, 1/2)$, and $\mathbb{E}|X_1|^\varkappa < \infty$ for some $\varkappa > 1/(1 - \beta)$, then (4) holds uniformly in $y \in \left[\sqrt{2ax} - \frac{R\sqrt{2ax}}{g(\sqrt{2ax})}, \sqrt{2ax} \right]$, $R > 0$.

This statement obviously implies the following lower bound for the tail of A_τ :

$$\liminf_{x \rightarrow \infty} \frac{\mathbb{P}(A_\tau > x)}{\bar{F}(\sqrt{2ax})} \geq \mathbb{E}\tau. \tag{5}$$

Furthermore, using this proposition one can give an alternative proof of (1) under the assumption of the regular variation of \bar{F} , which is much simpler than the original one in [2]. We first split the event $\{A_\tau > x\}$ into two parts,

$$\{A_\tau > x\} = \{A_\tau > x, M_\tau > y\} \cup \{A_\tau > x, M_\tau \leq y\}.$$

Clearly, $\{A_\tau > x, M_\tau \leq y\} \subseteq \{\tau > x/y\}$. Therefore,

$$\mathbb{P}(A_\tau > x, M_\tau > y) \leq \mathbb{P}(A_\tau > x) \leq \mathbb{P}(A_\tau > x, M_\tau > y) + \mathbb{P}(\tau > x/y). \tag{6}$$

When $\alpha > 1$, according to Theorem I in [9] or [7, Theorem 3.2], $\mathbb{P}(\tau > t) \sim \mathbb{E}\tau \bar{F}(at)$ as $t \rightarrow \infty$. Choosing $y = \varepsilon\sqrt{x}$ and recalling that \bar{F} is regularly varying, we get

$$\mathbb{P}(\tau > x/y) = \mathbb{P}(\tau > \sqrt{x}/\varepsilon) \sim \varepsilon^\alpha \mathbb{E}\tau \bar{F}(\sqrt{x}). \tag{7}$$

It follows from the first statement of Proposition 1 that

$$\mathbb{P}(A_\tau > x, M_\tau > \varepsilon\sqrt{x}) \sim \mathbb{E}\tau \bar{F}(\sqrt{2ax}).$$

Plugging this and (7) into (6), we get, as $x \rightarrow \infty$,

$$\mathbb{E}\tau \bar{F}(\sqrt{2ax})(1 + o(1)) \leq \mathbb{P}(A_\tau > x) \leq \mathbb{E}\tau \bar{F}(\sqrt{2ax}) \left(1 + \frac{\varepsilon^\alpha}{(2a)^{\alpha/2}} + o(1) \right).$$

Letting $\varepsilon \rightarrow 0$, we arrive at (1).

The case of heavy-tailed distributions, which satisfy the conditions of Proposition 1(b), is more complicated. In particular, it seems that in this case there is a regime when the asymptotics in (1) is no longer valid. We will treat this case by using exponential bounds similar to Section 2.2 in [14] and asymptotics for $\mathbb{P}(\tau > x)$ from [5] and [7].

First, we will introduce a subclass of subexponential distributions to consider. We will assume that $\mathbb{E}[X_1^2] = \sigma^2 < \infty$. Without loss of generality we may assume that $\sigma = 1$.

Assumption 1. Let

$$\bar{F}(x) \sim e^{-g(x)} x^{-2}, \quad x \rightarrow \infty, \tag{8}$$

where $g(x)$ is an eventually increasing function such that eventually

$$\frac{g(x)}{x^{\gamma_0}} \downarrow 0, \quad x \rightarrow \infty, \tag{9}$$

for some $\gamma_0 \in (0, 1]$.

Due to the asymptotic nature of equivalence in (8), without loss of generality we may assume that g is continuously differentiable and that (9) holds for all $x > 0$. Clearly, monotonicity in (9) implies that

$$g'(x) \leq \gamma_0 \frac{g(x)}{x} \quad (10)$$

for all sufficiently large x . Using the Karamata representation theorem we can show that this class of subexponential distributions includes regularly varying distributions $\bar{F}(x) \sim x^{-r}L(x)$ for $r > 2$. Also, it is not difficult to show that lognormal distributions and Weibull distributions ($\bar{F}(x) \sim e^{-x^\beta}$, $\beta \in (0, 1)$) belong to our class of distributions. This class previously appeared in [15] for the analysis of large deviations of sums of subexponential random variables on the whole axis.

Now we are able to give rough (logarithmic) asymptotics for $\gamma_0 \leq 1$.

Theorem 1. *Let $\mathbb{E}[X_1] = -a < 0$ and $\text{Var}(X_1) < \infty$. Assume that Assumption 1 holds with $\gamma_0 = 1$. Then, there exists a constant $C > 0$ such that*

$$\mathbb{P}(A_\tau > x) \leq Cx^{1/4} \exp \left\{ -g(\sqrt{2ax}) \sqrt{1 - \frac{2Cg(\sqrt{2ax})}{a\sqrt{2ax}}} \right\}.$$

Furthermore, for any $\varepsilon > 0$ there exists $C > 0$ such that

$$\liminf_{x \rightarrow \infty} \frac{\mathbb{P}(A_\tau > x)}{\bar{F}(\sqrt{2ax} + Cx^{1/4+\varepsilon})} \geq \mathbb{E}\tau.$$

In particular, if $\gamma_0 < 1$ then

$$\lim_{x \rightarrow \infty} \frac{\ln \mathbb{P}(A_\tau > x)}{\ln \bar{F}(\sqrt{2ax})} = 1.$$

To obtain the exact asymptotics we will impose a further requirement on the function g .

Assumption 2. *Let $g(x)$ satisfy*

$$xg'(x) \rightarrow \infty, \quad x \rightarrow \infty. \quad (11)$$

This assumption implies that

$$\frac{g(x)}{\log x} \rightarrow \infty. \quad (12)$$

In particular, it excludes all regularly varying distributions.

Theorem 2. *Let $\mathbb{E}[X_1] = -a < 0$ and $\text{Var}(X_1) < \infty$. Assume that Assumption 1 holds with $\gamma_0 < 1/2$ and, in addition, that Assumption 2 holds. Then*

$$\mathbb{P}(A_\tau > x) \sim \mathbb{E}\tau \bar{F}(\sqrt{2ax}), \quad x \rightarrow \infty.$$

1.2. Discussion and organisation of the paper

The main result of this note, Theorem 2, provides tail asymptotics for A_τ in the case when increments of the random walk have a Weibull-like distribution with the shape parameter $\gamma_0 < 1/2$. We believe that $\mathbb{P}(A_\tau > x)$ behaves differently in the case when $g(x) = x^{\gamma_0}$ with $\gamma_0 \geq 1/2$. This change in the asymptotic behaviour appears in the analysis of the exact asymptotics for $\mathbb{P}(\tau > n)$ and $\mathbb{P}(S_n > an)$; see, correspondingly, [5,7].

The conjecture in [13] was formulated for the workload process of a single-server queue rather than the area under the random walk excursion. However, one can prove analogous results for the Lévy processes by essentially the same arguments. It is well known that the workload of the M/G/1 queue can be represent as a Lévy process, and thus our results can be transferred to this setting almost immediately. We believe that the treatment of the workload of the general G/G/1 queue is not that different either.

The paper is organised as follows. We will start by proving Proposition 1 in Section 2. Then we will derive a useful exponential bound and prove Theorem 1 in Section 3. Finally, we derive exact asymptotics for $\mathbb{P}(A_\tau > x)$ and thus prove Theorem 2 in Section 4.

2. Proof of Proposition 1

Before giving the proof we collect some auxiliary results that we will need in this and the following sections.

We will require the following statement, the first part of which follows from Theorem 2 in [12] (see also [4] for a short proof), and the second part from [7, Theorem 3.2].

Proposition 2. *Let $\mathbb{E}[X_1] = -a$ and either (a) $\bar{F}(x) := \mathbb{P}(X_1 > x) = x^{-\alpha}L(x)$ with some $\alpha > 1$ or (b) $\bar{F}(x) \sim x^{-\varkappa}e^{-g(x)}$, where $g(x)$ is a monotone continuously differentiable function satisfying $\frac{g(x)}{x^\beta} \downarrow$ for $\beta \in (0, 1/2)$, and $\mathbb{E}|X_1|^\varkappa < \infty$ for some $\varkappa > 1/(1 - \beta)$; then, for any fixed k ,*

$$\mathbb{P}(M_k > y) \sim \mathbb{P}(S_k > y) \sim k\bar{F}(y), \quad y \rightarrow \infty, \tag{13}$$

$$\mathbb{P}\left(\max_{n \leq \tau \wedge k} S_n > y\right) \sim \mathbb{E}(\tau \wedge k)\bar{F}(y), \quad y \rightarrow \infty, \tag{14}$$

$$\mathbb{P}(M_\tau > y) \sim \mathbb{E}\tau\bar{F}(y), \quad y \rightarrow \infty \tag{15}$$

and

$$\mathbb{P}(\tau > n) \sim \mathbb{E}[\tau]\bar{F}(an), \quad n \rightarrow \infty. \tag{16}$$

In the proof we will need some properties of the function $\bar{F}(x) \sim x^{-\varkappa}e^{-g(x)}$ that we will summarise in the following lemma, which will also be used later in the paper.

Lemma 1. *Let the distribution function $\bar{F}(x)$ be such that $\bar{F}(x) \sim x^{-\varkappa}e^{-g(x)}$, where $g(x)$ is a monotone continuously differentiable function satisfying $\frac{g(x)}{x^\beta} \downarrow$ for $\beta \in (0, 1)$, and $\mathbb{E}|X_1|^\varkappa < \infty$ for some $\varkappa > 1/(1 - \beta)$. Then,*

$$g'(x) \leq \beta \frac{g(x)}{x}, \quad x > 0, \tag{17}$$

$$g(x) - g(y) \leq \beta g(y) \frac{x-y}{y}, \quad x > y > 0, \tag{18}$$

$$g(x) - g(x-y) \leq \beta g(y), \quad x \geq 2y > 0, \tag{19}$$

$$\sup_{y \leq x^{1/\varkappa}} \frac{\bar{F}(x-y)}{\bar{F}(x)} \rightarrow 1, \quad x \rightarrow \infty. \tag{20}$$

Proof. Since $g(x)$ is continuously differentiable and $\frac{g(x)}{x^\beta}$ is monotone decreasing then, with necessity,

$$0 \geq \left(\frac{g(x)}{x^\beta}\right)' = \frac{g'(x)x^\beta - \beta x^{\beta-1}g(x)}{x^{2\beta}},$$

implying (17). To prove (18), note that

$$g(x) - g(y) = \int_y^x g'(t) dt \leq \beta \int_y^x \frac{g(t)}{t} dt \leq \beta \frac{g(y)}{y^\beta} \int_y^x \frac{1}{t^{1-\beta}} dt = \beta g(y) \frac{x-y}{y}.$$

To prove (18), note that, since $x - y \geq y$,

$$\begin{aligned} g(x) - g(x-y) &= \int_{x-y}^x g'(t) dt \leq \beta \int_{x-y}^x \frac{g(t)}{t} dt \leq \beta \frac{g(y)}{y^\beta} \int_{x-y}^x \frac{1}{t^{1-\beta}} dt \\ &\leq \beta \frac{g(y)}{y^\beta} \int_y^{2y} \frac{1}{t^{1-\beta}} dt \leq \beta g(y) \frac{(2y)^\beta - y^\beta}{\beta y^\beta} \leq \beta g(y), \end{aligned}$$

since $2^\beta \leq 1 + \beta$ for $\beta \in [0, 1]$. To show (20), note that, uniformly in $y \leq x^{1/\varkappa}$, as $x \rightarrow \infty$,

$$\begin{aligned} 1 &\leq \frac{\bar{F}(x-y)}{\bar{F}(x)} \leq \frac{\bar{F}(x-x^{1/\varkappa})}{\bar{F}(x)} = (1 + o(1)) \exp \left\{ g(x) - g(x-x^{1/\varkappa}) \right\} \\ &\leq (1 + o(1)) \exp \left\{ \beta g(x-x^{1/\varkappa}) \frac{x^{\varkappa} - x^{1/\varkappa}}{x - x^{\varkappa}} \right\} \leq (1 + o(1)) \exp \left\{ C \frac{x^{1/\varkappa}}{(x - x^{1/\varkappa})^{1-\beta}} \right\} \rightarrow 1, \end{aligned}$$

since $1/\varkappa < 1 - \beta$. Here, we have also made use of (18). □

Proof of Proposition 2. To prove (13), (14), and (15), by Theorem 2 of [12] it is sufficient to show that (a) or (b) implies that $F \in \mathcal{S}^*$, that is, $\int_0^\infty \bar{F}(y) dy < \infty$ and

$$\int_0^x \bar{F}(y)\bar{F}(x-y) dy \sim 2\bar{F}(x) \int_0^\infty \bar{F}(y) dy, \quad x \rightarrow \infty.$$

The fact that (a) implies $F \in \mathcal{S}^*$ is well known and follows immediately from the dominated convergence theorem, since $\bar{F}(x) \sim \bar{F}(x-y)$ for all fixed y and

$$\int_0^x \frac{\bar{F}(y)\bar{F}(x-y)}{\bar{F}(x)} dy = 2 \int_0^{x/2} \frac{\bar{F}(y)\bar{F}(x-y)}{\bar{F}(x)} dy,$$

and $\bar{F}(x-y) \leq C\bar{F}(x)$ for some $C > 0$ when $y \leq x/2$. Now, assume that (b) holds and show that $F \in \mathcal{S}^*$. Consider

$$2 \int_0^{x/2} \frac{\bar{F}(y)\bar{F}(x-y)}{\bar{F}(x)} dy.$$

Uniformly in $y \in [\ln x, x/2]$ we have, by (19),

$$\frac{\bar{F}(y)\bar{F}(x-y)}{\bar{F}(x)} \leq C \frac{x^{\varkappa}}{(x-y)^{\varkappa} y^{\varkappa}} e^{g(x)-g(x-y)-g(y)} \leq C y^{-\varkappa} e^{\beta g(y)-g(y)}, \tag{21}$$

and therefore, since $\varkappa > 1$,

$$2 \int_{\ln x}^{x/2} \frac{\bar{F}(y)\bar{F}(x-y)}{\bar{F}(x)} dy \rightarrow 0.$$

Next, applying (20) we see that $\frac{\bar{F}(x-y)}{\bar{F}(x)} \rightarrow 1$ uniformly in $y \in [0, \ln x]$, which implies that $F \in \mathcal{S}^*$.

The proof of (16) can be done by verification that (8) and (9) imply that the conditions of Theorem 3.1 (and hence of Theorem 3.2) of [7] hold. We will provide the arguments in the more complicated case (b). First, $X_1 + a$ satisfies $\mathbb{E}[|X_1 + a|^\alpha] < \infty$ by the assumption of this proposition. Convergence (3.1) in Theorem 3.1 of [7] holds by (20). Let

$$\varepsilon(n) := \sup_{x \geq 2n^{1/\alpha}} \frac{\mathbb{P}(\xi_1 > n^{1/\alpha}, \xi_2 > n^{1/\alpha}, \xi_1 + \xi_2 > x)}{\mathbb{P}(\xi_1 > x)},$$

where $\xi_i = X_i + a, i = 1, 2$. To show (3.2) in Theorem 3.1 of [7] we need to prove that $\varepsilon(n) = o(1/n)$. For $x \geq 2n^{1/\alpha}$ we have

$$\begin{aligned} \frac{\mathbb{P}(\xi_1 > n^{1/\alpha}, \xi_2 > n^{1/\alpha}, \xi_1 + \xi_2 > x)}{\mathbb{P}(\xi_1 > x)} &= 2P_1 + P_2 \\ &:= 2 \int_{n^{1/\alpha}}^{x/2} \mathbb{P}(\xi_1 \in dy) \frac{\mathbb{P}(\xi_2 > x - y)}{\mathbb{P}(\xi_1 > x)} + \frac{\mathbb{P}(\xi_1 > x/2)^2}{\mathbb{P}(\xi_1 > x)}. \end{aligned}$$

Then, using (19),

$$\begin{aligned} P_1 &\leq \int_{n^{1/\alpha}}^{x/2} \mathbb{P}(\xi_1 \in dy) \frac{\mathbb{P}(\xi_2 > x - y)}{\mathbb{P}(\xi_1 > x)} \\ &\leq C \int_{n^{1/\alpha}}^{x/2} \mathbb{P}(\xi_1 \in dy) \frac{\bar{F}(x - y)}{\bar{F}(x)} \leq C \int_{n^{1/\alpha}}^{x/2} \mathbb{P}(\xi_1 \in dy) e^{\beta g(y)} \end{aligned}$$

for some C . Integrating by parts,

$$\begin{aligned} P_1 &\leq C\mathbb{P}(\xi_1 > n^{1/\alpha})e^{\beta g(n^{1/\alpha})} + C \int_{n^{1/\alpha}}^{x/2} dy g'(y) e^{\beta g(y)} \mathbb{P}(\xi_1 > y) \\ &\leq Cn^{-1}e^{(\beta-1)g(n^{1/\alpha})} + C \int_{n^{1/\alpha}}^{x/2} dy g'(y) y^{-\kappa} e^{(\beta-1)g(y)} \\ &\leq o(n^{-1}) + cn^{-1} \int_{n^{1/\alpha}}^{x/2} dy g'(y) e^{(\beta-1)g(y)} = o(n^{-1}) \end{aligned}$$

uniformly in $x \geq 2n^{1/\alpha}$. Using (21), $P_2 \leq Cx^{-\alpha}e^{(\beta-1)g(x)} = o(1/n)$ uniformly in $x \geq 2n^{1/\alpha}$, which proves that $\varepsilon(n) = o(1/n)$. □

Define $\sigma_y = \inf\{n < \tau : S_n > y\}$.

Lemma 2. *Under the conditions of Proposition 2,*

$$\lim_{y \rightarrow \infty} \mathbb{P}(\sigma_y = k \mid M_\tau > y) =: q_k, \quad k \geq 1; \quad \sum_{k=1}^\infty q_k = 1.$$

Proof. For every $k \geq 1$,

$$\begin{aligned} \mathbb{P}(\sigma_y = k \mid M_\tau > y) &= \frac{\mathbb{P}(\sigma_y = k)}{\mathbb{P}(M_\tau > y)} \\ &= \frac{\mathbb{P}(\max_{n \leq \tau \wedge k} S_n > y) - \mathbb{P}(\max_{n \leq \tau \wedge (k-1)} S_n > y)}{\mathbb{P}(M_\tau > y)}. \end{aligned}$$

It follows from (14) and (15) that

$$\begin{aligned} \lim_{y \rightarrow \infty} \mathbb{P}(\sigma_y = k \mid M_\tau > y) &= \frac{\mathbb{E}\tau \wedge k - \mathbb{E}\tau \wedge (k - 1)}{\mathbb{E}\tau} \\ &= \frac{\mathbb{P}(\tau > k - 1)}{\mathbb{E}\tau} =: q_k, \quad k \geq 1. \end{aligned} \tag{22}$$

It is clear that

$$\sum_{k=1}^{\infty} q_k = \frac{1}{\mathbb{E}\tau} \sum_{k=0}^{\infty} \mathbb{P}(\tau > k - 1) = 1. \quad \square$$

Lemma 3. *For every fixed k ,*

$$\sup_{v > y} \left| \frac{\mathbb{P}(S_k > v, \sigma_y = k)}{\bar{F}(v)} - \mathbb{P}(\tau > k - 1) \right| \rightarrow 0 \text{ as } y \rightarrow \infty.$$

Proof. Fix some $N > 0$ and define the events

$$D_{k,N} = \cup_{j=1}^k \{X_j > v + kN, |X_l| \leq N \text{ for all } l \neq j, l \leq k\}.$$

It is clear that $D_{k,N} \subseteq \{S_k > v\}$. Therefore,

$$\begin{aligned} \mathbb{P}(S_k > v, \sigma_y = k) &= \mathbb{P}(D_{k,N}, \sigma_y = k) + \mathbb{P}(S_k > v, D_{k,N}^c, \sigma_y = k) \\ &= \mathbb{P}(X_k > v + kN, |X_l| \leq N, \text{ for all } l < k, \sigma_y > k - 1) \\ &\quad + \mathbb{P}(S_k > v, D_{k,N}^c, \sigma_y = k). \end{aligned}$$

For the first term we have ($y > (k - 1)N$)

$$\begin{aligned} \mathbb{P}(X_k > v + kN, |X_l| \leq N, \text{ for all } l < k, \sigma_y > k - 1) \\ &= \mathbb{P}(\tau > k - 1, |X_l| \leq N, l < k) \bar{F}(v + kN) \\ &= \mathbb{P}(\tau > k - 1) \bar{F}(v) - \varepsilon_N^{(1)} \bar{F}(v) + o(\bar{F}(v)) \end{aligned} \tag{23}$$

uniformly in $v > y$, where $\varepsilon_N^{(1)} := \mathbb{P}(\tau > k - 1, |X_l| > N \text{ for some } l < k) \rightarrow 0$ as $N \rightarrow \infty$.

Furthermore,

$$\begin{aligned} \mathbb{P}(S_k > v, D_{k,N}^c, \sigma_y = k) &\leq \mathbb{P}(S_k > v, D_{k,N}^c) = \mathbb{P}(S_k > v) - \mathbb{P}(D_{k,N}) \\ &= \mathbb{P}(S_k > v) - k \mathbb{P}(X_1 > v + kN) (\mathbb{P}(|X_1| \leq N))^{k-1} \\ &= \varepsilon_N^{(2)} \bar{F}(v) + o(\bar{F}(v)), \end{aligned} \tag{24}$$

where $\varepsilon_N^{(2)} := k (1 - (\mathbb{P}(|X_1| \leq N))^{k-1}) \rightarrow 0$, as $N \rightarrow \infty$. Combining (23) and (24) and letting $N \rightarrow \infty$, we get the desired relation. \square

We now turn to the study of the tail behaviour of A_τ on the event $\{\sigma_y = k\}$. For the corresponding result we need the following property of \bar{F} .

Lemma 4. *Assume that the conditions of Proposition 1 are fulfilled. Then*

$$\lim_{y \rightarrow \infty} \frac{\bar{F}(y + h)}{\bar{F}(y)} = 1 \tag{25}$$

for any $h = o(y/g(y))$. Furthermore, for every $R > 0$ there exists a constant C such that

$$\bar{F}(y - h) \leq C\bar{F}(y) \text{ for all } h \leq R\frac{y}{g(y)}. \tag{26}$$

Proof. Since $\bar{F}(x) \sim x^{-\alpha}e^{-g(x)}$ and $\frac{y}{y+h} \rightarrow 1$, (25) will follow from

$$g(y + h) - g(y) \rightarrow 0. \tag{27}$$

Since $\frac{g(x)}{x^\beta}$ is monotone decreasing and g is differentiable,

$$g'(x) \leq \beta \frac{g(x)}{x}.$$

Then, for $h < 0$,

$$g(y) - g(y - h) = \int_y^{y-h} g'(t) dt \leq \beta \int_y^{y-h} \frac{g(t)}{t} dt \leq \beta \frac{g(y - h)}{y - h} h. \tag{28}$$

In the last step we have used the fact that $\frac{g(t)}{t}$ is decreasing. Similarly, for $h > 0$,

$$g(y + h) - g(y) \leq \beta \frac{g(y)}{y} h.$$

These estimates yield (27).

To prove the second claim we note that, by (28),

$$\frac{\bar{F}(y - h)}{\bar{F}(y)} \leq C \left(\frac{y}{y - h}\right)^\alpha e^{g(y) - g(y - h)} \leq C \exp \left\{ \beta \frac{g(y - h)}{y - h} h \right\}.$$

If $h \leq R\frac{y}{g(y)}$ then

$$\frac{\bar{F}(y - h)}{\bar{F}(y)} \leq C \exp \left\{ \beta R \frac{yg(y - h)}{(y - h)g(y)} \right\} \leq C \exp \left\{ \beta R \frac{1}{(1 - R/g(y))} \right\}.$$

This completes the proof. □

Lemma 5. Assume that the conditions of Proposition 1 hold. Then

$$\mathbb{P} \left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k \right) \sim q_k \mathbb{E} \tau \bar{F}(\sqrt{2az}), \quad k \geq 1,$$

uniformly in $y \in [\varepsilon\sqrt{z}, \sqrt{2az}]$ for regularly varying tails \bar{F} and in $y \in [\sqrt{2ax} - \frac{R\sqrt{2ax}}{g(\sqrt{2ax})}, \sqrt{2ax}]$ for tails satisfying the conditions of part (b) in Proposition 1.

Proof. By the Markov property, for every $z > 0$,

$$\mathbb{P} \left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k \right) = \int_y^\infty \mathbb{P}(S_k \in dv, \sigma_y = k) \mathbb{P}(A_\tau > z | S_0 = v).$$

Let $\varkappa \in (1/(1-\beta), 2)$ if \bar{F} satisfies the conditions of part (b), and let $\varkappa = 1$ in the case when \bar{F} is regularly varying. Fix some $\delta > 0$ and consider the set

$$B_v := \left\{ v - \delta v^{1/\varkappa} \leq S_n + na \leq v + \delta v^{1/\varkappa} \text{ for all } n \leq \frac{v + \delta v^{1/\varkappa}}{a} \right\}.$$

Since $\mathbb{E}|X_1|^\varkappa < \infty$, it follows from the Marcinkiewicz–Zygmund law of large numbers that

$$\lim_{R \rightarrow \infty} \mathbb{P} \left(-R - \frac{\delta}{2} n^{1/\varkappa} < S_n + na < R + \frac{\delta}{2} n^{1/\varkappa} \text{ for all } n \geq 1 \right) = 1.$$

Consequently, $\mathbb{P}(B_v | S_0 = v) \rightarrow 1$ as $v \rightarrow \infty$. This implies that, as $y \rightarrow \infty$,

$$\begin{aligned} & \mathbb{P} \left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k \right) \\ &= \int_y^\infty \mathbb{P}(S_k \in dv, \sigma_y = k) \mathbb{P}(A_\tau > z | S_0 = v) \\ &= \int_y^\infty \mathbb{P}(S_k \in dv, \sigma_y = k) \mathbb{P}(\{A_\tau > z\} \cap B_v | S_0 = v) + o(\mathbb{P}(\sigma_y = k)). \end{aligned}$$

On the event $B_v \cap \{S_0 = v\}$ one has

$$\frac{v - \delta v^{1/\varkappa}}{a} < \tau < \frac{v + \delta v^{1/\varkappa}}{a}.$$

Consequently,

$$A_\tau = \sum_{n=0}^{\tau-1} S_n \geq \sum_{n=0}^{\tau-1} (v - \delta v^{1/\varkappa} - na) \geq \tau \left(v - \delta v^{1/\varkappa} - \frac{a\tau}{2} \right) \geq \frac{(v - \delta v^{1/\varkappa})^2}{2a}$$

and

$$A_\tau = \sum_{n=0}^{\tau-1} S_n \leq v + \sum_{n=1}^{\tau-1} (v + \delta v^{1/\varkappa} - na) \leq \tau \left(v + \delta v^{1/\varkappa} - \frac{a\tau}{2} \right) \leq \frac{(v + \delta v^{1/\varkappa})^2}{2a}$$

on the same event. In other words, $\mathbb{P}(\{A_\tau > z\} \cap B_v | S_0 = v) = \mathbb{P}(B_v | S_0 = v)$ if $v - \delta v^{1/\varkappa} \geq \sqrt{2az}$, and $\mathbb{P}(\{A_\tau > z\} \cap B_v | S_0 = v) = 0$ if $v + \delta v^{1/\varkappa} < \sqrt{2az}$. Therefore, for all v large enough,

$$\begin{aligned} \mathbb{P} \left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k \right) &\leq \int_{\sqrt{2az} - \delta(2az)^{1/2\varkappa}}^\infty \mathbb{P}(S_k \in dv, \sigma_y = k) + o(\mathbb{P}(\sigma_y = k)) \\ &= \mathbb{P} \left(S_{\sigma_y} > \sqrt{2az} - \delta(2az)^{1/2\varkappa}, \sigma_y = k \right) + o(\mathbb{P}(\sigma_y = k)) \end{aligned}$$

and

$$\begin{aligned} \mathbb{P} \left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k \right) &\geq \int_{\sqrt{2az} + 2\delta(2az)^{1/2\varkappa}}^\infty \mathbb{P}(S_k \in dv, \sigma_y = k) \mathbb{P}(B_v) + o(\mathbb{P}(\sigma_y = k)) \\ &= \mathbb{P} \left(S_{\sigma_y} > \sqrt{2az} + 2\delta(2az)^{1/2\varkappa}, \sigma_y = k \right) + o(\mathbb{P}(\sigma_y = k)). \end{aligned}$$

By Lemma 3, $\mathbb{P}(S_{\sigma_y} > v, \sigma_y = k) \sim \bar{F}(v)\mathbb{P}(\tau > k - 1)$ as $y \rightarrow \infty$ uniformly in $v \geq y$ and, consequently,

$$\mathbb{P}\left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k\right) \leq \bar{F}\left((\sqrt{2az} - \delta(2az)^{1/2\kappa}) \vee y\right) (\mathbb{P}(\tau > k - 1) + o(1)) + o(\mathbb{P}(\sigma_y = k))$$

and

$$\mathbb{P}\left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k\right) \geq \bar{F}\left(\sqrt{2az} + 2\delta(2az)^{1/2\kappa}\right) (\mathbb{P}(\tau > k - 1) + o(1)) + o(\mathbb{P}(\sigma_y = k)).$$

Under our assumptions on \bar{F} , we have

$$\lim_{\delta \rightarrow 0} \lim_{z \rightarrow \infty} \frac{\bar{F}\left(\sqrt{2az} + 2\delta(2az)^{1/2\kappa}\right)}{\bar{F}\left((\sqrt{2az} - \delta(2az)^{1/2\kappa}) \vee y\right)} = 1.$$

Indeed, this relation is obvious for regularly varying tails, and under the conditions of part (b) it is a particular case of (25). Therefore,

$$\mathbb{P}\left(\sum_{j=k}^{\tau-1} S_j > z, \sigma_y = k\right) = \bar{F}\left(\sqrt{2az}\right) (\mathbb{P}(\tau > k - 1) + o(1)) + o(\mathbb{P}(\sigma_y = k)).$$

Combining (15) and (22), we get $\mathbb{P}(\sigma_y = k) \sim q_k \mathbb{E}\tau \bar{F}(y)$. Thus, it remains to show that $\bar{F}(y) = O(\bar{F}(\sqrt{2az}))$. This is obvious for regularly varying tails and $y \geq \varepsilon\sqrt{z}$. For distributions satisfying the conditions of part (b), it suffices to apply (26) with $y = \sqrt{2az}$. \square

Proof of Proposition 1. For every fixed $N \geq 1$ we have

$$\begin{aligned} &\mathbb{P}(A_\tau > x, M_\tau > y) \\ &= \sum_{k=1}^N \mathbb{P}(A_\tau > x, \sigma_y = k, M_\tau > y) + \mathbb{P}(A_\tau > x, \sigma_y > N, M_\tau > y). \end{aligned} \tag{29}$$

For the last term on the right-hand side we have

$$\begin{aligned} \mathbb{P}(A_\tau > x, \sigma_y > N, M_\tau > y) &\leq \mathbb{P}(\sigma_y > N, M_\tau > y) \\ &= \mathbb{P}(M_\tau > y)\mathbb{P}(\sigma_y > N \mid M_\tau > y). \end{aligned}$$

It follows from (22) that $\mathbb{P}(\sigma_y > N \mid M_\tau > y) \rightarrow \sum_{j=N+1}^\infty q_j$ as $y \rightarrow \infty$. Then, using (15), we get

$$\mathbb{P}(A_\tau > x, \sigma_y > N, M_\tau > y) \leq \varepsilon_N \bar{F}(y), \tag{30}$$

where $\varepsilon_N \rightarrow 0$ as $N \rightarrow \infty$.

For every fixed k we have $\mathbb{P}(A_\tau > x, \sigma_y = k, M_\tau > y) = \mathbb{P}(A_\tau > x, \sigma_y = k)$. Since $S_j \in (0, y)$ for all $j < k$, we obtain

$$\mathbb{P}(A_\tau > x, \sigma_y = k) \leq \mathbb{P}\left(\sum_{j=k}^{\tau-1} S_j > x - (k-1)y, \sigma_y = k\right)$$

and

$$\mathbb{P}(A_\tau > x, \sigma_y = k) \geq \mathbb{P}\left(\sum_{j=k}^{\tau-1} S_j > x, \sigma_y = k\right).$$

Using Lemma 5 with $z = x$ and with $z = x - ky$, we conclude that $\mathbb{P}(A_\tau > x, \sigma_y = k) \sim q_k \mathbb{E}\tau \bar{F}(\sqrt{2ax})$. Consequently,

$$\sum_{k=1}^N \mathbb{P}(A_\tau > x, \sigma_y = k, M_\tau > y) \sim \bar{F}(\sqrt{2ax}) \mathbb{E}\tau \sum_{k=1}^N q_k. \tag{31}$$

Plugging (30) and (31) into (29) and letting $N \rightarrow \infty$, we obtain

$$\mathbb{P}(A_\tau > x, M_\tau > y) = (\mathbb{E}\tau + o(1))\bar{F}(\sqrt{2ax}) + o(\bar{F}(y)).$$

Recalling that $\bar{F}(y) = O(\bar{F}(\sqrt{2ax}))$, we finish the proof. □

3. Proof of Theorem 1

We start by proving an exponential estimate for the area A_n when random variables X_j are truncated. Let $\bar{X}_n = \max(X_1, \dots, X_n)$. The next result is our main technical tool to investigate trajectories without big jumps.

Lemma 6. *Let $\mathbb{E}[X_1] = -a$ and $\sigma^2 := \text{Var}(X_1) < \infty$. Assume that the distribution function F of X_j satisfies (8) and that (9) holds with $\gamma_0 = 1$. Then there exists a constant $C_0 > 0$ such that*

$$\mathbb{P}(A_n > x, \bar{X}_n \leq y) \leq \exp\left\{-\lambda \frac{x}{n} - \lambda \frac{an}{2} + C_0 \lambda^2 n\right\},$$

where $\lambda = \frac{g(y)}{y}$.

Proof. We will prove this lemma by using the exponential Chebyshev inequality. For that, we need to obtain estimates for the moment-generating function of A_n . First,

$$\mathbb{E}\left[e^{\frac{\lambda}{n} A_n}; \bar{X}_n \leq y\right] = \mathbb{E}\left[e^{\frac{\lambda}{n} \sum_{j=1}^n (n-j+1)X_j}; \bar{X}_n \leq y\right] = \prod_{j=1}^n \varphi_y(\lambda_{n,j}),$$

where $\varphi_y(t) := \mathbb{E}[e^{tX_j}; X_j \leq y]$ and $\lambda_{n,j} := \lambda \frac{(n-j+1)}{n}$. Then,

$$\varphi_y(\lambda_{n,j}) = \mathbb{E}[e^{\lambda_{n,j} X_j}; X_j \leq 1/\lambda_{n,j}] + \mathbb{E}[e^{\lambda_{n,j} X_j}; 1/\lambda_{n,j} < X_j \leq y] =: E_1 + E_2.$$

Using the elementary bound $e^x \leq 1 + x + x^2$ for $x \leq 1$, we obtain

$$E_1 \leq 1 + \lambda_{n,j} \mathbb{E}[X_j] + \lambda_{n,j}^2 \mathbb{E}[X_j^2] = 1 - a\lambda_{n,j} + (a^2 + \sigma^2)\lambda_{n,j}^2.$$

Next, using integration by parts and the assumption in (8),

$$\begin{aligned}
 E_2 &= \int_{1/\lambda_{n,j}}^y e^{\lambda_{n,j}t} dF(t) = -\bar{F}(t)e^{\lambda_{n,j}t} \Big|_{t=1/\lambda_{n,j}}^{t=y} + \lambda_{n,j} \int_{1/\lambda_{n,j}}^y e^{\lambda_{n,j}t}\bar{F}(t) dt \\
 &\leq e\bar{F}(1/\lambda_{n,j}) + C\lambda_{n,j} \int_{1/\lambda_{n,j}}^y e^{\lambda_{n,j}t-g(t)}t^{-2} dt.
 \end{aligned}$$

Now note that, for $t \leq y$,

$$\lambda_{n,j}t - g(t) = t \left(\lambda_{n,j} - \frac{g(t)}{t} \right) \leq t \left(\lambda_{n,j} - \frac{g(y)}{y} \right),$$

due to the condition in (9). Then,

$$\lambda_{n,j} - \frac{g(y)}{y} \leq \lambda - \frac{g(y)}{y} = 0$$

and, therefore,

$$E_2 \leq e\bar{F}(1/\lambda_{n,j}) + C\lambda_{n,j} \int_{1/\lambda_{n,j}}^y t^{-2} dt \leq (C + e)\lambda_{n,j}^2,$$

where we also used the Chebyshev inequality. As a result, for some constant C ,

$$\varphi_y(\lambda_{n,j}) = E_1 + E_2 \leq 1 - a\lambda_{n,j} + C\lambda_{n,j}^2.$$

Consequently,

$$\begin{aligned}
 \mathbb{E} \left[e^{\frac{\lambda}{n}A_n}; \bar{X}_n \leq y \right] &\leq \prod_{j=1}^n \left(1 - a\lambda_{n,j} + C\lambda_{n,j}^2 \right) \\
 &= \exp \left\{ \sum_{j=1}^n \ln \left(1 - a\lambda_{n,j} + C\lambda_{n,j}^2 \right) \right\} \\
 &\leq \exp \left\{ \sum_{j=1}^n \left(-a\lambda_{n,j} + C\lambda_{n,j}^2 \right) \right\} \\
 &= \exp \left\{ \sum_{j=1}^n \left(-a\lambda \frac{n-j+1}{n} + C \left(\lambda \frac{n-j+1}{n} \right)^2 \right) \right\} \\
 &\leq \exp \left\{ -\frac{a\lambda}{2}n + C\lambda^2n \right\}.
 \end{aligned}$$

Finally,

$$\mathbb{P}(A_n > x, \bar{X}_n \leq y) \leq e^{-\lambda \frac{x}{n}} \mathbb{E} \left[e^{\frac{\lambda}{n}A_n}; \bar{X}_n \leq y \right] \leq \exp \left\{ -\lambda \frac{x}{n} - \frac{a\lambda}{2}n + C\lambda^2n \right\}. \quad \square$$

We can now obtain upper bounds for the tail of A_τ using the exponential bound in Lemma 6.

Lemma 7. Let $\mathbb{E}[X_1] = -a < 0$ and $\text{Var}(X_1) < \infty$. Assume that the distribution function F of X_j satisfies (8), and that (9) holds with $\gamma_0 = 1$. Then there exists a constant $C > 0$ such that

$$\mathbb{P}(A_\tau > x, \bar{X}_\tau \leq y) \leq Cx^{1/4} \exp \left\{ -2 \frac{g(y)}{y} \sqrt{\left(\frac{a}{2} - \frac{2C_0g(y)}{y} \right) x} \right\} \quad (32)$$

for all y satisfying $C_0g(y) \leq ay/4$, where C_0 is the constant given by Lemma 6. Moreover,

$$\mathbb{P}(A_\tau > x) \leq Cx^{1/4} \exp \left\{ -g(\sqrt{2ax}) \sqrt{\left(1 - \frac{2C_0g(\sqrt{2ax})}{a\sqrt{2ax}} \right)^+} \right\} x \geq 1,$$

Proof. Using Lemma 6 with $y = \sqrt{2ax}$ we obtain

$$\begin{aligned} \mathbb{P}(A_\tau > x, \bar{X}_\tau \leq y) &\leq \sum_{n=0}^{\infty} \mathbb{P}(A_n \geq x, \bar{X}_n \leq \sqrt{2ax}, \tau = n+1) \\ &\leq \sum_{n=1}^{\infty} \exp \left\{ -\lambda \frac{x}{n} - \frac{a\lambda}{2} n + C\lambda^2 n \right\} = \sum_{n=1}^{\infty} \exp \left\{ -\lambda \frac{x}{n} - \lambda In \right\}, \end{aligned}$$

where $\lambda = \frac{g(\sqrt{2ax})}{\sqrt{2ax}}$ and $I = \frac{a}{2} - C\lambda$. The assumption $C_0g(y) \leq y\frac{a}{4}$ implies that $I > \frac{a}{4}$. Since I is positive, we have the inequality

$$\int_{n-1}^n \exp \left\{ -\lambda \frac{x}{y} - \lambda I(y+1) \right\} dy \geq \exp \left\{ -\lambda \frac{x}{n} - \lambda In \right\}, \quad n \geq 1.$$

With formula (25) on page 146 of [1], we have

$$\begin{aligned} \sum_{n=1}^{\infty} \exp \left\{ -\lambda \frac{x}{n} - \lambda In \right\} &\leq \int_0^{\infty} \exp \left\{ -\lambda \frac{x}{y} - \lambda I(y+1) \right\} dy \\ &= e^{-\lambda I} \sqrt{\frac{4x}{I}} K_1(2\lambda\sqrt{Ix}). \end{aligned}$$

Now, using the asymptotics for the modified Bessel function

$$K_1(z) \sim \sqrt{\frac{\pi}{2z}} e^{-z},$$

we obtain

$$\sum_{n=1}^{\infty} \exp \left\{ -\lambda \frac{x}{n} - \lambda In \right\} \leq Cx^{1/4} \exp\{-2\lambda\sqrt{Ix}\}.$$

Therefore, (32) is proven.

The second claim in the lemma obviously holds for all x such that $C_0g(\sqrt{2ax}) \geq a\sqrt{2ax}$. Assume that x is so large that $C_0g(\sqrt{2ax}) < a\sqrt{2ax}$. Clearly,

$$\mathbb{P}(A_\tau > x) \leq \mathbb{P}(A_\tau > x, \bar{X}_\tau \leq \sqrt{2ax}) + \mathbb{P}(A_\tau > x, \bar{X}_\tau > \sqrt{2ax}) =: P_1 + P_2.$$

By (32) with $y = \sqrt{2ax}$,

$$P_1 \leq Cx^{1/4} \exp \left\{ -g(\sqrt{2ax}) \sqrt{1 - \frac{2C_0g(\sqrt{2ax})}{a\sqrt{2ax}}} \right\}.$$

Next,

$$\begin{aligned} P_2 &\leq \sum_{n=0}^{\infty} \mathbb{P}(A_\tau \geq x, M_n \leq \sqrt{2ax}, X_{n+1} > \sqrt{2ax}, \tau > n) \\ &\leq \sum_{n=0}^{\infty} \mathbb{P}(X_{n+1} > \sqrt{2ax}) \mathbb{P}(\tau > n) \leq \mathbb{E}[\tau] \bar{F}(\sqrt{2ax}) = o(P_1). \end{aligned}$$

Then, the claim follows. □

Now we will give a lower bound.

Lemma 8. *Let $\mathbb{E}[X_1] = -a < 0$ and $\text{Var}(X_1) < \infty$. Then, for any $\varepsilon > 0$ there exists $C > 0$ such that*

$$\liminf_{x \rightarrow \infty} \frac{\mathbb{P}(A_\tau > x)}{\bar{F}(\sqrt{2ax} + Cx^{1/4+\varepsilon/2})} \geq \mathbb{E}\tau.$$

Proof. Fix $N \geq 1$. Put $y^+ = \sqrt{2ax} + Cx^{1/4+\varepsilon/2}$, where C will be picked later. Since $\mathbb{E}[X_1^2] < \infty$, by the strong law of large numbers,

$$\frac{S_l + al}{l^{1/2+\varepsilon}} \rightarrow 0, \quad l \rightarrow \infty \text{ almost surely.}$$

Hence, for any $\delta > 0$ we can pick $R > 0$ such that

$$\mathbb{P} \left(\min_{l \leq \sqrt{2x/a}} (S_l + al + R + l^{1/2+\varepsilon}) > 0 \right) > (1 - \delta).$$

Define

$$E_k^+ := \left\{ \min_{l \leq \sqrt{2x/a}} (S_{k+l} - S_k + al + R + l^{1/2+\varepsilon}) > 0, \tau > k, S_k > y^+ \right\}.$$

If $C > 1 + (2/a)$ then, for all x large enough, $al + l^{1/2+\varepsilon} + R \leq \sqrt{2ax} + (2x/a)^{1/4+\varepsilon/2} + R \leq y^+$ for all $l \leq \sqrt{2x/a}$. Therefore, for every $k \leq N$, $E_k^+ \subset \{\tau > k + \sqrt{2x/a}\}$. Furthermore, if $\tau > k + \sqrt{2x/a}$ then, on the event E_k^+ ,

$$\begin{aligned} A_\tau &> \sum_{l=0}^{k+\sqrt{2x/a}} S_{k+l} > y^+ \sqrt{2x/a} + \sum_{l=0}^{k+\sqrt{2x/a}} (S_{k+l} - S_l) \\ &> y^+ \sqrt{2x/a} - \sum_{l=0}^{k+\sqrt{2x/a}} (al + l^{1/2+\varepsilon} + R). \end{aligned}$$

Now, we can choose C so large that, for every $k \leq N$, $E_k^+ \subset \{A_\tau > x\}$. Hence,

$$\begin{aligned} \mathbb{P}(A_\tau > x) &\geq \sum_{k=1}^N \mathbb{P}(A_\tau > x, \bar{X}_{k-1} \leq y^+, X_k > y^+, \tau > k) \\ &\geq \sum_{k=1}^N \mathbb{P}(E_k^+, \bar{X}_{k-1} \leq y^+, X_k > y^+, \tau > k) \\ &\geq \sum_{k=1}^N \mathbb{P}\left(\bar{X}_{k-1} \leq y^+, \tau > k-1, X_k > y^+, \min_{l \leq \sqrt{2x/a}} (S_{l+k} - S_k + R + l^{1/2+\varepsilon}) > 0\right) \\ &\geq (1 - \delta) \sum_{k=1}^N \mathbb{P}(\bar{X}_{k-1} \leq y^+, \tau > k-1) \bar{F}(y^+). \end{aligned}$$

For every fixed k we have $\mathbb{P}(\bar{X}_{k-1} \leq y^+, \tau > k-1) \rightarrow \mathbb{P}(\tau > k-1)$ as $x \rightarrow \infty$. Furthermore, $\sum_{k=0}^N \mathbb{P}(\tau > k) \rightarrow \mathbb{E}\tau$ as $N \rightarrow \infty$. Therefore, we can pick sufficiently large N such that

$$\liminf_{x \rightarrow \infty} \sum_{k=1}^N \mathbb{P}(\bar{X}_{k-1} \leq y^+, \tau > k-1) \geq (1 - \delta)\mathbb{E}\tau.$$

Then, for all x sufficiently large, $\mathbb{P}(A_\tau > x) \geq (1 - \delta)^2 \mathbb{E}\tau \bar{F}(y^+)$. As $\delta > 0$ is arbitrarily small, we arrive at the conclusion. □

Proof of Theorem 1. The upper bound follows from Lemma 7. The lower bound follows from Lemma 8. The rough asymptotics follow immediately from the lower and upper bounds, and from the observation that

$$\sup_{|y| \leq x\rho(x)} \left| \frac{\log \bar{F}(x)}{\log \bar{F}(x+y)} - 1 \right| \rightarrow 0, \tag{33}$$

where $\rho(x) \rightarrow 0$. To prove (33) we note that by (9) and (10)

$$\begin{aligned} g(x+y) - g(x) &= \int_x^{x+y} g'(t) dt \leq \gamma_0 \int_x^{x+y} \frac{g(t)}{t} dt \leq \gamma_0 \frac{g(x)}{x^{\gamma_0}} \int_x^{x+y} \frac{1}{t^{1-\gamma_0}} dt \\ &\leq \gamma_0 \frac{g(x)}{x^{\gamma_0}} \frac{y}{x^{1-\gamma_0}} = \gamma_0 g(x) \frac{y}{x}, \quad y > 0. \end{aligned}$$

This implies that, as $x \rightarrow \infty$,

$$\sup_{|y| \leq x\rho(x)} \left| \frac{g(x+y)}{g(x)} - 1 \right| \rightarrow 0. \tag{34}$$

Recalling that $\log \bar{F}(x) \sim -g(x) - 2 \log x$, one easily obtains (33). □

4. Proof of Theorem 2

Set

$$h(x) := \frac{\sqrt{2ax}}{g(\sqrt{2ax})}, \quad y = \sqrt{2ax} - Ch(x) \log x,$$

where $C > \frac{5/4}{1-\gamma_0}$. We first split the probability $\mathbb{P}(A_\tau > x)$ as follows:

$$\begin{aligned} \mathbb{P}(A_\tau > x) &= \mathbb{P}(A_\tau > x, \bar{X}_\tau \leq y) + \mathbb{P}\left(A_\tau > x, \bar{X}_\tau > \sqrt{2ax} - \frac{1}{\log x}h(x)\right) \\ &\quad + \mathbb{P}\left(A_\tau > x, \bar{X}_\tau \in \left[y, \sqrt{2ax} - \frac{1}{\log x}h(x)\right]\right) =: P_1 + P_2 + P_3. \end{aligned}$$

The first term will be estimated using the exponential bound proved in Lemma 6.

Lemma 9. *Let $\mathbb{E}[X_1] = -a$ and $\text{Var}(X_1) < \infty$. Assume that (8) and (9) hold with some $\gamma_0 < 1/2$, together with (11). Then, $P_1 = o(\bar{F}(\sqrt{2ax}))$.*

Proof. According to (32),

$$P_1 \leq Cx^{1/4} \exp \left\{ -2\frac{g(y)}{y} \sqrt{\left(\frac{a}{2} - 2C_0\frac{g(y)}{y}\right)x} \right\}.$$

Since (9) holds for some $\gamma_0 < 1/2$, $g^2(y)/y \rightarrow 0$, and hence

$$P_1 \leq Cx^{1/4} \exp \left\{ -\frac{g(y)}{y} \sqrt{2ax} \right\}.$$

Then,

$$\frac{P_1}{\bar{F}(\sqrt{2ax})} \leq Cx^{5/4} \exp \left\{ g(\sqrt{2ax}) - \frac{g(y)}{y} \sqrt{2ax} \right\}.$$

To finish the proof, it is sufficient to show that

$$g(\sqrt{2ax}) - \frac{g(y)}{y} \sqrt{2ax} + \frac{5}{4} \log x \rightarrow -\infty, \quad x \rightarrow \infty. \tag{35}$$

We first note that

$$\begin{aligned} d(x) &:= g(\sqrt{2ax}) - \frac{g(y)}{y} \sqrt{2ax} = g(\sqrt{2ax}) - \frac{g(y)}{1 - C\frac{\log x}{g(\sqrt{2ax})}} \\ &= g(\sqrt{2ax}) - g(y) - (C + o(1)) \log x \frac{g(y)}{g(\sqrt{2ax})}. \end{aligned}$$

Using (18), we can see that

$$g(\sqrt{2ax}) - g(y) \leq \gamma_0 \frac{g(y)}{y} (\sqrt{2ax} - y) = \gamma_0 C \frac{g(y)}{y} \log x \frac{\sqrt{2ax}}{g(\sqrt{2ax})}.$$

Hence,

$$d(x) \leq \left(\gamma_0 \frac{\sqrt{2ax}}{y} - 1 \right) (C + o(1)) \frac{g(y)}{g(\sqrt{2ax})} \log x.$$

According to (34), $g(y) \sim g(\sqrt{2ax})$. Therefore, (35) is valid for any C satisfying $C(\gamma_0 - 1) + \frac{5}{4} < 0$. □

The next lemma gives the term that dominates in $\mathbb{P}(A_\tau > x)$.

Lemma 10. *Under the assumptions of Lemma 9 we have the following estimate:*

$$P_2 \leq (\mathbb{E}\tau + o(1))\bar{F}(\sqrt{2ax}), \quad x \rightarrow \infty.$$

Proof. Put

$$y^* = \sqrt{2ax} - \frac{h(x)}{\log x}.$$

By the total probability formula,

$$\begin{aligned} P_2 &\leq \sum_{n=0}^{\infty} \mathbb{P}(A_\tau \geq x, \bar{X}_n \leq y^*, X_{n+1} > y^*, \tau > n) \\ &\leq \sum_{n=0}^{\infty} \mathbb{P}(X_{n+1} > y^*)\mathbb{P}(\tau > n) = \mathbb{E}[\tau]\bar{F}(y^*). \end{aligned}$$

Now, note that, by (18) and (34),

$$\begin{aligned} \frac{\bar{F}(y^*)}{\bar{F}(\sqrt{2ax})} &\leq (1 + o(1))e^{g(\sqrt{2ax}) - g(y^*)} \leq (1 + o(1)) \exp \left\{ \frac{\gamma_0 g(y^*)}{y^*} (\sqrt{2ax} - y^*) \right\} \\ &\leq (1 + o(1)) \exp \left\{ \frac{\gamma_0 g(y^*)}{y^*} \frac{1}{\log x} \frac{\sqrt{2ax}}{g(\sqrt{2ax})} \right\} = 1 + o(1). \end{aligned}$$

Then the statement immediately follows. □

We proceed to the analysis of P_3 . Fix some $\delta > 0$ and set $z = \frac{1}{a} (\sqrt{2ax} + \delta\sqrt{x})$. We split P_3 further as follows:

$$\begin{aligned} P_3 \leq P_{31} + P_{32} + P_{33} &:= \mathbb{P} \left(A_\tau > x, \bar{X}_\tau \in \left[y, \sqrt{2ax} - \frac{1}{\log x} h(x) \right]; J_1; \tau \leq z \right) \\ &\quad + \mathbb{P} \left(A_\tau > x, \bar{X}_\tau \in \left[y, \sqrt{2ax} - \frac{1}{\log x} h(x) \right]; J_{\geq 2}, \tau \leq z \right) \\ &\quad + \mathbb{P}(\tau > z), \end{aligned}$$

where $J_1 = \{\text{there exists } k(1, \tau) \text{ such that } X_k > y \text{ and } \max_{1 \leq i \leq \tau, i \neq k} X_i \leq y\}$ and, correspondingly, $J_{\geq 2} = \{\text{there exist } k, l \in (1, \tau) \text{ such that } X_k > y \text{ and } X_l > y\}$.

We start with the easier terms P_{32} and P_{33} . To deal with these terms we will use Proposition 2.

Lemma 11. *Let assumptions (8) and (9) hold for $\gamma_0 < 1/2$. Assume also that (11) holds as well. Then $P_{33} = o(\bar{F}(\sqrt{2ax}))$ as $x \rightarrow \infty$.*

Proof. We have, by Proposition 2, $P_{33} = \mathbb{P}(\tau > z) \leq (\mathbb{E}\tau + o(1))\bar{F}(az) = O(\bar{F}(\sqrt{2ax} + \delta\sqrt{x}))$. Therefore,

$$\frac{P_{33}}{\bar{F}(\sqrt{2ax})} \leq C \exp \left\{ g(\sqrt{2ax}) - g(\sqrt{2ax} + \delta\sqrt{x}) \right\}.$$

By the mean value theorem and by the assumption (11), $g(cx) - g(x) \rightarrow \infty$ as $x \rightarrow \infty$ for every $c > 1$. This completes the proof. □

Lemma 12. *Let $\mathbb{E}[X_1] = -a$ and $\text{Var}(X_1) < \infty$. Assume that (8) and (9) hold with some $\gamma_0 < 1/2$, together with (11). Then $P_{32} = o(\bar{F}(\sqrt{2ax}))$.*

Proof. We can use the formula of total probability to write

$$P_{32} \leq \sum_{k=1}^z \mathbb{P}(\tau = k, J_{\geq 2}) \leq \sum_{k=1}^z \frac{k^2}{2} \bar{F}(y)^2.$$

Then,

$$\frac{P_{32}}{\bar{F}(\sqrt{2ax})} \leq Cx^{3/2} \frac{\bar{F}(y)^2}{\bar{F}(\sqrt{2ax})} \leq Cx^{1/2} e^{g(\sqrt{2ax})-2g(y)}.$$

Using (18) we can see that, in view of (12),

$$\frac{P_{32}}{\bar{F}(\sqrt{2ax})} \leq Cx^{1/2} e^{C \ln x - g(y)} \rightarrow 0. \quad \square$$

P_{31} remains to be analysed. For that, introduce $\mu(y) := \min\{n \geq 1 : X_k > y\}$. Now we will complete the proof with the following lemma.

Lemma 13. *Let assumptions (8), (9), and (11) hold for $\gamma_0 < 1/2$. Then $P_{31} = o(\bar{F}(\sqrt{2ax}))$ as $x \rightarrow \infty$.*

Proof. First, represent event J_1 as $J_1 = J_{11} \cup J_{12}$, where

$$J_{11} := \{X_k > y \text{ for exactly one } k \in (0, \tau) \text{ and } X_i \leq x^\varepsilon \text{ for all other } i < \tau, \}$$

$$J_{12} := \{X_k > y \text{ for exactly one } k \in (0, \tau) \text{ and } X_i > x^\varepsilon \text{ for some } i \neq k, i < \tau\}.$$

Then,

$$\begin{aligned} Q_2 &:= \mathbb{P} \left(A_\tau > x, \bar{X}_\tau \in \left[y, \sqrt{2ax} - \frac{1}{\log x} h(x) \right]; J_{12}, \tau \leq z \right) \\ &\leq \sum_{j=1}^z \mathbb{P}(\tau = j, J_{12}) \leq \sum_{j=1}^z \frac{j^2}{2} \bar{F}(y) \bar{F}(x^\varepsilon) \leq z^3 \bar{F}(y) \bar{F}(x^\varepsilon), \end{aligned}$$

so

$$\frac{Q_2}{\bar{F}(\sqrt{2ax})} \leq Cx^{3/2-2\varepsilon} e^{g(\sqrt{2ax})-g(y)-g(x^\varepsilon)}$$

By (18), $g(\sqrt{2ax}) - g(y) \leq C \ln x$. Then, in view of the relation (12), we have $g(\sqrt{2ax}) - g(y) - g(x^\varepsilon) \leq -4 \ln x$, which implies that $Q_2 = o(\bar{F}(\sqrt{2ax}))$.

To estimate

$$Q_1 := \mathbb{P} \left(A_\tau > x, \bar{X}_\tau \in \left[y, \sqrt{2ax} - \frac{1}{\log x} h(x) \right]; J_{11}, \tau \leq z \right)$$

we make use of the exponential bound given in Lemma 6. Put

$$x^+(k) = x - k \left(\sqrt{2ax} - \frac{h(x)}{\log x} \right).$$

Then, we have

$$\begin{aligned} Q_1 &= \sum_{k=0}^{z-1} \sum_{j=1}^k \mathbb{P} \left(A_k > x, \max_{i \neq j, i \leq k} X_i \leq x^\varepsilon, X_j \in \left[y, \sqrt{2ax} - \frac{h(x)}{\log x} \right], \tau = k + 1 \right) \\ &\leq \sum_{k=1}^z (k+1) \mathbb{P}(A_k > x^+(k), \bar{X}_k \leq x^\varepsilon) \bar{F}(y) \\ &\leq Cx^{1/2} \bar{F}(y) \sum_{k=1}^z \exp \left\{ -\lambda \frac{x^+(k)}{k} - \frac{a\lambda}{2} k + C\lambda^2 k \right\}, \end{aligned}$$

where $\lambda = \frac{g(x^\varepsilon)}{x^\varepsilon}$. Now note that

$$-\lambda \frac{x^+(k)}{k} - \frac{a\lambda}{2} k = -\lambda \left(-\sqrt{2ax} + \frac{h(x)}{\log x} + \frac{x}{k} + \frac{ak}{2} \right).$$

Since

$$\frac{x}{k} + \frac{ak}{2} \geq \sqrt{2ax}, \quad k \geq 1,$$

we obtain

$$-\lambda \frac{x^+(k)}{k} - \frac{a\lambda}{2} k \leq -\lambda \frac{h(x)}{\log x}, \quad k \geq 1.$$

Thus, $Q_1 \leq Cx e^{-\lambda h(x)/\log x + \lambda^2 z} \bar{F}(y)$. Next, we can pick $\varepsilon = \frac{1}{4(1-\gamma_0)}$ to achieve

$$\begin{aligned} \lambda^2 z &\leq C \left(\frac{g(x^\varepsilon)}{x^\varepsilon} \right)^2 x^{1/2} = C \left(\frac{g(x^\varepsilon)}{x^{\varepsilon(1-1/(4\varepsilon))}} \right)^2 = C \left(\frac{g(x^\varepsilon)}{x^{\gamma_0 \varepsilon}} \right)^2 \\ &< C \sup_t \left(\frac{g(t)}{t^{\gamma_0}} \right)^2 < \infty \end{aligned}$$

by the condition (9). Note that the assumption $\gamma_0 < 1/2$ implies that $\varepsilon = \frac{1}{4(1-\gamma_0)} < 1/2$. Then, using (8), we obtain

$$\frac{Q_1}{\bar{F}(\sqrt{2ax})} \leq Cx e^{g(\sqrt{2ax}) - g(y) - \lambda h(x)/\log x},$$

and, using (18),

$$\frac{Q_1}{\bar{F}(\sqrt{2ax})} \leq Cx^C e^{-\lambda h(x)/\log x}.$$

Finally, noting that

$$\lambda h(x) = \frac{g(x^\varepsilon)}{x^\varepsilon} \frac{\sqrt{2ax}}{g(\sqrt{2ax})}$$

grows polynomially, we obtain the required convergence to 0. The polynomial growth can be immediately seen for $g(x) = x^{\gamma_0}$. However, a proper proof goes as follows:

$$\begin{aligned} g(C\sqrt{x}) &= g(x^\varepsilon) + \int_{x^\varepsilon}^{C\sqrt{x}} g'(t) dt \leq g(x^\varepsilon) + \gamma_0 \int_{x^\varepsilon}^{C\sqrt{x}} \frac{g(t)}{t} dt \\ &\leq g(x^\varepsilon) + \gamma_0 \int_{x^\varepsilon}^{C\sqrt{x}} \frac{g(t)}{t^{\gamma_0}} t^{\gamma_0-1} dt \leq g(x^\varepsilon) + \frac{g(x^\varepsilon)}{x^{\varepsilon\gamma_0}} \int_{x^\varepsilon}^{C\sqrt{x}} t^{\gamma_0-1} dt \\ &\leq g(x^\varepsilon) + C \frac{g(x^\varepsilon)}{x^{\varepsilon\gamma_0}} x^{\gamma_0/2} \leq Cg(x^\varepsilon)x^{\gamma_0(1/2-\varepsilon)}. \end{aligned}$$

Therefore $\lambda h(x) \geq x^{1/2-\varepsilon} x^{-\gamma_0(1/2-\varepsilon)} = x^{(1-\gamma_0)/2-1/4}$, where we have used the equality $\varepsilon = \frac{1}{4(1-\gamma_0)}$. \square

Proof of Theorem 2. Combining the preceding lemmas give us the upper bound. The lower bound has been shown in (5) under even weaker conditions. \square

Acknowledgement

We would like to thank the anonymous referees for a number of useful comments and suggestions that helped us to improve the paper.

References

- [1] BATEMAN, H. (1954). *Tables of Integral Transforms*, Vol. 1. McGraw-Hill, New York.
- [2] BOROVKOV, A. A., BOXMA, O. J. AND PALMOWSKI, Z. (2003). On the integral of the workload process of the single server queue. *J. Appl. Prob.* **40**, 200–225.
- [3] CARMONA, P. AND PÉTRÉLIS, N. (2019). A shape theorem for the scaling limit of the IPDSAW at criticality. *Ann. Appl. Prob.* **29**, 875–930.
- [4] DENISOV, D. (2005). A note on the asymptotics for the maximum on a random time interval of a random walk. *Markov Process. Relat. Fields* **11**, 165–169.
- [5] DENISOV, D., DIEKER, A. B. AND SHNEER, V. (2008). Large deviations for random walks under subexponentiality: the big-jump domain. *Ann. Prob.* **36**, 1946–1991.
- [6] DENISOV, D., KOLB M. AND WACHTEL, V. (2015). Local asymptotics for the area of random walk excursions. *J. London Math. Soc.* **91**, 495–513.
- [7] DENISOV, D. AND SHNEER, V. (2013). Asymptotics for the first passage times of Lévy processes and random walks. *J. Appl. Prob.* **50**, 64–84.
- [8] DOBRUSHIN, R. AND HRYNIV, O. (1996). Fluctuations of shapes of large areas under paths of random walks. *Prob. Theory Relat. Fields* **105**, 423–458.
- [9] DONEY, R. A. (1989). On the asymptotic behaviour of first passage times for transient random walks. *Prob. Theory Relat. Fields* **81**, 239–246.
- [10] DUFFY, K. AND MEYN, S. (2014). Large deviation asymptotics for busy periods. *Stoch. Syst.* **4**, 300–319.
- [11] FELLER, W. (1971). *An Introduction to Probability Theory and Its Applications*, Vol. 2. Wiley, New York.
- [12] FOSS, S., PALMOWSKI, Z. AND ZACHARY, S. (2005). The probability of exceeding a high boundary on a random time interval for a heavy-tailed random walk. *Ann. Appl. Prob.* **15**, 1936–1957.
- [13] KULIK, R. AND PALMOWSKI, Z. (2011). Tail behaviour of the area under a random process, with applications to queueing systems, insurance and percolations. *Queueing Systems* **68**, 275–284.
- [14] PERFILEV, A. AND WACHTEL, V. (2018). Local asymptotics for the area under the random walk excursion. *Adv. Appl. Prob.* **50**, 600–620.
- [15] ROZOVSKII, L. V. (1993). Probabilities of large deviations on the whole axis. *Theory Prob. Appl.* **38**, 53–79.
- [16] TAKACS, L. (1991). A Bernoulli excursion and its various applications. *Adv. Appl. Prob.* **23**, 557–585.
- [17] TAKACS, L. (1992). On the total heights of random rooted trees. *J. Appl. Prob.* **29**, 543–556.
- [18] TAKACS, L. (1994). On the total heights of random rooted binary trees. *J. Combinatorial Theory B* **61**, 155–166.