

Main Article

Dr O Reichel takes responsibility for the integrity of the content of the paper

Cite this article: Reichel O, Taxeidis M. Use of an image-guided navigation system for routine endonasal endoscopic dacryocystorhinostomy. *J Laryngol Otol* 2019;**133**:685–690. <https://doi.org/10.1017/S0022215119001567>

Accepted: 18 April 2019
First published online: 24 July 2019

Key words:

Dacryocystorhinostomy; Neuronavigation; Lacrimal Apparatus Diseases; Tears

Author for correspondence:

Dr Oliver Reichel,
Department of Otolaryngology,
Head and Neck Surgery,
Siloah St. Trudpert Hospital,
Wilferdinger Str. 67, Pforzheim 75179,
Germany
E-mail: o.reichel@siloh.de

Use of an image-guided navigation system for routine endonasal endoscopic dacryocystorhinostomy

O Reichel and M Taxeidis

Department of Otolaryngology, Head and Neck Surgery, Siloah St. Trudpert Hospital, Pforzheim, Germany

Abstract

Objective. This study aimed to evaluate the results of routine endonasal endoscopic dacryocystorhinostomy combined with computed tomography guided navigation in patients with nasolacrimal duct obstruction.

Method. This was a retrospective study of all patients with lacrimal drainage obstruction undergoing stereotactic endoscopic dacryocystorhinostomy between 1st January 2016 and 1st April 2018. Computed tomography dacryocystography was used for intra-operative navigation. Patients with a presaccal obstruction site location were excluded from the study.

Results. Endoscopic dacryocystorhinostomy with computed tomography guided navigation was successfully performed in all 17 cases without complications. Early post-operative dislocation of the inserted bicanalicular silicone stent occurred in two patients. Two other patients developed post-operative bacterial infection within the lacrimal sac. Otherwise, the silicone tube was removed three months after surgery, and after further follow up of 8 weeks, 94 per cent of the study population reported complete remission of epiphora.

Conclusion. The use of computed tomography guidance in routine endoscopic dacryocystorhinostomy enhanced safety for the patient and avoided unnecessary damage of bone and mucosa surrounding the lacrimal drainage system. Therefore, routine endoscopic dacryocystorhinostomy with additional stereotactic guidance by computed tomography navigation can contribute to high success rates with endoscopic dacryocystorhinostomy.

Introduction

With the introduction of rigid endoscopes, the endonasal approach to the saccus lacrimalis has become established as a real alternative to external surgical treatment of lacrimal drainage obstruction.¹ Studies of endoscopically performed dacryocystorhinostomy (DCR) report success rates of 81 to 94 per cent, comparable to those of external DCR.^{2,3} Therefore, both the endonasal and extranasal approach are well established and widely accepted surgical procedures for the treatment of nasolacrimal duct obstruction. In the hands of otolaryngologists, however, the endonasal endoscopic technique has several advantages, such as allowing paranasal sinus surgery or septoplasty to be performed simultaneously.⁴ The avoidance of a facial scar and preservation of the medial canthal ligament are further arguments for choosing an endonasal approach for surgical management of nasolacrimal duct obstruction.⁵

Even though ENT surgeons are very familiar with navigation-guided endoscopic surgery for paranasal sinus disorders, no studies of image-guided routine endoscopic DCR performed by otolaryngologists exist. Most studies on this topic are by ophthalmologists who have chosen an endonasal endoscopic approach to the saccus lacrimalis with simultaneous use of computed tomography (CT) navigation.^{6–11} Therefore, we conducted a retrospective study of patients with nasolacrimal duct obstruction undergoing routine endoscopic DCR with CT-guided navigation performed by one ENT surgeon in order to evaluate our results from the otolaryngologist's point of view.

Materials and methods

A retrospective study was performed on 16 patients with unilateral or bilateral ($n = 1$) epiphora who underwent endoscopic DCR with CT-guided navigation between January 2016 and April 2018.

Endoscopic DCR is well accepted as the most suitable treatment for patients with obstructions of the nasolacrimal system, particularly at the level of the lacrimal sac or in the nasolacrimal duct. Thus, in our ENT department endoscopic DCR with CT guidance is the standard procedure for all patients who have a localised obstruction in this area.¹² Seven patients were male, and the mean age of the study population was 54 years (range, 14–82 years). In most patients, no previous DCR had been performed. Only one patient suffered from recurrence of epiphora after endoscopic DCR without

Table 1. Patient characteristics

Patient	Age	Sex	Side	Obstruction site	Follow up after stent removal or dislocation (months)	Complete remission	Complication
1	41	Male	Left	Junction of lacrimal sac and nasolacrimal duct	9	Yes	
2	52	Female	Right	Lacrimal sac	4	No	
3	37*	Female	Right	Lacrimal sac	17	Yes	
4	65	Male	Right	Lacrimal sac	14	Yes	Dislocation after 3 weeks
5	67	Male	Left	Nasolacrimal duct	11	Yes	
6	68*	Female	Left	Lacrimal sac	8	Yes	
7	68 [†]	Female	Right	Junction of lacrimal sac and nasolacrimal duct	4	Yes	<i>Staphylococcus aureus</i> infection
8	77	Female	Bilateral	Lacrimal sac right, junction of lacrimal sac and nasolacrimal duct left	6	Yes	
9	52*	Male	Left	Lacrimal sac	5	Yes	
10	54	Female	Right	Lacrimal sac	12	Yes	<i>Streptococcus pneumoniae</i> infection
11	44	Male	Left	Lacrimal sac	7	Yes	Dislocation after 2 weeks
12	52	Male	Right	Nasolacrimal duct	11	Yes	
13	82	Male	Left	Lacrimal sac	4	Yes	
14	66	Female	Left	Lacrimal sac	6	Yes	
15	14	Female	Right	Nasolacrimal duct	2	Yes	
16	24*	Female	Right	Junction of lacrimal sac and nasolacrimal duct	2	Yes	

*use of a curved diamond drill; [†]revision case after previous surgery at another location

image-guidance that was performed elsewhere. Patients' characteristics are summarised in Table 1.

All patients or parents provided written informed consent and patients were referred for surgical therapy by ophthalmologists after verification of saccal or post-saccal obstruction by diagnosing reflux during irrigation of the efferent lacrimal duct system. In order to determine the exact location of nasolacrimal duct obstruction and for pre-operative analysis of concomitant nasal pathologies, all enrolled patients underwent low-dose helical CT-dacryocystography with three-dimensional reconstruction. This low-radiation imaging tool, first described in 2002 by Freitag *et al.*, was well tolerated by all patients in our study without pain or complications and provided detailed imaging of the nasolacrimal system including the surrounding structures.¹³

All surgical procedures were performed by one surgeon and under general anaesthesia as an in-patient procedure. During surgery, image-guided dacryolocalisation was performed by means of the BrainLab Kolibri navigation system (BrainLAB, Heimstetten, Germany) after setup and registration with the z-touch system (Figure 1). The calculated registration error was within 1 mm.

For mucosal decongestion, the nasal cavity was packed with gauze soaked with naphazoline for 5 minutes. After removal of the gauze, an infiltration of the mucosa in the area of the crista lacrimalis in front of the middle turbinate was performed with xylocaine (1 per cent) with adrenalin (1 per cent). Then, a bicanalicular silicone stent (Canaliculus Intubation Set of 15.5 × 0.051 cm; Beaver-Visitec International, Sydney, Australia) with stainless steel probes at both ends was inserted through the upper and lower puncta.

During the next step, a handpiece was used to accurately determine the location of the bony medial wall of the saccus lacrimalis using a 30° rigid endoscope together with the navigation system (Figures 1 and 2). In this area, a small and cranial stemmed mucosal flap was elevated using a sickled knife and a freer elevator in order to minimally expose the bone of the frontal process of the maxilla facing the lacrimal sac (Figure 3). In most cases, a Kerrison punch was used to remove the bone above the medial wall of the lacrimal sac. In three cases, a curved diamond burr was used because of very thick bone in this area. By deeper insertion of one of the steel probes, the medial wall of the lacrimal sac was medialised towards the nasal cavity and incised with a sickled knife. Then, the medial wall of the lacrimal sac was resected and the silicone stent with the two steel probes was pushed into the nasal cavity.

In the next step, the two ends of the silicone stent were grasped endonasally with small forceps, and the stainless-steel probes were removed. The two ends of the silicone tube were knotted and sutured to the lateral nasal wall. Finally, the mucosal flap was replaced to cover the resilient exposed bone above the lacrimal sac. No nasal package was used, but topical steroid ointment was applied onto the mucosal flap at the end of surgery.

No additional septoplasty or extended paranasal sinus surgery needed to be performed in any patient. However, low-dose CT dacryocystography showed anatomic variations such as hyperpneumatisation of agger nasi cells or concha bullosa in six patients (Figure 4).

After surgery, patients underwent regular check-ups for removal of endonasal crusting, which took place either in



Fig. 1. Image showing dacrylocalisation surgery using the BrainLab Kolibri navigation system.

our department or by an ENT specialist in a private practice. No eye drops or nasal steroid sprays were prescribed. After three months, the silicone tube was removed in our department and patients had two further check-ups at four-week intervals. After removal of the bicanalicular silicone stent, every patient underwent at least two control examinations before final assessment of the result after surgical intervention (minimal follow up of 5 months after endoscopic DCR). We considered our surgery to be successful if the patient was completely symptom-free after the second check-up.

Ethical approval

The present study is retrospective with no changes to any patient's care. Only members of the team caring for included patients had access to non-anonymised data. Data collection was only performed as part of the care of these patients, and all data were collated by the authors in an anonymous form. Therefore, research ethics committee approval was not sought for this retrospective analysis of our results.

Results

By using pre-operative CT-dacrycystography, nasolacrimal system obstruction was precisely located in the lacrimal sac in 10 patients (60 per cent), at the junction of lacrimal sac and nasolacrimal duct in 4 patients (23 per cent) and within the nasolacrimal duct in 3 patients (17 per cent). During all surgical procedures, the location of the lacrimal sac could be accurately determined by navigation (Figure 5), and all 17 endoscopic DCRs were carried out successfully and without intra-operative complications.

Pre-operative setup of the navigation system together with registration was completed within a maximum of 20 minutes in all cases. In two patients, a dislocation of the silicone stent occurred during the early post-operative period (patient 4 in week 3 and patient 11 in week 2; Table 1). In these two cases, the silicone stent was removed transnasally earlier than three months after endoscopic DCR. This procedure succeeded without complications after incising the stent in the area of the lacrimal puncta.

In two patients, a post-operative bacterial infection within the lacrimal sac occurred, causing sanious secretion at the lacrimal puncta (Table 1). Microbiological analysis showed there was infection with *Staphylococcus aureus* in patient 7 (8 weeks

post-operatively) and infection with *Streptococcus pneumoniae* in patient 10 (5 weeks post-operatively). In both patients, the infection was effectively treated within 1 week using eye drops containing ofloxacin.

In 15 cases, the silicone tube was removed as planned after 3 months (bilateral removal in patient 8). The result of surgical therapy was assessed after the second check-up at 8 weeks after surgery. Patients 4 and 11 also underwent a check-up after 8 weeks, despite earlier removal of the silicone stent. Five months after endoscopic DCR, all patients reported complete remission of epiphora, except for patient 2 (94 per cent). In this case the epiphora continued, and the patient was offered a second surgical intervention. However, the patient refused revision surgery and was referred to her ophthalmologist. Further follow up was performed every 4 weeks in all patients, and after a mean follow up of 7.6 months, the overall success rate was still 94 per cent (complete remission of epiphora).

Discussion

In 1913, West first reported an endonasal technique for DCR, describing resection of the medial wall of the lacrimal sac to treat nasolacrimal duct obstruction.¹⁴ In the next development in 1989, McDonogh *et al.* introduced endoscopic DCR, a minimally invasive surgical procedure routinely performed by otolaryngologists in particular and now well accepted as an alternative for external DCR.¹ Ophthalmologists have increasingly appreciated the advantages of endoscopic DCR and started to combine endoscopic DCR with CT-guided navigation, particularly in complex endoscopic lacrimal surgical procedures.⁶⁻¹¹ However, no studies by otolaryngologists examining the benefit of additional CT-guided navigation in routine endoscopic DCR exist.

When clinical examination suggests a mechanical obstruction of the nasolacrimal duct system, the preferred radiological tool for imaging the nasolacrimal system in our ENT department is low-dose CT dacrycystography. Thus, the most important requirement for endoscopic DCR with CT-guided navigation was satisfied in most of our patients suffering from epiphora. Although some authors assume that CT-guidance is a useful adjunct to endoscopic DCR only in challenging or secondary acquired nasolacrimal duct obstruction,^{6,9,10} we use this technique in all patients undergoing routine endoscopic DCR.

One reason against the use of image-guidance in routine endoscopic DCR might be an increase in operating room time. In our study, set up and registration of navigation took a maximum of 20 minutes, which is comparable to other studies.¹⁵ However, those 20 additional minutes significantly enhanced both safety for the patient and comfort for the surgeon. The location of the lacrimal sac could be determined with an accuracy of 1 mm by means of image guidance in all our patients, despite anatomical variations in the nasal cavity, turbinates and nasal septum (Figure 4). As a consequence, maximum preservation of nasal mucosa and bone was possible, contributing to the high success rate of 94 per cent. According to a systematic review by Leong *et al.*, success rates of endoscopic DCR vary between 84 and 94 per cent.¹⁶ Factors contributing to a high success rate in endoscopic DCR include adequate opening of the bony nasolacrimal canal and adequate removal of the medial wall of the lacrimal sac.¹⁷ In our study, navigation guidance significantly facilitated the accomplishment of these two tasks.

Given that stent insertion is helpful for post-operative care and impedes fibrous closure after endoscopic DCR,¹⁸ we




Fig. 2. Image showing precise identification of the bony medial wall of the saccus lacrimalis on the right side using the navigation handpiece.




Fig. 3. Image showing minimal exposure of the bone of the frontal process of the maxilla (white arrow) after elevation of a cranial stemmed mucosal flap (white star).

routinely inserted a bicanalicular silicone stent in all patients to be left for three months. However, earlier removal of the stent did not negatively influence the outcome of endoscopic DCR in two of our patients after dislocation of the tube (patients 4 and 11; [Table 1](#)). This result is consistent with a systematic review and meta-analysis by Sarode *et al.*, who found no advantage of using a silicone stent in primary endoscopic DCR.¹⁹ Thus, the efficacy of stent insertion in patients undergoing endoscopic DCR remains uncertain.²⁰ Our rate of stent dislocation (12 per cent) is significantly higher than in a study by Horn *et al.*, which reported this post-operative complication after endoscopic DCR in 3 of 102 cases (3 per cent).⁴ Both cases of stent dislocation in our study were caused mechanically by the patient drying their face with a towel; therefore, we can rule out an association between our operating technique and the high dislocation rate.

Most of our patients showed a proximal obstruction site in the lacrimal sac (10 patients, 60 per cent of cases), which correlates with a lower success rate for endoscopic DCR in some studies.^{20,21} In contrast, Yung *et al.* reported a success rate of 95 per cent for endoscopic DCR with an obstruction site in the lacrimal sac or duct.²² In our study, the only patient




Fig. 4. Image from low-dose computed tomography dacryocystography showing a nasolacrimal drainage obstruction located in the right lacrimal sac. During endoscopic dacryocystorhinostomy, a hyperpneumatized agger nasi cell had to be removed (white arrow).

with persistent epiphora after navigation-guided endoscopic DCR suffered from an obstruction in the lacrimal sac (patient 2; [Table 1](#)). Thus, for this location of obstruction, the success rate in our study was 90 per cent. However, in the studies mentioned above, endoscopic DCR was not performed with CT-guided navigation. Therefore, the results cannot adequately be compared with the results of our study, and future studies with a larger study population are needed to analyse the influence of the obstruction site on the success rate of endoscopic DCR with CT-guided navigation.

In a recent retrospective chart review published by Nomura *et al.*, another factor associated with poor prognosis for endoscopic DCR, in addition to a proximal obstruction site, was patient age being 65 years or older.²⁰ In our study population, 44 per cent of the patients were 65 years or older. However, in this group of patients our success rate was 100 per cent. As we did not augment the function of the orbicularis muscle that can be weakened due to age, our CT-guided surgical technique




Fig. 5. Image showing image-guided navigation to precisely locate the left lacrimal sac before removal of the bone.

cannot be the reason for the high success rate in this subgroup. However, the use of image guidance in our study enabled the surgeon to achieve minimal exposure of bone and maximal preservation of mucosa in all patients, resulting in less bone neogenesis and scarring. Thus, in our opinion, the additional use of navigation guidance can significantly reduce unnecessary collateral damage of tissue surrounding the nasolacrimal duct system, resulting in excellent results after endoscopic DCR in patients of all ages.

- Long-term failure after endoscopic dacryocystorhinostomy (DCR) is most commonly caused by inaccurately locating the lacrimal sac and inappropriate resection of the medial wall of the sac
- Navigation guidance in routine endoscopic DCR accurately targets the precise location and minimises excessive mucosal resection
- This is the first study to evaluate the results of image-guided endoscopic DCR performed by an otolaryngologist
- In this study, the additional use of a navigation system resulted in excellent long-term results after routine endoscopic DCR
- Navigation-guidance can give superior long-term results for surgical treatment of nasolacrimal obstruction, even in ENT departments performing only a few endoscopic DCRs per year

Another factor contributing to the favourable outcomes in our study population may have been the use of a mucosal flap, which has been suggested to decrease cicatricial stenosis after endoscopic DCR.²³ The positive effect of this mucosa-sparing technique has been described in previous studies.^{23,24} In our study, an ideal configuration of the flap was maximally facilitated by image-guided dacrylocalisation (Figure 3).

Without routine post-operative application of antibiotic eye drops, two patients in our study developed an infection within the lacrimal sac causing sanious secretion at the lacrima puncta. In a study by Horn *et al.*, all patients routinely received ofloxacin eye drops post-operatively and no infections occurred.⁴ Thus, routine post-operative use of antibiotic eye drops might reduce post-operative infection rates. However, whether this procedure has a statistically significant effect needs to be evaluated in future studies with a larger number of patients.

To our knowledge, this is the first study evaluating the results of routine endoscopic DCR with CT-guided navigation performed by otolaryngologists. Our data shows 0 per cent intra-operative complications and 94 per cent success after a minimum follow-up of 5 months, though we only performed 17 endoscopic DCRs over a period of 28 months. A weakness of this study is the lack of a control group without image-guided assistance. However, the success rate of 94 per cent in our patients may indicate that additional use of navigation has a positive effect on the long-term outcome after routine endoscopic DCR.

The strengths of the current study are that we had one single operating surgeon using the same technique in all surgical procedures and a mean follow up of nearly eight months after removal of the silicone intubation. However, this retrospective study is limited by the relatively small number of patients. Therefore, our promising results require confirmation by future prospective studies with larger study populations and a study design that includes patient randomisation to endoscopic DCR with or without the assistance of navigation.

Conclusion

Endonasal endoscopic routine DCR with CT-guided navigation provides additional comfort for the surgeon, maximum safety for the patient and avoids unnecessary damage of bone and mucosa surrounding the lacrimal drainage system. Thus, additional image-guidance may contribute to excellent post-operative results after routine surgical therapy of nasolacrimal duct obstruction.

Competing interests. None declared

References

- 1 McDonogh M, Meiring JH. Endoscopic transnasal dacryocystorhinostomy. *J Laryngol Otol* 1989;**103**:585–7
- 2 Ben Simon GJ, Joseph J, Lee S, Schwarcz RM, McCann JD, Goldberg RA. External versus endoscopic dacryocystorhinostomy for acquired nasolacrimal duct obstruction in a tertiary referral center. *Ophthalmology* 2005;**112**:1463–8
- 3 Cho WK, Paik JS, Yang SW. Surgical success rate comparison in functional nasolacrimal duct obstruction: simple lacrimal stent versus endoscopic versus external dacryocystorhinostomy. *Eur Arch Otorhinolaryngol* 2013;**270**:535–40
- 4 Horn IS, Tittmann M, Fischer M, Otto M, Dietz A, Mozet C. Endonasal lacrimal duct surgery: a comparative study of two techniques. *Eur Arch Otorhinolaryngol* 2014;**271**:1923–31

- 5 Patel BC. Management of acquired nasolacrimal duct obstruction: external and endonasal dacryocystorhinostomy. Is there a third way? *Br J Ophthalmol* 2009;**93**:1416–19
- 6 Day S, Hwang TN, Pletcher SD, Bhatki A, McCulley TJ. Interactive image-guided endoscopic dacryocystorhinostomy. *Ophthalmic Plast Reconstr Surg* 2008;**24**:338–40
- 7 Morley AM, Collyer J, Malhotra R. Use of image-guided navigation system for insertion of a Lester-Jones tube in a patient with disturbed orbito-nasal anatomy. *Orbit* 2009;**28**:439–41
- 8 Ali MJ, Naik MN. Image-guided dacryolocalization (IGDL) in traumatic secondary acquired lacrimal drainage obstructions (SALDO). *Ophthalmic Plast Reconstr Surg* 2015;**31**:406–9
- 9 Ali MJ, Singh S, Naik MN, Kaliki S, Dave TV. Interactive navigation-guided ophthalmic plastic surgery: navigation enabling of telescopes and their use in endoscopic lacrimal surgeries. *Clin Ophthalmol* 2016;**10**:2319–24
- 10 Ali MJ, Singh S, Naik MN, Kaliki S, Dave TV. Interactive navigation-guided ophthalmic plastic surgery: the utility of 3D CT-DCG-guided dacryolocalization in secondary acquired lacrimal duct obstructions. *Clin Ophthalmol* 2016;**11**:127–33
- 11 Ali MJ, Singh S, Naik MN. Image-guided lacrimal drainage surgery in congenital arhinia-microphthalmia syndrome. *Orbit* 2017;**36**:137–43
- 12 Penttilä E, Smirnov G, Tuomilehto H, Kaarniranta K, Seppä J. Endoscopic dacryocystorhinostomy as treatment for lower lacrimal pathway obstructions in adults: Review article. *Allergy Rhinol (Providence)* 2015;**6**:12–19
- 13 Freitag SK, Woog JJ, Kousoubris PD, Curtin HD. Helical computed tomographic dacryocystography with three-dimensional reconstruction. *Ophthalmic Plast Reconstr Surg* 2002;**18**:121–32
- 14 West JM. The transnasal technique for opening the nasolacrimal sac. *Arch Laryngol Rhinol* 1913;**24**:62–4
- 15 Metson R, Cosenza M, Gliklich RE, Montgomery WW. The role of image-guidance systems for head and neck surgery. *Arch Otolaryngol Head Neck Surg* 1999;**125**:1100–4
- 16 Leong SC, MacEwen CJ, White PS. A systematic review of outcomes after dacryocystorhinostomy in adults. *Am J Rhinol Allergy* 2010;**24**:81–90
- 17 Onerci M, Orhan M, Ogretmenoglu O, Irkec M. Long-term results and reason for failure of intranasal endoscopic dacryocystorhinostomy. *Acta Otolaryngol* 2000;**120**:319–22
- 18 Bernal-Sprekelsen M, Alobid I, Tomas-Barberan M, Della Rocca RC, Schaefer SD. Dacryocystorhinostomy surgical technique. In: Weber RK, Keerl R, Schaefer SD, Della Rocca RC, eds. *Atlas of Lacrimal Surgery*. New York: Springer, 2007;61–8
- 19 Sarode D, Bari DA, Cain AC, Syed MI, Williams AT. The benefit of silicone stents in primary endonasal dacryocystorhinostomy: a systematic review and meta-analysis. *Clin Otolaryngol* 2017;**42**:307–14
- 20 Nomura K, Arakawa K, Sugawara M, Hidaka H, Suzuki J, Katori Y. Factors influencing endoscopic dacryocystorhinostomy outcome. *Eur Arch Otorhinolaryngol* 2017;**274**:2773–7
- 21 Choi JC, Jin HR, Moon YE, Kim MS, Oh JK, Kim HA *et al*. The surgical outcome of endoscopic dacryocystorhinostomy according to the obstruction levels of lacrimal drainage system. *Clin Exp Otorhinolaryngol* 2009;**2**:141–4
- 22 Yung MW, Hardmann-Lea S. Analysis of the results of surgical endoscopic dacryocystorhinostomy: effect of the level of obstruction. *Br J Ophthalmol* 2002;**86**:792–4
- 23 Mueller SK, Freitag SK, Lefebvre DR, Bleier BS. Endoscopic DCR using bipediced interlacing mucosal flaps. *Laryngoscope* 2018;**128**:794–7
- 24 Tsirbas A, Wormald PJ. Endonasal dacryocystorhinostomy with mucosal flaps. *Am J Ophthalmol* 2003;**135**:76–83