

ORIGINAL ARTICLE

An Evaluation of Food as a Potential Source for *Clostridium difficile* Acquisition in Hospitalized Patients

Jennie H. Kwon, MSCI;¹ Cristina Lanzas, DVM, PhD;² Kimberly A. Reske, MPH;¹ Tiffany Hink, BS;¹ Sondra M. Seiler, BS;¹ Kerry M. Bommarito, PhD;¹ Carey-Ann D. Burnham, PhD;³ Erik R. Dubberke, MD, MSPH¹

OBJECTIVE. To determine whether *Clostridium difficile* is present in the food of hospitalized patients and to estimate the risk of subsequent colonization associated with *C. difficile* in food.

METHODS. This was a prospective cohort study of inpatients at a university-affiliated tertiary care center, May 9, 2011–July 12, 2012. Enrolled patients submitted a portion of food from each meal. Patient stool specimens and/or rectal swabs were collected at enrollment, every 3 days thereafter, and at discharge, and were cultured for *C. difficile*. Clinical data were reviewed for evidence of infection due to *C. difficile*. A stochastic, discrete event model was developed to predict exposure to *C. difficile* from food, and the estimated number of new colonization events from food exposures per 1,000 admissions was determined.

RESULTS. A total of 149 patients were enrolled and 910 food specimens were obtained. Two food specimens from 2 patients were positive for *C. difficile* (0.2% of food samples; 1.3% of patients). Neither of the 2 patients was colonized at baseline with *C. difficile*. Discharge colonization status was available for 1 of the 2 patients and was negative. Neither was diagnosed with *C. difficile* infection while hospitalized or during the year before or after study enrollment. Stochastic modeling indicated contaminated hospital food would be responsible for less than 1 newly colonized patient per 1,000 hospital admissions.

CONCLUSIONS. The recovery of *C. difficile* from the food of hospitalized patients was rare. Modeling suggests hospital food is unlikely to be a source of *C. difficile* acquisition.

Infect Control Hosp Epidemiol 2016;37:1401–1407

Clostridium difficile infection (CDI) is the most common healthcare-associated infection in the United States and is associated with significant patient morbidity, mortality, and high attributable acute care hospital costs.^{1–3} Given the continued high incidence and severity of clinical outcomes associated with CDI, measures to prevent CDI are an area of ongoing interest. Current strategies for CDI prevention are focused on interrupting the cycle of transmission from individuals with CDI; however, it is important to evaluate other potential modes for *C. difficile* acquisition.

Although CDI is primarily associated with healthcare facilities, the precise source of *C. difficile* exposure is unknown. Recent studies have found that only 15%–25% of CDI cases could be attributed to ward-based or patient-to-patient transmission, indicating that there may be other sources of *C. difficile* acquisition in the hospital.^{4,5} A potential reservoir and source for *C. difficile* acquisition is the food of hospitalized patients. *C. difficile* has been isolated from retail foods worldwide, including ground meats, poultry, and vegetables.^{6–17} The spores

of *C. difficile* are heat-resistant and thus may have the potential to survive cooking temperatures.¹⁰ Given the presence of *C. difficile* in food and its heat-resistant qualities, it is theoretically possible that hospitalized patients could be exposed to *C. difficile* from their food. We conducted a prospective cohort study with the objectives of determining whether *C. difficile* was present in the food of hospitalized patients and of estimating the risk of colonization associated with the presence of *C. difficile* in the food of hospitalized patients.

METHODS

Setting

This prospective cohort study was conducted at Barnes-Jewish Hospital (BJH), a 1,250-bed tertiary care center in St. Louis, MO, from May 9, 2011, through July 12, 2012, in conjunction with a study of *C. difficile* colonization in hospitalized patients.^{18,19} The study was approved by the Washington University Human Research Protection Office.

Affiliations: 1. Division of Infectious Diseases, Washington University School of Medicine, St. Louis, Missouri; 2. Department of Population Health and Pathobiology, North Carolina State University, Raleigh, North Carolina; 3. Departments of Pathology & Immunology, Molecular Microbiology, and Pediatrics, Washington University School of Medicine, St. Louis, Missouri.

Received June 15, 2016; accepted August 25, 2016; electronically published October 3, 2016

© 2016 by The Society for Healthcare Epidemiology of America. All rights reserved. 0899-823X/2016/3712-0002. DOI: 10.1017/ice.2016.218

Subjects

Subjects at least 18 years old admitted to the medical and surgical wards with a projected length of stay (LOS) of at least 3 days and no diarrhea were invited to participate; all provided written, informed consent.

Data Sources and Statistical Analyses

Data collected included demographic characteristics, comorbidities, and CDI diagnoses from 1 year prior to enrollment to 1 year after enrollment. Data sources included patient interviews, medical chart review, and Medical Informatics queries. Data analyses were descriptive. SPSS, version 21 (IBM), was used. The model was implemented in NetLogo, version 5.1. R (R Foundation for Statistical Computing) was used for model parameterization and output analysis.

Specimen Collection

Stool or rectal swab specimens were collected from patients upon study enrollment, every 3 days, and at discharge. Rectal swab samples (ESwab; Becton, Dickinson) were obtained from patients unable to provide a stool specimen within 48 hours of admission or 24 hours of a postadmission specimen collection time.^{20–22}

Each patient was provided with a cooler and 4 sterile specimen cups labeled breakfast, lunch, supper, or snack. Patients were instructed to place a piece of food from everything they ate into the corresponding container. If they did not eat a particular meal, no food was collected. As patients placed all components of their meal into the same container, there were multiple types of food per container.

Food specimens were transported to the laboratory and frozen at -30°C . Prior to culture, food specimens were thawed and the food types were documented. The food specimen was combined with 10 mL of sterile water and homogenized for approximately 1 minute.

Microbiological Analysis

Next, 1 mL of food homogenate was added to cycloserine-cefoxitin mannitol broth with taurocholate, lysozyme, and cysteine (CCMB-TAL; Anaerobe Systems), and the broth was subcultured to pre-reduced blood agar (Becton, Dickinson) as previously described.²⁰ *C. difficile* in food was quantified by weighing initial specimens, processing via heat shock, plating onto a pre-reduced blood agar plate, then streaking for isolation. Colonies per gram of food were calculated. Additionally, food was diluted in CCMB-TAL broth in a series of five 10-fold dilutions to approximate the burden of *C. difficile*. *C. difficile* negative and positive controls were included with every set of cultures to monitor for contamination. Ribotyping was performed on all *C. difficile* isolates as previously described.²³

Stool and rectal swabs were cultured for *C. difficile* using CCMB-TAL according to methods previously published.²⁰ Isolates were evaluated for the presence of *tcdA*, *tcdB*, and

binary toxin genes (*cdtA/cdtB*) by multiplex polymerase chain reaction as previously described.¹⁸ Isolates were also characterized by polymerase chain reaction ribotyping for strain comparison.²³

Model Overview

To estimate the risk of *C. difficile* acquisition associated with exposure to *C. difficile*-contaminated food during a hospital stay, we developed a stochastic, individual-based model that simulated the flow of patients admitted to BJH, antimicrobial exposures, number of meals eaten per day, and concentration of *C. difficile* in food (Figure 1). A formal description of the model, code, and parameters is available at <http://www.lanzaslab.org/research/cdifficile#food>. The hospital model simulated the 171 BJH general medicine hospital ward beds. Each patient was followed from admission to discharge. On admission, each patient was assigned a LOS. LOS depended on whether the patient received antibiotic treatment based on the distribution of LOS from 11,046 admissions from these wards to several distributions using maximum likelihood methods.²⁴ On the basis of these data, antibiotic use was used as a marker for longer LOS. Antibiotic use was also a marker for susceptibility to *C. difficile* colonization. The log-normal distribution, which provided the best fit, was used to parameterize LOS (Table 1).

The number of meals consumed by a patient daily and the probability that a meal was contaminated were based on this study's results (Table 1). A Poisson log-normal distribution was used to simulate the number of spores per contaminated meal. This distribution is often used to describe microbial counts.²⁵ The parameters of the distribution were chosen to generate a mean number of spores of approximately 10 colony-forming units/gm because this was the limit of detection of the culture methods. Data from a clinical trial in which healthy adults received escalating doses of nontoxigenic *C. difficile* spores were used to estimate the probability of *C. difficile* colonization upon dose exposure.²⁶ We used logistic regression to model the data from study subjects in cohort




FIGURE 1. Exposure assessment in study of *Clostridium difficile* in the food of hospitalized patients. *Cumulative dose calculated by total number of spores present in all contaminated meals eaten in a hospital stay.

4 who received 5 days of pretreatment with oral vancomycin before receiving a daily dose of 1×10^4 , 1×10^6 , or 1×10^8 spores for 14 days. Because of the repeated measurements on the same subjects, the binary correlated data were analyzed by means of the generalized estimating equation²⁷ as implemented in package *geepack* in R.²⁸

The model was iterated 5,000 times to assure output convergence. Each iteration simulated the patients' admissions to the wards for 1 year. In each iteration, model inputs described as probability distributions were sampled and fed to the model. The model outcomes were the number of patients exposed to *C. difficile* through food and the number of colonization events due to food exposure per 1,000 admissions.

RESULTS

Enrollment and Demographic Characteristics

A total of 149 patients were enrolled, and food specimens from 910 meals were obtained and cultured for *C. difficile*. Patient characteristics are in Table 2. Most patients had healthcare exposures within the previous 90 days (136 [91%]), but only 2 patients had a history of CDI within the previous year (none within 60 days prior to enrollment).

Food Cultures

Toxigenic *C. difficile* was recovered from 2 food specimens from 2 separate patients, representing 0.2% of food cultures and 1.3% of patients (Table 3). The food items that tested positive were a gelatin dessert (ribotype 001) and a sample consisting of vegetables/bread/grains (ribotype 027). The concentration of *C. difficile* spores recovered from the positive food samples was less than or equal to 10 colony-forming units/mL. *C. difficile* was successfully recovered from all positive controls, and there was no growth in any of the negative controls.

C. difficile Colonization and CDI

Neither of the 2 patients exposed to *C. difficile* in food was colonized at baseline with *C. difficile*. A discharge stool specimen was available for 1 of the 2 patients and was negative. Neither was diagnosed with CDI during their hospitalization or during the year before or after study enrollment. No patients in the study developed CDI within a year of discharge.

Exposure Modeling

A summary of the functions and probability distributions for the exposure model are detailed in Table 1. On 44.1% of days, no meals were eaten; on 17.5% of days 1 meal was eaten; on 26.4% of days 2 meals were eaten; on 11.2% of days 3 meals were eaten; and on 0.8% of days 4 meals were eaten. Reasons for missing meals were variable but included instructions to take nothing by mouth in preparation for an upcoming procedure(s) or lack of appetite. The mean number of patients who were exposed to *C. difficile* through food was 12.70 per 1,000 admissions (95% CI, 12.542–12.858). The minimum and maximum simulated values were 2.34 and 25.85 exposed patients per 1,000 admissions, respectively (Figure 2). The mean number of predicted colonization events was 0.609 per 1,000 admissions (95% CI, 0.600–0.618), and the median number was 0.57. The minimum and maximum simulated colonization events were 0.04 and 1.73 per 1,000 admissions, respectively (Figure 2).

Both the predicted number of exposed and the predicted number of colonized patients were highly influenced by the probability of meal contamination (Figure 3). A 0.1% increase in the probability of meal contamination resulted in an increase of 5.5 exposures and 0.26 colonization events per 1,000 admissions (Figure 3). Overall, the simulated number of spores in contaminated food was low, reflecting the low counts recovered from hospital food. As a result, on average fewer than 5% of the patients exposed to *C. difficile* became colonized. Antibiotic prescription probability had a marginal effect

TABLE 1. Summary of Functions and Probability Distributions for the Exposure Model

| Description | Functions and probability distributions |
|---|--|
| Antibiotic prescription probability | Uniform (0.3, 0.7) |
| Distribution of length of stay for patients who do not receive antibiotic | Log-normal (0.84, 0.77) |
| Distribution of length of stay for patients who receive antibiotic | Log-normal (1.54, 0.87) |
| N of meals (X) received by a patient daily | $P(X=0) = 0.441$, $P(X=1) = 0.175$, $P(X=2) = 0.264$, $P(X=3) = 0.112$, $P(X=4) = 0.008$ |
| Probability that a meal is contaminated with <i>Clostridium difficile</i> spores | Uniform (0.0005, 0.0035) |
| Distribution of the number of spores found on a contaminated meal | Poisson-lognormal (2, 0.8) |
| Relationship between cumulative dose in a hospital stay and probability of colonization | $\frac{e^{-3.073 + 0.539 \log(dose + 1)}}{1 + e^{-3.073 + 0.539 \log(dose + 1)}}$ |

NOTE. The parameters needed to characterize the probability distributions are indicated between brackets. For the uniform distribution (a,b): a is the minimum value and b is the maximum value. For the log-normal and Poisson-lognormal distributions (μ , σ): μ is the log mean and σ is the log standard deviation.

TABLE 2. Characteristics of 149 Patients in Study of *Clostridium difficile* in the Food of Hospitalized Patients

| Variable | Value |
|--|---------------|
| Age, median (range), y | 55 (23–90) |
| Length of stay, median (range), d | 4.0 (0.4–292) |
| Female sex | 80 (54) |
| Nonwhite race | 43 (29) |
| Healthcare worker | 20 (13) |
| Lives with a healthcare worker | 19 (13) |
| Spends ≥2 hours/week visiting a healthcare facility | 16 (11) |
| Admitted to medicine service | 145 (97) |
| Admitted from | |
| Home | 99 (66) |
| Healthcare facility | 50 (34) |
| Reason for admission | |
| Infection | 45 (30) |
| Exacerbation of chronic condition | 54 (36) |
| Elective surgery | 2 (1) |
| New medical or surgical problem | 48 (32) |
| Any healthcare exposures in previous 90 days | 136 (91) |
| Diabetes mellitus | 58 (39) |
| Congestive heart failure | 27 (18) |
| Liver disease | 28 (19) |
| Chronic renal insufficiency | 17 (11) |
| Chronic lung disease | 25 (17) |
| Human immunodeficiency virus | 3 (2) |
| Solid organ transplant | 8 (5) |
| Stem cell transplant | 0 |
| Solid malignant tumor | 24 (16) |
| Hematologic malignant tumor | 2 (1) |
| Other immunocompromised | 20 (13) |
| Inflammatory bowel disease | 5 (3) |
| Surgery in previous 90 days | 15 (10) |
| Upper endoscopy performed during hospitalization | 9 (6) |
| Lower endoscopy performed during hospitalization | 8 (5) |
| History of CDI in the year before enrollment | 2 (1) |
| CDI diagnosis within 1 year after enrollment | 0 |
| Colonized with <i>C. difficile</i> on admission | 34 (23) |
| Colonized with <i>C. difficile</i> on discharge ^a | 35/141 (25) |

NOTE. Data are no. (%) of patients unless otherwise indicated. CDI, *C. difficile* infection.

^aDischarge colonization status was unknown for 8 patients (5%).

on the number of predicted exposed and colonized patients compared with probability of meal contamination (Figure 3).

DISCUSSION

In this study of *C. difficile* in the food of hospitalized patients, recovery of toxigenic *C. difficile* was rare, with only 0.2% of food specimens testing positive for *C. difficile* with a low estimated concentration (≤10 colony-forming units/mL). Stated differently, 1.5% of patients ingested food from which *C. difficile* was recovered. On the basis of this finding,

TABLE 3. Types of Food Positive for *Clostridium difficile*, by Food Type, for 910 Meals

| Food item | Total | <i>C. difficile</i> , n (%) |
|--------------------|-------|-----------------------------|
| Meat | 308 | 0 |
| Poultry | 142 | 0 |
| Fruit | 179 | 0 |
| Vegetables | 455 | 1 (<1) ^a |
| Nuts | 1 | 0 |
| Dairy/eggs | 210 | 0 |
| Bread/grains | 376 | 1 (<1) ^a |
| Other ^b | 200 | 1 (1) ^c |

NOTE. Percentages are percent of positive samples / all food items of that type. As patients placed all components of their meal into the same container, there were multiple types of food per container.

^aThe positive specimens for vegetables and bread/grains were combined in a single specimen cup.

^bFor example, veggie burger, sauce/gravy, pudding, jelly, fish, cake.

^cGelatin dessert.




FIGURE 2. Histograms for the simulated number of patients exposed to *Clostridium difficile* in food and colonization events due to exposure to *C. difficile* spores in food. Counts are the number of exposed and colonized patients in each histogram bar.

theoretically hundreds of hospitalized patients could be exposed to *C. difficile* from food and develop CDI every year at BJH, which in 2014 alone had more than 53,300 inpatient admissions. Thus, our objective was to model the likelihood of *C. difficile* acquisition from food in the hospital setting. Using a similar modeling framework based on BJH data, we previously predicted that on average there were approximately 100 new colonization events per 1,000 admissions.²⁹ In this study, we found that at less than 1 new colonization event per 1,000 admissions, *C. difficile* acquisition linked to contaminated food was likely uncommon. The results of the modeling indicate




FIGURE 3. Scatterplots between the hospital-level variables (antibiotic prescription and meal contamination probability) and model outcomes (number of exposed and colonized patients per 1,000 admissions). The points indicate individual simulations (total number of simulations = 5,000) and the line indicates the linear trend between variables.

that acquisition of *C. difficile* from food is likely a rare event at BJH and that the food of hospitalized patients was not a significant source of new *C. difficile* colonization.

Most previously published studies of *C. difficile* in food have focused on retail meat products, with prevalence rates reported in Europe and Canada ranging from 2.7% in chicken and 4.3%–20% in beef/pork, and with rates in US studies greater than 40% (ground meats).^{8,10–12} The presence of *C. difficile* in foods of non-animal origin (eg, fruits, vegetables, grains) has not been fully explored, with a Scottish study indicating that 7.5% of salads and a French study reporting 2.9% of raw vegetables were contaminated with *C. difficile*.^{6–8} These previous studies were based on singular food types, rather than a mixture of foods that constitute a meal that a patient might eat. In our study, we included all food that the patient would be consuming during that meal, which would better represent a hospitalized patient's actual *C. difficile* exposure.

Although we were able to recover *C. difficile* from the food of hospitalized patients, this does not equate directly to *C. difficile* being a foodborne pathogen in the healthcare setting. The source of contamination is not known; contamination may have occurred at the food source (farm, factory), food handler, food transporter, and/or from the patient handling

the food. The results of our study are consistent with those of Rodriguez et al,¹³ who found *C. difficile* in less than 1% of food samples collected from the kitchens of a Belgian nursing home. *C. difficile* was isolated from only one meal sample composed of pork sausage, mustard sauce, and carrot salad.¹³ Together, our study and the Rodriguez study suggest that *C. difficile* is present in hospital foods but at lower rates compared with retail foods. The reason for this discrepancy is unclear. Although *C. difficile* spores can survive cooking temperatures, it is possible that soaking, washing, and/or cooking food reduces the *C. difficile* burden and may have accounted for this difference.¹⁰

This study had limitations: it was a single-center study, and results may not be generalizable to all institutions. Regarding specimen collection, patients placed food into containers for culture, potentially introducing variability. However, this provided a practical method of obtaining food samples that were actually consumed by the patient. There are no data available to indicate whether or not *C. difficile* is evenly distributed in food. Thus the amount of food included in each food culture may have impacted our findings, especially in the setting of a low contamination burden. In previously published studies the amount of food cultured varies widely,

from 1 g samples to complete pieces of meat.^{14,17} Previous studies were limited to specific food types; however, this study focused on the variety of foods that a patient ingests per meal, providing a more realistic estimate of a patient's entire meal.

A strength of our study was the collection of food that was actually consumed, rather than a single food type. In the studies of retail food products, the food would have likely been washed and/or cooked prior to consumption; therefore the prevalence of *C. difficile* detected may not represent what individuals would have consumed. Additional strengths include the collection of clinical data to examine CDI in the year after enrollment and culture of food specimens along with the culture of stool specimens throughout the patients' hospitalizations. This allowed us to link *C. difficile* food contamination with acquisition. Our laboratory standards were rigorous, and we included both positive and negative controls to ensure against laboratory contamination.

C. difficile is a ubiquitous organism, and only a minority of new *C. difficile* acquisitions in the hospital have been linked to another patient with CDI.^{4,5,30,31} Therefore, understanding all potential sources of *C. difficile* exposure in the hospital is necessary to inform prevention measures for CDI. Our findings indicate that food is unlikely to be a significant source of *C. difficile* acquisition in hospitalized patients. Towards the goal of CDI prevention, future studies aimed at understanding modes of *C. difficile* transmission and acquisition are necessary.

ACKNOWLEDGMENTS

Financial support. National Institute of Diabetes and Digestive and Kidney Diseases (grant P30DK52574); the Washington University Institute of Clinical and Translational Sciences (grant UL1TR000448, sub-award KL2TR000450, from the National Center for Advancing Translational Sciences of the National Institutes of Health, to J.H.K.); and the joint National Science Foundation/National Institute of General Medical Sciences Mathematical Biology Program (grant R01GM113239 to C.L.).

Potential conflicts of interest. C.-A.D.B. reports that she has received research support from bioMérieux, Cepheid, and Accelerate Diagnostics. E.R.D. reports that he has been a consultant to Sanofi Pasteur, Merck, Summit, Alere, Valenza, and Rebiotix. All other authors report no conflicts of interest relevant to this article.

Disclaimer: The content is solely the responsibility of the authors and does not necessarily represent the official view of the National Institutes of Health.

Address correspondence to Jennie H. Kwon, MSCI, Washington University School of Medicine, Division of Infectious Diseases, 660 S Euclid Ave, Campus Box 8051, St. Louis, MO 63110 (JKwon@wustl.edu).

REFERENCES

- Centers for Disease Control and Prevention (CDC). Antibiotic resistance threats in the United States, 2013. CDC website. <http://www.cdc.gov/drugresistance/threat-report-2013/>. Published 2014. Accessed January 9, 2016.
- Lessa FC, Mu Y, Bamberg WM, et al. Burden of *Clostridium difficile* infection in the United States. *N Engl J Med* 2015; 372:825–834.
- Kwon JH, Olsen MA, Dubberke ER. The morbidity, mortality, and costs associated with *Clostridium difficile* infection. *Infect Dis Clin North Am* 2015;29:123–134.
- Walker AS, Eyre DW, Wyllie DH, et al. Characterisation of *Clostridium difficile* hospital ward-based transmission using extensive epidemiological data and molecular typing. *PLOS Med* 2012;9:e1001172.
- Grundmann H, Barwolff S, Tami A, et al. How many infections are caused by patient-to-patient transmission in intensive care units? *Crit Care Med* 2005;33:946–951.
- Bakri MM, Brown DJ, Butcher JP, Sutherland AD. *Clostridium difficile* in ready-to-eat salads, Scotland. *Emerg Infect Dis* 2009;15:817–818.
- Eckert C, Burghoffer B, Barbut F. Contamination of ready-to-eat raw vegetables with *Clostridium difficile* in France. *J Med Microbiol* 2013;62:1435–1438.
- Gould LH, Limbago B. *Clostridium difficile* in food and domestic animals: a new foodborne pathogen? *Clin Infect Dis* 2010;51: 577–582.
- Jobstl M, Heuberger S, Indra A, Nepf R, Kofer J, Wagner M. *Clostridium difficile* in raw products of animal origin. *Int J Food Microbiol* 2010;138:172–175.
- Lund BM, Peck MW. A possible route for foodborne transmission of *Clostridium difficile*? *Foodborne Pathog Dis* 2015;12: 177–182.
- Rodriguez-Palacios A, Staempfli HR, Duffield T, Weese JS. *Clostridium difficile* in retail ground meat, Canada. *Emerg Infect Dis* 2007;13:485–487.
- Rodriguez-Palacios A, Reid-Smith RJ, Staempfli HR, et al. Possible seasonality of *Clostridium difficile* in retail meat, Canada. *Emerg Infect Dis* 2009;15:802–805.
- Rodriguez C, Korsak N, Taminiau B, et al. *Clostridium difficile* from food and surface samples in a Belgian nursing home: an unlikely source of contamination. *Anaerobe* 2015;32:87–89.
- Songer JG, Trinh HT, Killgore GE, Thompson AD, McDonald LC, Limbago BM. *Clostridium difficile* in retail meat products, USA, 2007. *Emerg Infect Dis* 2009;15:819–821.
- Von Abercron SM, Karlsson F, Wigh GT, Wierup M, Krovacek K. Low occurrence of *Clostridium difficile* in retail ground meat in Sweden. *J Food Prot* 2009;72:1732–1734.
- Weese JS, Avery BP, Rousseau J, Reid-Smith RJ. Detection and enumeration of *Clostridium difficile* spores in retail beef and pork. *Appl Environ Microbiol* 2009;75:5009–5011.
- Weese JS, Reid-Smith RJ, Avery BP, Rousseau J. Detection and characterization of *Clostridium difficile* in retail chicken. *Lett Appl Microbiol* 2010;50:362–365.
- Alasmari F, Seiler SM, Hink T, Burnham CA, Dubberke ER. Prevalence and risk factors for asymptomatic *Clostridium difficile* carriage. *Clin Infect Dis* 2014;59:216–222.
- Dubberke ER, Reske KA, Seiler S, Hink T, Kwon JH, Burnham CA. Risk factors for acquisition and loss of *C. difficile* colonization in hospitalized patients. *Antimicrob Agents Chemother* 2015;59:4533–4543.
- Hink T, Burnham CA, Dubberke ER. A systematic evaluation of methods to optimize culture-based recovery of *Clostridium difficile* from stool specimens. *Anaerobe* 2013;19:39–43.
- McFarland LV, Coyle MB, Kremer WH, Stamm WE. Rectal swab cultures for *Clostridium difficile* surveillance studies. *J Clin Microbiol* 1987;25:2241–2242.

22. Kundrapu S, Sunkesula VC, Jury LA, Sethi AK, Donskey CJ. Utility of perirectal swab specimens for diagnosis of *Clostridium difficile* infection. *Clin Infect Dis* 2012;55:1527–1530.
23. Westblade LF, Chamberland RR, Maccannell D, et al. Development and evaluation of a novel, semiautomated *Clostridium difficile* typing platform. *J Clin Microbiol* 2013;51:621–624.
24. Delignette-Muller ML, Dutang C. fitdistrplus: an R package for fitting distributions. *J Stat Softw* 2015;64.
25. Gonzales-Barron U, Butler F. A comparison between the discrete Poisson-gamma and Poisson-lognormal distributions to characterise microbial counts in foods. *Food Control* 2011;22:1279–1286.
26. Villano SA, Seiberling M, Tatarowicz W, Monnot-Chase E, Gerding DN. Evaluation of an oral suspension of VP20621, spores of nontoxigenic *Clostridium difficile* strain M3, in healthy subjects. *Antimicrob Agents Chemother* 2012;56:5224–5229.
27. Hosmer DW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*. 3rd ed. Hoboken, NJ: John Wiley & Sons; 2013.
28. Halekoh U, Hojsgaard S, Yan J. The R package geepack for generalized estimating equations. *J Stat Softw* 2016;15.
29. Lanzas C, Dubberke ER. Effectiveness of screening hospital admissions to detect asymptomatic carriers of *Clostridium difficile*: a modeling evaluation. *Infect Control Hosp Epidemiol* 2014;35:1043–1050.
30. Fekety R, Kim KH, Brown D, Batts DH, Cudmore M, Silva J Jr. Epidemiology of antibiotic-associated colitis; isolation of *Clostridium difficile* from the hospital environment. *Am J Med* 1981;70:906–908.
31. McFarland LV, Mulligan ME, Kwok RY, Stamm WE. Nosocomial acquisition of *Clostridium difficile* infection. *N Engl J Med* 1989;320:204–210.