

## ISOMETRIES ON EXTREMELY NON-COMPLEX BANACH SPACES

PIOTR KOSZMIDER<sup>1</sup>, MIGUEL MARTÍN<sup>2</sup> AND JAVIER MERÍ<sup>2</sup>

<sup>1</sup>*Instytut Matematyki Politechniki Łódzkiej, ul. Wólczańska 215,  
90-924 Łódź, Poland (pkoszmider.politechnika@gmail.com)*

<sup>2</sup>*Departamento de Análisis Matemático, Facultad de Ciencias,  
Universidad de Granada, 18071 Granada, Spain  
(mmartins@ugr.es; jmeri@ugr.es)*

(Received 14 April 2009; revised 18 January 2010; accepted 18 January 2010)

*Abstract* Given a separable Banach space  $E$ , we construct an extremely non-complex Banach space (i.e. a space satisfying that  $\|\text{Id} + T^2\| = 1 + \|T^2\|$  for every bounded linear operator  $T$  on it) whose dual contains  $E^*$  as an  $L$ -summand. We also study surjective isometries on extremely non-complex Banach spaces and construct an example of a real Banach space whose group of surjective isometries reduces to  $\pm\text{Id}$ , but the group of surjective isometries of its dual contains the group of isometries of a separable infinite-dimensional Hilbert space as a subgroup.

*Keywords:* few operators; complex structure; Daugavet equation; Banach spaces of continuous functions; extremely non-complex; surjective isometries

AMS 2010 *Mathematics subject classification:* Primary 46B04

Secondary 46B10; 46B20; 46E15; 47A99; 54G99

### 1. Introduction

All the Banach spaces in this paper will be real. Given a Banach space  $X$ , we write  $X^*$  for the topological dual,  $L(X)$  for the space of all bounded linear operators,  $W(X)$  for the space of weakly compact operators and  $\text{Iso}(X)$  for the group of surjective isometries.

A Banach space  $X$  is said to be *extremely non-complex* if the norm equality

$$\|\text{Id} + T^2\| = 1 + \|T^2\|$$

holds for every  $T \in L(X)$ . This concept was introduced very recently by the authors in [20], where several different examples of  $C(K)$  spaces are shown to be extremely non-complex. For instance, this is the case for some perfect compact spaces  $K$  constructed by the first author [19] such that  $C(K)$  has few operators (in the sense that every operator is a weak multiplier). There are other examples of extremely non-complex  $C(K)$  spaces which contain complemented copies of  $\ell_\infty$  or  $C(2^\omega)$  (and so, they do not have few operators). It is trivial that  $X = \mathbb{R}$  is extremely non-complex. The existence of infinite-dimensional extremely non-complex Banach spaces had been asked in [16, Question 4.11],

where possible generalizations of the Daugavet equation were investigated. We recall that an operator  $S$  defined on a Banach space  $X$  satisfies the *Daugavet equation* [4] if

$$\|\text{Id} + S\| = 1 + \|S\|$$

and that the space  $X$  has the *Daugavet property* [17] if the Daugavet equation holds for every rank-1 operator on  $X$ . We refer the reader to [1, 2, 17, 27] for background on the Daugavet property. Let us observe that  $X$  is extremely non-complex if the Daugavet equation holds for the square of every (bounded linear) operator on  $X$ . Spaces  $X$  in which the square of every rank-1 operator on  $X$  satisfies the Daugavet equation are studied in [23], where it is shown that their unit balls do not have strongly exposed points. In particular, the unit ball of an extremely non-complex Banach space (of dimension greater than one) does not have strongly exposed points and, therefore, the space does not have the Radon–Nikodým property, even the more it is not reflexive.

The name of extremely non-complex comes from the fact that a real Banach space  $X$  is said to have a *complex structure* if there exists  $T \in L(X)$  such that  $T^2 = -\text{Id}$ , so extremely non-complex spaces lack of complex structures in a very strong way. Let us also comment that no hyperplane (actually no finite-codimensional subspace) of an extremely non-complex Banach space admits a complex structure. The existence of infinite-dimensional real Banach spaces admitting no complex structure is known since the 1950s, when Dieudonné [6] showed that this is the case of the James space  $\mathcal{J}$ . We refer the reader to the very recent papers by Ferenczi [10] and Ferenczi and Medina Galego [11] and references therein for a discussion about complex structures on spaces and on their hyperplanes.

Our first goal in this paper is to present examples of extremely non-complex Banach spaces which are not isomorphic to  $C(K)$  spaces. Namely, it is proved in §3 that if  $K$  is a compact space such that all operators on  $C(K)$  are weak multipliers,  $L$  is a closed nowhere-dense subset of  $K$  and  $E$  is a subspace of  $C(L)$ , then the space

$$C_E(K\|L) = \{f \in C(K) : f|_L \in E\}$$

is extremely non-complex (see §2 for the definitions and basic facts about this kind of spaces). It is also shown that there are extremely non-complex  $C_E(K\|L)$  spaces which are not isomorphic to  $C(K')$  spaces. On the other hand, some spaces  $C_E(K\|L)$  are isometric to spaces  $C(K')$  for some compact  $K'$ . This can be used to note that the extremely non-complex spaces of the form  $C_E(K\|L)$  may have many operators besides weak multipliers (see Remark 3.14).

The next aim is to show that  $\text{Iso}(X)$  is a ‘discrete’ Boolean group when  $X$  is extremely non-complex. Namely, we show that  $T^2 = \text{Id}$  for every  $T \in \text{Iso}(X)$  (i.e.  $\text{Iso}(X)$  is a Boolean group and is therefore commutative) and that  $\|T_1 - T_2\| \in \{0, 2\}$  for every  $T_1, T_2 \in \text{Iso}(X)$ . Next, we discuss the relationship with the set of all unconditional projections on  $X$  and the possibility of this set to be a Boolean algebra. This is the content of §4. In §5 we particularize these results to spaces  $C_E(K\|L)$  which are extremely non-complex, getting necessary conditions on the elements of  $\text{Iso}(C_E(K\|L))$ . In particular, if  $C(K)$  is extremely non-complex, we show that the only homeomorphism from  $K$  to  $K$

is the identity, obtaining that  $\text{Iso}(C(K))$  is isomorphic to the Boolean algebra of clopen sets of  $K$ .

Section 6 is devoted to apply all the results above to get an example showing that the behaviour of the group of isometries with respect to duality can be extremely bad. Namely, we show that for every separable Banach space  $E$ , there is a Banach space  $\tilde{X}(E)$  such that  $\text{Iso}(\tilde{X}(E)) = \{\text{Id}, -\text{Id}\}$  and  $\tilde{X}(E)^* = E^* \oplus_1 Z$ , so  $\text{Iso}(\tilde{X}(E)^*)$  contains  $\text{Iso}(E^*)$  as a subgroup. To do so, we have to modify a construction of a connected compact space  $K$  with few operators given by the first named author in [19, §5] and use our construction of  $C_E(K||L)$  for a nowhere dense  $L \subset K$ . For the special case  $E = \ell_2$ , we have  $\text{Iso}(\tilde{X}(\ell_2)) = \{\text{Id}, -\text{Id}\}$ , while  $\text{Iso}(\tilde{X}(\ell_2)^*)$  contains infinitely many uniformly continuous one-parameter semigroups of surjective isometries. Let us comment that, in sharp contrast with the examples above, when a Banach space  $X$  is *strongly unique predual*, the group  $\text{Iso}(X^*)$  consists exactly of the conjugate operators to the elements of  $\text{Iso}(X)$ . Quite a lot of spaces are actually strong unique preduals. We refer the reader to [14] for more information.

We finish this introduction with some needed notation. If  $X$  is a Banach space, we write  $B_X$  to denote the closed unit ball of  $X$  and given a convex subset  $A \subseteq X$ ,  $\text{ext}(A)$  denotes the set of extreme points of  $A$ . A closed subspace  $Z$  of  $X$  is an  $L$ -summand if  $X = Z \oplus_1 W$  for some closed subspace  $W$  of  $X$ , where  $\oplus_1$  denotes the  $\ell_1$ -sum. A closed subspace  $Y$  of a Banach space  $X$  is said to be an  $M$ -ideal of  $X$  if the annihilator  $Y^\perp$  of  $Y$  is an  $L$ -summand of  $X^*$ . We refer the reader to [15] for background on  $L$ -summands and  $M$ -ideals.

**2. Notation and preliminary results on the spaces  $C_E(K||L)$**

Throughout the paper,  $K$  will be a (Hausdorff) compact (topological) space and  $L \subseteq K$  will stand for a nowhere-dense closed subset. Given a closed subspace  $E$  of  $C(L)$ , we will consider the subspace of  $C(K)$  given by

$$C_E(K||L) = \{f \in C(K) : f|_L \in E\}.$$

This notation is compatible with the Semadeni’s book [25, II.4] notation of

$$C_0(K||L) = \{f \in C(K) : f|_L = 0\}.$$

This last space can be identified with the space  $C_0(K \setminus L)$  of those continuous functions  $f : K \setminus L \rightarrow \mathbb{R}$  vanishing at infinity.

By the Riesz representation theorem, the dual space of  $C(K)$  is isometric to the space  $M(K)$  of Radon measures on  $K$ , i.e. signed, Borel, scalar-valued, countably additive and regular measures. More precisely, given  $\mu \in M(K)$  and  $f \in C(K)$ , the duality is given by

$$\mu(f) = \int f \, d\mu.$$

We recall that  $C_0(K||L)$  is an  $M$ -ideal of  $C(K)$  [15, Example I.1.4(a)], meaning that  $C_0(K||L)^\perp$  is an  $L$ -summand in  $C(K)^*$ . This fact allows to show the following well-known result.

**Lemma 2.1.**  $C_0(K\|L)^* \equiv \{\mu \in M(K) : |\mu|(L) = 0\}$ .

**Proof.** Since  $C_0(K\|L)$  is an  $M$ -ideal in  $C(K)$ , Proposition 1.12 and Remark 1.13 of [15] allow us to identify  $C_0(K\|L)^*$  with the subspace of  $C(K)^* = M(K)$  given by

$$C_0(K\|L)^\# = \{\mu \in M(K) : |\mu|(K) = |\mu|(K \setminus L)\} = \{\mu \in M(K) : |\mu|(L) = 0\}.$$

□

When we consider the space  $C_E(K\|L)$ , it still makes sense to talk about functionals corresponding to the measures on  $K$ , namely one understands them as the restriction of the functional from  $C(K)$  to  $C_E(K\|L)$ . However, given a functional belonging to  $C_E(K\|L)^*$  one may have several measures on  $K$  associated with it. The next result describes the dual of a  $C_E(K\|L)$  space for an arbitrary  $E \subseteq C(L)$ . It is worth mentioning that its proof is an extension of that appearing in [21, Theorem 3.3].

**Lemma 2.2.** *Let  $K$  be a compact space,  $L$  a closed subset of  $K$  and  $E \subseteq C(L)$ . Then,*

$$C_E(K\|L)^* \equiv C_0(K\|L)^* \oplus_1 C_0(K\|L)^\perp \equiv C_0(K\|L)^* \oplus_1 E^*.$$

**Proof.** We write  $P : C(K) \rightarrow C(L)$  for the restriction operator, i.e.

$$[P(f)](t) = f(t) \quad (t \in L, f \in C(K)).$$

Then,  $C_0(K\|L) = \ker P$  and  $C_E(K\|L) = \{f \in C(K) : P(f) \in E\}$ . Since  $C_0(K\|L)$  is an  $M$ -ideal in  $C(K)$ , it is *a fortiori* an  $M$ -ideal in  $C_E(K\|L)$  by [15, Proposition I.1.17], meaning that

$$C_E(K\|L)^* \equiv C_0(K\|L)^* \oplus_1 C_0(K\|L)^\perp \equiv C_0(K\|L)^* \oplus_1 [C_E(K\|L)/C_0(K\|L)]^*.$$

Now, it suffices to prove that the quotient  $C_E(K\|L)/C_0(K\|L)$  is isometrically isomorphic to  $E$ . To do so, we define the operator  $\Phi : C_E(K\|L) \rightarrow E$  given by  $\Phi(f) = P(f)$  for every  $f \in C_E(K\|L)$ . Then  $\Phi$  is well defined,  $\|\Phi\| \leq 1$ , and  $\ker \Phi = C_0(K\|L)$ . To see that the canonical quotient operator  $\tilde{\Phi} : C_E(K\|L)/C_0(K\|L) \rightarrow E$  is a surjective isometry, it suffices to show that

$$\Phi(\{f \in C_E(K\|L) : \|f\| < 1\}) = \{g \in E : \|g\| < 1\}.$$

Indeed, the left-hand side is contained in the right-hand side since  $\|\Phi\| \leq 1$ . Conversely, for every  $g \in E \subseteq C(L)$  with  $\|g\| < 1$ , we just use Tietze’s extension theorem to find  $f \in C(K)$  such that  $\Phi(f) = f|_L = g$  and  $\|f\| = \|g\|$ . □

If  $\phi \in C_E(K\|L)^*$ , by the above lemma we have  $\phi = \phi_1 + \phi_E$  with  $\phi_1 \in C_0(K\|L)^*$  and  $\phi_E \in C_0(K\|L)^\perp \equiv E^*$ . Observe that  $\phi_1$  can be isometrically associated with a measure on  $K \setminus L$  by Lemma 2.1, which will be denoted  $\phi|_{K \setminus L}$ . Given a subset  $A \subseteq K$  satisfying  $A \cap L = \emptyset$ ,  $\phi|_A$  will stand for the measure  $\phi|_{K \setminus L}$  restricted to  $A$ .

The next result is a straightforward consequence of the above two lemmas.

**Lemma 2.3.** *Let  $\phi \in C_E(K\|L)^*$  and  $x \in K \setminus L$ . Then*

$$\|\phi\| = \|\phi|_{\{x\}}\| + \|\phi|_{K \setminus (L \cup \{x\})}\| + \|\phi_E\|.$$

The next easy lemma describes the set of extreme points in the unit ball of the dual of  $C_E(K\|L)$  and gives a norming set for  $C_E(K\|L)$ . We recall that a subset  $A$  of the unit ball of the dual of a Banach space  $X$  is said to be *norming* if

$$\|x\| = \sup\{|\phi(x)| : \phi \in A\} \quad (x \in X).$$

**Lemma 2.4.** *Let  $K$  be a compact space,  $L$  a nowhere-dense closed subset and  $E \subseteq C(L)$ . We consider the set*

$$\mathcal{A} = \{\theta\delta_y : y \in K \setminus L, \theta \in \{-1, 1\}\} \subset C_E(K\|L)^*.$$

Then

- (a)  $\text{ext}(B_{C_E(K\|L)^*}) = \mathcal{A} \cup \text{ext}(B_{E^*})$ ,
- (b)  $\mathcal{A}$  is norming for  $C_E(K\|L)$  and
- (c) therefore,  $\mathcal{A}$  is weak\*-dense in  $\text{ext}(B_{C_E(K\|L)^*})$ .

**Proof.** By Lemma 2.2 and the description of the extreme points of the unit ball of an  $\ell_1$ -sum of Banach spaces [15, Lemma I.1.5], we have

$$\text{ext}(B_{C_E(K\|L)^*}) = \text{ext}(B_{C_0(K \setminus L)^*}) \cup \text{ext}(B_{E^*}).$$

It suffices to recall that  $\text{ext}(B_{C_0(K \setminus L)^*}) = \mathcal{A}$  (see [12, Theorem 2.3.5] for example) to get (a). The fact that  $\mathcal{A}$  is norming for  $C_E(K\|L)$  is a direct consequence of the fact that  $K \setminus L$  is dense in  $K$ . Finally, every norming set is weak\*-dense in  $\text{ext}(B_{C_E(K\|L)^*})$  by the Hahn–Banach theorem and the reversed Krein–Milman theorem.  $\square$

We introduce one more ingredient which will play a crucial role in our arguments. Given an operator  $U \in L(C_E(K\|L)^*)$ , we consider the function

$$g_U : K \setminus L \rightarrow [-\|U\|, \|U\|], \quad g_U(x) = U(\delta_x)(\{x\}) \quad (x \in K \setminus L).$$

This obviously extends to operators on  $C_E(K\|L)$  by passing to the adjoint, that is, for  $T \in L(C_E(K\|L))$  one can consider  $g_{T^*} : K \setminus L \rightarrow [-\|T\|, \|T\|]$ . This is a generalization of a tool used in [26] under the name ‘stochastic kernel’. One of the results in that paper can be generalized to the following.

**Lemma 2.5.** *Let  $K$  be a compact space,  $L$  a nowhere-dense closed subset of  $K$ ,  $E \subseteq C(L)$ , and  $T \in L(C_E(K\|L))$ . If the set  $\{x \in K \setminus L : g_{T^*}(x) \geq 0\}$  is dense in  $K \setminus L$ , then  $T$  satisfies the Daugavet equation.*

**Proof.** We use Lemmas 2.2–2.4 to get

$$\begin{aligned}
 \|\text{Id} + T^*\| &\geq \sup_{x \in K \setminus L} \|\delta_x + T^*(\delta_x)\| \\
 &= \sup_{x \in K \setminus L} |1 + T^*(\delta_x)(\{x\})| + \|T^*(\delta_x)|_{(K \setminus (L \cup \{x\}))}\| + \|T^*(\delta_x)|_E\| \\
 &= \sup_{x \in K \setminus L} |1 + T^*(\delta_x)(\{x\})| - |T^*(\delta_x)(\{x\})| + \|T^*(\delta_x)\| \\
 &\geq \sup_{x \in K \setminus L} 1 + T^*(\delta_x)(\{x\}) - |T^*(\delta_x)(\{x\})| + \|T^*(\delta_x)\|. \tag{2.1}
 \end{aligned}$$

Now, we claim that the set  $\{x \in K \setminus L : \|T^*(\delta_x)\| > \|T\| - \varepsilon\}$  is non-empty and open in  $K \setminus L$  for every  $\varepsilon > 0$ . Indeed, take a norm-1 function  $f \in C_E(K\|L)$  such that  $\|T(f)\| > \|T\| - \varepsilon$  and use the fact that  $K \setminus L$  is dense in  $K$  to find  $x \in K \setminus L$  satisfying

$$\|T^*(\delta_x)\| \geq |T^*(\delta_x)(f)| = |T(f)(x)| > \|T\| - \varepsilon.$$

To show that  $\{x \in K \setminus L : \|T^*(\delta_x)\| > \|T\| - \varepsilon\}$  is open we prove that the mapping

$$x \mapsto \|T^*(\delta_x)\| \quad (x \in K \setminus L)$$

is lower semicontinuous. To do so, since  $T^*$  is weak\* continuous and  $\|\cdot\|$  is weak\* lower semicontinuous, it suffices to observe that the mapping which sends  $x$  to  $\delta_x$  is continuous with respect to the weak\* topology on  $C_E(K\|L)^*$ . But this is so since, for  $a \in \mathbb{R}$  and  $f \in C_E(K\|L)$ , the preimage of the sub-basic set  $\{\phi \in C_E(K\|L)^* : \phi(f) < a\}$  in this topology is  $\{x \in K \setminus L : f(x) < a\}$  which is open in  $K \setminus L$ .

To finish the proof, we use the hypothesis to find  $x_0 \in K \setminus L$  satisfying

$$\|T^*(\delta_{x_0})\| > \|T\| - \varepsilon \quad \text{and} \quad g_{T^*}(x_0) = T^*(\delta_{x_0})(\{x_0\}) \geq 0$$

and we use it in (2.1) to obtain

$$\|\text{Id} + T^*\| \geq 1 + \|T^*(\delta_{x_0})\| > 1 + \|T\| - \varepsilon,$$

which implies that  $\|\text{Id} + T\| = \|\text{Id} + T^*\| \geq 1 + \|T\|$  since  $\varepsilon$  was arbitrary. □

### 3. Spaces $C_E(K\|L)$ when $C(K)$ has few operators

Let us start fixing some notation and terminology that will be used throughout the section. If  $g : K \rightarrow \mathbb{R}$  is a bounded Borel function, we will consider the operator  $g\text{Id} : C(K)^* \rightarrow C(K)^*$  which sends the functional which is the integration of a function  $f \in C(K)$  with respect to a measure  $\mu$  to the functional which is the integration of the product  $fg$  with respect to  $\mu$ .

**Definition 3.1.** Let  $K$  be a compact space and  $T \in L(C(K))$ . We say that  $T$  is a *weak multiplier* if  $T^* = g\text{Id} + S$  where  $g : K \rightarrow \mathbb{R}$  is a bounded Borel function on  $K$  and  $S \in W(C(K)^*)$ .

This definition was given in [19] in an equivalent form (see [19, Definition 2.1] and [19, Theorem 2.2]).

**Definition 3.2.** We say that an open set  $V \subseteq K$  is *compatible* with  $L$  if and only if  $L \subseteq V$  or  $L \cap \bar{V} = \emptyset$ . In the first case, the notation  $C_E(\bar{V}||L)$  has the same meaning as in the previous section. If  $L \cap \bar{V} = \emptyset$ , we will write  $C_E(\bar{V}||L)$  just to denote  $C(\bar{V})$ . Let us also observe that if  $L \subseteq V$ , then

$$C_E(\bar{V}||L)^* \equiv C_0(\bar{V}||L)^* \oplus_1 C_0(\bar{V}||L)^\perp \equiv C_0(\bar{V}||L)^* \oplus_1 E^*$$

since Lemma 2.2 applies to  $\bar{V}$ .

Given an open set  $V \subseteq K$  compatible with  $L$ , we consider the restriction operator  $P^{\bar{V}} : C_E(K||L)^* \rightarrow C_E(\bar{V}||L)^*$  given by

$$P^{\bar{V}}(\phi) = \phi|_{V \setminus L} + \phi_E$$

for  $\phi = \phi|_{K \setminus L} + \phi_E$  where  $\phi|_{K \setminus L} \in C_0(K||L)^*$  and  $\phi_E \in C_0(K||L)^\perp \equiv E^*$ . Observe that  $\phi_E$  can also be viewed as an element of  $C_0(\bar{V}||L)^\perp$  since the spaces  $C_0(\bar{V}||L)^\perp$  and  $C_0(K||L)^\perp$  are isometrically isomorphic (both coincide with  $E^*$ ).

Given an open set  $V \subseteq K$  compatible with  $L$  and  $h : K \rightarrow [0, 1]$  a continuous function constant on  $L$  with support included in  $V$ , we denote by  $P_{\bar{V}} : C_E(K||L) \rightarrow C_E(\bar{V}||L)$  and  $I_{h, \bar{V}} : C_E(\bar{V}||L) \rightarrow C_E(K||L)$  the operators defined by

$$P_{\bar{V}}(f) = f|_{\bar{V}} \quad \text{and} \quad I_{h, \bar{V}}(\tilde{f}) = h\tilde{f}$$

for  $f \in C_E(K||L)$  and  $\tilde{f} \in C_E(\bar{V}||L)$  respectively. We observe that  $I_{h, \bar{V}}$  is well defined, that is,  $h\tilde{f}$  is a function in  $C(K)$  with  $h\tilde{f}|_L \in E$  (indeed,  $h\tilde{f}$  is continuous in  $V$  as a product of two continuous functions and it is continuous in  $K \setminus \text{Supp}(h)$  as a constant function, since these two sets form an open cover of  $K$  we have that  $h\tilde{f}$  is continuous in  $K$ ; being  $h$  constant on  $L$ , it is clear that  $h\tilde{f}|_L \in E$ ). Finally, If  $V_1 \subseteq K$  is an open set compatible with  $L$  satisfying  $\bar{V} \subseteq V_1$  we will also use the notation  $P_{\bar{V}}$  for the restriction operator from  $C_E(\bar{V}_1||L)$  to  $C_E(\bar{V}||L)$ . In the next result we gather some easy facts concerning these operators.

**Lemma 3.3.** *Let  $V$  and  $h$  be as above. Then, the following hold.*

- (a)  $P_{\bar{V}}^*(\phi)(f) = \phi(f|_{\bar{V}})$  for  $\phi \in C_E(\bar{V}||L)^*$  and  $f \in C_E(K||L)$ .
- (b)  $I_{h, \bar{V}}^*(\phi)(\tilde{f}) = \phi(\tilde{f}h)$  for  $\phi \in C_E(K||L)^*$  and  $\tilde{f} \in C_E(\bar{V}||L)$ .
- (c) If  $V_0$  is an open set such that  $\bar{V}_0 \subseteq V$  and  $h|_{\bar{V}_0} \equiv 1$ , then  $(I_{h, \bar{V}}^* P_{\bar{V}}^*)(\mu) = \mu$  for every  $\mu \in C(K)^*$  with  $\text{Supp}(\mu) \subseteq V_0$ .
- (d) If  $E = C(L)$ , then  $C_E(K||L) = C(K)$ ,  $C_E(\bar{V}||L) = C(\bar{V})$ , and  $P^{\bar{V}} P_{\bar{V}}^*(\mu) = \mu$  for every  $\mu \in C(\bar{V})^*$ .

**Proof.** Parts (a) and (b) of the lemma are obvious from the definitions of the operators. To prove (c) we fix  $\tilde{f} \in C_E(\bar{V}||L)$ ,  $\mu \in C(K)^*$  with  $\text{Supp}(\mu) \subseteq V_0$  and we observe that

$$\begin{aligned} (I_{h,\bar{V}}^* P_{\bar{V}}^*)(\mu)(\tilde{f}) &= P_{\bar{V}}^*(\mu)(I_{h,\bar{V}}(\tilde{f})) \\ &= P_{\bar{V}}^*(\mu)(\tilde{f}h) \\ &= \mu((\tilde{f}h)|_{\bar{V}}) \\ &= \int (\tilde{f}h)|_{\bar{V}} \, d\mu \\ &= \int_{V_0} \tilde{f} \, d\mu \\ &= \mu(\tilde{f}). \end{aligned}$$

The first two assertions of (d) are obvious. For the third one, given  $\mu \in C(\bar{V})^*$  and  $\tilde{f} \in C(\bar{V})$ , use the regularity of the measure  $P_{\bar{V}}^*(\mu)$  to find an open set  $V_n \subseteq K$  satisfying

$$\bar{V} \subseteq V_n \quad \text{and} \quad |P_{\bar{V}}^*(\mu)|(V_n \setminus \bar{V}) < \frac{1}{n} \tag{3.1}$$

for every  $n \in \mathbb{N}$ . Next, take  $f_n \in C(K)$  satisfying

$$f_n|_{\bar{V}} \equiv \tilde{f}, \quad f_n|_{K \setminus V_n} \equiv 0 \quad \text{and} \quad \|f_n\| = \|\tilde{f}\|$$

for every  $n \in \mathbb{N}$ , and observe that

$$\begin{aligned} P^{\bar{V}} P_{\bar{V}}^*(\mu)(\tilde{f}) &= (P_{\bar{V}}^*(\mu))|_{\bar{V}}(\tilde{f}) \\ &= \int_{\bar{V}} \tilde{f} \, d(P_{\bar{V}}^*(\mu))|_{\bar{V}} \\ &= \int_K f_n \, dP_{\bar{V}}^*(\mu) - \int_{V_n \setminus \bar{V}} f_n \, dP_{\bar{V}}^*(\mu) \\ &= P_{\bar{V}}^*(\mu)(f_n) - \int_{V_n \setminus \bar{V}} f_n \, dP_{\bar{V}}^*(\mu) \\ &= \mu(\tilde{f}) - \int_{V_n \setminus \bar{V}} f_n \, dP_{\bar{V}}^*(\mu). \end{aligned}$$

Therefore, using (3.1) and letting  $n \rightarrow \infty$ , it follows that  $P^{\bar{V}} P_{\bar{V}}^*(\mu)(\tilde{f}) = \mu(\tilde{f})$ . □

Our first application uses the above operators in the simple case in which  $E = C(L)$ .

**Proposition 3.4.** *Let  $K$  be a compact space, let  $V_0, V_1$  and  $V_2$  be open non-empty subsets of  $K$  such that  $\bar{V}_0 \subseteq V_1$ , and let  $R : C(\bar{V}_2) \rightarrow C(\bar{V}_1)$  be a linear operator. Suppose that all operators on  $C(K)$  are weak multipliers. Then, there are a Borel function  $g : \bar{V}_1 \rightarrow \mathbb{R}$  with support included in  $\bar{V}_1 \cap \bar{V}_2$  and a weakly compact operator  $S : C(\bar{V}_1)^* \rightarrow C(\bar{V}_2)^*$  such that*

$$R^*(\mu) = g\mu + S(\mu)$$

for every  $\mu \in C(K)^*$  with  $\text{Supp}(\mu) \subseteq V_0$ .

**Proof.** We fix a continuous function  $h : K \rightarrow [0, 1]$  satisfying  $h|_{\bar{V}_0} \equiv 1$  and  $h|_{(K \setminus \bar{V}_1)} \equiv 0$  and we define  $R_0 \in L(C(K))$  by

$$R_0(f) = I_{h, \bar{V}_1} R P_{\bar{V}_2}^*(f) \tag{3.2}$$

for  $f \in C(K)$ . Hence, there are a bounded Borel function  $\hat{g} : K \rightarrow \mathbb{R}$  and a weakly compact operator  $\hat{S} \in L(C(K)^*)$  such that  $R_0^*(\mu) = \hat{g}\mu + \hat{S}(\mu)$  for  $\mu \in C(K)^*$ , which allows us to write

$$P^{\bar{V}_2} R_0^* P_{\bar{V}_1}^* = P^{\bar{V}_2} \hat{g} \text{Id}_{C(K)^*} P_{\bar{V}_1}^* + P^{\bar{V}_2} \hat{S} P_{\bar{V}_1}^*. \tag{3.3}$$

We claim that, considering the weakly compact operator given by  $S = P^{\bar{V}_2} \hat{S} P_{\bar{V}_1}^*$  and defining the functions  $\check{g} : K \rightarrow \mathbb{R}$  and  $g : \bar{V}_1 \rightarrow \mathbb{R}$  by

$$\check{g}(x) = \begin{cases} \hat{g}(x) & \text{if } x \in \bar{V}_1 \cap \bar{V}_2, \\ 0 & \text{if } x \notin \bar{V}_1 \cap \bar{V}_2 \end{cases} \quad \text{and} \quad g = \check{g}|_{\bar{V}_1}$$

the following holds for  $\mu \in C(\bar{V}_1)^*$ :

$$(P^{\bar{V}_2} R_0^* P_{\bar{V}_1}^*)(\mu) = g\mu + S(\mu). \tag{3.4}$$

Indeed, for  $\mu \in C(\bar{V}_1)^*$  and  $f \in C(\bar{V}_2)$  we observe that

$$\begin{aligned} (P^{\bar{V}_2} \hat{g} \text{Id}_{C(K)^*} P_{\bar{V}_1}^*)(\mu)(f) &= (\hat{g} P_{\bar{V}_1}^*(\mu))|_{\bar{V}_2}(f) \\ &= \int_{\bar{V}_2} \hat{g} f \, dP_{\bar{V}_1}^*(\mu) \\ &= \int_{\bar{V}_1 \cap \bar{V}_2} \hat{g} f \, dP_{\bar{V}_1}^*(\mu) \\ &= \int_{\bar{V}_1 \cap \bar{V}_2} \check{g} f \, dP_{\bar{V}_1}^*(\mu) \\ &= \int_K \check{g} f \, dP_{\bar{V}_1}^*(\mu) \end{aligned}$$

and, for  $n \in \mathbb{N}$ , we use Lusin’s Theorem (see, for example, [22, Theorem 21.4]) to find a compact set  $K_n \subseteq K$  such that

$$\check{g}|_{K_n} \text{ is continuous on } K_n, \quad |P_{\bar{V}_1}^*(\mu)|(K \setminus K_n) < \frac{1}{n} \quad \text{and} \quad |\mu|(\bar{V}_1 \setminus (\bar{V}_1 \cap K_n)) < \frac{1}{n}. \tag{3.5}$$

Using Tietze’s Extension Theorem we may find a continuous function  $g_n : K \rightarrow \mathbb{R}$  satisfying

$$g_n|_{K_n} = \check{g}|_{K_n} \quad \text{and} \quad \|g_n\| = \|\check{g}|_{K_n}\| \leq \|\check{g}\|$$

for every  $n \in \mathbb{N}$ . Now it is easy to check that

$$\begin{aligned} \int_K \check{g}f \, dP_{\check{V}_1}^*(\mu) &= \int_K g_n f \, dP_{\check{V}_1}^*(\mu) + \int_{K \setminus K_n} (\check{g} - g_n) f \, dP_{\check{V}_1}^*(\mu) \\ &= P_{\check{V}_1}^*(\mu)(g_n f) + \int_{K \setminus K_n} (\check{g} - g_n) f \, dP_{\check{V}_1}^*(\mu) \\ &= \mu((g_n f)|_{\check{V}_1}) + \int_{K \setminus K_n} (\check{g} - g_n) f \, dP_{\check{V}_1}^*(\mu) \\ &= \int_{\check{V}_1} g_n f \, d\mu + \int_{K \setminus K_n} (\check{g} - g_n) f \, dP_{\check{V}_1}^*(\mu) \\ &= \int_{\check{V}_1} g f \, d\mu + \int_{\check{V}_1 \setminus (\check{V}_1 \cap K_n)} (g_n - \check{g}) f \, d\mu + \int_{K \setminus K_n} (\check{g} - g_n) f \, dP_{\check{V}_1}^*(\mu) \\ &= \mu(gf) + \int_{\check{V}_1 \setminus (\check{V}_1 \cap K_n)} (g_n - \check{g}) f \, d\mu + \int_{K \setminus K_n} (\check{g} - g_n) f \, dP_{\check{V}_1}^*(\mu) \end{aligned}$$

which, letting  $n \rightarrow \infty$  and using (3.5), implies that

$$\int_K \check{g}f \, dP_{\check{V}_1}^*(\mu) = \mu(gf)$$

and, therefore,

$$(P^{\check{V}_2} \hat{g} \text{Id}_{C(K)^*} P_{\check{V}_1}^*)(\mu)(f) = \mu(gf).$$

This, together with (3.3) and the definition of  $S$ , finishes the proof of the claim. On the other hand by (3.2), we can write

$$P^{\check{V}_2} R_0^* P_{\check{V}_1}^* = P^{\check{V}_2} P_{\check{V}_2}^* R^* I_{h, \check{V}_1}^* P_{\check{V}_1}^*.$$

So, if the support of  $\mu$  is included in  $V_0$ , by parts (c) and (d) of Lemma 3.3 we obtain

$$P^{\check{V}_2} R_0^* P_{\check{V}_1}^*(\mu) = R^*(\mu)$$

and, consequently,  $R^*(\mu) = g\mu + S(\mu)$  follows from (3.4). □

**Remark 3.5.** The result above shows that if every operator on  $C(K)$  is a weak multiplier then, in the above sense, there are also few operators on  $C(\bar{V})$  for  $V$  open (since for such a closed set it is possible to define an appropriate function  $h$  as in the proof). In general, one cannot replace closures of open sets by general closed sets: it is shown in [9] that under the Continuum Hypothesis, there are compact  $K$ s as above which contain  $\beta\mathbb{N}$  (and, of course, there are many operators on  $C(\beta\mathbb{N}) \equiv \ell_\infty$ ). On the other hand, using the set-theoretic principle  $\diamond$ , it is also shown in [9] that there are  $K$ s such that for every infinite closed  $K' \subseteq K$ , all operators on the space  $C(K')$  are weak multipliers.

**Corollary 3.6.** *Let  $K$  be a compact space, let  $V_0, V_1$  and  $V_2$  be open non-empty subsets of  $K$  such that  $\bar{V}_0 \subseteq V_1$  and  $\bar{V}_1 \cap \bar{V}_2 = \emptyset$ , and let  $R : C(\bar{V}_2) \rightarrow C(\bar{V}_1)$  be a linear operator. Suppose that all operators on  $C(K)$  are weak multipliers. Then,  $P_{V_0} R$  is weakly compact.*

**Proof.** By Proposition 3.4 there is a weakly compact operator  $S : C^*(\bar{V}_1) \rightarrow C^*(\bar{V}_2)$  such that  $R^*(\mu) = S(\mu)$  for every measure  $\mu$  with support included in  $\bar{V}_0$ . In other words,  $(P_{\bar{V}_0}R)^* = R^*P_{\bar{V}_0}^*$  is weakly compact and so, by Gantmacher theorem,  $P_{\bar{V}_0}R$  is weakly compact.  $\square$

The following result is an easy consequence of the Dieudonné–Grothendieck Theorem which we state for the sake of clearness.

**Lemma 3.7.** *Let  $K$  be a compact space,  $X$  a Banach space and  $S : X^* \rightarrow C(K)^*$  a weakly compact operator. Then, for every bounded subset  $B \subseteq X^*$ , the set*

$$\{x \in K : \exists \phi \in B \text{ so that } S(\phi)(\{x\}) \neq 0\}$$

is countable.

**Proof.** Suppose that the set

$$\{x \in K : \exists \phi \in B \text{ so that } S(\phi)(\{x\}) \neq 0\}$$

is uncountable for some bounded set  $B \subseteq X^*$ . Then, there is  $\varepsilon > 0$  so that the set

$$\{x \in K : \exists \phi \in B \text{ so that } |S(\phi)(\{x\})| \geq \varepsilon\}$$

is infinite, which contradicts the fact of being  $S(B)$  relatively weakly compact by the Dieudonné–Grothendieck Theorem (see, for example, [5, Theorem VII.14]).  $\square$

**Lemma 3.8.** *Let  $K$  be a compact space, let  $V_0, V_1$  and  $V_2$  be open non-empty subsets of  $K$  compatible with  $L$  such that  $\bar{V}_0 \subseteq V_1$  and  $\bar{V}_1 \cap L = \emptyset$ , and let  $T : C_E(K||L) \rightarrow C_E(K||L)$  be a linear operator. Then, there exists an operator  $R : C(\bar{V}_1) \rightarrow C_E(\bar{V}_2||L)$  such that*

$$R^*(\phi)|_{\bar{V}_0} = (T^*P_{V_2}^*)(\phi)|_{\bar{V}_0}$$

for all  $\phi \in C_E(\bar{V}_2||L)^*$ .

**Proof.** Take a continuous function  $h : K \rightarrow [0, 1]$  satisfying  $h|_{\bar{V}_0} \equiv 1$  and  $h|_{(K \setminus V_1)} \equiv 0$ , and define the operator  $R = P_{\bar{V}_2}TI_{h, \bar{V}_1}$ . Given  $\phi \in C_E(\bar{V}_2||L)^*$  and  $f \in C(\bar{V}_1)$  with  $\text{Supp}(f) \subseteq \bar{V}_0$ , by parts (b) and (a) of Lemma 3.3 and using the facts  $h|_{\bar{V}_0} \equiv 1$  and  $\text{Supp}(f) \subseteq \bar{V}_0$ , we can write

$$R^*(\phi)(f) = I_{h, \bar{V}_1}^*T^*P_{\bar{V}_2}^*(\phi)(f) = T^*P_{\bar{V}_2}^*(\phi)(fh) = T^*P_{\bar{V}_2}^*(\phi)(f),$$

which finishes the proof.  $\square$

We are ready to state and prove the main result of the section.

**Theorem 3.9.** *Let  $K$  be a perfect compact space such that all operators on  $C(K)$  are weak multipliers, let  $L \subseteq K$  be closed and nowhere dense, and  $E$  a closed subspace of  $C(L)$ . Then,  $C_E(K||L)$  is extremely non-complex.*

**Proof.** Fixed  $T \in L(C_E(K\|L))$ , we have to show that its square satisfies the Daugavet equation. By Lemma 2.5, it is enough to prove that the set  $\{x \in K \setminus L : g_{(T^2)^*}(x) \geq 0\}$  is dense in  $K \setminus L$ . To do so, we proceed ad absurdum: suppose that there is an open set  $U_1 \subseteq K$  such that  $\bar{U}_1 \cap L = \emptyset$  and  $g_{(T^2)^*}(x) < 0$  for each  $x \in U_1$ . By going to a subset, we may without loss of generality assume that  $\bar{U}_1$  is a  $G_\delta$  set. Therefore, we can find open sets  $W_n \subseteq K$  such that  $\bigcap_{n \in \mathbb{N}} W_n = \bar{U}_1$ ,  $\bar{W}_{n+1} \subseteq W_n$ , and  $K \setminus W_n$  is the closure of an open set containing  $L$  for every  $n \in \mathbb{N}$ . Next, we fix a non-empty open set  $U_0 \subseteq K$  with  $\bar{U}_0 \subseteq U_1$ , and we observe that it is uncountable (since  $K$  is perfect) so there is  $\varepsilon > 0$  such that the set

$$A = \{x \in U_0 : g_{(T^2)^*}(x) < -\varepsilon\}$$

is uncountable. Moreover, we claim that there is  $n_0 \in \mathbb{N}$  such that the set

$$B = \left\{ x \in A : |T^*(\delta_x)|(W_{n_0} \setminus \bar{U}_1) < \frac{\varepsilon}{2\|T\|} \right\}$$

is uncountable. Indeed, fixed  $x \in A$ , the regularity of the measure  $T^*(\delta_x)$  implies that there is  $n \in \mathbb{N}$  so that  $|T^*(\delta_x)|(W_n \setminus \bar{U}_1) < \varepsilon/2\|T\|$  which gives us

$$A = \bigcup_{n \in \mathbb{N}} \left\{ x \in A : |T^*(\delta_x)|(W_n \setminus \bar{U}_1) < \frac{\varepsilon}{2\|T\|} \right\}$$

and the uncountability of  $A$  finishes the argument.

For  $x \in B$ , we write

$$\phi_x = T^*(\delta_x)|_{K \setminus (L \cup W_{n_0})} + T^*(\delta_x)_E \in C_E((K \setminus W_{n_0})\|L)^*$$

and we can decompose  $T^*(\delta_x)$  as follows:

$$T^*(\delta_x) = T^*(\delta_x)|_{\bar{U}_1} + T^*(\delta_x)|_{W_{n_0} \setminus \bar{U}_1} + \phi_x.$$

Hence, for every  $x \in B$ , we get

$$-\varepsilon > [(T^*)^2(\delta_x)](\{x\}) = T^*[T^*(\delta_x)|_{\bar{U}_1} + T^*(\delta_x)|_{W_{n_0} \setminus \bar{U}_1} + \phi_x](\{x\}). \tag{3.6}$$

However, the following claims show that this is impossible.

**Claim 1.**  $\|T^*[T^*(\delta_x)|_{W_{n_0} \setminus \bar{U}_1}]\| < \frac{1}{2}\varepsilon$ .

**Proof of Claim 1.** It follows obviously from the choice of  $n_0$ .

**Claim 2.** The function  $x \mapsto T^*(T^*(\delta_x)|_{\bar{U}_1})(\{x\})$  is non-negative for all but countably many  $x \in B$ .

**Proof of Claim 2.** By Lemma 3.8 applied to  $V_0 = U_1$ ,  $V_1 = W_{n_0}$  and  $V_2 = U_1$ , we obtain an operator  $R : C(\bar{W}_{n_0}) \rightarrow C(\bar{U}_1)$  such that

$$R^*(\psi)|_{\bar{U}_1} = T^*P_{\bar{U}_1}^*(\psi)|_{\bar{U}_1}$$

for  $\psi \in C(\bar{U}_1)^*$ . On the other hand, by Proposition 3.4 applied to  $R$  and  $V_0 = U_0$ ,  $V_1 = U_1$ ,  $V_2 = W_{n_0}$ , we get a bounded Borel function  $g : \bar{U}_1 \rightarrow \mathbb{R}$  and a weakly compact operator  $S : C(\bar{U}_1)^* \rightarrow C(\bar{W}_{n_0})^*$  such that

$$R^*(\mu) = g\mu + S(\mu)$$

for every  $\mu$  with support in  $U_0$ . In particular, for  $x \in B$  we have

$$T^*(\delta_x)|_{\bar{U}_1} = T^*P_{\bar{U}_1}^*(\delta_x)|_{\bar{U}_1} = R^*(\delta_x)|_{\bar{U}_1} = g\delta_x + S(\delta_x)|_{\bar{U}_1} = g(x)\delta_x + S(\delta_x)|_{\bar{U}_1}$$

and using this twice we get

$$\begin{aligned} T^*(T^*(\delta_x)|_{\bar{U}_1})(\{x\}) &= T^*[g(x)\delta_x + S(\delta_x)|_{\bar{U}_1}](\{x\}) \\ &= g(x)T^*(\delta_x)(\{x\}) + T^*[S(\delta_x)|_{\bar{U}_1}](\{x\}) \\ &= g(x)T^*(\delta_x)|_{\bar{U}_1}(\{x\}) + R^*[S(\delta_x)|_{\bar{U}_1}](\{x\}) \\ &= g(x)^2 + g(x)S(\delta_x)|_{\bar{U}_1}(\{x\}) + (gS(\delta_x)|_{\bar{U}_1})(\{x\}) + S[S(\delta_x)|_{\bar{U}_1}](\{x\}) \\ &= g(x)^2 + (2gP^{\bar{U}_1}S)(\delta_x)(\{x\}) + (SP^{\bar{U}_1}S)(\delta_x)(\{x\}) \end{aligned}$$

for every  $x \in B$ . Finally, since  $gP^{\bar{U}_1}S$  and  $SP^{\bar{U}_1}S$  are weakly compact operators, we conclude by Lemma 3.7 that  $(gP^{\bar{U}_1}S)(\delta_x)(\{x\})$  and  $(SP^{\bar{U}_1}S)(\delta_x)(\{x\})$  are zero for all but countably many  $x$ , completing the proof of the claim.

**Claim 3.**  $T^*(\phi_x)(\{x\}) = 0$  for all but countably many  $x \in U_0$ .

**Proof of Claim 3.** By Lemma 3.8 applied to  $V_0 = U_0$ ,  $V_1 = U_1$ , and  $V_2 = K \setminus \bar{W}_{n_0+1}$  we obtain an operator  $R : C(\bar{U}_1) \rightarrow C_E((K \setminus W_{n_0+1})\|L)$  such that  $R^*(\phi)(\{x\}) = T^*P_{K \setminus W_{n_0}}^*(\phi)(\{x\})$  for  $x \in U_0$  and  $\phi \in C_E((K \setminus W_{n_0+1})\|L)^*$ . We denote

$$J : C_E((K \setminus W_{n_0+1})\|L) \rightarrow C(K \setminus W_{n_0+1})$$

the inclusion operator and we apply Corollary 3.6 for the operator  $JR$  and the open sets  $V_0 = K \setminus \bar{W}_{n_0}$ ,  $V_1 = K \setminus \bar{W}_{n_0+1}$ , and  $V_2 = U_1$  to obtain that the operator  $P_{K \setminus W_{n_0}}JR$  is weakly compact.

Besides, we recall that  $\phi_x \in C_E((K \setminus W_{n_0}\|L))^*$  and that it can be viewed as an element of  $C_E((K \setminus W_{n_0+1}\|L))^*$  by just extending it by zero outside  $K \setminus W_{n_0}$ . For  $x \in U_0$  we take  $\tilde{\phi}_x$  a Hahn–Banach extension of  $\phi_x$  to  $C(K \setminus W_{n_0})$  and we observe that  $J^*P_{K \setminus W_{n_0}}^*(\tilde{\phi}_x) = \phi_x$ . Indeed, for  $f \in C_E((K \setminus W_{n_0+1})\|L)$  we have that

$$\begin{aligned} J^*P_{K \setminus W_{n_0}}^*(\tilde{\phi}_x)(f) &= \tilde{\phi}_x(P_{K \setminus W_{n_0}}(Jf)) \\ &= \tilde{\phi}_x(P_{K \setminus W_{n_0}}(f)) \\ &= \tilde{\phi}_x(f|_{K \setminus W_{n_0}}) \\ &= \phi_x(f|_{K \setminus W_{n_0}}) \\ &= \phi_x(f). \end{aligned}$$

Therefore, for  $x \in U_0$ , we can write

$$\begin{aligned} T^*(\phi_x)(\{x\}) &= T^*P_{K \setminus W_{n_0}}^*(\phi_x)(\{x\}) \\ &= R^*(\phi_x)(\{x\}) \\ &= R^*J^*P_{K \setminus W_{n_0}}^*(\tilde{\phi}_x)(\{x\}) \\ &= (P_{K \setminus W_{n_0}}JR)^*(\tilde{\phi}_x)(\{x\}), \end{aligned}$$

where we are identifying  $\phi_x$  with its extension by zero to  $K$ . Now the proof of the claim is finished by just applying Lemma 3.7 to the operator  $P_{K \setminus W_{n_0}}JR$ .

Finally, the claims obviously contradict (3.6) completing the proof of the theorem.  $\square$

When  $E = \{0\}$ , we get a sufficient condition to get that a space of the form  $C_0(K \setminus L)$  is extremely non-complex.

**Corollary 3.10.** *Let  $K$  be a compact space such that all operators on  $C(K)$  are weak multipliers. Suppose  $L \subseteq K$  is closed and nowhere dense. Then,  $C_0(K \setminus L)$  is extremely non-complex.*

To show that there are extremely non-complex spaces of the form  $C_E(K||L)$  which are not isomorphic to the  $C(K')$  spaces, we need the following (well-known) result which allows us to construct spaces  $C_E(K||L)$  for every perfect separable compact space  $K$  and every separable Banach space  $E$ .

**Lemma 3.11.** *Let  $K$  be a perfect compact space. Then*

- (a) *there is a nowhere-dense closed subset  $L \subset K$  such that  $L$  can be continuously mapped onto the Cantor set;*
- (b) *therefore, every separable Banach space  $E$  is (isometrically isomorphic to) a subspace of  $C(L)$ .*

**Proof.** (a) As  $K$  is perfect, given a non-empty open subset  $U$  in  $K$  and  $x \in U$ , there are two non-empty open subsets  $V_1, V_2$  of  $U$  such that

$$\bar{V}_1 \cap \bar{V}_2 = \emptyset \quad \text{and} \quad x \notin \bar{V}_i \quad (i = 1, 2).$$

This allows us to construct a family of open sets  $U_s$  for  $s \in \{0, 1\}^{<\omega}$  such that

$$U_\emptyset = K, \quad \bar{U}_{s \smallfrown 0} \cap \bar{U}_{s \smallfrown 1} = \emptyset, \quad \bar{U}_{s \smallfrown 0}, \bar{U}_{s \smallfrown 1} \subseteq U_s \quad \text{and} \quad U_s \setminus [\bar{U}_{s \smallfrown 0} \cup \bar{U}_{s \smallfrown 1}] \neq \emptyset.$$

Take any point  $y_s$  in the above difference. Define  $L$  to be the set of all the accumulation points of the set  $\{y_s : s \in \{0, 1\}^{<\omega}\}$ .

For  $n \in \mathbb{N}$ , let  $f_n : K \rightarrow [0, 1]$  be such continuous functions that for all  $s \in \{0, 1\}^n$  we have  $f_n[\bar{U}_{s \smallfrown 0}] = \{0\}$  and  $f_n[\bar{U}_{s \smallfrown 1}] = \{1\}$  which can be easily obtained since  $\bar{U}_s \cap \bar{U}_{s'} = \emptyset$  if  $s, s' \in \{0, 1\}^n$  are distinct. Let  $f : K \rightarrow [0, 1]^{\mathbb{N}}$  be defined by  $f(x)(n) = f_n(x)$ . We claim that  $f|_L$  satisfies the lemma. One can easily check that  $f$  is continuous since the  $f_n$ s are continuous.

Because each  $\bar{U}_s$  contains infinitely many points  $y_t$ , we have that  $L \cap \bar{U}_s \neq \emptyset$  for each  $s \in \{0, 1\}^{<\omega}$ . Note that if  $x \in \bar{U}_s$  and  $s \in \{0, 1\}^n$ , then  $f(x)|\{0, \dots, n-1\} = s$ . So, as the image of  $L$  under  $f$ , is closed, it contains  $\{0, 1\}^{\mathbb{N}}$ .

On the other hand, if  $x \in L$ , then for each  $n \in \mathbb{N}$  there is  $s \in \{0, 1\}^n$  such that  $x \in \bar{U}_s$ , this is because  $K \setminus \bigcup\{\bar{U}_s : |s| = n\}$  contains only finitely many points  $y_t$  and hence no element of  $L$ . Thus  $f_n(x) \in \{0, 1\}$  if  $x \in L$  which gives that  $f[L] \subseteq \{0, 1\}^{\mathbb{N}}$ , which together with the previous observation gives that  $f[L] = \{0, 1\}^{\mathbb{N}}$ .

Finally, let us prove that  $L$  has empty interior, and so, as a closed set, it is nowhere dense. It is enough to see that  $L$  has empty interior in the subspace  $\{y_s : s \in \{0, 1\}^{<\omega}\} \cup L$ . This is true since  $\{y_s : s \in \{0, 1\}^{<\omega}\}$  is dense and open in  $\{y_s : s \in \{0, 1\}^{<\omega}\} \cup L$ , as each point  $y_s$  is isolated by  $U_s \setminus [\bar{U}_{s-0} \cup \bar{U}_{s-1}]$  from the remaining points.

(b) Since the function  $f|_L : L \rightarrow 2^\omega$  of the above item is continuous and surjective,  $C(2^\omega)$  embeds isometrically into  $C(L)$  by just composing every element in  $C(2^\omega)$  with  $f|_L$ . Since every separable Banach space  $E$  embeds isometrically into  $C(2^\omega)$  (Banach–Mazur theorem), we get  $E \subseteq C(L)$  isometrically. □

**Remark 3.12.** If  $K$  is a compact space such that all operators on  $C(K)$  are weak multipliers, then it is easier to prove the existence of  $L \subseteq K$  closed nowhere dense which maps onto  $[0, 1]$  giving also (b) above. Indeed,  $C(K)$  is a Grothendieck space by [19, Theorem 2.4], so  $K$  has no convergent sequence (otherwise it would give rise to a complemented copy of  $c_0$  contradicting the Grothendieck property). Now, take any discrete sequence  $\{x_n : n \in \mathbb{N}\} \subseteq K$  and consider the set  $L$  of all its accumulation points. Then,  $L$  is perfect because an isolated point would produce a convergent subsequence of  $\{x_n : n \in \mathbb{N}\}$ , so  $L$  continuously maps onto  $[0, 1]$  [25, Theorem 8.5.4]. To see that  $L$  is nowhere dense we use the discreteness of  $\{x_n : n \in \mathbb{N}\}$ . If  $U \subseteq K$  is open and intersects  $L$ , then there is  $n_0 \in \mathbb{N}$  such that  $x_{n_0} \in U$ ; but by the discreteness of  $\{x_n : n \in \mathbb{N}\}$ , there is an open neighbourhood  $V$  of  $x_{n_0}$  not containing the remaining  $x_n$ s and hence, disjoint from  $L$ . Therefore,  $V \cap U$  is an open subset of  $U$  disjoint with  $L$ , proving that  $L$  is nowhere dense.

Now, we take a perfect compact space  $K$  such that every operator on  $C(K)$  is a weak multiplier [19], and we use Lemma 3.11 to find a nowhere-dense closed subset  $L$  such that  $C(L)$  contains isometric copies of every separable Banach space. Then, for every  $E \subset C(L)$ ,  $C_E(K||L)$  is extremely non-complex by Theorem 3.9 and  $C_E(K||L)^* = C_0(K||L)^* \oplus_1 E^*$  by Lemma 2.2. If  $E$  is infinite dimensional and reflexive,  $C_E(K||L)$  is not isomorphic to a  $C(K')$  space, since  $C(K')^*$  never contains complemented infinite-dimensional reflexive subspaces (see, for example, [3, Proposition 5.6.1]). Let us state all what we have proved.

**Example 3.13.**

- (a) For every separable Banach space  $E$ , there is an extremely non-complex Banach space  $C_E(K||L)$  such that  $E^*$  is an  $L$ -summand in  $C_E(K||L)^*$ .
- (b) If  $E$  is infinite dimensional and reflexive, then such  $C_E(K||L)$  is not isomorphic to any  $C(K')$  space.

- (c) Therefore, *there are extremely non-complex Banach spaces which are not isomorphic to  $C(K)$  spaces.*

We finish the section commenting that some  $C(K')$  spaces with many operators can be viewed as  $C_E(K\|L)$  spaces where  $C(K)$  has few operators and for which our previous results apply.

**Remark 3.14.** Let  $L \subseteq K$  be a nowhere-dense subset of a compact  $K$  as before. Consider the topological quotient map  $q : K \rightarrow K_L$ , where  $K_L$  is obtained from  $K$  by identifying all points of  $L$  to one point. The canonical isometric embedding  $I_q$  of  $C(K_L)$  into  $C(K)$  defined by  $I_q(f) = f \circ q$  has the image equal to the subspace of  $C(K)$  consisting of all functions constant on  $L$ . Thus  $C(K_L)$  is isometric to  $C_E(K\|L)$ , where  $E$  is the subspace of  $C(L)$  of all constant functions. Hence, by the results above, if all operators on  $C(K)$  are weak multipliers, then all spaces of the form  $C(K_L)$  are extremely non-complex. It turns out that the spaces of [20] can be realized as spaces of this form. In particular, there are extremely non-complex spaces of the form  $C_E(K\|L)$  which have many operators besides weak multipliers. For example, take  $K$  such that all operators on  $C(K)$  are weak multipliers. Choose a discrete sequence  $(x_n)$  in  $K$  and let  $L$  be the set of its accumulation points. Then the sequence  $(x_n)$  has a unique accumulation point in  $K_L$ , that is, it is a convergent sequence. By a well-known fact, this means that  $C(K_L) \cong C_E(K\|L)$  has a complemented copy of  $c_0$  and so is not Grothendieck, hence it has more operators than weak multipliers by results of [19] (actually, many operators which are not weak multipliers can be directly obtained from automorphisms of the complemented copy of  $c_0$  generated by permutations of the natural numbers).

#### 4. Isometries on extremely non-complex spaces

The following result shows that the group of isometries of an extremely non-complex Banach space is a discrete Boolean group.

**Theorem 4.1.** *Let  $X$  be an extremely non-complex Banach space. Then*

- (a) *if  $T \in \text{Iso}(X)$ , then  $T^2 = \text{Id}$ ,*
- (b) *as a consequence, for every  $T_1, T_2 \in \text{Iso}(X)$ ,  $T_1T_2 = T_2T_1$ ,*
- (c) *for every  $T_1, T_2 \in \text{Iso}(X)$ ,  $\|T_1 - T_2\| \in \{0, 2\}$ .*

**Proof.** (a) Given  $T \in \text{Iso}(X)$ , we define the operator  $S = (1/\sqrt{2})(T - T^{-1})$  and we observe that  $S^2 = \frac{1}{2}T^2 - \text{Id} + \frac{1}{2}T^{-2}$ . Since  $X$  is extremely non-complex, we get

$$1 + \|S^2\| = \|\text{Id} + S^2\| = \|\frac{1}{2}T^2 + \frac{1}{2}T^{-2}\| \leq 1$$

and, therefore,  $S^2 = 0$ . This gives us that  $\text{Id} = \frac{1}{2}T^2 + \frac{1}{2}T^{-2}$ . Finally, since  $\text{Id}$  is an extreme point of  $L(X)$  (see, for example, [24, Proposition 1.6.6]) and  $\|T^2\| \leq 1$ ,  $\|T^{-2}\| \leq 1$ , we get  $T^2 = \text{Id}$ .

(b) Commutativity comes routinely from the first part since  $T_1T_2 \in \text{Iso}(X)$ , so

$$\text{Id} = (T_1T_2)^2 = T_1T_2T_1T_2,$$

which finishes the proof by just multiplying by  $T_1$  from the left and by  $T_2$  from the right.

(c) We start by observing that  $\|\text{Id} - T\| \in \{0, 2\}$  for every  $T \in \text{Iso}(X)$ . Indeed, from (a) we have

$$(\text{Id} - T)^2 = \text{Id} + \text{Id} - 2T = 2(\text{Id} - T),$$

which gives us that

$$2\|\text{Id} - T\| = \|(\text{Id} - T)^2\| \leq \|\text{Id} - T\|^2.$$

Therefore, we get either  $\|\text{Id} - T\| = 0$  or  $\|\text{Id} - T\| \geq 2$ . Now, if  $T_1, T_2 \in \text{Iso}(X)$  we observe that

$$\|T_1 - T_2\| = \|T_1(\text{Id} - T_1T_2)\| = \|\text{Id} - T_1T_2\| \in \{0, 2\}.$$

□

As an immediate consequence we obtain the following result. Let us observe that there is no topological consideration on the semigroup.

**Corollary 4.2.** *If  $X$  is an extremely non-complex Banach space and  $\Phi : \mathbb{R}_0^+ \rightarrow \text{Iso}(X)$  is a one-parameter semigroup, then  $\Phi(\mathbb{R}_0^+) = \{\text{Id}\}$ .*

**Proof.** Just observe that  $\Phi(t) = \Phi(\frac{1}{2}t + \frac{1}{2}t) = \Phi(\frac{1}{2}t)^2 = \text{Id}$  for every  $t \in \mathbb{R}_0^+$ . □

Let  $X$  be a Banach space. A projection  $P \in L(X)$  is said to be *unconditional* if  $2P - \text{Id} \in \text{Iso}(X)$  (equivalently,  $\|2P - \text{Id}\| = 1$ ). We write  $\text{Unc}(X)$  for the set of unconditional projections on  $X$ . It is straightforward to show that  $P \in \text{Unc}(X)$  if and only if  $P = \frac{1}{2}(\text{Id} + T)$  for some  $T \in \text{Iso}(X)$  with  $T^2 = \text{Id}$ . It is then immediate that  $\text{Unc}(X)$  identifies with  $\{T \in \text{Iso}(X) : T^2 = \text{Id}\}$  and both sets are Boolean groups: the group operation in  $\{T \in \text{Iso}(X) : T^2 = \text{Id}\}$  is just the composition and so the group operation in  $\text{Unc}(X)$  is

$$(P_1, P_2) \mapsto P_1 + P_2 - P_1P_2.$$

It also follows that all unconditional projections commute.

If  $X$  is extremely non-complex, the set  $\{T \in \text{Iso}(X) : T^2 = \text{Id}\}$  is the whole  $\text{Iso}(X)$  (Theorem 4.1). We summarize all of this in the next result, where we will also discuss when these Boolean groups are actually Boolean algebras. The proof is completely straightforward. We refer the reader to the book [18, § 1.8] for background on Boolean algebras of projections.

**Proposition 4.3.** *Let  $X$  be an extremely non-complex Banach space.*

- (a)  $\text{Iso}(X)$  is a Boolean group for the composition operation.
- (b)  $\text{Unc}(X)$  is (equivalently,  $\text{Iso}(X)$  is isomorphic to) a Boolean algebra if, and only if,  $P_1P_2 \in \text{Unc}(X)$  for every  $P_1, P_2 \in \text{Unc}(X)$  if, and only if,  $\|\text{Id} + T_1 + T_2 - T_1T_2\| = 2$  for every  $T_1, T_2 \in \text{Iso}(X)$ .

We will show later that for many examples of extremely non-complex Banach spaces the set of unconditional projections is a Boolean algebra, but we do not know if this always happens.

**5. Surjective isometries on extremely non-complex  $C_E(K\|L)$  spaces**

Our aim in this section is to describe the group of isometries of the spaces  $C_E(K\|L)$  when they are extremely non-complex. We will deduce all the results from the following theorem.

**Theorem 5.1.** *Suppose that the space  $C_E(K\|L)$  is extremely non-complex. Then, for every  $T \in \text{Iso}(C_E(K\|L))$  there is a continuous function  $\theta : K \setminus L \rightarrow \{-1, 1\}$  such that*

$$[T(f)](x) = \theta(x)f(x)$$

for all  $x \in K \setminus L$  and  $f \in C_E(K\|L)$ .

**Proof.** We divide the proof into several claims.

**Claim 1.** The set  $D_0 = \{x \in K \setminus L : \exists y \in K \setminus L, \theta_0 \in \{-1, 1\} \text{ with } T^*(\delta_x) = \theta_0\delta_y\}$  is dense in  $K$ .

**Proof of Claim 1.** Let  $W$  be a non-empty open subset of  $K$ . Since  $K \setminus L$  is open and dense in  $K$ , there is  $V$  non-empty and open satisfying  $V \subseteq W \cap (K \setminus L)$ . Now,  $\{\delta_x : x \in V\}$  is a subset of  $\text{ext}(B_{C_E(K\|L)})$  by Lemma 2.4, and it is easy to check that it is weak\* open in  $\text{ext}(B_{C_E(K\|L)^*})$  (indeed, for  $x_0 \in V$  take a non-negative  $f \in C_0(K\|L)$  such that  $f(x_0) = 1$  and  $f(K \setminus V) = 0$ , and observe that  $\delta_{x_0} \in \{\delta_x : \delta_x(f) > \frac{1}{2}\} \subset \{\delta_x : x \in V\}$ ). Now, since  $T^*$  is a weak\* continuous surjective isometry, the mapping

$$T^* : (\text{ext}(B_{C_E(K\|L)^*}), w^*) \rightarrow (\text{ext}(B_{C_E(K\|L)^*}), w^*)$$

is a homeomorphism and so the set  $\{T^*(\delta_x) : x \in V\}$  is weak\* open in  $\text{ext}(B_{C_E(K\|L)^*})$ . Since, by Lemma 2.4, the set  $\{\theta\delta_y : y \in K \setminus L, \theta \in \{-1, 1\}\}$  is weak\* dense in  $\text{ext}(B_{C_E(K\|L)^*})$ , there are  $x \in V, y \in K \setminus L$ , and  $\theta_0 \in \{-1, 1\}$  such that  $T^*(\delta_x) = \theta_0\delta_y$ , which implies  $x \in V \cap D_0 \subseteq W \cap D_0$ , finishing the proof of Claim 1.

Now, we can consider functions  $\phi : D_0 \rightarrow D_0$  and  $\theta : D_0 \rightarrow \{-1, 1\}$  such that

$$T^*(\delta_x) = \theta(x)\delta_{\phi(x)} \tag{5.1}$$

for all  $x \in D_0$ . Since  $T^2 = \text{Id}$  by Theorem 4.1, if  $x \in D_0$  and  $T^*(\delta_x) = \pm\delta_y$ , then  $T^*(\delta_y) = \pm\delta_x$  and so  $y \in D_0$ . Therefore,  $\phi$  is a well-defined function from  $D_0$  into itself. Moreover, it also follows that

$$\phi^2(x) = x \quad \text{and} \quad \theta(x)\theta(\phi(x)) = 1 \quad (x \in D_0). \tag{5.2}$$

Indeed, given  $x \in D_0$ , we use the fact that  $(T^*)^2 = \text{Id}$  to get

$$\delta_x = T^*(T^*(\delta_x)) = T^*(\theta(x)\delta_{\phi(x)}) = \theta(x)\theta(\phi(x))\delta_{\phi^2(x)}.$$

**Claim 2.**  $\phi$  is a homeomorphism of  $D_0$ .

**Proof of Claim 2.** As  $\phi^2$  is the identity on  $D_0$ , it is enough to prove that  $\phi$  is continuous. To do so, fixed  $x_0 \in D_0$  and an open subset  $W$  of  $K \setminus L$  with  $\phi(x_0) \in W$ , we have to show that  $\phi^{-1}(W \cap D_0)$  is a neighbourhood of  $x_0$  in  $D_0$ . Indeed, we consider a continuous function  $f_0 \in C_0(K\|L) \subseteq C_E(K\|L)$  such that

$$f_0(\phi(x_0)) = 1 = \|f_0\| \quad \text{and} \quad f_0 \equiv 0 \quad \text{in } K \setminus W.$$

Since the mapping

$$x \mapsto [T^*(\delta_x)](f_0) = \theta(x)f_0(\phi(x)) \quad (x \in D_0)$$

is continuous at  $x_0$ , there is an open neighbourhood  $U_0$  of  $x_0$  such that

$$\| |f_0(\phi(x))| - 1 | \leq | \theta(x)f_0(\phi(x)) - \theta(x_0)f_0(\phi(x_0)) | < \frac{1}{2} \quad (x \in U_0 \cap D_0).$$

Since  $f_0 \equiv 0$  outside  $W$ , we get that  $U_0 \cap D_0 \subseteq \phi^{-1}(W \cap D_0)$ .

**Claim 3.**  $\phi(x) = x$  for all  $x \in D_0$ .

**Proof of Claim 3.** Suppose otherwise that there are  $x_0, y_0 \in D_0$  such that  $\phi(x_0) = y_0 \neq x_0$ . Let  $V_i \subseteq \bar{V}_i \subseteq K \setminus L$  with  $i = 1, 2$  be open subsets of  $K$  satisfying

$$x_0 \in V_1, \quad y_0 \in V_2, \quad \bar{V}_1 \cap \bar{V}_2 = \emptyset \quad \text{and} \quad V_1 \cap D_0 \subseteq \phi^{-1}(V_2 \cap D_0).$$

The last condition obviously implies  $\phi(V_1 \cap D_0) \subseteq V_2$  and, since  $\phi$  is a homeomorphism of  $D_0$ , it follows that  $\phi(V_1 \cap D_0)$  is open in  $D_0$ . Therefore, we can find  $V_0 \subseteq V_2$  an open subset of  $K$  such that  $V_0 \cap D_0 = \phi(V_1 \cap D_0)$ . Then, we may find  $g \in C_0(K\|L) \subseteq C_E(K\|L)$  satisfying  $g(x_0) = 1, g(y_0) = -1, g(x) \in [-1, 0]$  for  $x \in V_0, g(x) \in [0, 1]$  for  $x \in V_1$ , and  $g(x) = 0$  for  $x \notin V_1 \cup V_0$ . In particular, for  $x \in D_0$ , we have that

$$g(x)g(\phi(x)) \in [-1, 0].$$

Next, we define the operator  $T_g : C_E(K\|L) \rightarrow C_E(K\|L)$  by  $T_g(f) = gf$ , which is well defined (since  $g|_L \equiv 0$ ) and satisfies  $T_g^*(\delta_x) = g(x)\delta_x$  for each  $x \in K \setminus L$ . Finally, we consider the composition  $S = T_g T$  and, for  $x \in D_0$ , we use (5.1) and (5.2) to write

$$(S^*)^2(\delta_x) = T^*(T_g^*(T^*(T_g^*(\delta_x)))) = \theta(x)\theta(\phi(x))g(x)g(\phi(x))\delta_{\phi^2(x)} = g(x)g(\phi(x))\delta_x.$$

This, together with our choice of  $g$ , tells us that

$$\| [\text{Id} + (S^*)^2](\delta_x) \| \leq 1 \quad (x \in D_0).$$

As  $D_0$  is dense in  $K$  by Claim 1 and  $\text{Id} + (S^*)^2$  is weak\*-continuous, we deduce that  $\| \text{Id} + (S^*)^2 \| \leq 1$ . Now, the fact that  $C_E(K\|L)$  is extremely non-complex implies that  $S^2 = 0$ , which is a contradiction since  $(S^*)^2(\delta_{x_0}) = -\delta_{x_0} \neq 0$ .

**Claim 4.**  $D_0 = K \setminus L$ .

**Proof of Claim 4.** Let us fix  $x_0 \in K \setminus L$ . Since  $D_0$  is dense in  $K \setminus L$ , we may find a net  $(x_\lambda)_{\lambda \in \Lambda}$  in  $D_0$  such that  $(x_\lambda)_{\lambda \in \Lambda} \rightarrow x_0$ , so  $(T^*(\delta_{x_\lambda}))_{\lambda \in \Lambda} \rightarrow T^*(\delta_{x_0})$ . But  $T^*(\delta_{x_\lambda}) = \theta(x_\lambda)\delta_{x_\lambda}$ , so the only possible accumulation points of the net  $(T^*(\delta_{x_\lambda}))_{\lambda \in \Lambda}$  are  $+\delta_{x_0}$  and  $-\delta_{x_0}$ . Therefore,  $x_0 \in D_0$  as claimed.

**Claim 5.**  $\theta$  is continuous on  $K \setminus L$ .

**Proof of Claim 5.** We fix  $x_0 \in K \setminus L$  and an open subset  $W$  of  $K$  such that  $x_0 \in W \subseteq \bar{W} \subseteq K \setminus L$  and we take a function  $f \in C_0(K\|L) \subseteq C_E(K\|L)$  satisfying  $f|_{\bar{W}} \equiv 1$ . Since the mapping

$$x \mapsto \psi(x) = [T^*(\delta_x)](f) = \theta(x)f(x) \quad (x \in K \setminus L)$$

is continuous and  $\psi|_W \equiv \theta|_W$ , we get the continuity of  $\theta$  at  $x_0$ . □

We are now able to completely describe the set of surjective isometries in some special cases. The first one covers the case when  $K$  and  $K \setminus L$  are connected.

**Corollary 5.2.** *Let  $K$  be a connected compact space such that  $K \setminus L$  is also connected. Suppose that  $C_E(K\|L)$  is extremely non-complex. Then,  $\text{Iso}(C_E(K\|L)) = \{\text{Id}, -\text{Id}\}$ .*

**Proof.** Given  $T \in \text{Iso}(C_E(K\|L))$ , Theorem 5.1 gives a continuous function  $\theta : K \setminus L \rightarrow \{-1, 1\}$  such that  $[T(f)](x) = \theta(x)f(x)$  for every  $x \in K \setminus L$  and every  $f \in C_E(K\|L)$ . If  $K \setminus L$  is connected, there are only two possible functions  $\theta$ . Being  $L$  nowhere dense, the values of  $[T(f)](x)$  for  $x \in K \setminus L$  determine completely the function  $T(f)$  for every  $f \in C_E(K\|L)$ . This gives only two possible surjective isometries,  $\text{Id}$  and  $-\text{Id}$ . □

**Corollary 5.3.** *Suppose  $E$  is a subspace of  $C(L)$  such that  $C_E(K\|L)$  is extremely non-complex and for every  $x \in L$ , there is  $f \in E$  such that  $f(x) \neq 0$ . If  $T \in \text{Iso}(C_E(K\|L))$ , then there is a continuous function  $\theta : K \rightarrow \{-1, 1\}$  such that  $T(f) = \theta f$  for all  $f \in C_E(K\|L)$ .*

**Proof.** By Theorem 5.1, we may find  $\theta' : K \setminus L \rightarrow \{-1, 1\}$  continuous such that

$$[T(f)](x) = \theta'(x)f(x) \quad (x \in K \setminus L, f \in C_E(K\|L)).$$

First, we note that  $\theta'$  can be extended to a continuous function  $\theta$  on  $K$  (indeed, if  $x \in L$ , there is an open neighbourhood  $U$  of  $x$  on  $K$  and an  $f \in C_E(K\|L)$  such that  $f(y) \neq 0$  for every  $y \in U$  and  $T(f)|_U/f|_U$  is a continuous function on  $U$  which extends  $\theta'|_{U \setminus L}$ ). Now, for each  $f \in C_E(K\|L)$  we have

$$[T(f)](x) = \theta(x)f(x) \quad (x \in K \setminus L),$$

so  $T(f) = \theta f$  since they are two continuous functions which agree on a dense set. □

By just taking  $E = C(L)$  in the above result, we get a description of all surjective isometries on an extremely non-complex  $C(K)$  space. One direction is the above corollary, the converse result is just a consequence of the classical Banach–Stone Theorem (see, for example, [12, Theorem 2.1.1]).

**Corollary 5.4.** *Let  $K$  be a perfect Hausdorff space such that  $C(K)$  is extremely non-complex. If  $T \in \text{Iso}(C(K))$ , then there is a continuous function  $\theta : K \rightarrow \{-1, 1\}$  such that  $T(f) = \theta f$  for every  $f \in C(K)$ . Conversely, for every continuous function  $\theta' : K \rightarrow \{-1, 1\}$ , the operator given by  $T(f) = \theta' f$  for every  $f \in C(K)$  is a surjective isometry. In other words,  $\text{Iso}(C(K))$  is isomorphic to the Boolean algebra of clopen subsets of  $K$ .*

It follows from the above result and the Banach–Stone Theorem on the representation of surjective isometries on  $C(K)$  (see, for example, [12, Theorem 1.2.2]) that the only homeomorphism of  $K$  is the identity.

**Corollary 5.5.** *Let  $K$  be a perfect Hausdorff space such that  $C(K)$  is extremely non-complex. Then, the unique homeomorphism from  $K$  onto  $K$  is the identity.*

We finish the section with the study of the opposite extreme case, i.e. when  $E = \{0\}$ . Then, the hypotheses of Corollary 5.3 are not satisfied, but we obtain a description of the surjective isometries of the spaces  $C_0(K\|L) \equiv C_0(K \setminus L)$  directly from Theorem 5.1. Again, the converse result comes from the Banach–Stone Theorem (see, for example, [12, Corollary 2.3.12]).

**Corollary 5.6.** *Let  $K$  be a compact Hausdorff space,  $L \subset K$  closed nowhere dense, and suppose that  $C_0(K \setminus L)$  is extremely non-complex. If  $T \in \text{Iso}(C_0(K \setminus L))$ , then there is a continuous function  $\theta : K \setminus L \rightarrow \{-1, 1\}$  such that  $T(f) = \theta f$  for every  $f \in C_0(K \setminus L)$ . Conversely, for every continuous function  $\theta' : K \setminus L \rightarrow \{-1, 1\}$ , the operator*

$$[T(f)](x) = \theta'(x)f(x) \quad (x \in K \setminus L, f \in C_0(K \setminus L))$$

*is a surjective isometry. In other words,  $\text{Iso}(C_0(K \setminus L))$  is isomorphic to the Boolean algebra of clopen subsets of  $K \setminus L$ .*

**Proof.** The first part is a direct consequence of Theorem 5.1 for  $E = \{0\}$ . For the converse result, just observe that given any extension of  $\theta'$  to  $L$ , the product  $\theta' f : K \rightarrow \mathbb{R}$  does not depend on the extension, belongs to  $C_0(K\|L)$  and  $\|\theta' f\|_\infty = \|f\|_\infty$ . □

### 6. The construction of the main example

Our goal here is to construct a compact space  $K$  and a nowhere-dense subset  $L \subseteq K$  with very special properties which will allow us to provide the main example on surjective isometries and duality.

**Theorem 6.1.** *There exist a compact space  $K$  and a closed nowhere-dense subset  $L \subseteq K$  with the following properties:*

- (a)  $K$  and  $K \setminus L$  are connected;
- (b) there is a continuous mapping  $\phi$  from  $L$  onto the Cantor set; and
- (c) every operator on  $C(K)$  is a weak multiplier.

**Proof.**  $K$  is the compact space constructed in [19, § 5]. The fact that all operators on  $C(K)$  are weak multipliers is given in [19, Lemma 5.2].

We just need to find the appropriate  $L$ . We will assume the familiarity of the reader with the above construction of  $K$  in [19, § 5]. In particular, that  $K \subseteq [0, 1]^{2^\omega}$  is the inverse limit of  $K_\alpha \subseteq [0, 1]^\alpha$  for  $\alpha \leq 2^\omega$  where  $K_1 = [0, 1]^2$ . For  $\beta \leq \alpha \leq 2^\omega$  the projection from  $[0, 1]^\alpha$  onto  $[0, 1]^\beta$  is denoted  $\pi_{\beta, \alpha}$ .

Choose any  $N \subseteq [0, 1]^2$  which is a copy of a Cantor set included in some subinterval of  $[0, 1]^2$ . In particular, it is compact nowhere-dense perfect and such that  $[0, 1]^2 \setminus N$  is connected. Let  $N_\alpha = \pi_{1, \alpha}[N]$ . We claim that  $L = N_{2^\omega}$  works, i.e. is nowhere dense in  $K$ ,  $K \setminus L$  is connected and there is a continuous mapping of  $L$  onto the Cantor set. The last part is clear as  $\pi_{1, 2^\omega}$  sends  $L$  onto  $N$  which is a homeomorphic copy of the Cantor set.

One proves by induction on  $\alpha \leq 2^\omega$  that  $N_\alpha$  is nowhere dense in  $K_\alpha$  and  $K_\alpha \setminus N_\alpha$  is connected. This is essentially a generalization of [19, Lemma 4.6] from a finite set to a nowhere-dense set with a connected complement in  $[0, 1]^2$ .

Lemma 4.3.a of [19] says that being nowhere dense is preserved when we pass by preimages from  $K_\alpha$  to  $K_{\alpha+1}$  so, as the limit stage is trivial, it follows that every  $N_\alpha$  is nowhere dense in  $K_\alpha$ . Therefore,  $L$  is nowhere dense in  $K$ .

So we are left with showing that  $K_\alpha \setminus N_\alpha$  are connected. As in [19, Lemma 4.6], we prove by induction on  $\alpha$  that there are  $M_\alpha^n \subseteq K_\alpha$  such that

- (1)  $\pi_{\alpha', \alpha}[M_\alpha^n] = M_{\alpha'}^n$  for  $\alpha' \leq \alpha \leq \beta$ ,
- (2) the  $M_\alpha^n$ s are compact and connected,
- (3)  $M_\alpha^n \cap N_\alpha = \emptyset$ ,  $M_\alpha^n \subseteq M_\alpha^{n+1}$ ,
- (4)  $\bigcup_{n \in \mathbb{N}} M_\alpha^n$  is dense in  $K_\alpha \setminus N_\alpha$ .

We start by choosing  $M_1^n$  to satisfy (2)–(4) and such that  $[0, 1]^2 \setminus \bigcup_{n \in \mathbb{N}} M_\alpha^n$  is  $N = N_1 \subseteq [0, 1]^2$ . The rest of the argument is exactly as in the last part of the proof of [19, Lemma 4.6]. □

We are now ready to present the main application of the results of the paper.

**Theorem 6.2.** *For every separable Banach space  $E$ , there is a Banach space  $\tilde{X}(E)$  such that  $\text{Iso}(\tilde{X}(E)) = \{\text{Id}, -\text{Id}\}$  and  $\tilde{X}(E)^* = E^* \oplus_1 Z$  for a suitable space  $Z$ . In particular,  $\text{Iso}(\tilde{X}(E)^*)$  contains  $\text{Iso}(E^*)$  as a subgroup.*

**Proof.** Consider the compact space  $K$  and the nowhere-dense closed subset  $L \subset K$  given in Theorem 6.1. As there is a surjective continuous function from  $L$  to the Cantor set, every separable Banach space  $E$  is a subset of  $C(L)$ . Let  $\tilde{X}(E)$  be  $C_E(K\|L)$ . Then,  $\tilde{X}(E)$  is extremely non-complex since every operator on  $C(K)$  is a weak multiplier and we may use Theorem 3.9. Now, since  $K \setminus L$  is connected, we may apply Corollary 5.2 to get that  $\text{Iso}(\tilde{X}(E)) = \{\text{Id}, -\text{Id}\}$ . Finally, Lemma 2.2 gives us that  $\tilde{X}(E)^* = E^* \oplus_1 C_0(K\|L)^*$  and so  $\text{Iso}(\tilde{X}(E)^*)$  contains  $\text{Iso}(E^*)$  as a subgroup (see, for example, [21, Proposition 2.4]). □

Let us comment that all the spaces  $\tilde{X}(E)$  constructed above are non-separable. We do not know whether separable spaces with the same properties can be constructed.

The case  $E = \ell_2$  in Theorem 6.2 gives the following specially interesting example.

**Example 6.3.** There is a Banach space  $\tilde{X}(\ell_2)$  such that  $\text{Iso}(\tilde{X}(\ell_2)) = \{\text{Id}, -\text{Id}\}$  but  $\text{Iso}(\tilde{X}(\ell_2)^*)$  contains  $\text{Iso}(\ell_2)$  as a subgroup. Therefore,  $\text{Iso}(\tilde{X}(\ell_2))$  is trivial, while  $\text{Iso}(\tilde{X}(\ell_2)^*)$  contains infinitely many uniformly continuous one-parameter semigroups of surjective isometries.

Recently, the second author of this paper constructed a Banach space  $X(\ell_2)$  such that  $\text{Iso}(X(\ell_2))$  does not contain any *uniformly continuous* one-parameter semigroup of surjective isometries, while  $\text{Iso}(X(\ell_2)^*)$  contains infinitely many of them [21, Example 4.1]. But it is not difficult to show that  $\text{Iso}(X(\ell_2))$  does not reduce to  $\{\text{Id}, -\text{Id}\}$  and, actually, it contains infinitely many *strongly continuous* one-parameter semigroups of surjective isometries. We refer the reader to the books of Engel and Nagel [7, 8] for background on one-parameter semigroups of operators and to the monographs of Fleming and Jamison [12, 13] for more information on isometries on Banach spaces.

**Acknowledgements.** P.K. was partially supported by Polish Ministry of Science and Higher Education Research Grant N N201 386234. M.M. and J.M. were partially supported by Spanish MEC Project MTM2006-04837 and Junta de Andalucía Grants FQM-185 and P06-FQM-01438.

The authors would like to thank Vladimir Kadets, Rafael Ortega, Rafael Payá and Armando Villena for fruitful conversations on the subject of the paper. Part of this work was done while M.M. was visiting the Technical University of Łódź in July 2008 and also while P.K. was visiting the University of Granada in September 2008. Both authors would like to thank the hosting institutions for partial financial support.

## References

1. Y. ABRAMOVICH AND C. ALIPRANTIS, *An invitation to operator theory*, Graduate Texts in Mathematics, Volume 50 (American Mathematical Society, Providence, RI, 2002).
2. Y. ABRAMOVICH AND C. ALIPRANTIS, *Problems in operator theory*, Graduate Texts in Mathematics, Volume 51 (American Mathematical Society, Providence, RI, 2002).
3. F. ALBIAC AND N. J. KALTON, *Topics in Banach space theory*, Graduate Texts in Mathematics, Volume 233 (Springer, 2006).
4. I. K. DAUGAVET, On a property of completely continuous operators in the space  $C$ , *Usp. Mat. Nauk* **18** (1963), 157–158 (in Russian).
5. J. DIESTEL, *Sequences and series in Banach spaces*, Graduate Texts in Mathematics, Volume 92 (Springer, 1984).
6. J. DIEUDONNÉ, Complex structures on real Banach spaces, *Proc. Am. Math. Soc.* **3** (1952), 162–164.
7. K. J. ENGEL AND R. NAGEL, *One-parameter semigroups for linear evolution equations*, Graduate Texts in Mathematics, Volume 194 (Springer, 2000).
8. K. J. ENGEL AND R. NAGEL, *A short course on operator semigroups*, Universitext (Springer, 2006).
9. R. FAJARDO, Consistent constructions of Banach spaces  $C(K)$  with few operators, PhD Thesis, Universidade de São Paulo (2007).

10. V. FERENCZI, Uniqueness of complex structure and real hereditarily indecomposable Banach spaces, *Adv. Math.* **213** (2007), 462–488.
11. V. FERENCZI AND E. MEDINA GALEGO, Even infinite-dimensional real Banach spaces, *J. Funct. Analysis* **253** (2007), 534–549.
12. R. FLEMING AND J. JAMISON, *Isometries on Banach spaces: function spaces*, Monographs and Surveys in Pure and Applied Mathematics, Volume 129 (Chapman & Hall/CRC, Boca Raton, FL, 2003).
13. R. FLEMING AND J. JAMISON, *Isometries on Banach spaces: vector-valued function spaces and operator spaces*, Monographs and Surveys in Pure and Applied Mathematics, Volume 132 (Chapman & Hall/CRC, Boca Raton, FL, 2008).
14. G. GODEFROY, Existence and uniqueness of isometric preduals: a survey, in *Banach Space Theory, Iowa City, IA, 1987*, Contemporary Mathematics, Volume 85, pp. 131–193 (American Mathematical Society, Providence, RI, 1989).
15. P. HARMAND, D. WERNER AND W. WERNER, *M-ideals in Banach spaces and Banach algebras*, Lecture Notes in Mathematics, Volume 1547 (Springer, 1993).
16. V. KADETS, M. MARTÍN AND J. MERÍ, Norm equalities for operators on Banach spaces, *Indiana Univ. Math. J.* **56** (2007), 2385–2411.
17. V. KADETS, R. SHVIDKOY, G. SIROTKIN AND D. WERNER, Banach spaces with the Daugavet property, *Trans. Am. Math. Soc.* **352** (2000), 855–873.
18. S. KOPPELBERG, *Handbook of Boolean algebras*, Volume 1 (North-Holland, Amsterdam, 1989).
19. P. KOSZMIDER, Banach spaces of continuous functions with few operators, *Math. Annalen* **330** (2004), 151–183.
20. P. KOSZMIDER, M. MARTÍN AND J. MERÍ, Extremely non-complex  $C(K)$  spaces, *J. Math. Analysis Applic.* **350** (2009), 584–598.
21. M. MARTÍN, The group of isometries of a Banach space and duality, *J. Funct. Analysis* **255** (2008), 2966–2976.
22. M. E. MUNROE, *Measure and integration*, 2nd edn (Addison-Wesley, Reading, MA, 1971).
23. T. OIKHBERG, Some properties related to the Daugavet property, in *Banach spaces and their applications in analysis*, pp. 399–401 (Walter de Gruyter, Berlin, 2007).
24. S. SAKAI,  *$C^*$ -algebras and  $W^*$ -algebras* (Springer, 1971).
25. Z. SEMADENI, *Banach spaces of Continuous functions* Volume I, Monografie Matematyczne, Tom 55 (PWN/Polish Scientific Publishers, Warsaw, 1971).
26. D. WERNER, An elementary approach to the Daugavet equation, in *Interaction between functional analysis, harmonic analysis and probability* (ed. N. Kalton, E. Saab and S. Montgomery-Smith), Lecture Notes in Pure and Applied Mathematics, Volume 175, pp. 449–454 (CRC Press, Boca Raton, FL, 1994).
27. D. WERNER, Recent progress on the Daugavet property, *Irish Math. Soc. Bull.* **46** (2001), 77–97.