

Original Article

*Contributed equally to the manuscript.

Cite this article: Bécaries L, Dewey ME, Das-Munshi J (2018). Ethnic density effects for adult mental health: systematic review and meta-analysis of international studies. *Psychological Medicine* **48**, 2054–2072. <https://doi.org/10.1017/S0033291717003580>

Received: 6 July 2017

Revised: 3 October 2017

Accepted: 9 November 2017

First published online: 14 December 2017

Key words:

race; ethnicity; ethnic density; common mental disorders; depression; anxiety; psychosis; psychotic experiences; suicidal ideas; suicide; mental health; systematic review; meta-analysis

Author for correspondence:

Laia Bécaries, E-mail: Laia.becaries@manchester.ac.uk

Ethnic density effects for adult mental health: systematic review and meta-analysis of international studies

Laia Bécaries^{1*}, Michael E. Dewey² and Jayati Das-Munshi^{2*}

¹Department of Social Statistics, University of Manchester, Manchester, UK and ²Department of Health Services & Population Research, King's College London, Institute of Psychiatry, Psychology & Neuroscience, London, UK

Abstract

Background. Despite increased ethnic diversity in more economically developed countries it is unclear whether residential concentration of ethnic minority people (ethnic density) is detrimental or protective for mental health. This is the first systematic review and meta-analysis covering the international literature, assessing ethnic density associations with mental health outcomes.

Methods. We systematically searched Medline, PsychINFO, Sociological Abstracts, Web of Science from inception to 31 March 2016. We obtained additional data from study authors. We conducted random-effects meta-analysis taking into account clustering of estimates within datasets. Meta-regression assessed heterogeneity in studies due to ethnicity, country, generation, and area-level deprivation. Our main exposure was ethnic density, defined as the residential concentration of own racial/ethnic minority group. Outcomes included depression, anxiety and the common mental disorders (CMD), suicide, suicidality, psychotic experiences, and psychosis.

Results. We included 41 studies in the review, with meta-analysis of 12 studies. In the meta-analyses, we found a large reduction in relative odds of psychotic experiences [odds ratio (OR) 0.82 (95% confidence interval (CI) 0.76–0.89)] and suicidal ideation [OR 0.88 (95% CI 0.79–0.98)] for each 10 percentage-point increase in own ethnic density. For CMD, depression, and anxiety, associations were indicative of protective effects of own ethnic density; however, results were not statistically significant. Findings from narrative review were consistent with those of the meta-analysis.

Conclusions. The findings support consistent protective ethnic density associations across countries and racial/ethnic minority populations as well as mental health outcomes. This may suggest the importance of the social environment in patterning detrimental mental health outcomes in marginalized and excluded population groups.

Introduction

Mental health disorders account for 8.6 million years of life lost to premature mortality, and are the leading cause of years lived with disability worldwide (Whiteford *et al.* 2013), ranking second only to cardiovascular disease in their impact on disability in high-income countries (Murray & Lopez, 1996). There is some heterogeneity in the incidence and prevalence of mental disorders across racial/ethnic populations, and long-standing controversies in the etiological origins of these differences. Whereas a body of research has focused on the intrinsic biologic variations among human subpopulations (Kaufman & Cooper, 2001), a parallel line of research has examined the social determinants of mental health and has indicated that social factors rooted in the experience of belonging to a racial/ethnic minority, marginalized group, has a significant effect on adverse mental health outcomes (Pickett & Wilkinson, 2008).

A burgeoning body of research suggests that for racial/ethnic minority populations, there might be benefits to living in neighborhoods with a greater concentration of other racial/ethnic minority residents, a so-called ethnic density effect (Shaw *et al.* 2012; Bécaries *et al.* 2012b). Studies suggest that ethnic density – the residential concentration of racial/ethnic minorities in a neighborhood – protects racial/ethnic minorities from the stress of racial discrimination, low status stigma, and socioeconomic disadvantage, while encouraging stronger social support and community cohesion. Understanding the contribution of ethnic density effects for mental health is important because it highlights the role of the environment, including racialization and exclusion, in maintaining and perpetuating the increased risk of mental health disorders among racial/ethnic minority populations.

However, this body of work is contentious, as protective effects are offset by the detrimental effects of concentrated poverty and disinvestment in areas with larger concentrations of racial/ethnic minority people. Previous research has been unable to disentangle the competing effects for mental health of ethnic density and area deprivation, and there are challenges in

attempting to draw conclusions from diverse, international settings with distinct historical patterns of settlement and diversity. A previous narrative review was not able to quantify the magnitude of differences for mental health by ethnic density (Shaw & Pickett, 2011), and sources of potential heterogeneity have hitherto remain unexplored, although understanding these may lead to important clues around underlying mechanisms for mental disorder. These issues are particularly salient given the current scale of global migration (United Nations, 2016) and policies around the forced dispersal of migrant populations to areas without previous histories of migrant settlement.

To address these controversies, we conducted a systematic review with meta-analysis and meta-regression of observational studies of mental disorders where ethnic density effects have been examined. The aim of the review and meta-analysis was to establish: (1) if ethnic density associations are evident, whether they are consistent across mental health outcomes including depression, anxiety and the common mental disorders (CMD), suicide, suicidality, psychotic experiences, and psychosis; (2) the nature and magnitude of ethnic density effects on mental disorders across and within countries, racial/ethnic groups, generational status, and by area-level deprivation.

Methods

Search strategy

Search strategies were based on and updated from a narrative review of ethnic density and mental health (Shaw & Pickett, 2011) that searched the following databases: Medline, PsychINFO, Sociological Abstracts, and the Social Science Citation and Science citations from the Web of Science. We performed searches to March 2016. All reference lists for retrieved papers with potential studies for inclusion were hand-searched and first and/or last authors of all papers were contacted for further additional and unpublished data.

Eligibility

We excluded ecological studies, and studies that did not focus on adult (aged 16+) populations. We included all cross-sectional survey, case-control, or cohort-design studies that used representative population data, in order to minimize biases from differences in help-seeking from using patient-/service-level data. In order to further minimize ascertainment bias due to differences in health-seeking behavior, we restricted to studies which used community samples. As schizophreniform/psychotic disorders are rare and not usually detected in community samples for these disorders, we also included first contact studies which utilized leakage studies to identify all cases of schizophreniform disorders/psychosis, or alternatively studies which employed databases of whole populations with linkage. We also included studies of suicide which employed death certificate information, since there is no other way to ascertain this outcome from community survey-level data.

Main outcomes and measures

Ethnic density was measured as 10% increase in the residential concentration of own racial/ethnic group, for the purposes of meta-analysis. Studies had to have a measure of ethnic density and had to specify the geographical area of measurement, which had to be at the level of neighborhoods. Studies could be

in a journal or book, but we also included doctoral dissertations and unpublished estimates obtained from contacting authors. Outcomes were depression, anxiety, suicidality and suicide, psychotic experiences, and schizophreniform/psychotic disorders. For some studies, it was not possible to examine depression and anxiety separately, in these instances we opted to analyze the outcome of CMD (anxiety or depression).

Search and extraction

Searches were conducted by two of the authors (JD-M and LB). Retrieved studies were compared with inclusion/exclusion criteria. Uncertainties around whether to include studies were settled by discussions between the authors. For each of the studies which met criteria for inclusion, relevant fields including medical outcomes, were extracted and cross-checked between two of the investigators (JD-M and LB). Where possible, we extracted estimates for the association of own ethnic density and mental health outcome, adjusted for age, sex, individual-level and area-level deprivation. For studies where data on ethnic density measured as a 10 percentage-point increase in the residential concentration of own ethnic group were not available, study authors were contacted for additional estimates, and in some cases, for unpublished estimates. We considered multiple publications using the same dataset as one set of analyses. There were no restrictions on language of manuscript. We were aware of two other studies for which we were co-investigators, but we could not include the data due to the publications being embargoed at the time of analysis.

Assessment of methodological quality

A small pilot study was conducted to assess the reliability of a scale devised to assess the quality of retrieved studies [based on quality assessment scales of observational studies (Khan *et al.* 2001)]. LB and JD-M independently assessed the quality of a random sample of 20 papers using this scale, blind to each other's quality ratings (see online Supplementary Table S1 for criteria in quality assessment tool). The κ for inter-rater reliability of the scale was 67%. Efforts to avoid publication biases were taken by contacting all first authors and experts in the field for unpublished data. Funnel plots were derived to formally assess this (see online Supplementary Fig. S1). We registered the review protocol with PROSPERO (registration number: CRD42014012992).

Statistical analysis

We conducted meta-analysis followed by meta-regression to examine the association between own ethnic density and the following mental health outcomes: depression, anxiety and the CMD, suicide, suicidality, psychotic experiences, and psychosis. We obtained raw data from some of the studies (Bécares *et al.* 2009; Das-Munshi *et al.* 2010; Das-Munshi *et al.* 2012; Bécares *et al.* 2012a; Bécares & Das-Munshi, 2013; Bécares, 2014; Bécares *et al.* 2014) which permitted us to obtain estimates adjusted for *a priori* confounders, including age and gender, and area-level deprivation, or if area-level deprivation was not available, adjusted only for age and gender. These were combined with estimates extracted from publications or provided by authors of publications, also adjusted for the same confounders. We then used meta-regression to assess the effect of moderators, which included: race/ethnicity, country of study, generational status. We assessed the moderating effect of area-level deprivation

using datasets which had estimates adjusted for age and sex. Meta-regression models were run with fixed effects for moderators, and random effects for study. Studies which used symptom scales for depression were converted into mean differences and their standard errors using the method outlined by Chinn (Chinn, 2000). In each of our models, we derived an I^2 -like statistic for each level of random variability. To assess for bias introduced through small study effects, we used funnel plots. Meta-analysis and meta-regression were conducted in R (Team, 2015), version 3.2.2, using the package *metafor*, version 1.9-8 (Viechtbauer, 2010). To supplement findings from the meta-analysis and meta-regression, we also conducted narrative review synthesis to all papers, which relied on vote counting of studies (Popay *et al.* 2006), based on statistical significance of results.

Results

Our search retrieved 2288 articles, 135 were assessed for eligibility, and 41 studies were included in the synthesis. Of these, we were only able to include 12 studies in meta-analysis because, despite our efforts, most authors either did not respond to our requests, were unable to provide data in suitable format for meta-analysis, or no longer had access to the data (Fig. 1). In total, there were 29 studies assessing CMD, depression, and anxiety (Table 1); 12 studies assessing ethnic density associations for psychotic experiences and psychosis; and four studies assessing suicidality and completed suicide. Table 1 is ordered according to outcome and quality of studies; asterisked studies in the table provided estimates for meta-analysis. All of the studies that were included in

the meta-analysis were of a moderate- to high-quality rating. No studies were excluded based on quality. We also obtained two other datasets from the UK, publicly available under end-user license agreements, which had not been published as part of any study but which we included in meta-analysis and meta-regression. For the meta-analysis and meta-regressions, we focus on datasets and not on studies, reflecting the fact that several studies may have used the same data sources, to avoid duplication. Visual assessment of funnel plots did not indicate publication biases (see online Supplementary Fig. S1).

In general, qualitative assessment of included studies (Table 1) provides support for a protective association between increased own ethnic density and reduced risk of depression, anxiety and the CMD, suicide, suicidality, psychotic experiences, and psychosis; for all outcomes over half of studies supported protective associations. This was particularly evident for psychotic experiences (3/4 studies) and suicide (2/2 studies). Negative associations were identified only in US studies. Out of 30 international studies that assessed the association between ethnic density and CMD, depression, and anxiety, six reported a detrimental effect in African American and Latino populations; all of these studies were from the USA. One of these studies examined the non-linearity of ethnic density and depression in African American people and found that the detrimental association of own ethnic density and depression was only observed at the highest level of ethnic density (85% or greater); at levels below this the association was protective (Bécaries *et al.* 2014).

One UK-based study (out of four) reported detrimental associations between Pakistani ethnic density and psychotic




Fig. 1. PRISMA flow chart of studies.

Table 1. Studies investigating ethnic density effects with mental health outcomes. Studies ranked according to quality assessment by outcome

Reference	Dataset, country	Sample	Instrument	Area unit	Ethnic density measure	Covariates	Method	Results (threshold/non-linearity)	QA score
Depression									
*Arévalo <i>et al.</i> (2015)	Boston Puerto Rican Health Study (USA)	1142 Puerto Ricans from initial sample of 1504 (76%) with complete baseline and follow-up assessment. Aged 45–75	Center for Epidemiologic Studies-Depression (CES-D)	Census tract	Percentage Puerto Ricans	Gender, age, education, depressive symptoms at baseline (for follow-up assessment), Neighborhood poverty level, pre-migration urbanicity, reasons for migration, length of stay, acculturation, discrimination, recreational/friend/neighbor/family/church activities	Multilevel multivariate linear regression	At 2-year follow-up, after adjusting for all covariates and baseline CES-D, men living in high v. low ethnic density neighborhoods had lower CES-D scores but not women	14
Booth (2014)	Chicago Community Adult Health Survey (CCAHS) 2001–2003 (USA)	802 Latino people; 64% Mexican American, 17% Puerto Rican, 19% other Hispanic/Latino	CES-D	343 neighborhood 'clusters' consisting of one or more contiguous census tracts, from 865 aggregated census tracts	Percentage of Hispanic people, continuous according to 2000 US census	Gender, age, education Neighborhood-level disadvantage, generational status	Multilevel regression models	Protective association of increasing own Hispanic density in crude models and models adjusting for individual and area-level covariates excluding generational status and acculturation	14
Alegria <i>et al.</i> (2014)	2001–2003 Collaborative Psychiatric Epidemiology Surveys (CPES) – these included National Comorbidity Survey Replication (NCS-R), the National Survey of American Life (NSAL), the National Latino and Asian American Study (NLAAS) (USA)	4180 Non-Latino white, 2554 Latino, 2095 Asian, 3570 African American and 1438 African Caribbean residents	World Health Organization World Mental Health Composite International Diagnostic Interview (WHO WMH-CIDI)	Census tracts	Percentage Latino and foreign-born population – interacted with race/ethnicity	Individual – race/ethnicity, age, gender, marital status, education, poverty status, work status, and nativity Neighborhood – disadvantage, Latino/immigrant concentration, affluence, and residential instability	Multi-level logistic regression models	In an interaction between individual-level ethnicity (Latino v. non-Latino white), there was not a statistically significant association between increased Latino/immigrant ethnic density and depressive disorders	14
*Bécares <i>et al.</i> (2014)	NSAL; (USA)	3570 African Americans	CES-D	Census tract	Percentage black residents	Age, gender, marital status, household income, education, employment Area deprivation score composed of 4 indicators	Multi-level logistic regression models	Increased black ethnic density up to 85% associated with decreased depressive symptoms Increased black ethnic density from 85% onwards associated with increased depressive symptoms	14

(Continued)

Table 1. (Continued.)

Reference	Dataset, country	Sample	Instrument	Area unit	Ethnic density measure	Covariates	Method	Results (threshold/ non-linearity)	QA score
Mair et al. (2010)	Multi-Ethnic Study of Atherosclerosis (MESA); (USA)	837 African American women, 669 African American men, 608 Hispanic women, 571 Hispanic men, 339 Chinese women, 321 Chinese men	CES-D	Census tract	Percent non-Hispanic African American, Hispanic, non-Hispanic white, and non-Hispanic Asian	Age, income, marital status, education and nativity. Analyses stratified by gender Neighborhood social cohesion, safety, problems, aesthetic quality and socioeconomic factors	Multilevel regression models	Living in a neighborhood with a higher ethnic density was associated with increased CES-D scores in African American men <i>When adjusted for area deprivation, Hispanic ethnic density was association with decreased CES-D scores among Hispanic men</i> No association between ethnic density and depression for the other racial/ethnic and gender categories	13
Wight et al. (2009)	Study of Assets and Health Dynamics Among the Oldest Old (AHEAD); (USA)	3442 older adults at time 1 (17% African American, 6% Hispanic), 2632 at time 2 (16% African American, 6% Hispanic), 1871 at time 3 (16% African American, 6% Hispanic)	8-item CES-D	Census tracts	Proportion of African American residents; proportion of Hispanic residents Numerator is proportion of residents who are over the age of 65 years	Educational attainment, household income, household wealth, gender, age, ethnicity, marital status, religion, assistance with activities of daily living (ADL), heart problems, stroke, a count of other major medical conditions, and cognitive function. Neighborhood-level SES factor based on proportion of residents aged 25+ without a high school degree, households receiving public assistance income, residents living below the poverty level, and unemployed residents aged 16+ Affluence indicator based on proportion of households with incomes of \$ 50 000 or more	Multilevel regression models	There was a trend for a protective, but not statistically significant association between Hispanic ethnic density and a reduction in depressive symptoms from time 1 to time 2, and from time 2 to time 3 There were no statistically significant associations for African American ethnic density, although the direction of the effect was detrimental from time 1 to time 2, and protective from time 2 to time 3	13
Shell et al. (2013)	Texas City Stress and Health Study; (USA)	1238 US-born and Mexican-born Hispanics of Mexican descent	Center for Epidemiologic Studies Depression Revised scale (CES-DR)	48 neighborhoods, with boundaries defined using street patterns, residential patterns	Percentage of Hispanic people in the neighborhood based on US Census 2000 data, in 6 groups: ≥16% >16% to ≤20%	Gender, age, education, income, marital status, language spoken, life events, physical health, social support, discrimination, stress	Multilevel regression models	<i>Living in the highest category of Hispanic ethnic density is associated with a reduction in depressive symptoms in models that adjust for individual-level</i>	12.5

					>20 to ≤25% >25 to ≤35% >35 to ≤45% >45%	Residential stability, median house value		<i>demographic covariates and area level covariates</i> This association loses statistical significance after adjusting for social support	
*Ostir <i>et al.</i> (2003)	Hispanic established population for the epidemiologic study of the elderly (HEPSE); (USA)	2710 Mexican Americans aged 65 years or older	CES-D. Subjects with a score of 16 or more were classified as having high depressive symptomatology	Census tracts	Percentage Mexican American	Age, gender, marital status, highest grade of schooling completed, per capita household income, nativity, chronic disease and ADL limitations. Neighborhood socioeconomic status	Multilevel regression models	There was a non-significant association between Mexican ethnic density and depression in models adjusted for age and sex only <i>After adjusting for individual-level covariates and for area deprivation, a 10% increase in percentage Mexican American was associated with reduced CES-D scores</i>	12
Wight <i>et al.</i> (2011)	Health and Retirement Study (HRS); (USA)	4805 adults aged 52– 63. 77% white, 13.18% African American, 7.37% Hispanic, 2.45% Other	CES-D	Census tracts	Percentage of African American residents, and percentage of Hispanic residents	Education, household wealth, household income, religion, gender, age, ethnicity, marital status, ADL, medical conditions. Socioeconomic disadvantage factor, neighborhood affluence	Multilevel regression models	No associations were found between ethnic density and depression	11.5
English <i>et al.</i> (2014)	Healthy Aging in Neighborhoods of Diversity across the Life Span study (HANDLS); (USA)	2197 African American and 1523 white American adults aged 30–64 years	CES-D	Census tract	Proportion African American and white American residents	Age, gender, poverty status (family income)	Structural Equation Modelling (SEM)	No association was found between African American ethnic density and depression, after adjusting for covariates	11
Kwag <i>et al.</i> (2011)	Own data; (USA)	567 Korean American adults aged 60+	Geriatric Depression Scale-Short Form (GDS-SF)	Census block group	Proportion of racial/ethnic minorities-comprising 'Hispanic, non-Hispanic white, non-Hispanic black, Asian, native Hawaiian and Pacific islander, American Indian, Alaskan native, and non-Hispanic multi-race'	Age, gender, marital status, education, length of stay (USA), chronic conditions, functional disability, health perceptions Area-level proportion people living below poverty-line, proportion people >65 years, proportion ethnic minorities	Multilevel multivariate linear regression	No association between proportion racial/ethnic minorities and mean depression after adjusting for covariates	11
Vogt Yuan (2007)	1995 Community, Crime, and Health Survey Illinois; (USA)	2292 black and Hispanic respondents	CES-D	Census tract	Percentage black and percentage Hispanic (logged)	Age, gender, education, household income, employment status, number of children, and marital status. Neighborhood-level: % households below federal poverty threshold and % female headed households with children	Multilevel multiple linear regression models	<i>Black Americans living in areas of higher own group density had lower mean depression in models adjusted for neighborhood disadvantage</i>	11

(Continued)

Table 1. (Continued.)

Reference	Dataset, country	Sample	Instrument	Area unit	Ethnic density measure	Covariates	Method	Results (threshold/non-linearity)	QA score
Aneshensel et al. (2007)	AHEAD; (USA)	3442 older adults (10% African American, 4% Hispanic)	8-item CES-D	Census tracts	Proportion of African American residents; proportion of Hispanic residents	Gender, age, marital status, educational attainment, household wealth, household income, religion, assistance with ADL, number of medical conditions, heart problems, stroke, cognition Neighborhood-level SES factor based on proportion of residents aged 25+ without a high school degree, households receiving public assistance income, residents living below the poverty level, and unemployed residents aged 16+ Affluence indicator based on proportion of households with incomes of \$50 000 or more	Multilevel regression models Tested census-tract variables for non-linear associations, especially threshold effects, but found no significant departures from linearity	No associations between ethnic density and depressive symptoms	11
Stafford et al. (2011)	2000–2001 Canadian Community Health Survey (CCHS); (Canada)	108 064 participants	Composite Diagnostic Interview Schedule Short Form for major depression	Health region	Percentage immigrant population	Immigrant status, length of residence, visible minority status, gender, age, marital status, tenure, education	Multilevel regression models	<i>An increase in immigrant ethnic density was associated with lower likelihood of depression among immigrants and visible minority participants</i>	9
Anxiety									
Alegria et al. (2014)	2001–2003 CPES, NCS-R, NSAL, NLAAS; (USA)	4180 Non-Latino white, 2554 Latino, 2095 Asian, 3570 African American and 1438 African Caribbean residents	WHO WMH-CIDI	Census tracts	Percentage Latino and foreign-born population – interacted with race/ethnicity	Individual – race/ethnicity, age, gender, marital status, education, poverty status, work status, and nativity Neighborhood – disadvantage, Latino/immigrant concentration, affluence, and residential instability	Multi-level logistic regression models	In the interaction model of Latino v. non-Latino white, there was an adverse association between increased Latino/immigrant ethnic density and any-past year anxiety disorders, compared to non-Latino whites	14
Menezes et al. (2011)	2002 CCHS; (Canada)	35 708 individuals of which 21.8% immigrant and 78.2% non-immigrant	WHO WMH-CIDI	Dissemination area (DA)	Percentage immigrants	Age, gender, income, marital status, education. Mean income, % moving in the past 5 years, % rental, and % below low income cut-off thresholds	Multilevel multivariate logistic regression models	<i>Immigrants in the sample had reduced relative odds of mood and anxiety disorders with increasing immigrant density</i>	12

Common mental disorders									
*Das-Munshi <i>et al.</i> (2010); *Bécares & Das-Munshi (2013)	Ethnic Minorities Psychiatric Illness Rates In the Community (EMPIRIC); (UK)	837 white British, 733 Irish, 694 black Caribbean people, 650 Bangladeshi, 643 Indian and 724 Pakistani people	Clinical Interview Schedule-Revised (CIS-R)	Middle super output level	Percentage people of same ethnic group	Age, gender, social class, education, marital status, area-level deprivation	Multilevel multivariate logistic regression	<i>Protective associations for common mental disorders in Irish and Bangladeshi people</i> No associations were found for other ethnic groups	15
*Bécares (2014)	NLAAS; (USA)	577 Cubans, 495 Puerto Ricans, 868 Mexican Americans	6-item Kessler Psychological Distress scale (K6)	US County	Percentage Cuban residents, percentage Puerto Rican residents, percentage Mexican American residents, percentage Latin American immigrant residents, percentage Latino residents	Age, gender, household income, education, work status, marital status, language of interview, nativity. Area deprivation score composed of 3 indicators	Structural equation models accounting for clustering	Increased Cuban ethnic density was not associated with psychological distress for Cuban respondents Increased Puerto Rican ethnic density was not associated with psychological distress for Puerto Rican respondents, but increased immigrant and Latino ethnic density were associated with decreased psychological distress for Puerto Rican respondents. These associations were stronger for the second generation Increased Mexican American ethnic density was not associated with psychological distress among Mexican American participants. Among the first generation, an increase in own, immigrant and overall Latino ethnic density was associated with increased psychological distress	15
Alegria <i>et al.</i> (2014)	2001–2003 CPES, NCS-R, NSAL, NLAAS; (USA)	4180 Non-Latino white, 2554 Latino, 2095 Asian, 3570 African American and 1438 African Caribbean residents	WHO WMH-CIDI	Census tracts	Percentage Latino and foreign-born population	Individual – race/ethnicity, age, gender, marital status, education, poverty status, work status, and nativity Neighborhood – disadvantage, Latino/immigrant concentration, affluence, and residential instability	Multi-level logistic regression models	Latinos residing in neighborhoods with higher Latino/immigrant ethnic density had increased risk of any past year common mental disorder, compared to non-Latino whites.	14
*Bécares <i>et al.</i> (2013)	New Zealand Health Survey (NZHS); (New Zealand)	3160 Maori people	Doctor-diagnosed common mental disorders and 10-item Kessler Psychological Distress scale (K10)	Census area units	Percentage Maori residents	Age, gender, education, household income and work status, area-level deprivation	Multilevel multivariate logistic and linear regression	<i>Increasing ethnic density was associated with a reduction in reporting doctor-diagnosed common mental disorders</i> Ethnic density not associated with non-specific psychological distress (K10)	13

(Continued)

Table 1. (Continued.)

Reference	Dataset, country	Sample	Instrument	Area unit	Ethnic density measure	Covariates	Method	Results (threshold/ non-linearity)	QA score
Choi et al. (2016)	2004–2007 Medical Expenditure Panel Survey linked to county-level data from the Area Health Resources Files; (USA)	29 011 individuals categorized into US-born individuals ($n = 24\,225$); earlier immigrants (≥ 15 years in the USA; $n = 3866$); and recent immigrants (< 15 years in the USA; $n = 920$)	6-item Kessler Scale (K-6) and the Patient Health Questionnaire (PHQ-2)	County	Percentage of foreign-born individuals in the county (continuous variable)	Age, gender, education, income, race/ethnicity, marital status, limiting illnesses % foreign born in county; % uninsured in county; % unemployed in county; population density per square mile; % in poverty in county; % families with female head in county	Multilevel linear regression with interactions between immigrant status and the percentage of foreign-born individuals in the county	<i>Increased foreign-born density associated with worse mental health among earlier immigrants when mental health was measured with the K6</i> <i>When mental health was measured with the PHQ-2, increased foreign-born density was associated with worse mental health for recent and earlier immigrants</i>	12
*Schrier et al. (2014)	G4 Monitor 2008; (The Netherlands)	1206 Surinamese-Dutch, 978 Turkish-Dutch and 784 Moroccan-Dutch citizens	K10	356 neighborhoods. Mean population size of 5853 inhabitants, ranging from 30 to 26 960.	Percentage native Dutch, Percentage Surinamese-Dutch, Percentage Turkish-Dutch Percentage Moroccan-Dutch	Length of stay in the Netherlands, marital status, highest educational qualification, financial situation. Percentage of households in with an income under the national minimum poverty threshold, average house value in the neighborhood	Multilevel regression models	Ethnic density was not associated with non-specific psychological distress in any of the three ethnic minority groups	12
*Jarrin et al. (2013)	Own data; (Spain)	568 Ecuadorians living in Spain, aged 18–55	General Health Questionnaire (GHQ-28), dichotomized at 5+	17 city neighborhoods and 16 municipalities (largely rural) within 4 regions in Spain (Alicante, Almeria, Madrid, Murcia)	Proportion of people with Ecuadorian nationality among all subjects recorded in the municipal council registry within the 33 areas. Ethnic density ranged from 0.9 to 19.5% with a median of 4.7% and a mean of 6.1%. Categorized as: low $> 5.9\%$ high $6\%+$	Gender, age, civil status, number of children, education, social support, social network diversity, emotional support, financial strain, income, employment, time since migration, discrimination. Proportion of people with less than primary education	Multilevel regression models	In unadjusted models, residence in areas of high ethnic density was associated with increased odds of common mental disorders No association between ethnic density and common mental disorders in fully adjusted models	11
Syed & Juan (2012)	NLAAS; (USA)	Vietnamese Americans ($n = 478$), Chinese Americans ($n = 566$), Filipino Americans ($n = 493$)	K10	US county	Percentage Vietnamese, percentage Chinese, percentage Filipino, percentage any Asian American, percentage non-white	Age, gender, nativity, English fluency, schooling, racial discrimination, social cohesion	Multilevel regression models	Vietnamese, Chinese and Filipino ethnic densities were not associated with psychological distress	11
Leu et al. (2011)	NLAAS; (USA)	1583 respondents including Vietnamese, Chinese, Filipino, and 'other Asian' people	WHO WMH-CIDI; 12-month prevalence common mental disorders	Census tract	Number of Asians divided by total population in census tract	Education, household income, subjective social status, employment, family size, marital status, age, ethnicity, citizenship, language, time in the USA, ethnic identification	Multivariate logistic regression	<i>Protective association for common mental disorders of Asian ethnic density in men only</i>	11
Propper et al. (2005)	10 waves of the British Household Panel Survey (BHPS) (1991–2000); (UK)	8184 individuals	General Health Questionnaire (GHQ-12)	Bespoke neighborhoods, composed of approximately	Percent Indian, percent black, percent Pakistani or Bangladeshi	Age, gender, ethnicity, and education, household income, number of adults, number of	Multilevel regression models. Ethnic density analyses examined with	There was a significant interaction for ethnicity of area and ethnicity of the individual. <i>Non-white</i>	10.5

				the nearest 500–800 people to the respondent's home address		children, tenure, employment status of the head of household. Composite index of area characteristics	interaction between individual and area-level ethnicity	<i>participants in more ethnically mixed areas experience less negative mental ill health trajectories than whites</i>	
*Halpern & Nazroo (2000)	Fourth National Survey of ethnic minorities (FNS); (UK)	2867 white British, 1205 black Caribbean, 1273 Indian, 728 African Asian, 1185 Pakistani, 591 Bangladeshi, 214 Chinese people	CIS-R	Ward	Percentage people of same ethnic group	Age, gender, hardship, language, age at migration	Multivariate linear regression (re-analyzed using robust standard errors for meta-analysis)	<i>Ethnic minorities had a lower mean score on CIS-R in areas of higher ethnic density</i> <i>Indian and Caribbean people had lower mean scores on CIS-R in areas of higher ethnic density</i>	10
*Feng <i>et al.</i> (2013)	45 and up study; (Australia)	226 487 adults from 19 ethnic groups	K10	Census collection districts	Percentage of people of same ethnic group	Age and gender	Multilevel multivariate logistic regression	<i>Protective associations for common mental disorders with increasing ethnic density for Australians not born in Australia, English born in the UK, Scottish born in Australia, Irish born in Australia and Chinese born in Australia</i>	9
Ecob & Williams (1991)	Self-collected data in Glasgow from 1986 to 1987; (UK)	173 Asians aged 30–40 years old	General Health Questionnaire (GHQ-12), dichotomized at 4+	Postcode sectors	Proportion of Asian born. Three categories, high (>6%), medium (3 <6%), and low (<3%)	Age, gender, religion, household size, car ownership, house needing repair, number of household durables	Single-level regressions	<i>Protective effect of ethnic density found for mental health (GHQ) and feeling sad or low</i>	7
<i>Psychosis and schizophrenia</i>									
Mezuk <i>et al.</i> (2015)	Swedish population-based registries; (Sweden)	1 442 931 adults age 15–60 years followed from 2005 to 2010	Incident cases of non-affective and affective psychosis identified from psychiatric in-patient and out-patient registries	Small Area Market Statistics (SAMS) from Sweden used to determine ethnic enclaves	Proportion Iraqi people 'enclave' v. 'non-enclave' areas	Age, gender, family income, education attainment, generation status	Multi-level logistic regression models	<i>Iraqi people living in Iraqi enclaves did not have an increased risk of psychosis</i>	16
Kirkbride <i>et al.</i> (2007, 2008)	Aetiology and Ethnicity in Schizophrenia and Other Psychoses study (AESOP); (UK)	218 cases aged 16–64 identified during 5 65 000 person-years of follow-up over a 2-year period (1997–1999)	Schedules for Clinical Assessment in Neuropsychiatry (SCAN) and a modified Personal and Psychiatric History Schedule. Non-affective psychoses (F20–29), schizophrenia (F20), other non-affective psychoses (F21–29)	Census statistical wards (CAS wards)	Proportion of ethnic minorities in ward	Age, gender Population density, ethnic fragmentation, Index of Multiple Deprivation	Multilevel Poisson regression with random effects	<i>Trend for negative association between overall ethnic minority density and risk of schizophrenia for ethnic minority individuals. Association not statistically significant</i>	12
	AESOP and postal survey conducted January and March 2004, and January and March 2006; (UK)	4231 survey participants living in South London (8.2% black Caribbean and 7% black African)						<i>Ethnic minorities living in neighborhoods with the lowest proportion of ethnic minorities had higher risk of schizophrenia than white people. This increased risk was not statistically significant in areas of</i>	

(Continued)

Table 1. (Continued.)

Reference	Dataset, country	Sample	Instrument	Area unit	Ethnic density measure	Covariates	Method	Results (threshold/ non-linearity)	QA score
								medium and high ethnic density	
Menezes et al. (2011)	CCHS 2002; (Canada)	35 708 individuals of which 21.8% immigrant and 78.2% non-immigrant	WMH-CIDI; respondents given screening questions on whether they had a life time diagnosis of schizophrenia or any other psychosis	DA	Percentage immigrants	Age, gender, income, marital status, education. mean income, % moving in the past 5 years, % rental, and % below low income cut-off thresholds	Multilevel multivariate logistic regression models	Very weak evidence supportive of immigrants in the sample having a reduced relative odds of reporting life-time schizophrenia or psychosis diagnosis with increasing immigrant density	12
Kirkbride et al. (2014)	East London First Episode Study (ELFEP); (UK)	484 people with first-episode psychosis, aged 18–65; 68 white British, 38 non-British white, 55 black Caribbean, 49 black African, 53 Bangladeshi people	SCAN: non-affective psychoses (schizophrenia, schizophreniform disorder, and schizoaffective disorder) and affective psychoses	'Statistical wards' based on information from LSOA	SD increase in own group density	Age at first contact, social class, gender, ethnicity	Bayesian hierarchical modeling of standardized incidence ratios	<i>Reduced risk of psychosis in black African people with increasing own group density</i>	10.5
Veling et al. (2008)	Prospective first-contact incidence study of psychotic disorders in the Hague over 7-year period (from 1 April 1997 to 1 April 1999 and 1 October 2000 to 1 October 2005); (The Netherlands)	463 subjects of Moroccan, Surinamese, Turkish, and native Dutch origin	Comprehensive Assessment of Symptoms and History	Neighborhoods classified according to postal codes ($n = 44$). Maximum of 38 000 inhabitants per neighborhood	Percentage immigrant group	Age, gender, marital status, area-level deprivation (average income, housing quality, proportion long-term unemployed, mean educational level)	Multilevel Poisson regression	<i>Compared with native Dutch, immigrants had a higher incidence rate of schizophrenia in low ethnic density neighborhoods, but not in high ethnic density neighborhoods</i> <i>Interactions between ethnic density and ethnicity showed a significant ethnic density effect for Moroccans only, so that elevated incidence rates of psychotic disorders was higher among Moroccan people living in low ethnic density neighborhoods</i>	10
Psychotic experiences									
*Das-Munshi et al. (2012); *Bécáres & Das-Munshi (2013)	EMPIRIC; (UK)	837 white British, 733 Irish, 694 black Caribbean people, 650 Bangladeshi, 643 Indian and 724 Pakistani people	Psychosis Screening Questionnaire (PSQ)	Middle super output level	Percentage people of same ethnic group	Age, gender, social class, education, marital status, area-level deprivation	Multilevel multivariate logistic regression	<i>Strong evidence of increased OR of psychotic experiences with decreasing ethnic density for Indian people</i> <i>Weaker evidence of</i>	15

								increased OR psychotic experiences with decreasing ethnic density for Irish and Bangladeshi people. No association for black Caribbean and Pakistani people	
*Bécares <i>et al.</i> (2009)	FNS; (UK)	2867 white British, 1205 black Caribbean, 1273 Indian, 728 African Asian, 1185 Pakistani, 591 Bangladeshi, 214 Chinese people	PSQ	Electoral ward	Percentage people of same ethnic group	Age, gender, socioeconomic position. A Classification of Residential Neighbourhoods (ACORN) measure of area deprivation	Regression models with robust standard errors	<i>Protective association between ethnic density and psychotic symptomatology for Indian and Bangladeshi respondents</i> Detrimental association for Pakistani respondents	14
*Halpern & Nazroo (2000)	FNS; (UK)	2867 white British, 1205 black Caribbean, 1273 Indian, 728 African Asian, 1185 Pakistani, 591 Bangladeshi, 214 Chinese people	PSQ	Electoral ward	Percentage people of same ethnic group	Age, gender, hardship, language, age at migration	Multivariate linear regression	<i>Ethnic minorities had a lower mean score on PSQ in areas of higher ethnic density</i> <i>Indian, Caribbean, and Bangladeshi people had lower mean PSQ scores in areas of higher ethnic density</i>	10
Suicide									
Termorshuizen <i>et al.</i> (2015)	Population registry linked to the causes of death registry of Statistics Netherlands, 2000–2011; (The Netherlands)	20 240 668 person years of observation (59% Dutch, 13% Other western, 5% Turkish, 5.5% Moroccan, 9.7% Surinamese/Antillean, 7.4% other non-Western)	ICD-10 X60-84, ICD-10 Y10-34	384 neighborhoods in Amsterdam, Rotterdam, Den Haag, and Utrecht	Percentage of observation time in 2000–2011 related to a specific ethnic group	Age, gender, mean neighborhood income	Multilevel Poisson regression	<i>Increasing minority density was associated with a decreased rate ratio (RR) of suicide among immigrants (both Western and non-Western), when compared to Dutch people</i>	8
Neeleman & Wessely (1999)	Coroners' records of suicides in Lewisham, Lambeth, Southwark, Greenwich between 1 January 1991 and 31 December 1993; (UK)	902 008 cases	Deaths considered as suicides if suicide verdicts given, suicide notes found, methods unambiguously indicated suicide and/or communications of suicidal intent had been recorded	109 electoral wards of the London boroughs of Lewisham, Lambeth, Southwark, and Greenwich	Proportions of all ethnic minority groups combined in ward; proportion Afro-Caribbean residents in ward; proportion Asians in ward	Age, gender, local Jarman underprivileged area (UPA) scores excluding ethnicity	Random-effects Poisson regression	<i>Increased ethnic density was associated with lower suicide rates among ethnic minority groups</i>	7
Suicidal ideation									
*Bécares & Das-Munshi (2013)	EMPIRIC; (UK)	837 white British, 733 Irish, 694 black Caribbean people, 650 Bangladeshi, 643	Suicidal ideation	Middle super output level	Percentage people of same ethnic group	Age, gender, social class, education, marital status, generation, area-level deprivation	Multilevel multivariate logistic regression	<i>Protective associations for suicidal ideation in Bangladeshi and Irish people</i> No associations between	15

(Continued)

Table 1. (Continued)

Reference	Dataset, country	Sample	Instrument	Area unit	Ethnic density measure	Covariates	Method	Results (threshold/non-linearity)	QA score
*Bécaries et al. (2012a)	NSAL, (USA) and EMPIRIC, (UK)	Indian, and 724 Pakistani people 1568 black Caribbean American and 661 black English people Aged 18–74	Suicidal ideation	Census tracts, middle super output level	Percentage born in the Caribbean Percentage black Caribbean Percentage black African + black Caribbean + black other	Age, gender, household income, educational qualifications, employment, nativity, marital status, racism and discrimination, area-level deprivation	Multilevel multivariate logistic regression	ethnic density and suicidal ideation for the other ethnic minority groups No associations between any of the two ethnic density measures and suicidal ideation for black Caribbean American people No associations between any of the two ethnic density measures and suicidal ideation for black Caribbean English people	13 15

Bold italic references refer to estimates included in meta-analyses; italic text indicates protective ethnic density effects; bold text indicates detrimental ethnic density effects.

*Studies contributed data to the meta-analyses.

experiences. There were no studies of psychosis and suicidality in the narrative review that reported detrimental associations with ethnic density (Table 1).

In the meta-analysis, we found that for CMD, depression, and anxiety, associations were indicative of protective effects of own ethnic density; however, 95% confidence intervals (CIs) just spanned the null (Figs 2–4). We also observed a large reduction in relative odds of psychotic experiences [odds ratio (OR) 0.82 (95% CI 0.76–0.89)] and suicidal ideation [OR 0.88 (95% CI 0.79–0.98)] for each 10 percentage-point increase in own ethnic density (Figs 5–6).


We assessed the role of race/ethnicity, country, generational status, and area-level deprivation as moderators in the association between own ethnic density and each of the mental health outcomes. Assessment for *a priori* moderators indicated that these did not account for any observed heterogeneity across estimates.

From the studies reviewed, several mechanisms emerged as providing possible explanations behind ethnic density effects. Racism and social support were the most commonly examined mechanisms. Studies across different national settings suggested that reports of experienced racial discrimination were lower in areas of higher own ethnic density, and reports of social support were higher (Bécaries et al. 2009; Das-Munshi et al. 2010; Das-Munshi et al. 2012; Bécaries et al. 2012a; Bécaries & Das-Munshi, 2013; Bécaries et al. 2013; Shell et al. 2013; Bécaries, 2014; Arévalo et al. 2015). Studies provided support for the role of perceived racism and social support in either moderating (Bécaries et al. 2009; Syed & Juan, 2012; Shell et al. 2013), or mediating (Bécaries et al. 2009; Das-Munshi et al. 2012; Shell et al. 2013; Bécaries, 2014; English et al. 2014) the association between own ethnic density and mental health outcome. Other less explored mechanisms included social cohesion (Syed & Juan, 2012; Bécaries, 2014), acculturation (Arévalo et al. 2015), and stress (Das-Munshi et al. 2012; Shell et al. 2013).

In many of the studies, area-level deprivation was adjusted for as a confounder. Only a few studies permitted the comparison of models adjusted for/not adjusted for area-level deprivation (Bécaries et al. 2013; Bécaries et al. 2014; Schrier et al. 2014; Mezuk et al. 2015). Where it was possible to assess adjustment for area-level deprivation, protective ethnic density associations became stronger or more apparent, and detrimental ethnic density associations reversed direction into a positive association (Bécaries et al. 2013; Bécaries et al. 2014; Schrier et al. 2014; Mezuk et al. 2015). Formal assessment of area-level deprivation as a moderator in the meta-analysis did not provide any conclusive results.

Discussion

Our meta-analysis and review finds evidence indicating protective ethnic density effects across suicidal ideation and psychotic experiences. For CMD, depression, and anxiety, associations were indicative of protective effects but were of a smaller magnitude. For psychosis, although we were unable to perform meta-analysis on any of the studies, the narrative review indicated protective density effects across three studies (Kirkbride et al. 2008; Veling et al. 2008; Kirkbride et al. 2014), with no evidence of detrimental associations across the other studies. The results of the meta-analysis for depression and anxiety were all from the USA, and so should be considered in light of this limitation. Nonetheless, findings relating to all the mental health outcomes were broadly consistent with the narrative review, which tended




1: Hispanic Established Populations for the Epidemiologic Study of the Elderly (HEPESE); 2: National Survey of American Life (NSAL); 3: National Latino and Asian American Study (NLAAS); F: estimates between datasets 76.9%, estimates within datasets 13.2%

Fig. 2. Forest plot of the association between 10% increase in own ethnic density and depression.

to suggest positive associations, except for some negative findings reported in US studies.


Because the ethnic density variable was scaled as per 10% increase, these estimates appear to be small effects, however

given the range of actual own ethnic density (e.g. Bangladeshi people in the UK ranging up to 73% own ethnic density, Māori people in New Zealand up to 86%, and black Caribbean people in the USA up to 53%) (Bécares *et al.* 2009; Bécares *et al.*



1: National Survey of American Life (NSAL); 2: National Latino and Asian American Study (NLAAS); F: between datasets 58.3%

Fig. 3. Forest plot of the association between 10% increase in own ethnic density and anxiety.



1: Health Survey for England (HSE) 1999; 2: Ethnic Minority Psychiatric Illness Rates in the Community (EMPRIC); 3: National Survey of American Life (NSAL); 4: HSE 2004; 5: New Zealand Health Survey (NZHS); 6: National Latino and Asian American Study (NLAAS); 7: G4 Monitor 2008; F: estimates between datasets 0%, estimates within datasets 82.2%

Fig. 4. Forest plot of the association between 10% increase in own ethnic density and common mental disorders.


2012a; Bécates *et al.* 2013) in absolute terms; this means that for people from minority groups living in higher ranges of own ethnic density, this could translate into larger protective effects for mental health. Conversely for people from minority groups living in areas of the lowest own ethnic density, these associations potentially represent larger detrimental effects. Results from the meta-analysis did not find any evidence of detrimental effects of ethnic density on any mental health outcomes.

The mechanisms underlying ethnic density effects are not yet fully understood although the evidence suggests that in areas of higher own ethnic density experiences of racial discrimination are fewer, and the association between racism and adverse mental health outcomes is weaker. Studies show that in these areas, levels of social support are higher. Our assessment of moderation by race/ethnicity, country, generational status, and area-level deprivation did not find any support in moderation by these variables. This may suggest that ethnic density associations are not particular to one racial/ethnic group or national context but rather may be a consequence of being a marginalized racial/ethnic minority. We were only able to empirically assess heterogeneity across 12 studies which may limit conclusions. Other factors such as social cohesion, stigma in reporting mental health problems in areas of high ethnic density, stress, and acculturation may play a role; however, further research is needed.

Strengths and limitations

This is the first systematic review with meta-analysis and detailed meta-regression to explore ethnic density effects on several mental

health outcomes. Our inclusion criteria enables us to be certain that the findings are not a function of help-seeking differences. This is important as it is known that there are potential differences by race/ethnicity in health-seeking behaviors and more complex pathways into care for mental disorders (Bhui *et al.* 2003). Our inclusion of 'leakage' studies for rare conditions such as psychosis in this review meant that we also included people with serious mental illness who would be missed in general population surveys. The use of meta-analysis with meta-regression allowed us to explore potential sources of heterogeneity which have previously been unexplored. The breadth of the systematic reviews meant that we were able to include estimates across national settings and diverse populations, which took into account characteristics such as generational status, area-level deprivation, and individual-level covariates such as age, gender, and socioeconomic position. The review also includes previously unpublished data, which will have reduced publication biases and increased the validity of findings. This was confirmed by formal statistical assessment of publication biases. One of the limitations is that due to data availability, we were only able to meta-analyze 12 studies and most of these data came from observational, cross-sectional studies, which did not allow us to ascertain the direction of causality or temporality of associations and also limited conclusions with respect to sources of heterogeneity. We aimed to address this through the narrative review which demonstrates difference across and within countries. Our review encompasses a number of countries, and although we did not restrict studies by location, generalizability of our findings is restricted to those countries included. In addition, we restricted inclusion criteria




1: Ethnic Minority Psychiatric Illness Rates in the Community (EMPIRIC); 2: Fourth National Survey of Ethnic Minorities (FNS); F: estimates between datasets 8.4%

Fig. 5. Forest plot of the association between 10% increase in own ethnic density and psychotic experiences.

to studies of ethnic density that focused on adults, but for some of these mental health conditions, the risk for disorders may emerge earlier in the life course. Future research should consider the development of mental health disorders and the exposure to socio-environmental stressors earlier in the life course.

We were not able to assess the existence of threshold effects in the meta-analysis, but two studies in the literature document a non-linear association between ethnic density and mental health. Both studies noted this in African American populations in the USA, for whom levels of own ethnic density are higher (Mair *et al.* 2010; Bécares *et al.* 2014). In one of the two studies, the investigators noted that detrimental associations of own ethnic density with depression were only observed at a higher threshold of 85% or greater own ethnic density; at levels below this the association was protective (Bécares *et al.* 2014). These findings highlight that in the USA, extreme levels of own ethnic density as found among African American populations are not always beneficial for mental health, as these areas are characterized by long-standing disinvestment and concentrated poverty (Williams & Collins, 2001). It is this concentration of poverty, and not of ethnic minority people, which drives the harmful associations for mental health. Although we were unable to assess this further due to lack of data, this could be relevant to other international contexts.

There are long-standing concerns that pre-existing mental health problems may cause people to 'drift' into less affluent neighborhoods (Halpern & Nazroo, 2000). High levels of ethnic density correlate with high levels of deprivation (Bécares *et al.* 2009; Bécares *et al.* 2012a; b; Bécares *et al.* 2013; Bécares *et al.* 2014), and these two phenomena operate in opposite directions – whereas area deprivation is associated with worse health (Pickett & Pearl, 2001), our meta-analysis and narrative review show that ethnic density has protective associations for the mental health of racial/ethnic minority people. If the drift hypothesis was true, one would anticipate that racial/ethnic minority people living in areas of high deprivation, and high ethnic density, would have higher rates of mental disorders, but in fact the opposite is true. Once area deprivation is adjusted for, living in areas of high ethnic density is protective for mental health. It is also possible that people with mental disorders move away from family and friends. However, in the context of ethnic density, it is unlikely that this move would happen towards an area of reduced ethnic density, as these areas tend to be more affluent. This would mean that individuals with mental disorders would have the social and economic resources to be able to move into a less dense, and more affluent area, which is unlikely as the evidence suggests that individuals with mental disorders tend to have lower socio-economic resources (Lorant *et al.* 2003).



1: Ethnic Minority Psychiatric Illness Rates in the Community (EMPIRIC); 2: National Latino and Asian American Study (NLAAS); 3: National Survey of American Life (NSAL); F: estimates between datasets 54.5%

Fig. 6. Forest plot of the association between 10% increase in own ethnic density and suicidal ideation.

Implications

A previous narrative review found consistent associations for ethnic density and psychosis, and was only to make tentative conclusions about the association between ethnic density and other mental disorders (Shaw *et al.* 2012). Despite the methodological challenges of the present study, which included limited availability and heterogeneity of data, differential adjustment for confounders, and the differing contexts of racial/ethnic groups across different national settings, the findings of the current study are able to indicate, for the first time, protective ethnic density associations across countries and racial/ethnic minority populations, as well as across mental health outcomes. This is indicative of the importance of the social environment in patterning detrimental mental health outcomes in marginalized and excluded population groups, in particular the role of social exclusion (Selten & Cantor-Graae, 2005), racism and racial discrimination (Bécaries *et al.* 2009), and minority stigma (Pickett & Wilkinson, 2008) for health.

Attempts to engineer patterns of racial/ethnic minority and migrant settlement have led to public policy initiatives to address the tendency for new migrants to concentrate in diverse cities by actively seeking to disperse people more widely, across a range of international settings, including the USA, Canada, Australia, Europe, and the UK, among others. These dispersal programs

place immigrants in areas with limited previous history of accommodating new migrants and lower ethnic density (Robinson & Reeve, 2006; Stewart & Shaffer, 2015). Although our study was not designed to examine this, results of our review and meta-analysis alert us to the possible unintended mental health consequences of these policies for new migrants, and provide evidence for positive outcomes emerging from the residential concentration of racial/ethnic minority populations, as long as the deprivation and concentrated poverty in these areas are adequately addressed.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291717003580>

Acknowledgements. The authors are grateful to all authors who responded to their inquiries. In particular, the authors are grateful to the following authors for providing additional data: James Kirkbride, Sandra Arévalo, Devon English, Jamie Booth, Inma Jarrin, Thomas Jucik, Fabian Temorshuizen, Ozcan Erdem, and Kristine Molina. The authors are also grateful to Craig Morgan for advice on leakage studies for psychosis. They are grateful to Richard Shaw for sharing search strategies from an earlier review. They are also grateful to the British Library for use of their reading rooms and other facilities. LB was funded by the ESRC (grant number ES/K001582/1) and a Hallsworth Research Fellowship. JD-M is funded by the Health Foundation/Academy of Medical Sciences.

Declaration of interest. None.

References

- Alegria M, Molina KM and Chen CN** (2014) Neighborhood characteristics and differential risk for depressive and anxiety disorders across racial/ethnic groups in the United States. *Depression & Anxiety* **31**, 27–37.
- Aneshensel CS, Wight RG, Miller-Martinez D, Botticello AL, Karlamangla AS and Seeman TE** (2007) Urban neighborhoods and depressive symptoms among older adults. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences* **62**, S52–S59.
- Arévalo SP, Tucker KL and Falcón LM** (2015) Beyond cultural factors to understand immigrant mental health: neighborhood ethnic density and the moderating role of pre-migration and post-migration factors. *Social Science & Medicine* **138**, 91–100.
- Bécares L** (2014) Ethnic density effects on psychological distress among Latino ethnic groups: an examination of hypothesized pathways. *Health & Place* **30**, 177–186.
- Bécares L, Cormack D and Harris R** (2013) Ethnic density and area deprivation: neighbourhood effects on Maori health and racial discrimination in Aotearoa/New Zealand. *Social Science & Medicine* **88**, 76–82.
- Bécares L and Das-Munshi J** (2013) Ethnic density, health care seeking behaviour and expected discrimination from health services among ethnic minority people in England. *Health & Place* **22**, 48–55.
- Bécares L, Nazroo J and Jackson J** (2014) Ethnic density and depressive symptoms among African Americans: threshold and differential effects across social and demographic subgroups. *American Journal of Public Health* **104**, 2334–2341.
- Bécares L, Nazroo J, Jackson J and Heuvelman H** (2012a) Ethnic density effects on health and experienced racism among Caribbean people in the US and England: a cross-national comparison. *Social Science & Medicine* **75**, 2107–2115.
- Bécares L, Nazroo J and Stafford M** (2009) The buffering effects of ethnic density on experienced racism and health. *Health & Place* **15**, 670–678.
- Bécares L, Shaw R, Nazroo J, Stafford M, Albor C, Atkin K et al.** (2012b) Ethnic density effects on physical morbidity, mortality, and health behaviors: a systematic review of the literature. *American Journal of Public Health* **102**, e33–e66.
- Bhui K, Stansfeld SA, Hull S, Priebe S, Mole F and Feder G** (2003) Ethnic variations in pathways to and use of specialist mental health services in the UK. *British Journal of Psychiatry* **182**, 105–116.
- Booth JM** (2014) *The protective effect of community organization on distress in disadvantaged neighbourhoods: Considering the Latino experience in Chicago* (Doctoral dissertation). Arizona State University. Published by ProQuest LLC. UMI 3619567.
- Chinn S** (2000) A simple method for converting an odds ratio to effect size for use in meta-analysis. *Statistics in Medicine* **19**, 3127–3131.
- Choi S, Kim G and Lee S** (2016) Effects of nativity, length of residence, and county-level foreign-born density on mental health among older adults in the U.S. *The Psychiatric Quarterly* **87**, 675–688.
- Das-Munshi J, Bécares L, Boydell JE, Dewey ME, Morgan C, Stansfeld SA et al.** (2012) Ethnic density as a buffer for psychotic experiences: findings from a national survey (EMPIRIC). *British Journal of Psychiatry* **201**, 282–290.
- Das-Munshi J, Bécares L, Dewey ME, Stansfeld SA and Prince MJ** (2010) Understanding the effect of ethnic density on mental health: multi-level investigation of survey data from England. *BMJ* **341**, c5367.
- Ecob R and Williams R** (1991) Sampling Asian minorities to assess health and welfare. *Journal of Epidemiology and Community Health* **45**, 93–101.
- English D, Lambert SF, Evans MK and Zonderman AB** (2014) Neighborhood racial composition, racial discrimination, and depressive symptoms in African Americans. *American Journal of Community Psychology* **54**, 219–228.
- Feng X, Astell-Burt T and Kolt GS** (2013) Do social interactions explain ethnic differences in psychological distress and the protective effect of local ethnic density? A cross-sectional study of 226 487 adults in Australia. *BMJ Open* **3**, e002713. doi: 10.1136/bmjopen-2013-002713.
- Halpern D and Nazroo J** (2000) The ethnic density effect: results from a national community survey of England and Wales. *International Journal of Social Psychiatry* **46**, 34–46.
- Jarrin I, Garcia-Fulgueiras A, Ibanez-Rojo V, Alvarez D, Garcia-Pina R, Fernandez-Liria A et al.** (2013) Absence of protective ethnic density effect on Ecuadorian migrants' mental health in a recent migration setting: a multi-level analysis. *Social Psychiatry and Psychiatric Epidemiology* **48**, 95–103.
- Kaufman J and Cooper R** (2001) Considerations for use of racial/ethnic classification in etiologic research. *American Journal of Epidemiology* **154**, 291–298.
- Khan KS, Riet GT, Popay J, Nixon J and Kleijnen J** (2001) STAGE II: conducting the review; PHASE 5: study quality assessment. In Khan KS, Riet GT, Glanville H, Sowden AJ and Kleijnen J (eds). *CRD Report Number 4*, pp. 1–20. York, UK: NHS Centres for Reviews and Dissemination (CRD).
- Kirkbride JB, Boydell J, Ploubidis GB, Morgan C, Dazzan P, McKenzie K et al.** (2008) Testing the association between the incidence of schizophrenia and social capital in an urban area. *Psychological Medicine* **38**, 1083–1094.
- Kirkbride JB, Jones PB, Ullrich S and Coid JW** (2014) Social deprivation, inequality, and the neighborhood-level incidence of psychotic syndromes in East London. *Schizophrenia Bulletin* **40**, 169–180.
- Kirkbride JB, Morgan C, Fearon P, Dazzan P, Murray RM and Jones PB** (2007) Neighbourhood-level effects on psychoses: re-examining the role of context. *Psychological Medicine* **37**, 1413–1425.
- Kwag KH, Jang Y, Rhew SH and Chiriboga DA** (2011) Neighborhood effects on physical and mental health: a study of Korean American older adults. *Asian American Journal of Psychology* **2**, 91–100.
- Leu J, Walton E and Takeuchi D** (2011) Contextualizing acculturation: gender, family, and community reception influences on Asian immigrant mental health. *American Journal of Community Psychology* **48**, 168–180.
- Lorant V, Deliège D, Eaton W, Robert A, Philippot P and Anseau M** (2003) Socioeconomic inequalities in depression: a meta-analysis. *American Journal of Epidemiology* **157**, 98–112.
- Mair C, Roux AVD, Osypuk TL, Rapp SR, Seeman T and Watson KE** (2010) Is neighborhood racial/ethnic composition associated with depressive symptoms? The multi-ethnic study of atherosclerosis. *Social Science & Medicine* **71**, 541–550.
- Menezes NM, Georgiades K and Boyle MH** (2011) The influence of immigrant status and concentration on psychiatric disorder in Canada: a multi-level analysis. *Psychological Medicine* **41**, 2221–2231.
- Mezuk B, Li X, Cedering K, Concha J, Kendler KS, Sundquist J et al.** (2015) Ethnic enclaves and risk of psychiatric disorders among first- and second-generation immigrants in Sweden. *Social Psychiatry and Psychiatric Epidemiology* **50**, 1713–1722.
- Murray C and Lopez A** (1996). *The Global Burden of Disease*. Cambridge, MA: Harvard University Press.
- Neeleman JAN and Wessely S** (1999) Ethnic minority suicide: a small area geographical study in south London. *Psychological Medicine* **29**, 429–436.
- Ostir G, Eschbach K, Markides K and Goodwin J** (2003) Neighbourhood composition and depressive symptoms among older Mexican Americans. *Journal of Epidemiology and Community Health* **57**, 987–992.
- Pickett K and Pearl M** (2001) Multilevel analyses of neighbourhood socioeconomic context and health outcomes: a critical review. *Journal of Epidemiology & Community Health* **55**, 111–122.
- Pickett K and Wilkinson R** (2008) People like us: ethnic group density effects on health. *Ethnicity & Health* **13**, 321–334.
- Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M et al.** (2006) *Guidance on the Conduct of Narrative Synthesis in Systematic Reviews: A Product from the ESRC Methods Programme*. Lancaster University. Available online at <http://www.lancs.ac.uk/shm/research/nssr/research/dis-semination/publications.php> (3 November 2017).
- Propper C, Jones K, Bolster A, Burgess S, Johnston R and Sarker R** (2005) Local neighbourhood and mental health: evidence from the UK. *Social Science & Medicine* **61**, 2065–2083.
- Robinson D and Reeve K** (2006) *Neighbourhood Experiences of new Immigration. Reflections from the Evidence Base*. York: Joseph Rowntree Foundation.
- Schrier AC, Peen J, de Wit MAS, van Ameijden EJC, Erdem Ö, Verhoeff AP et al.** (2014) Ethnic density is not associated with psychological distress in Turkish-Dutch, Moroccan-Dutch and Surinamese-Dutch ethnic minorities

- in the Netherlands. *Social Psychiatry and Psychiatric Epidemiology* **49**, 1557–1567.
- Selten JP and Cantor-Graae E** (2005) Social defeat: risk factor for schizophrenia? *British Journal of Psychiatry* **187**, 101–102.
- Shaw RJ, Atkin K, Bécades L, Albor CB, Stafford M, Kiernan KE et al.** (2012) Impact of ethnic density on adult mental disorders: narrative review. *British Journal of Psychiatry* **201**, 11–19.
- Shaw RJ and Pickett KE** (2011) The association between ethnic density and poor self-rated health among US black and Hispanic people. *Ethnicity & Health* **16**, 225–244.
- Shell AM, Peek MK and Eschbach K** (2013) Neighborhood Hispanic composition and depressive symptoms among Mexican-descent residents of Texas city, Texas. *Social Science & Medicine* **99**, 56–63.
- Stafford M, Newbold BK and Ross NA** (2011) Psychological distress among immigrants and visible minorities in Canada: a contextual analysis. *International Journal of Social Psychiatry* **57**, 428–441.
- Stewart E and Shaffer M** (2015) *Moving on? Dispersal Policy, Onward Migration and Integration of Refugees in the UK*. Glasgow: University of Strathclyde.
- Syed M and Juan MJD** (2012) Discrimination and psychological distress: examining the moderating role of social context in a nationally representative sample of Asian American adults. *Asian American Journal of Psychology* **3**, 104–120.
- Team RC** (2015) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Termorshuizen F, Braam AW and van Ameijden EJC** (2015) Neighborhood ethnic density and suicide risk among different migrant groups in the four big cities in the Netherlands. *Social Psychiatry and Psychiatric Epidemiology* **50**, 951–962.
- United Nations, Department of Economic and Social Affairs, Population Division** (2016) International Migration Report 2015 (ST/ESA/SER.A/384). United Nations.
- Viechtbauer W** (2010) Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software* **36**, 1–48.
- Veling W, Susser E, van Os J, Mackenbach JP, Selten J-P and Hoek HW** (2008) Ethnic density of neighborhoods and incidence of psychotic disorders Among immigrants. *American Journal of Psychiatry* **165**, 66–73.
- Vogt Yuan AS** (2007) Racial composition of neighborhood and emotional well-being. *Sociological Spectrum* **28**, 105–129.
- Whiteford H, Degenhardt L, Rehm J, Baxter A, Ferrari A, Erskine H et al.** (2013) Global burden of disease attributable to mental and substance use disorders: findings from the global burden of disease study 2010. *Lancet* **382**, 1575–1586.
- Wight RG, Cummings JR, Karlamangla AS and Aneshensel CS** (2009) Urban neighborhood context and change in depressive symptoms in late life. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences* **64B**, 247–251.
- Wight RG, Ko MJ and Aneshensel CS** (2011) Urban neighborhoods and depressive symptoms in late middle age. *Research on Aging* **33**, 28–50.
- Williams DR and Collins C** (2001) Racial residential segregation: a fundamental cause of racial disparities in health. *Public Health Reports* **116**, 404–416.