




RESEARCH ARTICLE

# Nucleic acids and melanin pigments after exposure to high doses of gamma rays: a biosignature robustness test

A. Cassaro<sup>1</sup> , C. Pacelli<sup>1,2</sup>, M. Baqué<sup>3</sup> , A. Maturilli<sup>3</sup>, U. Boettger<sup>4</sup>, R. Moeller<sup>5,6</sup>, A. Fujimori<sup>7</sup>, J.-P.P. de Vera<sup>8</sup>  and S. Onofri<sup>1</sup>

<sup>1</sup>Department of Ecological and Biological Sciences, University of Tuscia, Largo dell'Università snc, 01100 Viterbo, Italy

<sup>2</sup>Human Spaceflight and Scientific Research Unit, Italian Space Agency, via del Politecnico snc, Rome, Italy

<sup>3</sup>Planetary Laboratories Department, German Aerospace Center (DLR e.V.), Institute of Planetary Research, Rutherfordstraße 2, Berlin, Germany

<sup>4</sup>German Aerospace Center (DLR e.V.), Institute of Optical Sensor Systems, Rutherfordstraße 2, Berlin, Germany

<sup>5</sup>Radiation Biology Department, Aerospace Microbiology, German Aerospace Center (DLR e.V.), Institute of Aerospace Medicine, Cologne (Köln), Germany

<sup>6</sup>Department of Natural Sciences, University of Applied Sciences Bonn-Rhein-Sieg (BRSU), Rheinbach, Germany

<sup>7</sup>Molecular and Cellular Radiation Biology Group, Department of Basic Medical Sciences for Radiation Damages, NIRS/QST, Chiba, Japan

<sup>8</sup>German Aerospace Center (DLR e.V.), MUSC, Space Operations and Astronaut Training, Linder Höhe, Cologne, Germany

**Author for correspondence:** C. Pacelli, E-mail: [claudia.pacelli@asi.it](mailto:claudia.pacelli@asi.it)

**Received:** 3 December 2021; **Revised:** 16 April 2022; **Accepted:** 1 May 2022; **First published online:** 06 July 2022

**Key words:** Biosignatures, ExoMars, life-detection, Mars, nucleic acids, pigments

## Abstract

The question about the stability of certain biomolecules is directly connected to the life-detection missions aiming to search for past or present life beyond Earth. The extreme conditions experienced on extraterrestrial planet surface (e.g. Mars), characterized by ionizing and non-ionizing radiation, CO<sub>2</sub>-atmosphere and reactive species, may destroy the hypothetical traces of life. In this context, the study of the biomolecules behaviour after ionizing radiation exposure could provide support for the onboard instrumentation and data interpretation of the life exploration missions on other planets. Here, as a part of STARLIFE campaign, we investigated the effects of gamma rays on two classes of fungal biomolecules—nucleic acids and melanin pigments – considered as promising biosignatures to search for during the ‘*in situ* life-detection’ missions beyond Earth.

## Contents

<b>Introduction</b>	<b>296</b>
<b>Materials and methods</b>	<b>298</b>
Fungal colonies preparation and irradiation conditions . . . . .	298
Nucleic acid extractions . . . . .	298
Quantitative PCR (qPCR) analysis of <i>C. antarcticus</i> nucleic acids . . . . .	298
Statistical analyses . . . . .	298
Spectrophotometric analysis . . . . .	299
Confocal Raman spectroscopy analyses . . . . .	299
Fourier transform infrared resonance spectroscopy analysis . . . . .	299
<b>Results</b>	<b>300</b>
Dna analyses by qPCR assay after gamma rays exposure . . . . .	300
Spectrophotometric analysis of <i>C. antarcticus</i> melanin pigments . . . . .	300
Confocal Raman spectroscopy analyses . . . . .	300
Fourier transformed infrared resonance spectroscopy analysis . . . . .	302
<b>Discussion</b>	<b>302</b>

## Introduction

Mars is one of the main targets for the ongoing and in development life-detection missions beyond Earth, due to its potential past habitability. It is known that Mars actually hosts carbon, energy sources and different water reservoirs (Malin and Edgett, 2000; Rasmussen *et al.*, 2009). Nevertheless, the current atmospheric and surface conditions are characterized by a CO<sub>2</sub>-rich atmosphere (Owen *et al.*, 1977), oxidizing substances, such as perchlorates (Lasne *et al.*, 2016), oxygen peroxide (H<sub>2</sub>O<sub>2</sub>) and a hostile surface environment due to the presence of radiation (ionizing and non-ionizing radiation; Hassler *et al.*, 2014). This highly radiative environment is one of the main challenges both for life survival and biomolecule integrity, reducing the chance for robotic exploration to detect traces of life. Ionizing radiation can penetrate several metres into the Martian soil causing structural and chemical damages to biological molecules (Dartnell *et al.*, 2007). Indeed, no organic matter has been found in Martian regolith by the previously Viking missions on Mars (Biemann *et al.*, 1977), even though the high infall meteorites should have accumulated detectable amounts of organics (Drake *et al.*, 1988). The exploration missions focus their investigation on the Martian sub-surface being the Martian surface hostile for life as we know it.

Traces of life beyond Earth, the so-called biosignatures are defined as ‘an object, molecules or pattern of exclusively biological origin’ (des Marais *et al.*, 2008). Among biosignatures, different biomolecules could be considered as direct and unambiguous signs of extant or extinct life, due to chemical and structural complexity and to the specificity that allows to assume that they can be only synthesized by living organisms (Pace, 2001; Summons *et al.*, 2008; Davila and McKay, 2014). Terrestrial biomolecules that store and transfer genetic information (nucleic acids and proteins) are considered the most unambiguous signs of life, given the implausibility that these can be produced abiotically in a natural environment (Neveu *et al.*, 2018). Besides, intra and extra-cellular molecules (*e.g.* pigments) that acts as protective molecules against external stresses are considered good biosignatures.

Considering the previously mentioned radiation environment beyond Earth, which biosignatures are effectively stable and, therefore, recognizable during life-detection missions? The answer to this question is investigated in the frame of the STARLIFE project that aimed at understanding the effect of radiation on selected microorganisms and their associated molecules, by exposing them to different space-relevant radiations (Moeller *et al.*, 2017). The evaluation of the stability/degradation of organics in conditions similar to those experienced beyond Earth’s magnetic field protection, is of outmost importance to support the *in situ* life-detection missions. Among the microorganisms and biomolecules tested in the STARLIFE project, we have chosen the cryptoendolithic black fungus *Cryomyces antarcticus*, isolated from McMurdo Dry Valleys (Antarctica), one of the best terrestrial analogues of the Martian environment (Cassaro *et al.*, 2021a).

The fungus has been chosen for its proven ability to resist radiation stresses (Selbmann *et al.*, 2018). It survived gamma rays (up to 55.61 kGy, <sup>60</sup>Co, Pacelli *et al.*, 2017a), X-rays (up to 300 Gy, Pacelli *et al.*, 2017b), heavy ions in de-hydrated and hydrated conditions (iron ions up to 1000 and 2000 Gy, respectively; Aureli *et al.*, 2020; Pacelli *et al.*, 2021a) and helium ions (up to 1000 Gy, Pacelli *et al.*, 2020a). Melanin, as constituent of the fungal cell-wall, was previously identified as the main responsible for the fungal ability to resist radiation stresses. In particular, it was recently demonstrated that the fungus *C. antarcticus* has two types of melanins produced from two different metabolic pathways: L-DOPA and DHN melanin. This peculiarity could contribute to its enhanced resistance to radiation (Pacelli *et al.*, 2020b).

The melanin was previously reported as highly resistant to the damaging conditions encountered in space and therefore proposed as a promising biosignature for life detection (Pacelli *et al.*, 2021b). Furthermore, its spectroscopic stability after ionizing and non-ionizing radiation (UV, X and gamma rays, iron and helium ions) exposure was largely demonstrated (Pacelli *et al.*, 2017a, 2018, 2019, 2021a, 2021b, 2021c; Aureli *et al.*, 2020). Melanic pigments are ancient pigments widespread in all living kingdoms. Many extremophilic organisms, living in high altitude habitats and Arctic and

Antarctic regions, synthesize melanic pigments gaining protection against UV and solar radiation (Nosanchuk and Casadevall, 2003). In some cases, fungi, even in living condition, have been isolated from highly radiative environments (e.g. cooling pools of nuclear reactors, the stratosphere, the International Space Station (ISS), and inside the damaged nuclear reactor at Chernobyl; Dadachova and Casadevall, 2008), suggesting the role of pigments in radioprotection and microbial survivability mechanisms. In this context, de-hydrated colonies of the fungus *C. antarcticus* have been exposed to increasing doses of gamma rays ( $^{60}\text{Co}$  irradiation doses, up to  $\sim 113$  kGy) as one of the constituents of the Galactic Cosmic Rays (GCRs; Simpson, 1983) with the aim to investigate the stability of its associated biomolecules through quantitative amplification methods and spectroscopic analyses.

## Materials and methods

### *Fungal colonies preparation and irradiation conditions*

In this experiment, the meristematic black fungus *C. antarcticus* MNA-CCFEE (Italian National Antarctic Museum's Culture Collection of Fungi from Extreme Environment) 515 isolated from sandstone rocks collected by L. Vishniac at Linnaeus Terrace (McMurdo Dry Valleys, Southern Victoria Land, Antarctica) during the expedition in 1980–1981, was used as test organism. The fungus was stored in the Fungal Culture Collection of the Antarctic National Museum of Mycological Section of the University of Tuscia (Viterbo, Italy). Colonies for the experiment were obtained by spreading 2000 Colony-Forming Units (CFU) on MEA medium (malt extract, powdered  $30\text{ g L}^{-1}$ ; agar  $15\text{ g L}^{-1}$ ; Applichem, GmbH) on Petri dishes. Fungal samples were incubated at  $15\text{ }^{\circ}\text{C}$  for 3 months. After growth, the colonies were de-hydrated under a laminar flow in sterile conditions and exposed to gamma rays, generated from a  $^{60}\text{Co}$  source (gamma rays at  $1.17\text{ MeV}$ , low linear energy transfer of  $0.2\text{ keV }\mu\text{m}^{-\text{e}}$ ; Table 1), provided by Beta-Gamma-Service GmbH in Cologne (Germany). Irradiation received doses and the total irradiation time are reported in Table 1.

### *Nucleic acid extractions*

DNA was extracted from de-hydrated fungal colonies, by using the Nucleospin Plant kit (Macherey-Nagel, Düren, Germany) following the protocol optimized for black fungi as reported in Selbmann *et al.* (2005). Before amplification, DNA was quantified using QUBIT system and diluted at the concentration of  $0.1\text{ ng}\mu\text{L}^{-1}$  for the following analyses.

### *Quantitative PCR (qPCR) analysis of C. antarcticus nucleic acids*

After extraction and quantification process two gene clusters, a long-repeated fragment (LSU gene of 939 bp) and a short and non-repeated fragment in the genome ( $\beta$ -actin gene of 330 bp); respectively, were amplified. qPCR analysis was performed with a BioRad CFX96 real-time PCR detection system (BioRad, Hercules, CA) using primers targeting the fungal LSU rRNA gene and the  $\beta$ -actin genes: LR0R (ACCCGCTGAACTTAAGC) (Cubeta *et al.*, 1991) and LR5 (TCCTGAGGGAACTTC) (Vilgalys and Hester, 1990), ACT512-F (ATGTGCAAGGCCGTTTCGC) and ACT783-R (TACGAGTCCTTCTGGCCCAT) (Carbone and Kohn, 1999), each at 5 pmol final concentration, following the optimized protocol reported by Pacelli *et al.* (2021b).

### *Statistical analyses*

For multiple data points, mean and standard deviation were calculated. Statistical analyses were performed by one-way analysis of variance (Anova) and pair wise multiple comparison procedure (*t* test), carried out using the statistical software SigmaStat 2.0 (Jandel, USA).

**Table 1.** Samples description and radiation values experienced during STARLIFE experiments. Samples were exposed at the dose rate of  $100 \text{ Gymin}^{-1}$  (Moeller et al., 2017)

Sample	Received doses (kGy)	Total irradiation time (hours)
Lab Control (Ctr)	Non-irradiated	–
Irradiated de-hydrated colonies	6.66	1.11
Irradiated de-hydrated colonies	12.72	2.12
Irradiated de-hydrated colonies	19.04	3.17
Irradiated de-hydrated colonies	27.15	4.5
Irradiated de-hydrated colonies	55.81	9.30
Irradiated de-hydrated colonies	81.11	13.52
Irradiated de-hydrated colonies	113.25	18.9

### Spectrophotometric analysis

Melanin pigments were purified from irradiated and non-irradiated (Lab Ctr) fungal colonies as previously described in Pacelli *et al.* (2020a, 2020b). After three day of lyophilization, purified pigments were solubilized in  $500 \mu\text{L}$  of NaOH 1 M and its UV-Visible spectrum was measured by the use of a VWR-UV 1600 PC Spectrophotometer. NaOH 1 M was used as a zero and the instrument was set in a range of 200–800 nm for the analysis. The acquired data were analysed using Spectragryph software version 1.2.14. In order to quantify the extracted fungal melanin, a calibration line using the absorbance at 650 nm of the synthetic DHN (1,8-DiHydroxyNaphthalene, Thermo-Fisher scientific) melanin was performed, as stated in Raman and Ramasamy (2017).


### Confocal Raman spectroscopy analyses

Confocal Raman spectroscopy was performed at the German Aerospace Center in Berlin, using a 532 nm excitation laser, with a WITec alpha300 Confocal Raman microscope, at room temperature, under ambient atmospheric conditions. Raman spectrometer equipped with 532 nm green laser excitation line in the  $4000\text{--}100 \text{ cm}^{-1}$  region, with  $4\text{--}5 \text{ cm}^{-1}$  resolution with a diode laser of  $1.5 \mu\text{m}$  laser spot size. Magnification was  $10 \times$  via microscope. Image scans were performed at 0.7 mW using 1 accumulation and 1s for each measurement. In order to reduce saturation or damaging effects, measurement is repeated on three distinct areas of each sample up to  $100 \mu\text{m} \times 100 \mu\text{m}$  and up to 500 image points, thus collecting a minimum of 1000 measurements per sample. All obtained data were analysed with the WITec Project FIVE software, using the optimized protocol described by Pacelli *et al.* (2021b).

### Fourier transform infrared resonance spectroscopy analysis

Fourier Transform Infrared Resonance (FTIR) spectroscopy was used to identify the major functional groups, as a part of fungal biomolecules, through their characteristic absorption bands in defined regions of the spectrum.

A pool of colonies of non-irradiated and irradiated samples were placed into aluminium supports without any preparation and the analyses were carried out at room temperature under evacuated conditions. A FTIR spectrometer (Bruker Vertex80 V, Germany) at the Planetary Spectroscopy Laboratory (PSL) of the German Aerospace Center (DLR) in Berlin, was used to record the spectra of fungal samples region at resolution of  $4\text{--}5 \text{ cm}^{-1}$  over 250 scans in  $10\,000\text{--}400 \text{ cm}^{-1}$  wavenumbers range. The same protocol is followed for each measurement. To compare data, spectra were analysed using Spectragryph software version 1.2.14. Bands assignment were performed with a 3-threshold value; 3% of maximum intensity and a noise factor of 2.



**Fig. 1.** (a) Quantitative qPCR of a 939 bp target gene (LSU); (b) a 330 bp target gene ( $\beta$ -actin) of *C. antarcticus* DNA, after exposure to increasing doses of  $\gamma$ -rays. On the axis of the ordinates the number of amplified copies on a logarithmic scale is shown; on the abscissa axis the treatments, expressed in kGy. Lab Ctr: DNA of *C. antarcticus* de-hydrated colonies not exposed to the treatments. Pos Ctr: DNA of *C. antarcticus* hydrated colonies. The same letters above the bars indicate that the values are not significantly different according to the t test ( $P \leq 0.05$ ).

## Results

### Dna analyses by qPCR assay after gamma rays exposure


The integrity of fungal DNA after exposure to increasing doses of gamma rays was investigated amplifying two distinct regions (the ribosomal LSU region and the housekeeping  $\beta$ -actin genes) through qPCR assay. DNA amplification revealed a good nucleic acid integrity, with a difference in copy numbers amplification due to the gene repetition in the fungal genome. Overall, the amplification shows a similar trend for both LSU (Fig. 1A) and  $\beta$ -actin (Fig. 1B) genes, with a decrease in the number of amplified copy numbers as the radiation dose increases (up to ~113 kGy). Summarizing, we obtained an average of 930 and 370 DNA copies amplified; in particular, approximately 85 DNA copy numbers were obtained from samples at ~113 kGy whose LSU gene has been amplified (Fig. 1A) and 200 DNA copies from the same samples whose  $\beta$ -actin gene has been amplified (Fig. 1B).

### Spectrophotometric analysis of *C. antarcticus* melanin pigments

Spectrophotometric analyses were performed on extracted melanin pigments from *C. antarcticus* irradiated colonies in comparison with relative laboratory control (Lab Ctr). The obtained UV-Vis spectra (Fig. 2) showed a similarity among irradiated and control samples, with the highest absorption peak near 230 nm in the UV region and a decrease in absorption with increasing wavelength in the visible region (Pacelli *et al.*, 2020a, 2020b). However, a slight bulge was highlighted near 290–330 nm in all tested samples. In addition, a clear peak at 430 nm has been found in samples exposed to 19.04 kGy of gamma radiation. Overall, no evident changes in absorbed spectra have been reported (Fig. 2).

### Confocal Raman spectroscopy analyses


Raman spectroscopy is an optical technique that is one of the powerful techniques in biopolymer characterization (Krafft, 2004), largely used in space science experiments. Based on inelastic light scattering resulting from the vibration modes in a molecule, it provides specific biochemical information, which allows to discriminate different substances alone or in a mixture (Rebrošová *et al.*, 2017) and to investigate possible changes at the molecular level (Ferrara *et al.*, 2016).




**Fig. 2.** (a) UV-VIS absorbance spectra (200–800 nm) of extracted melanin of *C. antarcticus* after exposure to increasing doses of  $\gamma$ -rays; on x-axis, UV-VIS wavelengths (in nm); on y-axis, absorbance values (in arbitrary unit). (b) Amount of the extracted melanin from each sample.

In this work, we performed confocal Raman spectroscopy analyses in order to investigate similarities and differences among melanic pigments of fungal colonies exposed to increasing doses of  $\gamma$ -rays (results of 6.66 and 55.81 kGy are not show), in comparison with the laboratory controls (Lab Ctr).

The spectra acquired directly from fungal colonies showed two broad bands at 1590–1605  $\text{cm}^{-1}$  and a second lower intensity band around 1340  $\text{cm}^{-1}$ , which correspond to the melanin spectrum (Culka *et al.*, 2017; Pacelli *et al.*, 2021a) (Fig. 3). A third band near 1457  $\text{cm}^{-1}$ , which is also a characteristic Raman band of melanic pigments of the black fungus *C. antarcticus* (Pacelli *et al.*, 2021b), was found. Overall, no evident differences were reported among the different exposure conditions (Fig. S1A). As reported in Fig. S1B, a similar band position (near 1601  $\text{cm}^{-1}$ ) was revealed for control samples (Lab Ctr) and the lower irradiation dose (12.72 kGy). On the contrary, a slight shift (around 1603–1604  $\text{cm}^{-1}$ ) appeared for the higher irradiation doses (up to  $\sim$ 113 kGy). Moreover, a slight



**Fig. 3.** Raman spectra comparison of *C. antarcticus* colonies exposed to increasing doses of  $\gamma$ -rays (generated from a  $^{60}\text{Co}$  source). Melanin main bands are located at 1340 and 1603–1604  $\text{cm}^{-1}$ .



**Fig. 4.** FT-IR spectra in the 600–4000 nm spectral range of de-hydrated *C. antarcticus* colonies exposed to increasing doses of  $\gamma$ -rays (generated from a  $^{60}\text{Co}$  source).

depletion was revealed only in samples exposed to  $\sim 113$  kGy, but, nevertheless, the three bands that characterize the fungal melanin are still evident, even at the highest dose. In conclusion, the characteristic melanin Raman bands are detectable in all the exposure conditions.

#### Fourier transformed infrared resonance spectroscopy analysis

FTIR analysis is one of the most powerful techniques to detect the integrity of biomolecules since it provides an assignment of the spectral characteristic of different functional groups and the absence of a band in the spectrum of a treated sample may indicate a change in the structure of the molecule. In the analysed material, chemical bonds are able to vibrate at a characteristic frequency representative of the material structure and in a FTIR plot, for each absorption band individual chemical bonds can be identified and assigned. In this work, the useful spectral information was obtained between 600 and 4000  $\text{cm}^{-1}$ . Overall, each spectrum is characterized by a similar trend if compared with relative control (Lab Ctr), no evident changes were reported in the acquired spectra (Fig. 4). In particular, all the spectra show strong characteristic stretching bands for C=C bond [ $1750\text{ cm}^{-1}$  associated to lipids and fatty acids; Fig. 4 (Movasaghi *et al.*, 2008)]. The bands observed at  $1454\text{ cm}^{-1}$  and near  $1158\text{ cm}^{-1}$  revealed the detection of chitin characteristic vibrations (Salman *et al.*, 2014; Forfang *et al.*, 2017). The bands near  $1637$  and  $1044\text{ cm}^{-1}$  [due to the bending vibration modes of aromatic ring C=C or C=N (Mboniyirivuze *et al.*, 2015) and to carbon ring breathing vibrations in cyclic compounds (Shurvell, 2006; Pal *et al.*, 2013)], could be associated with the presence of melanin pigments (Fig. 4). The band at  $2857\text{ cm}^{-1}$  is clearly attributed to fungal melanins (Paim *et al.*, 1990; Pal *et al.*, 2013; Sun *et al.*, 2016; Raman and Ramasamy, 2017); while band at  $2926\text{ cm}^{-1}$  (Fig. 4) corresponds to *C. antarcticus* melanin (Pacelli *et al.*, 2020a, 2020b).

#### Discussion

Over the years, the search for life on other planets has undergone changes. During the Viking missions to Mars, research studies aimed at searching for microorganisms in Martian soils; a method that requires the viability of microorganisms and a good growth rate (McKay, 2020). Nowadays, life-detection missions encompass different measurements, including the search of biosignatures, which could represent unambiguously the sign of extant or recently extinct life (McKay, 2004; Neveu *et al.*, 2018). One of the main hurdles in the search for life is the highly radiative environment that could destroy organic molecules or their remains, not allowing their detection during *in situ*

life-detection mission. Among the GCRs, gamma-rays are the highest-energy form of light and have deleterious effects on biological molecules. Some are generated by transient events, such as solar flares and the huge star explosions known as supernovas. Others are produced by steady sources like the supermassive black holes at the hearts of galaxies.

Previous studies on the same samples showed a high resistance of the black fungus to this type of radiation in terms of cell survival and ultrastructure integrity (Pacelli *et al.*, 2017a). Here, we aimed at investigating the effects of gamma rays on two classes of fungal biomolecules, nucleic acids and pigments, in order to provide support for life-detection missions on Mars and for the future analyses of the Mars Samples Return. Based on the idea that life shares a similar biochemistry in the Universe and on the biogenicity requirement for a biosignature, we firstly focused our attention on the stability of nucleic acids after gamma rays' exposure. The choice of this macromolecule is determined by a multitude of factors. First of all, DNA is widespread in all life forms on Earth; the formation of double-stranded DNA does not occur under abiotic conditions, differently from other biological molecules, such as amino acids, polycyclic aromatic hydrocarbons, etc. (Trevors, 2003; Berger *et al.*, 2013). And secondly, PCR-based detection methods are highly specific and sensitive, allowing the detection of a single DNA molecule (Briggs *et al.*, 2009). By using a qPCR assay, we reported that the fungal DNA is always amplifiable, and therefore detectable, even after the ~113 kGy dose of exposure inside the fungal cells (Fig. 1A and B). In particular, we amplified 930 and 370 DNA copy numbers by amplifying LSU and  $\beta$ -actin genes; respectively.

Similar results have been obtained for DNA of the cyanobacterium *Chroococcidiopsis* exposed to gamma radiation, in the frame of the same irradiation campaign: a reduction of the amplifiable target gene was detected in hydrated and de-hydrated samples exposed to 113.25 kGy (Verseux *et al.*, 2017). Additional studies on hydrated colonies of *C. antarcticus* exposed to gamma rays will be carried out, in order to evaluate if protection against radiation may be enhanced from the desiccated condition. Preliminary studies on hydrated and metabolically active colonies of this fungus exposed to accelerated iron ions showed a good DNA amplification up to 2000 Gy (Pacelli *et al.*, 2021a).

The resistance of *C. antarcticus*' DNA under different stressors has been extensively reported (Onofri *et al.*, 2012, 2015, 2019; Pacelli *et al.*, 2017a, 2017b, 2019, 2020a, 2020b, 2021a, 2021c; Aureli *et al.*, 2020; Cassaro *et al.*, 2021b). Also, DNA was successfully extracted and amplified in *C. antarcticus* samples exposed to space conditions in Low Earth orbit (LEO) in the presence of lunar regolith analogue (Cassaro *et al.*, 2022) and of Martian regolith analogues (Pacelli *et al.*, 2021b).

These results suggest the need to accelerate the development of miniaturized amplification instruments, as part of the instrumentation onboard the next generation of rovers. For example, Oxford Nanopore Technologies MinION instrument, which can detect and sequence nucleic acids, has been already tested (Raymond-Bouchard *et al.*, 2022). One of the main challenges in using these devices on the surface of planetary bodies, could be represented by the disturbing compounds in the soil, that may prevent the amplification. However, recent studies reported that the sequencing of two different types of extract (purified and unpurified) DNA revealed a comparable community composition in Permafrost-associated soil samples (Raymond-Bouchard *et al.*, 2022).

Then, we focused on the stability of melanin pigments. This class of pigments is widespread in all living kingdoms and often dominating extremophilic species, suggesting that these pigments are used as a protective strategy in their lifecycle (Dadachova *et al.*, 2007). Here, we investigated potential melanin modifications through a multidisciplinary analysis (UV-Vis spectrophotometry, and FT-IR and Raman spectroscopies). The results of UV-Vis analysis revealed a good stability of melanic pigments even at the ~113 kGy dose of exposure: an absorption peak at 230 nm and a decrease in absorbance at higher wavelength was detected (Fig. 2, Pacelli *et al.*, 2020a, 2020b). No additional peaks or bulges were detected in the acquired spectra, in contrast with fungal samples exposed to Mars-like conditions (Cassaro *et al.*, 2021b). These results are in accordance with Meeßen *et al.* (2013), where UV/VIS-spectrometry data supported the identification of melanin in lichens of astrobiological interest, confirming its role in photoprotection.



Raman spectra showed the three characteristic bands of melanin (Fig. 3): a major band at higher wavenumber (1590–1605  $\text{cm}^{-1}$ ), associated to C-N bonds (Samokhvalov *et al.*, 2004) and a second band at 1340  $\text{cm}^{-1}$ , due to the stretching of the C-C bonds within the rings of the aromatic melanin monomers (Galván *et al.*, 2013). The third band (1457  $\text{cm}^{-1}$ , Fig. 3), distinctive of the fungal melanin pigments (Pacelli *et al.*, 2020a, 2020b) allows to discriminate melanin from (i) any thermal degradation or (ii) amorphous carbon. Our results, confirmed that fungal melanin maintains its stability also at the irradiation dose of 113 kGy, although with a slight thinning of the bands (Fig. 3, red line), and a slight shift of the main peak position at the higher doses (Fig. S1B). These results are in accordance with previous analyses on fungal melanin after exposure to space conditions in Low Earth Orbit (LEO) (Pacelli *et al.*, 2021b) and to simulated space conditions (Pacelli *et al.*, 2021a, 2021c; Cassaro *et al.*, 2021b).

On the contrary, the carotenoid Raman signals of the photobiont part of the lichen *Circinaria gyrosa* exposed to gamma radiation, showed alterations at the dose of 113 kGy (Meeßen *et al.*, 2017); while the carotenoid signal of de-hydrated cells of the cyanobacterium *Nostoc* sp., have been detected up to 27 kGy of gamma rays when exposed alone and to  $\sim$ 113 kGy, when protected by Martian regolith simulants (Baqué *et al.*, 2018). Photosynthetic pigments and nucleic acids of *Chroococcidiopsis* cells also resulted to be detectable even at the dose of 113.25 kGy of gamma rays (Verseux *et al.*, 2017).

Melanin also exhibits unaltered absorption bands in the 600–4000  $\text{cm}^{-1}$  region of FT-IR spectra (Fig. 4). In particular, bands referred to melanin pigments have been detected at different wavelength: the bands at 2857, 2926, 1637 and 1044  $\text{cm}^{-1}$  are indicative of phenols or carboxylic groups, which characterize the absorption of fungal melanin (Sun *et al.*, 2016; Pacelli *et al.*, 2020a, 2020b). The FT-IR spectra showed identical spectroscopic spectra among irradiated and control samples (Fig. 4).

The evaluation of the stability of melanins through three different and complementary techniques, supports its use as a promising biosignature.

The hypothetical finding of a spectral signature of pigment would be a possible sign of life, since no abiotic and uncatalyzed pigment synthesis is known so far in environmental conditions (Neveu *et al.*, 2018). The reported stability of melanin pigments against various types of ionizing and non-ionizing radiation (Robinson, 2001; Vember and Zhdanova, 2001; Nosanchuk and Casadevall, 2003; Cassaro *et al.*, 2021b; Pacelli *et al.*, 2021b, 2021c) could be exploited also in different scenarios of space exploration. As demonstrated in Turick *et al.* (2011), ionizing radiations interact with melanin altering its oxidation/reduction potential. This alteration results in electric current production and further study on the interaction between melanic pigments and ionizing radiation could offer new insights on the application of melanin pigments in the synthetic biology field. Besides, the role of melanin as energy generators could be useful also to generate melanin-based products, especially important to reduce global warming changes (Malo and Dadachova, 2019). When exposed to ionizing radiation, melanized fungal cells showed an increment of the growth rate in comparison to non-melanized cells, probably due to the capability of these pigments to capture electromagnetic radiation, which could be used by microorganisms as metabolic energy, suggesting a potential role in energy capture (Dadachova *et al.*, 2007).

The FT-IR analyses revealed additional bands related to lipids and fatty acids (near 1750  $\text{cm}^{-1}$ , Movasaghi *et al.*, 2008). The identification of chemical bonds belonging to chitin was detected near 1454 and 1158  $\text{cm}^{-1}$  (Fig. 4, Movasaghi *et al.*, 2008). Chitin is one of the main constituents of the fungal cell wall and one of the most abundant compounds on Earth (Hunt, 1970). It is a compound formed of  $\beta$  1–4-bonded N-acetylglucosamine. The hypothesis to consider chitin as a biosignature for the search of life elsewhere has been already reported in Pacelli *et al.* (2021c). In this work, we confirmed the stability of chitin after the exposure to  $\sim$ 113 kGy of gamma rays. The characterization of suitable candidates for a good biosignature could be difficult considering the preservation of certain molecules over geological time scale and the environmental conditions (Summons *et al.*, 2008). However, chitin may be preserved over geological time period: it has been detected in fungi preserved in Cretaceous amber (Speranza *et al.*, 2015) and in a 50-Ma-old marine sponge (Ehrlich *et al.*, 2013).

Recently, chitin has been also discovered in fungal fossil filaments in a Neoproterozoic shale rock (Bonneville *et al.*, 2020), supporting its utilization as a biosignature in space exploration.

In conclusion, this work aimed to investigate the persistence of fungal biomolecules to space-relevant radiation over time, considering that the maximum tested dose of ~113 kGy may be compared to an exposure of 1.5 million years on the Martian surface and of 13 million at two metres-depth in the subsurface (extrapolated from Hassler *et al.*, 2014). The study about the stability of biosignatures is of utmost importance for the *in-situ* life detection missions on other planets, such as Mars. Through this work, we confirmed the persistence of melanin pigments and nucleic acids and we highlighted chitin as a promising biosignature. Further investigations through the application of -omics analyses could represent a challenging task to deepen the molecular mechanisms at the basis of the nucleic acid stability of this fungus.

**Acknowledgements.** We thank the PNRA (Italian National Program for Antarctic Research) for supporting sample collection in Antarctica, and the Italian National Antarctic Museum ‘Felice Ippolito’, for funding MNA-CCFEE. We acknowledge the HIMAC team at the NIRS in Chiba, Japan for the irradiation. We thank the Italian Space Agency for the grant n. 2018-6- U.0 (BioSigN MicroFossils).

**Author contributions.** CP, SO, JPPdV, RM, designed the research; AC, CP, MB, AF performed the experiments. AC, CP, MB, AM analysed data. AC drafted the paper with inputs from all other authors. All authors approved the submitted version.

**Financial support.** This research was financially supported by the Italian Space Agency (ASI), BioSigN MicroFossils- ASI grant n. 2018-6-U.0). STARLIFE project was supported by the MEXT Grant-in-Aid for Scientific Research on Innovative Areas ‘Living in Space’ (Grant Numbers: 15H05935 and 15K21745). RM was supported by the DLR grant FuE-Projekt ‘ISS LIFE’ (Programm RF-FuW, TP 475). MB acknowledges the support of the Deutsche Forschungsgemeinschaft (DFG – German Research Foundation) for the project ‘Raman Biosignatures for Astrobiology Research’ (RaBioFAM; project number: 426601242).

**Conflict of interest.** The authors declare that they have no competing financial interests.

**Conflict of interest.** The authors declare they have not conflicts to disclose.

## References

- Aureli L, Pacelli C, Cassaro A, Fujimori A, Moeller R and Onofri S (2020) Iron ion particle radiation resistance of dried colonies of *Cryomyces antarcticus* embedded in Martian regolith analogues. *Life* **10**, 306.
- Baqué M, Hanke F, Böttger U, Leya T, Moeller R and de Vera JP (2018) Protection of cyanobacterial carotenoids’ Raman signatures by Martian mineral analogues after high-dose gamma irradiation. *Journal of Raman Spectroscopy* **49**, 1617–1627.
- Berger T, Bilski P, Hajek M, Puchalska M and Reitz G (2013) The MATROSHKA experiment: results and comparison from extravehicular activity (MTR-1) and intravehicular activity (MTR-2A/2B) exposure. *Radiation Research* **180**, 622–637.
- Biemann K, Oro JIIPT, Toulmin III P, Orgel LE, Nier AO, Anderson DM and Laffleur AL (1977) The search for organic substances and inorganic volatile compounds in the surface of Mars. *Journal of Geophysical Research* **82**, 4641–4658.
- Bonneville S, Delpomdor F, Préat A, Chevalier C, Araki T, Kazemian M and Benning LG (2020) Molecular identification of fungi microfossils in a Neoproterozoic shale rock. *Science Advances* **6**, eaax7599.
- Briggs AW, Good JM, Green RE, Krause J, Maricic T, Stenzel U and Pääbo S (2009) Primer extension capture: targeted sequence retrieval from heavily degraded DNA sources. *JoVE (Journal of Visualized Experiments)* **31**, e1573.
- Carbone I and Kohn LM (1999) A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* **91**, 553–556.
- Cassaro A, Pacelli C, Aureli L, Catanzaro I, Leo P and Onofri S (2021a) Antarctica As a reservoir of planetary analogue environments. *Extremophiles* **25**, 437–458.
- Cassaro A, Pacelli C, Baqué M, de Vera JPP, Böttger U, Botta L and Onofri S (2021b) Fungal biomarkers stability in Mars regolith analogues after simulated space and Mars-like conditions. *Journal of Fungi* **7**, 859.
- Cassaro A, Pacelli C, Baqué M, Cavalazzi B, Gasparotto G, Saladino R, Botta L, Böttger U, Rabbow E, de Vera JPP and Onofri S (2022) Investigation of fungal biomolecules after low earth orbit exposure: a testbed for the next moon missions. *Environmental Microbiology*.
- Cubeta MA, Echandi E, Abernethy T and Vilgalys R (1991) Characterization of anastomosis groups of binucleate *Rhizoctonia* species using restriction analysis of an amplified ribosomal RNA gene. *Phytopathology* **81**, 1395–1400.
- Culka A, Jehlička J, Ascaso C, Artieda O, Casero CM and Wierzchos J (2017) Raman Microspectrometric study of pigments in melanized fungi from the hyperarid Atacama desert gypsum crust. *Journal of Raman Spectroscopy* **48**, 1487–1493.
- Dadachova E and Casadevall A (2008) Ionizing radiation: how fungi cope, adapt, and exploit with the help of melanin. *Current Opinion* **11**, 525–531.

- Dadachova E, Bryan RA, Huang X, Moadel T, Schweitzer AD, Aisen P and Casadevall A (2007) Ionizing radiation changes the electronic properties of melanin and enhances the growth of melanized fungi. *PLoS ONE* **2**, e457.
- Dartnell LR, Desorgher L, Ward JM and Coates AJ (2007) Martian sub-surface ionising radiation: biosignatures and geology. *Biogeosciences* **4**, 545–558.
- Davila AF and McKay CP (2014) Chance and necessity in biochemistry: implications for the search for extraterrestrial biomarkers in earth-like environments. *Astrobiology* **14**, 534–540.
- Des Marais DJ, Nuth III JA, Allamandola LJ, Boss AP, Farmer JD, Hoehler TM and Spormann AM (2008) The NASA astrobiology roadmap. *Astrobiology* **8**, 715–730.
- Drake MJ, Greeley R, McKay GA, Blanchard DP, Carr MH, Gooding JL, McKay CP, Spudis PD and Squyres SW (eds) (1988) *Workshop on Mars Sample Return Science*. LPI Tech. Rpt. 88-07. Houston: Lunar and Planetary Institute.
- Ehrlich H, Kaluzhnaya OV, Brunner E, Tsurkan MV, Ereskovsky A, Ilan M and Wörheide G (2013) Identification and first insights into the structure and biosynthesis of chitin from the freshwater sponge *Spongilla lacustris*. *Journal of Structural Biology* **183**, 474–483.
- Ferrara MA, De Angelis A, De Luca AC, Coppola G, Dale B and Coppola G (2016) Simultaneous holographic microscopy and Raman spectroscopy monitoring of human spermatozoa photodegradation. *IEEE Journal of Selected Topics in Quantum Electronics* **22**, 27–34.
- Forfang K, Zimmermann B, Kosa G, Kohler A and Shapaval V (2017) FTIR spectroscopy for evaluation and monitoring of lipid extraction efficiency for oleaginous fungi. *PLoS ONE* **12**, e0170611.
- Galván I, Jorge A, Ito K, Tabuchi K, Solano F and Wakamatsu K (2013) Raman spectroscopy as a non-invasive technique for the quantification of melanins in feathers and hairs. *Pigment Cell & Melanoma Research* **26**, 917–923.
- Hassler DM, Zeitlin C, Wimmer-Schweingruber RF, Ehresmann B, Rafkin S, Eigenbrode JL and Berger G (2014) Mars' surface radiation environment measured with the Mars science laboratory's curiosity rover. *Science* **343**, 1244797.
- Hunt S (1970) *Polysaccharide-Protein Complexes in Invertebrates*. London: Academic Press.
- Krafft C (2004) Bioanalytical applications of Raman spectroscopy. *Analytical and Bioanalytical Chemistry* **378**, 60–62.
- Lasne J, Noblet A, Szopa C, Navarro-González R, Cabane M, Poch O and Coll P (2016) Oxidants at the surface of Mars: a review in light of recent exploration results. *Astrobiology* **16**, 977–996.
- Malin MC and Edgett KS (2000) Sedimentary rocks of early Mars. *Science* **290**, 1927–1937.
- Malo ME and Dadachova E (2019) Melanin as an energy transducer and a radioprotector in black fungi. In Tiquia-Arashiro S and Grube M (eds), *Fungi in Extreme Environments: Ecological Role and Biotechnological Significance*. Cham: Springer, pp. 175–184.
- Mbonyiriyvuzé A, Mwakikunga BW, Dhlamini SM and Maaza M (2015) Fourier transform infrared spectroscopy for sepia melanin. *Physics and Materials Chemistry* **3**, 25–29.
- McKay CP (2004) What is life—and how do we search for it in other worlds?. *PLoS Biology* **2**, e302.
- McKay CP (2020) What is life—and when do we search for it on other worlds. *Astrobiology* **20**, 163–166.
- Meeßen J, Sánchez FJ, Sadowsky A, de la Torre R, Ott S and de Vera JP (2013) Extremotolerance and resistance of lichens: comparative studies on five species used in astrobiological research II. Secondary lichen compounds. *Origins of Life and Evolution of Biospheres* **43**, 501–526.
- Meeßen J, Backhaus T, Brandt A, Raguse M, Böttger U, de Vera JP and de la Torre R (2017) The effect of high-dose ionizing radiation on the isolated photobiont of the astrobiological model lichen *Circinaria gyrosa*. *Astrobiology* **17**, 154–162.
- Moeller R, Raguse M, Leuko S, Berger T, Hellweg CE and Fujimori A and STARLIFE Research Group (2017) STARLIFE—An international campaign to study the role of galactic cosmic radiation in astrobiological model systems. *Astrobiology* **17**, 101–109.
- Movasaghi Z, Rehman S and Rehman DI (2008) Fourier transform infrared (FTIR) spectroscopy of biological tissues. *Applied Spectroscopy Reviews* **43**, 134–179.
- Neveu M, Hays LE, Voytek MA, New MH and Schulte MD (2018) The ladder of life detection. *Astrobiology* **18**, 1375–1402.
- Nosanchuk JD and Casadevall A (2003) The contribution of melanin to microbial pathogenesis. *Cellular Microbiology* **5**, 203–223.
- Onofri S, de la Torre R, de Vera JP, Ott S, Zucconi L, Selbmann L and Horneck G (2012) Survival of rock-colonizing organisms after 1.5 years in outer space. *Astrobiology* **12**, 508–516.
- Onofri S, de Vera JP, Zucconi L, Selbmann L, Scalzi G, Venkateswaran KJ and Horneck G (2015) Survival of Antarctic cryptoendolithic fungi in simulated Martian conditions on board the international space station. *Astrobiology* **15**, 1052–1059.
- Onofri S, Selbmann L, Pacelli C, Zucconi L, Rabbow E and de Vera JP (2019) Survival, DNA, and ultrastructural integrity of a cryptoendolithic Antarctic fungus in Mars and lunar rock analogs exposed outside the international space station. *Astrobiology* **19**, 170–182.
- Owen T, Biemann K, Rushneck DR, Biller JE, Howarth DW and Lafleur AL (1977) The composition of the atmosphere at the surface of Mars. *Journal of Geophysical Research* **82**, 4635–4639.
- Pace NR (2001) The universal nature of biochemistry. *PNAS* **98**, 805–808.
- Pacelli C, Selbmann L, Zucconi L, Raguse M, Moeller R, Shuryak I and Onofri S (2017a) Survival, DNA integrity, and ultrastructural damage in antarctic cryptoendolithic eukaryotic microorganisms exposed to ionizing radiation. *Astrobiology* **17**, 126–135.
- Pacelli C, Bryan RA, Onofri S, Selbmann L, Shuryak I and Dadachova E (2017b) Melanin is effective in protecting fast and slow growing fungi from various types of ionizing radiation. *Environmental Microbiology* **19**, 1612–1624.

- Pacelli C, Bryan RA, Onofri S, Selbmann L, Zucconi L, Shuryak I and Dadachova E (2018) The effect of protracted X-ray exposure on cell survival and metabolic activity of fast and slow growing fungi capable of melanogenesis. *Environmental Microbiology Reports* **10**, 255–263.
- Pacelli C, Selbmann L, Zucconi L, Coleine C, de Vera JP, Rabbow E and Onofri S (2019) Responses of the black fungus *Cryomyces antarcticus* to simulated Mars and space conditions on rock analogs. *Astrobiology* **19**, 209–220.
- Pacelli C, Cassaro A, Aureli L, Moeller R, Fujimori A and Onofri S (2020a) The responses of the black fungus *Cryomyces antarcticus* to high doses of accelerated helium ions radiation within Martian regolith simulants and their relevance for Mars. *Life* **10**, 130.
- Pacelli C, Cassaro A, Maturilli A, Timperio AM, Gevi F, Cavalazzi B and Onofri S (2020b) Multidisciplinary characterization of melanin pigments from the black fungus *Cryomyces antarcticus*. *AMAB* **104**, 6385–6395.
- Pacelli C, Alessia C, Siong LM, Lorenzo A, Moeller R, Fujimori A and Silvano O (2021a) Insights into the survival capabilities of *Cryomyces antarcticus* hydrated colonies after exposure to Fe particle radiation. *Journal of Fungi* **7**, 495.
- Pacelli C, Cassaro A, Baqué M, Selbmann L, Zucconi L, Maturilli A and Onofri S (2021b) Fungal biomarkers are detectable in Martian rock-analogues after space exposure: implications for the search of life on Mars. *International Journal of Astrobiology* **20**, 345–358.
- Pacelli C, Cassaro A, Catanzaro I, Baqué M, Maturilli A, Böttger U, Botta L, Saladino R, Rabbow E and Onofri S (2021c) The ground-based BIOMEX experiment verification tests for life detection on Mars. *Life* **11**, 1212.
- Paim S, Linhares LF, Mangrich AS and Martin JP (1990) Characterization of fungal melanins and soil humic acids by chemical analysis and infrared spectroscopy. *Biology and Fertility of Soils* **10**, 72–76.
- Pal AK, Gajjar DU and Vasavada AR (2013) DOPA And DHN pathway orchestrate melanin synthesis in *Aspergillus* species. *Medical Mycology* **52**, 10–18.
- Raman NM and Ramasamy S (2017) Genetic validation and spectroscopic detailing of DHN-melanin extracted from an environmental fungus. *Biochemistry and Biophysics Reports* **12**, 98–107.
- Rasmussen B, Blake TS, Fletcher IR and Kilburn MR (2009) Evidence for microbial life in synsedimentary cavities from 2.75 Ga terrestrial environments. *Geology* **37**, 423–426.
- Raymond-Bouchard I, Maggiori C, Brennan L, Altshuler I, Manchado JM, Parro V and Whyte LG (2022) Assessment of automated nucleic acid extraction systems in combination with MinION sequencing as potential tools for the detection of microbial biosignatures. *Astrobiology* **22**, 87–103.
- Rebrošová K, Šiler M, Samek O, Růžička F, Bernatová S, Holá V and Petráš P (2017) Rapid identification of staphylococci by Raman spectroscopy. *Scientific Reports* **7**, 1–8.
- Robinson CH (2001) Cold adaptation in Arctic and Antarctic fungi. *New phytologist* **151**, 341–353.
- Salman A, Shufan E, Tsror L, Moreh R, Mordechai S and Huleihel M (2014) Classification of *Colletotrichum coccodes* isolates into vegetative compatibility groups using infrared attenuated total reflectance spectroscopy and multivariate analysis. *Methods* **68**, 325–330.
- Samokhvalov A, Liu Y and Simon JD (2004) Characterization of the Fe (III)-binding site in sepia eumelanin by resonance Raman confocal microspectroscopy. *Journal of Photochemistry and Photobiology* **80**, 84–88.
- Selbmann L, De Hoog GS, Mazzaglia A, Friedmann EI and Onofri S (2005) Fungi at the edge of life: cryptoendolithic black fungi from Antarctic desert. *Studies in Mycology* **51**, 1–32.
- Selbmann L, Pacelli C, Zucconi L, Dadachova E, Moeller R, de Vera JP and Onofri S (2018) Resistance of an Antarctic cryptoendolithic black fungus to radiation gives new insights of astrobiological relevance. *Fungal Biology* **122**, 546–554.
- Shurvell HF (2006) Spectra–structure correlations in the mid- and far-infrared. In Chalmers JM and Griffiths PR (eds), *Handbook of Vibrational Spectroscopy*, vol. 1. Chichester, UK: John Wiley & Sons, Ltd, pp. 1–34.
- Simpson JA (1983) Elemental and isotopic composition of the galactic cosmic rays. *Annual Review of Nuclear and Particle Science* **33**, 323–382.
- Speranza M, Ascaso C, Delclòs Martínez X and Peñalver E (2015) Cretaceous mycelia preserving fungal polysaccharides: taphonomic and paleoecological potential of microorganisms preserved in fossil resins. *Geologica Acta: An International Earth Science Journal* **13**, 363–385.
- Summons RE, Albrecht P, McDonald G and Moldowan JM (2008) Molecular biosignatures. In Botta O, Bada JL, Gomez-Elvira J, Javaux E, Selsis F and Summons R (eds). *Strategies of Life Detection*, Space Sciences Series of ISSI, vol. 25. Boston, MA: Springer, pp. 133–159.
- Sun S, Zhang X, Sun S, Zhang L, Shan S and Zhu H (2016) Production of natural melanin by *Auricularia auricula* and study on its molecular structure. *Food Chemistry* **190**, 801–807.
- Trevors JT (2003) Possible origin of a membrane in the subsurface of the earth. *Cell Biology International* **27**, 451–457.
- Turick CE, Ekechukwu AA, Milliken CE, Casadevall A and Dadachova E (2011) Gamma radiation interacts with melanin to alter its oxidation–reduction potential and results in electric current production. *Bioelectrochem* **82**, 69–73.
- Vember VV and Zhdanova NN (2001) Peculiarities of linear growth of the melanin-containing fungi *Cladosporium sphaerospermum* Penz. and *Alternaria alternata* (Fr.) Keissler. *Journal of Microbiology* **63**, 3–12.
- Verseux C, Baqué M, Cifariello R, Faglarone C, Raguse M, Moeller R and Billi D (2017) Evaluation of the resistance of *Chroococcidiopsis* spp. to sparsely and densely ionizing irradiation. *Astrobiology* **17**, 118–125.
- Vilgalys R and Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**, 4238–4246.