



ON STOCHASTIC CONTROL UNDER POISSONIAN INTERVENTION: OPTIMALITY OF A BARRIER STRATEGY IN A GENERAL LÉVY MODEL

KEI NOBA *, *The Institute of Statistical Mathematics*

KAZUTOSHI YAMAZAKI ,** *The University of Queensland*

Abstract

We study a version of the stochastic control problem of minimizing the sum of running and controlling costs, where control opportunities are restricted to independent Poisson arrival times. Under a general setting driven by a general Lévy process, we show the optimality of a periodic barrier strategy, which moves the process upward to the barrier whenever it is observed to be below it. The convergence of the optimal solutions to those in the continuous-observation case is also shown.

Keywords: Stochastic control; inventory models; periodic observations; mathematical finance; Lévy processes

2020 Mathematics Subject Classification: Primary 60G51

Secondary 93E20; 90B05

1. Introduction

Stochastic control aims to obtain an optimal dynamic strategy in cases of uncertainty. In its typical formulation, the problem reduces to obtaining an adapted control process that maximizes/minimizes the expected total reward/cost, which depends on the paths of the controlling and controlled processes. The continuous-time stochastic control research, active in various fields such as financial/actuarial mathematics and research on inventory models, has been developed along with stochastic analysis and differential equations theory. In contrast to its discrete-time counterpart, for which numerical approaches are typically required, various analytical approaches, such as Itô calculus and first passage analysis, are available in continuous-time models to obtain explicit results.

Poissonian observation/intervention models have been developed to explore the interface between continuous-time and discrete-time models. The earliest papers on this model include those of Wang [33] and Dupuis and Wang [15] for Brownian motion models. More recently, these results have been extended to spectrally one-sided Lévy models, as discussed in, for example, [1–3], [23], [27–29], and [39–41]. For a comprehensive survey on this subject, see Saariinen [32] and the references therein. In the Poissonian model, instead of allowing the decision maker to observe the state process continuously and control it at all times, these

Received 21 October 2023; accepted 26 August 2024.

* Postal address: Department of Fundamental Statistical Mathematics, The Institute of Statistical Mathematics, 10-3 Midori-cho, Tachikawa-shi, Tokyo 190-8562, Japan. Email address: knoba@ism.ac.jp

** Postal address: School of Mathematics and Physics, The University of Queensland, St Lucia, Brisbane, QLD 4072, Australia. Email address: k.yamazaki@uq.edu.au

© The Author(s), 2025. Published by Cambridge University Press on behalf of Applied Probability Trust.

opportunities are given only at independent Poisson arrival times. Although this assumption of Poisson arrivals is indeed restrictive in real applications, it provides a more flexible approach for approximating the discrete-time counterpart (with deterministic interarrivals) than the classical continuous-time model. As confirmed numerically in studies such as [22], approximation via Poisson arrivals (as a special case of Erlangization [11, 14, 21]) often achieves accurate approximation of the discrete-time model in stochastic control problems.

This paper studies the classical stochastic control problem, described as follows. Given a stochastic process $X = \{X_t : t \geq 0\}$, the objective is to choose a strategy $\pi = \{R_t^\pi : t \geq 0\}$ to minimize the total expected values of the running cost $\int_0^\infty e^{-qt} f(U_t^\pi) dt$ and the controlling cost $\int_{[0, \infty)} e^{-qt} dR_t^\pi$, where $U^\pi := X + R^\pi$ is the controlled process when π is applied. More precisely, we want to minimize over π the expected sum

$$v_\pi(x) := \mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^\pi) dt + C \int_0^\infty e^{-qt} dR_t^\pi \right] \quad \text{for } C \in \mathbb{R}.$$

This framework enables the modeling of various optimization scenarios by suitably selecting the process X . See [5], [6], and [7] for inventory models, and [10], [18], and [24] for financial applications.

This problem has been studied in several papers when X is a spectrally negative Lévy process (i.e. a Lévy process with only negative jumps). Under the assumption that the running cost function is convex, the barrier strategy, with the lower barrier b^* selected to be a unique root of

$$\mathbb{E}_b \left[\int_0^\infty e^{-qt} f'_+(U_t^b) dt \right] + C = 0, \quad (1.1)$$

is optimal. Here, U^b is the reflected process starting at b . Interestingly, this optimality result continues to hold in different formulations with additional constraints on the admissible strategies. In a version where R^π is restricted to being absolutely continuous with respect to the Lebesgue measure with a given density bound [17], the same optimality result holds, with U^b being the so-called refracted processes [20, 26, 31, 34, 35, 38]. The Poissonian observation version we consider in this paper has been solved by Pérez, Yamazaki, and Bensoussan [30] for the spectrally negative case. In this case, U^b is a version of the reflected process that is pushed to b whenever it is observed to be below it. By selecting the barrier using (1.1), this version of the barrier strategy, which we call the periodic barrier strategy, has been shown to be optimal.

The results described above all rely on the so-called scale function (see [8], [16], [19], and [36]), which makes sense only for spectrally one-sided Lévy processes. However, the spectrally negative assumption is often unrealistic in real applications. For example, financial asset prices are empirically known to have both positive and negative jumps (see [12]); also, water storage levels of dams experience both positive and negative jumps, due to rainfall and surges in consumption. See also the introduction of [13] for the application of processes of two-sided jumps in modeling the surplus of an insurance company.

Although the existing results for a general Lévy process in stochastic control are significantly limited in comparison with diffusion and spectrally one-sided Lévy models, the problem described above has recently been solved for a general Lévy process in the continuous-observation setting. Noba and Yamazaki [25] have shown that the classical barrier strategy described in (1.1) continues to be optimal even in the presence of positive jumps. It is thus a natural conjecture that the form of optimal strategy is invariant to the existence of upward

jumps. The objective of this paper is to verify this conjecture. We solve the Poissonian observation case for a general Lévy process X with both positive and negative jumps, generalizing the results of [25] and [30] simultaneously, and provide a unified way of expressing the optimal strategy. Despite the obvious difficulty over the continuous-observation model, for which many analytical results are available for classical reflected processes, we provide a more concise proof than those given in [25].

The remainder of the paper is organized as follows. In Section 2, we formally define the problem under consideration. In Section 3, we define periodic barrier strategies and obtain their key properties. Then, in Section 4, we select the barrier and demonstrate its optimality. In Section 5, we show the convergence to the results in the classical setting as the rate of observation approaches infinity. These results are confirmed with numerical experiments in Section 6. Finally, we conclude the paper in Section 7. Some proofs are deferred to the appendix. Throughout the paper, we let $g'_+(\cdot)$ and $g'_-(\cdot)$ be the right-hand and left-hand derivatives of any function g whenever they make sense.

2. Problem

Let $X = \{X_t : t \geq 0\}$ be a (one-dimensional) Lévy process defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For $x \in \mathbb{R}$, we let \mathbb{P}_x denote the law of X when its initial value is x , and write $\mathbb{P} = \mathbb{P}_0$ for the case $x = 0$. Let Ψ be the characteristic exponent of X , i.e. $e^{-t\Psi(\lambda)} = \mathbb{E}[e^{i\lambda X_t}]$, $\lambda \in \mathbb{R}$ and $t \geq 0$. It is known to admit the form

$$\Psi(\lambda) := -i\gamma\lambda + \frac{1}{2}\sigma^2\lambda^2 + \int_{\mathbb{R} \setminus \{0\}} (1 - e^{i\lambda z} + i\lambda z 1_{\{|z| < 1\}})\Pi(dz), \quad \lambda \in \mathbb{R},$$

for some $\gamma \in \mathbb{R}$, $\sigma \geq 0$, and a Lévy measure Π on $\mathbb{R} \setminus \{0\}$ satisfying $\int_{\mathbb{R} \setminus \{0\}} (1 \wedge z^2)\Pi(dz) < \infty$.

We consider a version of the stochastic control problem defined as follows. The set of control opportunities

$$\mathcal{T}_\eta := \{T(k) : k \in \mathbb{N}\}$$

are given by the arrival times of a Poisson process $N^\eta = \{N_t^\eta : t \geq 0\}$ with intensity $\eta > 0$ which is independent of X . In other words, the interarrival times $\{T(k) - T(k - 1) : k \in \mathbb{N}\}$ are an i.i.d. sequence of exponential random variables with intensity η , where we let $T(0) = 0$ for notational convenience. Let $\mathbb{F} := \{\mathcal{F}_t : t \geq 0\}$ be the natural filtration generated by (X, N^η) . A strategy, representing the cumulative amount of controlling, $\pi = \{R_t^\pi : t \geq 0\}$, is a process of the form

$$R_t^\pi = \int_{[0,t]} v_s^\pi dN_s^\eta = \sum_{0 \leq s \leq t: N_s^\eta \neq N_{s-}^\eta} v_s^\pi, \quad t \geq 0, \tag{2.1}$$

for some càglàd (left-continuous with right limits) and non-negative \mathbb{F} -adapted process $v^\pi = \{v_t^\pi : t \geq 0\}$, where it is understood that $N_{0-}^\eta = 0$. The corresponding controlled process becomes

$$U_t^\pi = X_t + R_t^\pi, \quad t \geq 0.$$

We focus on the case where we can control the state process in one direction, and hence $v_s^\pi \geq 0$ a.s. for all $\pi \in \mathcal{A}$, which is standard as in [5–7] and [17]. Such an assumption is applicable in many inventory models where only replenishment is allowed, as well as in dam management scenarios where the water level can only be decreased by the decision maker.

For a given discount factor $q > 0$ and initial value $x \in \mathbb{R}$, the objective is to minimize

$$\begin{aligned} v_\pi(x) &:= \mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^\pi) dt + C \int_0^\infty e^{-qt} dR_t^\pi \right] \\ &= \mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^\pi) dt + C \sum_{0 \leq t < \infty: N_t^\pi \neq N_{t-}^\pi} e^{-qt} v_t^\pi \right], \end{aligned}$$

which is the sum of running costs for a given measurable function $f: \mathbb{R} \rightarrow \mathbb{R}$ and a controlling cost/reward for a unit cost/reward $C \in \mathbb{R}$ (cost if it is positive and reward if negative). Let \mathcal{A} be the set of all admissible strategies satisfying the constraints described above as well as the integrability condition:

$$\begin{aligned} &\mathbb{E}_x \left[\int_0^\infty e^{-qt} |f(U_t^\pi)| dt + \int_0^\infty e^{-qt} dR_t^\pi \right] \\ &= \mathbb{E}_x \left[\int_0^\infty e^{-qt} |f(U_t^\pi)| dt + \sum_{0 \leq t < \infty: N_t^\pi \neq N_{t-}^\pi} e^{-qt} v_t^\pi \right] \\ &< \infty. \end{aligned}$$

The aim of the problem is to obtain the (*optimal*) *value function*

$$v(x) := \inf_{\pi \in \mathcal{A}} v_\pi(x), \quad x \in \mathbb{R},$$

and an optimal strategy π^* such that $v_{\pi^*}(x) = v(x)$ (if such a strategy exists).

For the running cost function f , the unit cost/reward C and the Lévy process X , we impose the same conditions as those assumed in [25]; similar conditions are commonly assumed in the literature (see [5], [7], [17], and [37]).

Assumption 2.1. (Assumption on f and C .)

- (1) *The function f is convex.*
- (2) *There exist $k_1, k_2 > 0$ and $N \in \mathbb{N}$ such that $|f(x)| \leq k_1 + k_2|x|^N$ for all $x \in \mathbb{R}$.*
- (3) *We have $f'_+(-\infty) < -Cq < f'_+(\infty)$ where $f'_+(-\infty) := \lim_{x \rightarrow -\infty} f'_+(x) \in [-\infty, \infty)$ and $f'_+(\infty) := \lim_{x \rightarrow \infty} f'_+(x) \in (-\infty, \infty]$.*

Remark 2.1. Examples of f satisfying the above assumptions include classical examples such as $f(x) = x^2$ and $f(x) = |x|$, as well as asymmetric functions used for our numerical examples (6.1) in Section 6.

Note that the right- and left-hand derivatives $f'_+(x)$ and $f'_-(x)$, respectively, for all $x \in \mathbb{R}$ as well as their limits are well-defined by Assumption 2.1(1). Assumption 2.1(3) is necessary to avoid the optimality of a trivial strategy and the case optimal strategy does not exist; see [25, Remark 1]. More precisely, when this assumption is violated, the optimal strategy is never to modify the process, or to move the process to an arbitrarily large value.

Assumption 2.2. (Assumption on X .)

- (1) *X is not a (driftless) compound Poisson process.*
- (2) *For some $\bar{\theta} > 0$, $\int_{\mathbb{R} \setminus (-1, 1)} e^{\bar{\theta}|z|} \Pi(dz) < \infty$.*

Remark 2.2. By Assumption 2.2(1) and [8, Proposition I.15], the potential measure of X has no atoms. This also shows that

$$\mathbb{P}_x(X_{T(1)} = b) = \eta \mathbb{E}_x \left[\int_0^\infty e^{-\eta t} 1_{\{X_t=b\}} dt \right] = 0 \quad \text{for all } x, b \in \mathbb{R}.$$

Assumption 2.2(2) together with [19, Theorem 3.6] guarantees the finiteness of $\mathbb{E}[\exp(\bar{\theta}|X_1|)]$ and also that of $\mathbb{E}[|X_1|]$ (since $\exp(x) \geq x$ for $x \geq 0$).

Remark 2.3. From Assumptions 2.1(2) and 2.2(2) and by the proof of [37, Lemma 11], the expectation

$$\mathbb{E}_x \left[\int_0^\infty e^{-qt} |f(X_t)| dt \right]$$

is finite and it is at most of polynomial growth as $x \uparrow \infty$ and $x \downarrow -\infty$.

3. Periodic barrier strategies

Our objective is to show the optimality of a periodic barrier strategy π^b for a suitable selection of the barrier $b \in \mathbb{R}$ in the considered stochastic control problem.

Fix $b \in \mathbb{R}$. A periodic barrier strategy π^b pushes the process upward to b whenever it is observed to be below b (see Figure 1). The epochs of controlling $\{T_b^{(n)} : n \in \mathbb{N}\} \subset \mathcal{T}_\eta$ are given by a sequence of \mathbb{F} -stopping times, recursively defined as follows: with $T_b^{(0)} := 0$,

$$T_b^{(n)} = \inf\{t \in \mathcal{T}_\eta : t > T_b^{(n-1)}, \tilde{X}_t^{(n-1),b} < b\}, \quad n \geq 1,$$

where

$$\tilde{X}_t^{(m),b} := \begin{cases} X_t, & m = 0, \\ b + (X_t - X_{T_b^{(m)}}), & m \geq 1, \end{cases}$$

is a parallel shift of X so that it starts from b at $T_b^{(m)}$ when $m \geq 1$. The strategy π^b modifies X by adding at $T_b^{(n)}$ the shortage $b - \tilde{X}_{T_b^{(n)}}^{(n-1),b}$ so that the path of the controlled process is the concatenation of $(\tilde{X}^{(m),b})_{m \geq 1}$. The corresponding control and controlled processes, respectively, can be written as

$$R_t^b := R_t^{\pi^b} = \sum_{n=1}^\infty (b - \tilde{X}_{T_b^{(n)}}^{(n-1),b}) 1_{\{T_b^{(n)} \leq t\}},$$

$$U_t^b := U_t^{\pi^b} = X_t + R_t^b = \sum_{n=1}^\infty \tilde{X}_t^{(n-1),b} 1_{\{t \in [T_b^{(n-1)}, T_b^{(n)})\}}.$$

Note that X and N^η do not jump at the same time.

Alternatively, in terms of the càglàd \mathbb{F} -adapted process $v^b = \{v_t^b : t \geq 0\}$ with

$$v_t^b = \begin{cases} (b - X_{t-})^+, & t \in [0, T_b^{(1)}], \\ (X_{T_b^{(n-1)}} - X_{t-})^+, & t \in (T_b^{(n-1)}, T_b^{(n)}) \text{ with } n \geq 2, \end{cases}$$

where $x^+ := x \vee 0$ and it is understood that $X_{0-} = X_0$, it can be also written as

$$R_t^b = \int_{[0,t]} v_s^b dN_s^\eta, \quad t \geq 0. \tag{3.1}$$




FIGURE 1. Sample path of U^b . Control opportunities \mathcal{T}_η are shown by dotted vertical lines. The control times $\{T_b^{(n)} : n \in \mathbb{N}\}$ and control sizes ΔR^b are indicated by the vertical red lines.

Remark 3.1. In terms of the minimum of X observed until time t , we can also write

$$R_t^b = \max_{1 \leq k \leq N_t^\eta} (b - X_{T(k)})^+, \quad t \geq 0,$$

and $T_b^{(n)}$ as the n th jump time of R^b . This expression will be used to show the convergence to the classical case in Section 5.

For the rest of the paper, we denote the expected total cost under the periodic barrier strategy π^b by

$$v_b(x) := v_{\pi^b}(x), \quad x \in \mathbb{R}.$$

We now show the admissibility of periodic barrier strategies along with related results. The proof of the following lemma is deferred to Appendix A.1.

Lemma 3.1 For $x, b \in \mathbb{R}$,

- (i) $\mathbb{E}_x[\int_0^\infty e^{-qt} |f(U_t^b)| dt] < \infty$,
- (ii) $\mathbb{E}_x[\int_0^\infty e^{-qt} dR_t^b] < \infty$,
- (iii) $x \mapsto v_b(x)$ is at most of polynomial growth.

As a corollary of the above, we also have the following. Thanks to Assumption 2.1(1), this can be shown exactly in the same way as the proof of [25, Lemma 4] and thus we omit the proof.

Corollary 3.1. *For $x, b \in \mathbb{R}$, we have*

$$\mathbb{E}_x \left[\int_0^\infty e^{-qt} |f'_+(U_t^b)| dt \right] < \infty.$$

Lemma 3.1 together with (3.1) shows the following.

Proposition 3.1. *For $b \in \mathbb{R}$, the strategy π^b is admissible.*

Let $T_b := T_b^{(1)}$ be the first control time under the policy π^b . We conclude this section with the expression of the slope of v_b written in terms of T_b and the uncontrolled Lévy process X .

Proposition 3.2. *For $b \in \mathbb{R}$, the function v_b is continuously differentiable with its derivative*

$$v'_b(x) = \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_+(X_t) dt \right] - C \mathbb{E}_x [e^{-qT_b}], \quad x \in \mathbb{R}. \tag{3.2}$$

The proof of Proposition 3.2 requires the following continuity result of T_b ; its proof is deferred to Appendix A.2.

Lemma 3.2. *For fixed $b \in \mathbb{R}$, we have $\lim_{b' \rightarrow b} T_{b'} = T_b$ on $\{T_b < \infty\}$, almost surely.*

Proof of Proposition 3.2. By Lemma 3.1, we can decompose the expected costs as follows:

$$v_b(x) = v_b^{(1)}(x) + Cv_b^{(2)}(x), \quad x \in \mathbb{R},$$

where we write

$$v_b^{(1)}(x) := \mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^b) dt \right], \quad v_b^{(2)}(x) := \mathbb{E}_x \left[\int_0^\infty e^{-qt} dR_t^b \right].$$

For $y \in \mathbb{R}$, we write $X_t^{[y]} := X_t + y, t \geq 0$, and let $U_t^{[y],b}, R_t^{[y],b}, T_b^{[y]}$ be those corresponding to this shifted process.

(i) Fix $b \in \mathbb{R}$ and $\varepsilon > 0$. We show that $t \mapsto U_t^{[\varepsilon],b} - U_t^b = \varepsilon + R_t^{[\varepsilon],b} - R_t^b$ is non-increasing and always lies on $[0, \varepsilon]$. Because this difference is a step function in t with jump times contained in the set \mathcal{T}_η , it suffices to show that

$$\zeta(k) := U_{T(k)}^{[\varepsilon],b} - U_{T(k)}^b = \varepsilon + R_{T(k)}^{[\varepsilon],b} - R_{T(k)}^b, \quad k \geq 0,$$

is non-increasing in k and takes values only on $[0, \varepsilon]$. We show this claim by induction.

First it holds trivially when $k = 0$ with $\zeta(0) = \varepsilon \in [0, \varepsilon]$.

Now, suppose it holds that $\zeta(k) \in [0, \varepsilon]$ for some $k \geq 0$. With the set of indices of controlling,

$$A^{[\delta]} := \{k \geq 1 : \Delta R_{T(k)}^{[\delta],b} > 0\} = \{k \geq 1 : U_{T(k-1)}^{[\delta],b} + (X_{T(k)} - X_{T(k-1)}) < b\}, \quad \delta = 0, \varepsilon,$$

we have

$$\begin{aligned} \{k + 1 \in A^{[\varepsilon]}\} &= \{U_{T(k)}^b + \zeta(k) + (X_{T(k+1)} - X_{T(k)}) < b\} \\ &\subset \{U_{T(k)}^b + (X_{T(k+1)} - X_{T(k)}) < b\} \\ &= \{k + 1 \in A^{[0]}\}. \end{aligned} \tag{3.3}$$

(A) Suppose $k + 1 \in A^{[\varepsilon]}$ so that $U_{T(k+1)}^{[\varepsilon],b} = b$. By (3.3), this also implies $k + 1 \in A^{[0]}$ or equivalently $U_{T(k+1)}^b = b$. Hence, $\zeta(k + 1) = 0$.

(B) Suppose $k + 1 \notin A^{[\varepsilon]}$ so that

$$U_{T(k)}^{[\varepsilon],b} + (X_{T(k+1)} - X_{T(k)}) \geq b. \tag{3.4}$$

(a) Suppose $k + 1 \notin A^{[0]}$, then because $\Delta R_{T(k+1)}^{[\varepsilon],b} = \Delta R_{T(k+1)}^b = 0$, we have $\zeta(k + 1) = \zeta(k) \in [0, \varepsilon]$.

(b) Suppose $k + 1 \in A^{[0]}$. Then, clearly $\zeta(k + 1) = \zeta(k) - \Delta R_{T(k+1)}^b < \zeta(k)$. In addition, by (3.4),

$$\zeta(k + 1) = (U_{T(k)}^{[\varepsilon],b} + (X_{T(k+1)} - X_{T(k)})) - b \geq 0.$$

In sum, in all cases we have $\zeta(k + 1) \leq \zeta(k)$, and in addition, $\zeta(k + 1) \in [0, \varepsilon]$. By mathematical induction we have that ζ is non-increasing and always lies in $[0, \varepsilon]$. In view of (3.3), this also shows $A^{[\varepsilon]} \subset A^{[0]}$.

At the moment $T_b^{[\varepsilon]} = \inf_{k \in A^{[\varepsilon]}} T(k)$ with $\inf \emptyset = \infty$, the difference between $U^{[\varepsilon],b}$ and U^b becomes 0 and must stay at 0 afterwards. On the other hand, before T_b there is no control for both and the difference is ε . In sum,

$$U_t^{[\varepsilon],b} - U_t^b = \begin{cases} \varepsilon, & t \in [0, T_b), \\ 0, & t \in [T_b^{[\varepsilon]}, \infty), \end{cases} \quad R_t^{[\varepsilon],b} - R_t^b = \begin{cases} 0, & t \in [0, T_b), \\ -\varepsilon, & t \in [T_b^{[\varepsilon]}, \infty). \end{cases} \tag{3.5}$$

(ii) By (3.5), we have

$$\begin{aligned} \frac{v_b^{(1)}(x + \varepsilon) - v_b^{(1)}(x)}{\varepsilon} &= \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} \frac{f(U_t^b + \varepsilon) - f(U_t^b)}{\varepsilon} dt \right] \\ &\quad + \mathbb{E}_x \left[\int_{T_b}^{T_b^{[\varepsilon]}} e^{-qt} \frac{f(U_t^{[\varepsilon],b}) - f(U_t^b)}{\varepsilon} dt \right]. \end{aligned}$$

By (i) (in particular that the process $\{U_t^{[\varepsilon],b} - U_t^b : t \geq 0\}$ is non-increasing), mean value theorem, and the convexity of f , for all $0 < \varepsilon < \bar{\varepsilon}$,

$$\begin{aligned} \left| \mathbb{E}_x \left[\int_{T_b}^{T_b^{[\varepsilon]}} e^{-qt} \frac{f(U_t^{[\varepsilon],b}) - f(U_t^b)}{\varepsilon} dt \right] \right| &\leq \mathbb{E}_x \left[\int_{T_b}^{T_b^{[\varepsilon]}} e^{-qt} \frac{|f(U_t^{[\varepsilon],b}) - f(U_t^b)|}{\varepsilon} dt \right] \\ &\leq \mathbb{E}_x \left[\int_{T_b}^{T_b^{[\varepsilon]}} e^{-qt} \sup_{y \in [U_t^b, U_t^b + \bar{\varepsilon}]} |f'_+(y)| dt \right] \\ &\leq \mathbb{E}_x \left[\int_{T_b}^{T_b - \varepsilon} e^{-qt} (|f'_+(U_t^b)| + |f'_+(U_t^b + \bar{\varepsilon})|) dt \right] \\ &\xrightarrow{\varepsilon \downarrow 0} 0, \end{aligned}$$

where $T_b^{[\varepsilon]} = T_{b-\varepsilon}$ holds because

$$\{t < T_b^{[\varepsilon]}\} = \left\{ \min_{1 \leq k \leq N_t^\eta} X_{T(k)}^{[\varepsilon]} \geq b \right\} = \left\{ \min_{1 \leq k \leq N_t^\eta} X_{T(k)} \geq b - \varepsilon \right\} = \{t < T_{b-\varepsilon}\}$$

(see Remark 3.1) and the last limit holds by monotone convergence and Lemma 3.2. Note that the finiteness of the expectations above hold by Corollary 3.1. Now, by the convexity of f , monotone convergence gives

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \frac{v_b^{(1)}(x + \varepsilon) - v_b^{(1)}(x)}{\varepsilon} &= \lim_{\varepsilon \downarrow 0} \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} \frac{f(U_t^b + \varepsilon) - f(U_t^b)}{\varepsilon} dt \right] \\ &= \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_+(U_t^b) dt \right]. \end{aligned}$$

In the same way, we compute the left derivative. By (i), with x changed to $x - \varepsilon$, we have

$$\frac{v_b^{(1)}(x) - v_b^{(1)}(x - \varepsilon)}{\varepsilon} = \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} \frac{f(U_t^b) - f(U_t^b - \varepsilon)}{\varepsilon} dt \right] + h(\varepsilon),$$

where

$$h(\varepsilon) := \mathbb{E}_x \left[\int_{T_b^{[-\varepsilon]}}^{T_b} e^{-qt} \frac{f(U_t^b) - f(U_t^{[-\varepsilon],b})}{\varepsilon} dt \right] - \mathbb{E}_x \left[\int_{T_b^{[-\varepsilon]}}^{T_b} e^{-qt} \frac{f(U_t^b) - f(U_t^b - \varepsilon)}{\varepsilon} dt \right].$$

For all $0 < \varepsilon < \bar{\varepsilon}$, the mean value theorem and the convexity of f give

$$\left| \frac{f(U_t^b) - f(U_t^b - \varepsilon)}{\varepsilon} \right| \vee \left| \frac{f(U_t^b) - f(U_t^{[-\varepsilon],b})}{\varepsilon} \right| \leq |f'_+(U_t^b - \bar{\varepsilon})| + |f'_+(U_t^b)|, \quad t \geq 0,$$

and thus

$$|h(\varepsilon)| \leq 2\mathbb{E}_x \left[\int_{T_{b+\varepsilon}}^{T_b} e^{-qt} (|f'_+(U_t^b - \bar{\varepsilon})| + |f'_+(U_t^b)|) dt \right] \xrightarrow{\varepsilon \downarrow 0} 0,$$

where we used $T_b^{[-\varepsilon]} = T_{b+\varepsilon}$ and Lemma 3.2. Therefore, as in the case of the right derivative, we have by monotone convergence

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \frac{v_b^{(1)}(x) - v_b^{(1)}(x - \varepsilon)}{\varepsilon} &= \lim_{\varepsilon \downarrow 0} \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} \frac{f(U_t^b) - f(U_t^b - \varepsilon)}{\varepsilon} dt \right] \\ &= \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_-(U_t^b) dt \right]. \end{aligned}$$

Because the right and left derivatives coincide thanks to Remark 2.2 and $U_t^b = X_t$ for $t < T_b$,

$$v_b^{(1)'}(x) = \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_+(U_t^b) dt \right] = \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_+(X_t) dt \right].$$

(iii) We now show

$$\lim_{\varepsilon \downarrow 0} \frac{v_b^{(2)}(x + \varepsilon) - v_b^{(2)}(x)}{\varepsilon} = \lim_{\varepsilon \downarrow 0} \frac{v_b^{(2)}(x) - v_b^{(2)}(x - \varepsilon)}{\varepsilon} = -\mathbb{E}_x [e^{-qT_b}]. \tag{3.6}$$

Indeed, since the process $\{R_t^{[\varepsilon],b} - R_t^b : t \geq 0\}$ is non-increasing by (i) and from (3.5), we have

$$-\mathbb{E}_x [e^{-qT_b}] \leq \frac{v_b^{(2)}(x + \varepsilon) - v_b^{(2)}(x)}{\varepsilon} \leq -\mathbb{E}_x [e^{-qT_{b+\varepsilon}}]. \tag{3.7}$$

By Lemma 3.2 and (3.7), we have that the first term in (3.6) is equal to the third term in (3.6). By changing from x to $x - \varepsilon$ in the above argument, we have the second equality in (3.6).

From (ii) and (iii), we obtain (3.2).

(iv) It remains to show that $x \mapsto v'_b(x)$ is continuous. We have

$$\begin{aligned} & |v'_b(x + \varepsilon) - v'_b(x)| \\ & \leq \left| \mathbb{E}_{x+\varepsilon} \left[\int_0^{T_b} e^{-qt} f'_+(X_t) dt \right] - \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_+(X_t) dt \right] \right| \\ & \quad + |C| \left| \mathbb{E}_{x+\varepsilon} [e^{-qT_b}] - \mathbb{E}_x [e^{-qT_b}] \right| \\ & = \left| \mathbb{E}_x \left[\int_0^{T_{b-\varepsilon}} e^{-qt} f'_+(X_t + \varepsilon) dt \right] - \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} f'_+(X_t) dt \right] \right| \\ & \quad + |C| \left| \mathbb{E}_x [e^{-qT_{b-\varepsilon}}] - \mathbb{E}_x [e^{-qT_b}] \right| \\ & \leq \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} |f'_+(X_t + \varepsilon) - f'_+(X_t)| dt \right] + \mathbb{E}_x \left[\int_{T_b}^{T_{b-\varepsilon}} e^{-qt} |f'_+(X_t + \varepsilon)| dt \right] \\ & \quad + |C| \left| \mathbb{E}_x [e^{-qT_{b-\varepsilon}}] - \mathbb{E}_x [e^{-qT_b}] \right|. \end{aligned}$$

As $\varepsilon \searrow 0$, the first expectation converges to zero by monotone convergence, the second expectation converges to zero by monotone convergence and Lemma 3.2. The last expectation converges to zero by Lemma 3.2. By replacing x with $x - \varepsilon$ and again using Remark 2.2, we also have the left continuity. \square

4. The optimal barrier b^* in the periodic barrier strategies

In this section, we show the optimality of a periodic barrier strategy. Define

$$\rho(b) := \mathbb{E}_b \left[\int_0^\infty e^{-qt} f'_+(U_t^b) dt \right], \quad b \in \mathbb{R}, \tag{4.1}$$

which takes real values by Corollary 3.1. Our candidate optimal barrier is

$$b^* := \inf\{b \in \mathbb{R} : \rho(b) + C \geq 0\}, \tag{4.2}$$

which is well-defined by Lemma 4.1 below; see Appendix A.3 for the proof.

Lemma 4.1. *The function ρ is non-decreasing and continuous. We also have $\lim_{b \uparrow \infty} \rho(b) = f'_+(\infty)/q > -C$ and $\lim_{b \downarrow -\infty} \rho(b) = f'_+(-\infty)/q < -C$.*

We now state the main result of the paper.

Theorem 4.1. *The periodic barrier strategy at b^* is an optimal strategy and thus we have $v(x) = v_{b^*}(x)$ for $x \in \mathbb{R}$.*

In the remaining part, we show Theorem 4.1. Acting on a measurable function $g: \mathbb{R} \rightarrow \mathbb{R}$ belonging to $C^1(\mathbb{R})$ (resp. $C^2(\mathbb{R})$) when X has bounded (resp. unbounded) variation paths with at most polynomial growth, define the operator

$$\begin{aligned} \mathcal{L}g(x) & := \gamma g'(x) + \frac{1}{2} \sigma^2 g''(x) \\ & \quad + \int_{\mathbb{R} \setminus \{0\}} (g(x+z) - g(x) - g'(x)z 1_{\{|z|<1\}}) \Pi(dz), \quad x \in \mathbb{R}. \end{aligned}$$

Let $(\mathcal{L} - q)g := \mathcal{L}g - qg$. Define also, for any measurable function $g: \mathbb{R} \rightarrow \mathbb{R}$,

$$\mathcal{M}g(x) := \inf_{l \geq 0} \{Cl + g(x + l)\}, \quad x \in \mathbb{R}.$$

The following verification lemma gives a sufficient condition for optimality. The proof is the same as that for the spectrally negative case in [30, Lemma 3.1] and hence we omit it.

Lemma 4.2. (Verification lemma.) *Let $w: \mathbb{R} \rightarrow \mathbb{R}$ be of polynomial growth and belong to $C^1(\mathbb{R})$ (resp. $C^2(\mathbb{R})$) when X has bounded (resp. unbounded) variation paths. If it satisfies*

$$(\mathcal{L} - q)w(x) + \eta(\mathcal{M}w(x) - w(x)) + f(x) = 0, \quad x \in \mathbb{R},$$

then we have $w(x) \leq v(x)$ for $x \in \mathbb{R}$.

Before confirming these sufficient conditions for $w = v_{b^*}$, we explicitly compute $\mathcal{M}v_{b^*}$. To this end, we show the following, which is a direct consequence of Proposition 3.2.

Lemma 4.3. *For $x \in \mathbb{R}$, we have*

$$v'_{b^*}(x) = \mathbb{E}_x \left[\int_0^\infty e^{-qt} f'_+(U_t^{b^*}) dt \right].$$

Proof. By Lemma 4.1 and the definition of b^* , we have $\rho(b^*) + C = 0$. This together with the strong Markov property gives

$$-C\mathbb{E}_x[e^{-qT_{b^*}}] = \mathbb{E}_x[e^{-qT_{b^*}}]\rho(b^*) = \mathbb{E}_x \left[\int_{T_{b^*}}^\infty e^{-qt} f'_+(U_t^{b^*}) dt \right],$$

where we recall that $T_{b^*} := T_{b^*}^{(1)}$ is the first control time under the policy π^{b^*} . Substituting this in (3.2) gives the result. \square

From part (i) of the proof of Proposition 3.2, for each $t \geq 0$, $U_t^{b^*}$ is monotonically increasing in the start value $X_0 = x$. By this and Lemma 4.3, the derivative v'_{b^*} is non-decreasing. In addition, by the definition of b^* and the continuity of ρ as in Lemma 4.1, we have $v'_{b^*}(b^*) = -C$. Thus we have

$$v'_{b^*}(x) \begin{cases} \leq -C, & x < b^*, \\ \geq -C, & x \geq b^*. \end{cases}$$

Since the derivative of the function $l \mapsto Cl + v_{b^*}(x + l)$ is equal to $l \mapsto C + v'_{b^*}(x + l)$, it is minimized when $l = (b^* - x)^+$, showing the following.

Proposition 4.1. *We have*

$$\mathcal{M}v_{b^*}(x) = \begin{cases} v_{b^*}(x), & x \geq b^*, \\ C(b^* - x) + v_{b^*}(b^*), & x < b^*. \end{cases}$$

Regarding the smoothness of v_{b^*} , it belongs to $C^1(\mathbb{R})$ by Proposition 3.2. This is sufficient for the case of bounded variation, but care is needed for the unbounded variation case.

We temporarily assume the following, to first consider the case when the C^2 property of v_{b^*} is guaranteed.

Condition 4.1. *When X has unbounded variation paths, the running cost function f belongs to $C^2(\mathbb{R})$ and f'' has polynomial growth in the tails.*

The proof of the following lemma is deferred to Appendix A.4.

Lemma 4.4. *Suppose Condition 4.1 holds. When X has unbounded variation paths, the function v_{b^*} belongs to $C^2(\mathbb{R})$.*

We assume Condition 4.1 temporarily for Lemma 4.5. However, Condition 4.1 can be completely relaxed by following the arguments in Section 4.2 of [25]. We later provide a brief remark on how Condition 4.1 can be removed in the proof of Theorem 4.1.

By Proposition 3.2 and Lemma 4.4, the function $x \mapsto v_{b^*}(x)$ is sufficiently smooth to apply \mathcal{L} (under Condition 4.1).

Lemma 4.5. *Suppose Condition 4.1 holds. For $x \in \mathbb{R}$, we have $(\mathcal{L} - q)v_{b^*}(x) + \eta(\mathcal{M}v_{b^*}(x) - v_{b^*}(x)) + f(x) = 0$.*

Proof. It suffices to show $\mathcal{L}v_{b^*}(x) - (q + \eta)v_{b^*}(x) + h(x) = 0$ with

$$h(x) := f(x) + \eta\mathcal{M}v_{b^*}(x) = f(x) + \eta C(b^* - x)^+ + \eta v_{b^*}(x \vee b^*),$$

where the second equality holds by Proposition 4.1. Because v_{b^*} is smooth enough to apply Ito's formula (see Proposition 3.2 and Lemma 4.4), it is enough to show that the process $\{M_t : t \geq 0\}$, where

$$M_t := e^{-(q+\eta)t}v_{b^*}(X_t) + \int_0^t e^{-(q+\eta)s}h(X_s) ds,$$

is a local martingale with respect to the natural filtration $\{\mathcal{F}_t^X : t \geq 0\}$ generated by X . See the proof of [9, (12)].

By the strong Markov property and because $U_t^{b^*} = X_t$ for $t < T(1)$, we have, for $x \in \mathbb{R}$,

$$\begin{aligned} v_{b^*}(x) &= \mathbb{E}_x \left[\int_0^{T(1)} e^{-qt}f(X_t) dt \right] + C\mathbb{E}_x \left[e^{-qT(1)}(b^* - X_{T(1)})^+ \right] \\ &\quad + \mathbb{E}_x \left[e^{-qT(1)}v_{b^*}(X_{T(1)} \vee b^*) \right] \\ &= \mathbb{E}_x \left[\int_0^\infty e^{-(q+\eta)t}f(X_t) dt \right] + \eta C\mathbb{E}_x \left[\int_0^\infty e^{-(q+\eta)t}(b^* - X_t)^+ dt \right] \\ &\quad + \eta\mathbb{E}_x \left[\int_0^\infty e^{-(q+\eta)t}v_{b^*}(X_t \vee b^*) dt \right] \\ &= \mathbb{E}_x \left[\int_0^\infty e^{-(q+\eta)t}h(X_t) dt \right]. \end{aligned}$$

This together with the strong Markov property gives, for $t \geq 0$ and $\tau_{[n]} := \inf\{t > 0 : |X_t| > n\}$ with $n \in \mathbb{N}$,

$$\begin{aligned} & \mathbb{E}_x \left[\int_0^\infty e^{-(q+\eta)s} h(X_s) ds \mid \mathcal{F}_{t \wedge \tau_{[n]}}^X \right] \\ &= \int_0^{t \wedge \tau_{[n]}} e^{-(q+\eta)s} h(X_s) ds + \mathbb{E}_x \left[\int_{t \wedge \tau_{[n]}}^\infty e^{-(q+\eta)s} h(X_s) ds \mid \mathcal{F}_{t \wedge \tau_{[n]}}^X \right] \\ &= \int_0^{t \wedge \tau_{[n]}} e^{-(q+\eta)s} h(X_s) ds + e^{-(q+\eta)(t \wedge \tau_{[n]})} v_{b^*}(X_{t \wedge \tau_{[n]}}) = M_{t \wedge \tau_{[n]}}. \end{aligned}$$

By the tower property of conditional expectations, M is a local martingale. □

Proof of Theorem 4.1. By Lemmas 3.1(iii), 4.4, 4.5, and Proposition 3.2, the function v_{b^*} satisfies the conditions in Lemma 4.2. Thus $v_{b^*}(x) \leq v(x)$ for all $x \in \mathbb{R}$. Because π^{b^*} is admissible as in Proposition 3.1, the reverse inequality also holds. This completes the proof for the case when Condition 4.1 holds.

For the case when Condition 4.1 is violated, we can write the cost function f in terms of the limit of a sequence of $C^2(\mathbb{R})$ functions for which Condition 4.1 is fulfilled and the optimality of a barrier strategy holds. We omit the details because the proof is exactly the same as those proofs given in Section 4.2 of [25]. □

5. Convergence as $\eta \rightarrow \infty$

In this section we verify the convergence of the optimal solutions to the classical case [25] as the rate of observation $\eta \rightarrow \infty$.

Recall that in the classical case strategy R^π is any adapted (with respect to the natural filtration of X) and non-decreasing process, which does not have to be of the form (2.1). The classical barrier strategy with barrier $b \in \mathbb{R}$ is given by $R_t^{b,\infty} = (b - \underline{X}_t)^+$, where \underline{X} is the running infimum process of X and the corresponding controlled process is $U_t^{b,\infty} = X_t + R_t^{b,\infty}$, $t \geq 0$. As obtained in [25], the barrier strategy $\{R_t^{b^*,\infty} : t \geq 0\}$ with barrier

$$b_\infty^* := \inf\{b \in \mathbb{R} : \rho_\infty(b) + C \geq 0\} \quad \text{for } \rho_\infty(b) := \mathbb{E}_b \left[\int_0^\infty e^{-qt} f'_+(U_t^{b,\infty}) dt \right], \quad b \in \mathbb{R},$$

is optimal; the value function becomes

$$v_\infty^*(x) := \mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^{b_\infty^*,\infty}) dt + C \int_0^\infty e^{-qt} dR_t^{b_\infty^*,\infty} \right] \quad \text{for } x \in \mathbb{R}.$$

Solely in this section, to spell out the dependence on the rate η , we add super/subscript η in an obvious way and add ∞ for the classical case.

The objective is to show the convergence $b_\eta^* \rightarrow b_\infty^*$ and $v_\eta^* \rightarrow v_\infty^*$, where b_η^* is as defined in (4.2). The results hold except for a very particular case when X is the negative of a subordinator (where the reflected process becomes a constant).

Theorem 5.1. *We have (i) $b_\eta^* \searrow b_\infty^*$ as $\eta \rightarrow \infty$ and (ii) $v_\eta^* \searrow v_\infty^*$ as $\eta \rightarrow \infty$ uniformly in x on any compact set, where we assume f' is strictly increasing at b_∞^* for the case when X is the negative of a subordinator.*

Proof. Let $\{\eta_n : n \in \{0\} \cup \mathbb{N}\}$ be a strictly increasing (deterministic) sequence such that $\eta_0 = 0$ and $\eta_n \xrightarrow{n \uparrow \infty} \infty$. Consider, for each n , a Poisson process M^n with rate

$\lambda_n := \eta_n - \eta_{n-1} > 0$ independent of X , and let

$$N_t^{\eta_n} = \sum_{k=1}^n M_t^k, \quad t \geq 0.$$

We assume $\{M^n: n \geq 1\}$ are mutually independent and also independent of X . Hence, their superposition N^{η_n} becomes a Poisson process with rate η_n independent of X . We consider the problems driven by these processes (defined on the same probability space) to show the convergence.

(i) Fix $u \geq 0$. Let $\bar{\sigma}_n(u) := \inf\{s > u: \Delta N_s^{\eta_n} \neq 0\}$ and $\underline{\sigma}_n(u) := \sup\{s < u: \Delta N_s^{\eta_n} \neq 0\}$, respectively, be the first arrival time after u and the last arrival time before u of N^{η_n} (with the understanding $\sup \emptyset = 0$). Then

$$\mathbb{P}(\bar{\sigma}_n(u) - u > \varepsilon) = \mathbb{P}(N_{u+\varepsilon}^{\eta_n} - N_u^{\eta_n} = 0) = e^{-\varepsilon \eta_n} \xrightarrow{n \uparrow \infty} 0, \quad \varepsilon > 0.$$

In other words $\bar{\sigma}_n(u) \xrightarrow{n \uparrow \infty} u$ in probability. Because it is decreasing, the convergence also holds in the a.s.-sense. Similarly, we also have $\underline{\sigma}_n(u) \nearrow u$ a.s. as $n \rightarrow \infty$.

Fix $t > 0$ and $G(t) := \sup\{s \in [0, t]: X_{s-} \wedge X_s = \underline{X}_t\}$ (with $X_{0-} = X_0$). Suppose $G(t) \in (0, t)$. If $X_{G(t)-} \geq X_{G(t)}$ (i.e. X is continuous or jumps downward at $G(t)$), because X is right-continuous a.s., then

$$X_{\bar{\sigma}^n(G(t))} \xrightarrow{n \uparrow \infty} X_{G(t)} = X_{G(t)} \wedge X_{G(t)-} = \underline{X}_t.$$

If $X_{G(t)} > X_{G(t)-}$ (i.e. X jumps upward at $G(t)$), then

$$X_{\underline{\sigma}^n(G(t))} \xrightarrow{n \uparrow \infty} X_{G(t)-} = X_{G(t)} \wedge X_{G(t)-} = \underline{X}_t.$$

These together with Remark 3.1 give, for any $b \in \mathbb{R}$,

$$R_t^{b, \eta_n} = \max_{1 \leq k \leq N_t^{\eta_n}} (b - X_{T(k)})^+ \geq (b - X_{\underline{\sigma}^n(G(t))})^+ \vee (b - X_{\bar{\sigma}^n(G(t))})^+ \xrightarrow{n \uparrow \infty} (b - \underline{X}_t)^+ = R_t^{b, \infty}.$$

For the case $G(t) = 0$ (i.e. $\underline{X}_t = X_0$), by slightly modifying the arguments,

$$R_t^{b, \eta_n} \geq (b - X_{\bar{\sigma}^n(0)})^+ \xrightarrow{n \uparrow \infty} (b - X_0)^+ = R_t^{b, \infty}.$$

If $G(t) = t$, we have $X_{\bar{\sigma}^n(t)} \xrightarrow{n \uparrow \infty} X_{t-}$ and hence $R_t^{b, \eta_n} \xrightarrow{n \uparrow \infty} (b - \underline{X}_{t-})^+$, which differs from $R_t^{b, \infty}$ only when X jumps downward at t .

By these and because $n \mapsto R_t^{b, \eta_n}$ is increasing, we have $R_t^{b, \eta_n} \nearrow R_t^{b, \infty}$ and consequently $U_t^{b, \eta_n} \nearrow U_t^{b, \infty}$ as $n \rightarrow \infty$ for a.e. $t > 0$ (more specifically all $t > 0$ except t at which X jumps downward) for all $b \in \mathbb{R}$.

By this, together with the fact that f'_+ is non-decreasing, we have

$$f'_-(U_t^{b, \infty}) \leq \lim_{n \rightarrow \infty} f'_+(U_t^{b, \eta_n}) \leq f'_+(U_t^{b, \infty}) \quad \text{for a.e. } t > 0,$$

and hence, by the monotone convergence theorem, $n \mapsto \rho_{\eta_n}(b)$ is non-decreasing and $\rho_\infty(b-) \leq \lim_{n \rightarrow \infty} \rho_{\eta_n}(b) \leq \rho_\infty(b)$ for all $b \in \mathbb{R}$.

This shows that $b^\dagger := \lim_{n \rightarrow \infty} b_{\eta_n}^*$ exists and $b^\dagger \geq b_\infty^*$. The monotonicity also suggests that $\rho_{\eta_n}(b^\dagger) \leq -C$ uniformly in n and hence $\rho_\infty(b^\dagger) \leq -C$. If $b^\dagger > b_\infty^*$, then we must have $\rho_\infty((b_\infty^* + b^\dagger)/2) \leq -C$. However, as shown in [25, Lemma 5], $\rho_\infty(b) > -C$ for $b > b_\infty^*$ for the case when X is not the negative of a subordinator. For the case when it is the negative of a subordinator, we have $\rho_\infty(b) = f'_+(b)/q$, which is strictly increasing at $b = b_\infty^*$ by assumption and hence the contradiction can be derived similarly. Hence we must have $b^\dagger = b_\infty^*$, as desired.

(ii) Fix $N \in \mathbb{N}$ and $x \in \mathbb{N}$. Because, for $n \geq N$, $b_\infty^* \leq b_{\eta_n}^* \leq b_{\eta_N}^*$ and hence $U_t^{b_\infty^*, \eta_N} \leq U_t^{b_{\eta_n}^*, \eta_n} \leq U_t^{b_{\eta_N}^*, \eta_N}$ and by the convexity of f ,

$$\begin{aligned} & \mathbb{E}_x \left[\int_0^\infty e^{-qt} \sup_{n \geq N} |f(U_t^{b_{\eta_n}^*, \eta_n})| dt \right] \\ & \leq \mathbb{E}_x \left[\int_0^\infty e^{-qt} (|f(U_t^{b_\infty^*, \eta_N})| + |f(U_t^{b_{\eta_N}^*, \eta_N})| + c) dt \right] \\ & < \infty, \end{aligned}$$

where c is a constant value defined in (A.2). On the other hand,

$$|U_t^{b_{\eta_n}^*, \eta_n} - U_t^{b_\infty^*, \infty}| \leq |U_t^{b_{\eta_n}^*, \eta_n} - U_t^{b_\infty^*, \eta_n}| + |U_t^{b_\infty^*, \eta_n} - U_t^{b_\infty^*, \infty}| \xrightarrow{n \uparrow \infty} 0$$

by Remark 3.1 and (i) for a.e. $t > 0$. Hence dominated convergence gives the pointwise convergence of

$$\mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^{b_{\eta_n}^*, \eta_n}) dt \right]$$

to

$$\mathbb{E}_x \left[\int_0^\infty e^{-qt} f(U_t^{b_\infty^*, \infty}) dt \right] \quad \text{for all } x \in \mathbb{R}.$$

On the other hand, by integration by parts,

$$\begin{aligned} & \left| \mathbb{E}_x \left[\int_0^\infty e^{-qt} dR_t^{b_{\eta_n}^*, \eta_n} \right] - \mathbb{E}_x \left[\int_{(0, \infty)} e^{-qt} dR_t^{b_\infty^*, \infty} \right] \right| \\ & = q \left| \mathbb{E}_x \left[\int_0^\infty e^{-qt} R_t^{b_{\eta_n}^*, \eta_n} dt \right] - \mathbb{E}_x \left[\int_0^\infty e^{-qt} R_t^{b_\infty^*, \infty} dt \right] \right| \\ & \leq q \mathbb{E}_x \left[\int_0^\infty e^{-qt} |R_t^{b_{\eta_n}^*, \eta_n} - R_t^{b_\infty^*, \infty}| dt \right]. \end{aligned}$$

Here,

$$\mathbb{E}_x \left[\int_0^\infty e^{-qt} \sup_{n \geq N} |R_t^{b_{\eta_n}^*, \eta_n} - R_t^{b_\infty^*, \infty}| dt \right] \leq 2 \mathbb{E}_x \left[\int_0^\infty e^{-qt} R_t^{b_{\eta_N}^*, \eta_N} dt \right] < \infty$$

thanks to

$$R_t^{b_{\eta_n}^*, \eta_n} \vee R_t^{b_\infty^*, \infty} \leq R_t^{b_{\eta_N}^*, \eta_N} \quad \text{for all } t > 0.$$

Thus, by the dominated convergence theorem together with

$$|R_t^{b_{\eta_n}^*, \eta_n} - R_t^{b_\infty^*, \infty}| = |U_t^{b_{\eta_n}^*, \eta_n} - U_t^{b_\infty^*, \infty}| \xrightarrow{n \uparrow \infty} 0 \quad \text{for a.e. } t > 0,$$

we have the pointwise convergence of

$$\mathbb{E}_x \left[\int_0^\infty e^{-qt} dR_t^{b_{\eta_n}^*, \eta_n} \right] \text{ to } \mathbb{E}_x \left[\int_{[0, \infty)} e^{-qt} dR_t^{b_{\infty}^*, \infty} \right] \text{ for all } x \in \mathbb{R}.$$

Finally, because $n \mapsto v_{\eta_n}^*(x)$ is monotone and each value function is continuous in x , $\lim_{n \uparrow \infty} v_{\eta_n}^*(x) = v_{\infty}^*(x)$ holds uniformly in x on any compact set by Dini’s theorem. \square

6. Numerical results

In this section we confirm the obtained results through numerical experiments via Monte Carlo simulation (classical Euler scheme). In order to confirm that the results hold for a wide class of Lévy processes, we choose a Lévy process X of the form

$$X_t = X_0 - 0.1t + 0.2B_t + \sum_{n=1}^{N_t^+} Z_n^+ - \sum_{n=1}^{N_t^-} Z_n^-, \quad 0 \leq t < \infty,$$

where $\{B_t : t \geq 0\}$ is a standard Brownian motion and $\{N_t^+ : t \geq 0\}$ and $\{N_t^- : t \geq 0\}$ are Poisson processes with arrival rates 0.4 and 0.6, respectively. The upward and downward jumps $\{Z_n^+ : n \in \mathbb{N}\}$ and $\{Z_n^- : n \in \mathbb{N}\}$ are i.i.d. sequences of (folded) normal random variables with mean zero and variance 1 and Weibull random variables with shape parameter 2 and scale parameter 1, respectively. These processes are assumed to be mutually independent.

For the running cost function f , we consider the following three cases:

$$\begin{aligned} f_1(x) &:= x^2, & f_2(x) &:= x^3 1_{\{x \geq 0\}} + x^2 1_{\{x < 0\}}, \\ f_3(x) &:= [x^2 + e^{-(x-1)}] 1_{\{x \geq 1\}} + \frac{x^2 + 3}{2} 1_{\{x < 1\}}, \end{aligned} \tag{6.1}$$

for $x \in \mathbb{R}$, which are convex and continuously differentiable on \mathbb{R} . For other parameters, we set $q = 0.05$ and $C = 1$. For each realization, we truncate the time horizon to $T = 100$ and discretize $[0, T]$ using $N = 10\,000$ equally spaced points with distance $\Delta_t := T/N$. Unless stated otherwise, we use $\eta = 1$.

For the approximation of the expectation, we first obtain a set of $M := 5\,000$ sample paths of X started at zero, say

$$\hat{\mathbf{X}} := (\hat{X}^{(1)}, \dots, \hat{X}^{(M)}) \text{ with } \hat{X}^{(m)} = \{\hat{X}_{n\Delta_t}^{(m)} : 1 \leq n \leq N\} \text{ for } 1 \leq m \leq M.$$

Control opportunities

$$\hat{\mathbf{N}}^\eta := (\hat{N}^{\eta, (1)}, \dots, \hat{N}^{\eta, (M)}) \text{ with } \hat{N}^{\eta, (m)} = \{\hat{N}_{n\Delta_t}^{\eta, (m)} : 1 \leq n \leq N\} \text{ for } 1 \leq m \leq M$$

are sampled by generating

$$\hat{N}_{(n+1)\Delta_t}^{\eta, (m)} - \hat{N}_{n\Delta_t}^{\eta, (m)} = 1_{\{e < \Delta_t\}} \text{ with i.i.d. } e \sim \exp(\eta)$$

and their corresponding reflected paths (with barrier zero)

$$\hat{U}^{0, (m)} = \{\hat{U}_{n\Delta_t}^{0, (m)} : 1 \leq n \leq N\}$$

are then computed. These sample paths can be used commonly for the approximation of the expectation in $\rho(b)$ as in (4.1). In other words, we approximate it by

$$\hat{\rho}_M(b) := M^{-1} \sum_{m=1}^M \Delta_t \sum_{n=0}^N e^{-qn\Delta_t} f'(\hat{U}_{n\Delta_t}^{0,(m)} + b).$$

As shown in Section 3, $\rho(b)$ is monotone and hence b^* can be obtained by classical bisection. While $\hat{\rho}_M(b)$ for each b is an approximated value, because we are using the same sample paths (\hat{X}, \hat{N}^η) , the monotonicity of $b \rightarrow \hat{\rho}_M(b)$ is still preserved, causing no problem in using bisection methods. Figure 2 shows the plots of $\hat{\rho}_M(b)$ for cases i for $i = 1, 2, 3$. It can be confirmed that it is indeed monotonically increasing, and the root becomes b^* . Note that for the case $i = 1$, $\rho_M(b)$ becomes a straight line.

With the approximated optimal barrier b^* , we shall now confirm the optimality by comparing the expected total costs v_{b^*} with v_b under suboptimal choices of b . In order to compute these, we continue using the set of paths (\hat{X}, \hat{N}^η) . Figure 3 shows the results. It can be confirmed that the selection b^* indeed minimizes the total expected cost for all starting points.

Finally, we confirm the convergence as $\eta \rightarrow \infty$. In Figure 4, we plot the value function when $\eta = 2, 5, 10, 20, 50, 100, 200, 500, 1000$ together with the classical case whose reflected path with lower barrier b under \mathbb{P}_x is approximated by

$$(\hat{X}_{n\Delta_t}^{(m)} + x) + \max_{0 \leq l \leq n} (b - (\hat{X}_{l\Delta_t}^{(m)} + x))^+.$$

It is observed in all cases that the optimal barrier and the value function converge decreasingly to those of the classical case, confirming Theorem 5.1.

7. Concluding remarks

In this paper we have solved the stochastic control problem of minimizing the sum of running and controlling costs under the constraint that control opportunities are restricted to independent Poisson arrival times. For a general Lévy process model, we showed the optimality of a simple barrier strategy, with its barrier analytically provided as a root of the equality (1). Furthermore, we demonstrated that the optimal solutions in the Poissonian setting converge to those in the continuous-observation setting. These results potentially provide a new approach to the classical case, using techniques developed in this paper for Poisson observation models.

One important extension is to consider the case where we can control the process in both directions. This scenario has been studied in the continuous-observation case driven by spectrally negative Lévy processes, as discussed in [4], where it is shown that it is optimal to reflect the process at both upper and lower barriers.

Another natural extension is to consider the case with a fixed intervention cost. In this case, the optimal strategy is expected to be of the two-barrier type. More specifically, it is expected to be a variant of the (s,S) -policy (see e.g. [5], [7]), which moves the process to a certain point, say \bar{b} , whenever it is observed to be below a different point, say \underline{b} , at Poisson observation times. This is a reasonable conjecture based on Yamazaki [37], who showed the optimality of such a policy for the continuous-observation case under a spectrally one-sided Lévy model.




FIGURE 2. Plot of $\hat{\rho}_M(b)$ for case i under the cost function f_i as in (6.1), for (a) $i = 1$, (b) $i = 2$, (c) $i = 3$. The root (indicated by a star) becomes an approximation of the optimal barrier b^* .




FIGURE 3. Plot of the approximated value functions v_{b^*} (solid) along with v_b (dotted) for $b = b^* - 1$, $b^* - 0.5$, $b^* + 0.5$, $b^* + 1.0$, for case i for (a) $i = 1$, (b) $i = 2$, (c) $i = 3$. The points at the barriers are indicated by stars and circles for $b = b^*$ and $b \neq b^*$, respectively.




FIGURE 4. Plot of the approximated value functions v_{b^*} (dotted) for $\eta = 2, 5, 10, 20, 50, 100, 200, 500, 1000$ along with that in the classical case (solid) for case i for (a) $i = 1$, (b) $i = 2$, (c) $i = 3$. The points at the barriers are indicated by stars.

Appendix A. Proofs

A.1. The proof of Lemma 3.1

We fix $b, x \in \mathbb{R}$. Let $U^{b,\infty}$ and $R^{b,\infty}$ be those defined in Section 5 for the controlled and control processes in the classical setting under the barrier strategy with barrier b . First we have a bound

$$X_t \leq U_t^b \leq U_t^{b,\infty}, \quad 0 \leq R_t^b \leq R_t^{b,\infty}. \tag{A.1}$$

By the convexity of f , we have $|f(U_t^b)| \leq |f(U_t^{b,\infty})| + |f(X_t)| + c$, where

$$c = \begin{cases} |\inf_{y \in \mathbb{R}} f(y)|, & \text{if it exists,} \\ 0, & \text{otherwise.} \end{cases} \tag{A.2}$$

By Remark 2.3 and [25, (A.3)] under Assumptions 2.1 and 2.2, we obtain (i). By (A.1) and since [25, Lemma 3] holds under Assumptions 2.1 and 2.2, we obtain (ii).

We have

$$|v_b(x)| \leq \mathbb{E}_x \left[\int_0^\infty e^{-qt} (|f(X_t)| + |f(U_t^{b,\infty})| + c) dt \right] + |C| \mathbb{E}_x \left[\int_{(0,\infty)} e^{-qt} dR_t^{b,\infty} \right],$$

which is of polynomial growth by Remark 2.3 and the proof of [25, Lemma 3], showing (iii). □

A.2. Proof of Lemma 3.2

Note that $T_b = T(k)$ for some $k \in \mathbb{N}$, almost surely on $\{T_b < \infty\}$. By the monotone convergence theorem and the strong Markov property, we have

$$\begin{aligned} & \lim_{\varepsilon \downarrow 0} \mathbb{P}_x(T_b = T_{b+\varepsilon}, T_b < \infty) \\ &= \lim_{\varepsilon \downarrow 0} \sum_{k \in \mathbb{N}} \mathbb{P}_x(T_b = T_{b+\varepsilon} = T(k)) \\ &= \sum_{k \in \mathbb{N}} \lim_{\varepsilon \downarrow 0} \mathbb{P}_x(X_{T(1)} \geq b + \varepsilon, X_{T(2)} \geq b + \varepsilon, \dots, X_{T(k-1)} \geq b + \varepsilon, X_{T(k)} < b), \\ &= \sum_{k \in \mathbb{N}} \mathbb{P}_x(X_{T(1)} > b, X_{T(2)} > b, \dots, X_{T(k-1)} > b, X_{T(k)} < b) \\ &= \sum_{k \in \mathbb{N}} E^{(k)}(x), \end{aligned} \tag{A.3}$$

where

$$E^{(1)}(x) = \mathbb{P}_x(X_{T(1)} < b), \quad E^{(l+1)}(x) = \mathbb{E}_x[1_{\{X_{T(1)} > b\}} E^{(l)}(X_{T(1)})], \quad l \in \mathbb{N}.$$

By Remark 2.2, with $\{X_{T(1)} > b\}$ replaced by $\{X_{T(1)} \geq b\}$ in the definition of $E^{(k)}$ and going backwards from (A.3), we have

$$\lim_{\varepsilon \downarrow 0} \mathbb{P}_x(T_b = T_{b+\varepsilon}, T_b < \infty) = \sum_{k \in \mathbb{N}} \mathbb{P}_x(T_b = T(k)) = \mathbb{P}_x(T_b < \infty).$$

On the other hand, by the monotone convergence theorem,

$$\begin{aligned} \lim_{\varepsilon \downarrow 0} \mathbb{P}_x(T_b = T_{b-\varepsilon}, T_b < \infty) &= \sum_{k \in \mathbb{N}} \lim_{\varepsilon \downarrow 0} \mathbb{P}_x(X_{T(1)} \geq b, X_{T(2)} \geq b, \dots, X_{T(k-1)} \geq b, X_{T(k)} < b - \varepsilon) \\ &= \sum_{k \in \mathbb{N}} \mathbb{P}_x(X_{T(1)} \geq b, X_{T(2)} \geq b, \dots, X_{T(k-1)} \geq b, X_{T(k)} < b) = \mathbb{P}_x(T_b < \infty). \end{aligned}$$

Finally, because the map $b \mapsto T_b$ is non-increasing, the proof is complete by monotone convergence.

A.3. Proof of Lemma 4.1

Since f'_+ is non-decreasing, the function

$$\rho(b) = \mathbb{E}_0 \left[\int_0^\infty e^{-qt} f'_+(U_t^0 + b) dt \right]$$

is non-decreasing. By Corollary 3.1, monotone convergence, and Assumption 2.1(3), we have $\lim_{b \uparrow \infty} \rho(b) = f'_+(\infty)/q > -C$ and $\lim_{b \downarrow -\infty} \rho(b) = f'_+(-\infty)/q < -C$. In what follows, we show the continuity of ρ .

(i) We first prove that the potential of the process U^b does not have mass. Recall the first control time $T_b = T_b^{(1)}$. For $x, y, b \in \mathbb{R}$, we have

$$\begin{aligned} \mathbb{E}_x \left[\int_0^\infty e^{-qt} 1_{\{y\}}(U_t^b) dt \right] &= \sum_{k \in \mathbb{N}} \mathbb{E}_x \left[\int_{T_b^{(k-1)}}^{T_b^{(k)}} e^{-qt} 1_{\{y\}}(U_t^b) dt \right] \\ &= \mathbb{E}_x \left[\int_0^{T_b} e^{-qt} 1_{\{y\}}(X_t) dt \right] \\ &\quad + \sum_{k \in \mathbb{N}} \mathbb{E}_x [e^{-qT_b}] (\mathbb{E}_b [e^{-qT_b}])^{k-1} \mathbb{E}_b \left[\int_0^{T_b} e^{-qt} 1_{\{y\}}(X_t) dt \right], \end{aligned}$$

which is equal to 0 by Remark 2.2.

(ii) Since f'_+ is right-continuous and by the dominated convergence theorem with Corollary 3.1, we have

$$\rho(b + \varepsilon) - \rho(b) = \mathbb{E}_0 \left[\int_0^\infty e^{-qt} (f'_+(U_t^0 + b + \varepsilon) - f'_+(U_t^0 + b)) dt \right] \xrightarrow{\varepsilon \downarrow 0} 0.$$

Let D be the set of discontinuous point of f'_+ on \mathbb{R} , which is at most countable set since f'_+ is non-decreasing. By Corollary 3.1 and the dominated convergence theorem, we have

$$\begin{aligned} \rho(b) - \rho(b - \varepsilon) &= \mathbb{E}_0 \left[\int_0^\infty e^{-qt} (f'_+(U_t^0 + b) - f'_+(U_t^0 + b - \varepsilon)) dt \right] \\ &\xrightarrow{\varepsilon \downarrow 0} \mathbb{E}_0 \left[\int_0^\infty e^{-qt} \sum_{y \in D} (f'_+(y) - f'_-(y)) 1_{\{y\}}(U_t^0 + b) dt \right] \\ &= \mathbb{E}_b \left[\int_0^\infty e^{-qt} \sum_{y \in D} (f'_+(y) - f'_-(y)) 1_{\{y\}}(U_t^b) dt \right], \end{aligned}$$

which is equal to 0 since the potential of U^b does not have mass as in (i). □

A.4. Proof of Lemma 4.4

The proof is essentially the same as that of [25, Lemma 9] by simply replacing the classical reflected process $U^{b^*,\infty}$ with the Poissonian version U^{b^*} . Following the same arguments, we obtain

$$v''_{b^*}(x) = \mathbb{E}_x \left[\int_0^{T_{b^*}} e^{-qt} f''(U_t^{b^*}) dt \right],$$

which can be shown to be continuous by the dominated convergence theorem using the assumption that f'' is of polynomial growth, (3.5), and Lemma 3.2. For more details, see [25, Section A.6]. \square

Acknowledgements

K. Noba stayed at Centro de Investigación en Matemáticas in Mexico as a JSPS Overseas Research Fellow and received support regarding the research environment there. K. Noba is grateful for their support during his visit.

Funding information

K. Noba was supported by JSPS KAKENHI grant no. JP21K13807 and JPJSBP120249936. K. Yamazaki was supported by JSPS KAKENHI grant no. JP20K03758, JP24K06844, and JP24H00328, and the start-up grant by the School of Mathematics and Physics of the University of Queensland. Both authors were supported by JSPS Open Partnership Joint Research Projects grant no. JPJSBP120209921.

Competing interests

There were no competing interests to declare which arose during the preparation or publication process of this article.

References

- [1] ALBRECHER, H., BÄUERLE, N. AND THONHAUSER, S. (2011). Optimal dividend-payout in random discrete time. *Statist. Risk Model.* **28**, 251–276.
- [2] ALBRECHER, H., IVANOV, J. AND ZHOU, X. (2016). Exit identities for Lévy processes observed at Poisson arrival times. *Bernoulli* **22**, 1364–1382.
- [3] AVANZI, B., CHEUNG, E. C., WONG, B. AND WOO, J. K. (2013). On a periodic dividend barrier strategy in the dual model with continuous monitoring of solvency. *Insurance Math. Econom.* **52**, 98–113.
- [4] BAURDOUX, E. J. AND YAMAZAKI, K. (2015). Optimality of doubly reflected Lévy processes in singular control. *Stoch. Process. Appl.* **125**, 2727–2751.
- [5] BENKHEROUF, L. AND BENSOUSSAN, A. (2009). Optimality of an (s, S) policy with compound Poisson and diffusion demands: a quasi-variational inequalities approach. *SIAM J. Control Optimization* **48**, 756–762.
- [6] BENSOUSSAN, A. AND TAPIERO, C. S. (1982). Impulsive control in management: prospects and applications. *J. Optimization Theory Appl.* **37**, 419–442.
- [7] BENSOUSSAN, A., LIU, R. H. AND SETHI, S. P. (2005). Optimality of an (s, S) policy with compound Poisson and diffusion demands: a quasi-variational inequalities approach. *SIAM J. Control Optimization* **44**, 1650–1676.
- [8] BERTOIN, J. (1996). *Lévy Processes*. Cambridge University Press.
- [9] BIFFIS, E. AND KYPRIANOU, A. E. (2010). A note on scale functions and the time value of ruin for Lévy insurance risk processes. *Insurance Math. Econom.* **46**, 85–91.
- [10] CAI, N. AND YANG, X. (2018). International reserve management: a drift-switching reflected jump-diffusion model. *Math. Finance* **28**, 409–446.
- [11] CARR, P. (1998). Randomization and the American put. *Rev. Financial Studies* **11**, 597–626.
- [12] CARR, P., GEMAN, H., MADAN, D. AND YOR, M. (2002). The fine structure of asset returns: an empirical investigation. *J. Business* **75**, 305–333.

- [13] CHEUNG, E. C., LIU, H. AND WILLMOT, G. E. (2018). Joint moments of the total discounted gains and losses in the renewal risk model with two-sided jumps. *Appl. Math. Comput.* **331**, 358–377.
- [14] DONG, H., YIN, C. AND DAI, H. (2019). Spectrally negative Lévy risk model under Erlangized barrier strategy. *J. Comput. Appl. Math.* **351**, 101–116.
- [15] DUPUIS, P. AND WANG, H. (2002). Optimal stopping with random intervention times. *Adv. Appl. Prob.* **34**, 141–157.
- [16] EGAMI, M. AND YAMAZAKI, K. (2014). Phase-type fitting of scale functions for spectrally negative Lévy processes. *J. Comput. Appl. Math.* **264**, 1–22.
- [17] HERNÁNDEZ-HERNÁNDEZ, D., PÉREZ, J. L. AND YAMAZAKI, K. (2016). Optimality of refraction strategies for spectrally negative Lévy processes. *SIAM J. Control Optimization* **54**, 1126–1156.
- [18] JEANBLANC-PICQUÉ, M. (1993). Impulse control method and exchange rate. *Math. Finance* **3**, 161–177.
- [19] KYPRIANOU, A. E. (2014). *Fluctuations of Lévy Processes with Applications*, second edition. Springer.
- [20] KYPRIANOU, A. E. AND LOEFFEN, R. L. (2010). Refracted Lévy processes. *Ann. Inst. H. Poincaré Prob. Statist.* **46**, 24–44.
- [21] KYPRIANOU, A. E. AND PISTORIUS, M. R. (2003). Perpetual options and Canadization through fluctuation theory. *Ann. Appl. Prob.* **13**, 1077–1098.
- [22] LEUNG, T., YAMAZAKI, K. AND ZHANG, H. (2015). An analytic recursive method for optimal multiple stopping: Canadization and phase-type fitting. *Int. J. Theor. Appl. Finance* **18**, 1550032.
- [23] LKABOUS, M. A. (2021). Poissonian occupation times of spectrally negative Lévy processes with applications. *Scand. Actuarial J.* **2021**, 916–935.
- [24] MUNDACA, G. AND ØKSENDAL, B. (1998). Optimal stochastic intervention control with application to the exchange rate. *J. Math. Econ.* **2**, 225–243.
- [25] NOBA, K. AND YAMAZAKI, K. (2023). On singular control for Lévy processes. *Math. Operat. Res.* **48**, 1213–1234.
- [26] NOBA, K., PÉREZ, J. L. AND YAMAZAKI, K. (2023). Refraction strategies in stochastic control: optimality for a general Lévy process model. To appear in *SIAM J. Control Optimization*. Available at arXiv:2308.08183.
- [27] PALMOWSKI, Z., PÉREZ, J. L., SURYA, B. AND YAMAZAKI, K. (2020). The Leland–Toft optimal capital structure model under Poisson observations. *Finance Stoch.* **24**, 1035–1082.
- [28] PALMOWSKI, Z., PÉREZ, J. L. AND YAMAZAKI, K. (2021). Double continuation regions for American options under Poisson exercise opportunities. *Math. Finance* **31**, 722–771.
- [29] PÉREZ, J. L. AND YAMAZAKI, K. (2020). Optimality of hybrid continuous and periodic barrier strategies in the dual model. *Appl. Math. Optimization* **82**, 105–133.
- [30] PÉREZ, J. L., YAMAZAKI, K. AND BENSOUSSAN, A. (2020). Optimal periodic replenishment policies for spectrally positive Lévy demand processes. *SIAM J. Control Optimization* **58**, 3428–3456.
- [31] PÉREZ, J. L., YAMAZAKI, K. AND YU, X. (2018). On the bail-out optimal dividend problem. *J. Optimization Theory Appl.* **179**, 553–568.
- [32] SAARINEN, H. (2023). On Poisson constrained control of linear diffusions. Doctoral Thesis, University of Turku, Finland.
- [33] WANG, H. (2001). Some control problems with random intervention times. *Adv. Appl. Prob.* **33**, 404–422.
- [34] WANG, W., WANG, N. AND CHEN, M. (2023). On a doubly reflected risk process with running maximum dependent reflecting barriers. *J. Comput. Appl. Math.* **422**, 114880.
- [35] WANG, Z., LKABOUS, M. A. AND LANDRIault, D. (2023). A refracted Lévy process with delayed dividend pullbacks. *Scand. Actuarial J.* **2023**, 885–906.
- [36] YAMAZAKI, K. (2015). Contraction options and optimal multiple-stopping in spectrally negative Lévy models. *Appl. Math. Optimization* **72**, 147–185.
- [37] YAMAZAKI, K. (2017). Inventory control for spectrally positive Lévy demand processes. *Math. Operat. Res.* **42**, 212–237.
- [38] YANG, C., SENDOVA, K. P. AND LI, Z. (2020). Parisian ruin with a threshold dividend strategy under the dual Lévy risk model. *Insurance Math. Econom.* **90**, 135–150.
- [39] ZHANG, Z. AND HAN, X. (2017). The compound Poisson risk model under a mixed dividend strategy. *Appl. Math. Comput.* **315**, 1–12.
- [40] ZHAO, Y., CHEN, P. AND YANG, H. (2017). Optimal periodic dividend and capital injection problem for spectrally positive Lévy processes. *Insurance Math. Econom.* **74**, 135–146.
- [41] ZHAO, Y., WANG, R., YAO, D., CHEN, P. (2015). Optimal dividends and capital injections in the dual model with a random time horizon. *J. Optimization Theory Appl.* **167**, 272–295.