

Social Media Review

SGEM Hot Off the Press: Computer provider order entry (CPOE) and emergency department flow

Katie Lin, MD*¹; Kenneth Chan, MD*²; Rohit Mohindra, MD[†]; Ken Milne, MD, MSc[‡]; Brent Thoma, MD, MA, MSc[§]; Chris Bond, MD*

As part of the Canadian Journal of Emergency Medicine (CJEM) developing social media strategy,¹ we are collaborating with the Skeptics' Guide to Emergency Medicine (SGEM) to summarize and critically appraise the current emergency medicine literature using evidence-based medicine principles. In the Hot Off the Press (HOP) series, we select original research manuscripts published in CJEM to be summarized and critically appraised on the SGEM website/podcast² and discussed by the study authors and the online EM community. A similar collaboration is underway between the SGEM and Academic Emergency Medicine. What follows is a summary of the selected article and the immediate post-publication critical appraisal from the SGEM podcast, as well as an overview of the subsequent discussion from the SGEM blog and other social media. Through this series, we hope to enhance the value, accessibility, and application of important, clinically relevant EM research. In this, the fourth SGEM HOP hosted collaboratively with CJEM, we discuss Gray et al.'s paper³ evaluating the impact of computerized provider order entry on patient flow through a quaternary emergency department in London, Ontario.

BACKGROUND

Computerized provider order entry (CPOE) has emerged in North America as a means to standardize and improve the delivery of health care services to patients.⁴ CPOE involves the utilization of electronic systems to track health care provider orders but can also include more advanced functionality such as clinical decision support.⁵

The debate surrounding CPOE has focused on the clinical benefits versus costs of implementation. Studied clinical benefits of CPOE include reduced prescribing errors and adverse medication interactions,⁶ improved

adherence to evidence-based protocols for specific presentations, such as renal colic and acute ischemic stroke,⁷⁻⁹ increased legibility and accessibility of documentation on record,¹⁰ and potential secondary uses of data by health care organizations for outcome tracking and quality assessment.¹⁰ CPOE's benefits, however, may come at the cost of decreased patient and physician satisfaction,^{11,12} impaired emergency physician productivity,¹³ and increased length of stay (LOS) in the emergency department (ED) for admitted patients.¹⁴ There is no strong evidence to suggest that CPOE improves patient mortality,¹⁵ and the literature has not yet provided consensus evidence in favour of CPOE when weighing its benefits and drawbacks.¹⁶

ARTICLE SUMMARY

Gray and colleagues conducted a retrospective cohort study to evaluate the impact of CPOE implementation on ED workflow at a quaternary care centre in London, Ontario, comprising two separate ED campuses. They adopted a pre-post implementation design to evaluate the effect of CPOE on three primary ED flow metrics: LOS, wait time (WT), and the proportion of patients who left without being seen (LWBS).

The study included all ED patients 18 years and older who were triaged during July and August 2013 (pre-implementation of CPOE) and July and August 2014 (post-implementation of CPOE). Data were extracted from the London Health Sciences Centre (LHSC) electronic database and health records. Any patients with incomplete or incorrect ED charts, negative WTs or LOS, extreme outliers with WT > 24 hours,

From the *Department of Emergency Medicine, University of Calgary, Calgary, AB; †Department of Emergency Medicine, McGill University, Montreal, QC; ‡Division of Emergency Medicine, University of Western Ontario, London, ON; and §Emergency Medicine, University of Saskatchewan, Saskatoon, SK.

Correspondence to: Drs. Katie Lin and Kenneth Chan, Department of Emergency Medicine, Room C231, Foothills Medical Centre, 1403 29 St. NW, Calgary, AB T2N 2T9; Email: kyllin@ucalgary.ca and kchan.med@gmail.com

or those missing vital statistics were excluded. The investigators also completed a subgroup analysis looking at variability by the Canadian Triage and Acuity Scale (CTAS) stratification and admitted patients to evaluate whether CPOE had differential effects based on acuity of illness or disposition status.

The authors analysed a combined data set of 36,758 ED visits: 18,872 visits in 2013 and 17,886 visits in 2014. Median age, gender distribution, CTAS stratification, and rate of admission were similar between the two groups at baseline. Statistical analyses were conducted to determine significant changes in WT, LOS, and LWBS between pre- and post-implementation of CPOE.

KEY RESULTS

The authors concluded that CPOE implementation detrimentally impacted patient flow in the two EDs that they studied. They found statistically significant increases in median WT (increased from 78 to 83 minutes), median LOS (increased from 254 to 264 minutes), proportion of patients who LWBS (increased from 7.2% to 8.1%), and the LOS for admitted patients (increased from 713 to 776 minutes) (Table 1).

QUALITY ASSESSMENT

The study by Gray et al.³ was a retrospective cohort design comparing ED patient flow pre- and post-implementation of CPOE using administrative data. The authors had clearly defined outcomes and collected data with objective

variables. The inclusion and exclusion criteria were appropriate and acceptable for cohort recruitment. Nevertheless, this study had several limitations.

The CPOE in this study was introduced to the two EDs in April 2014, only 3 months prior to the data sampling set for post-implementation. The results seen in this study could potentially be confounded by inexperience and lack of adequate training or uptake during the early stages of implementation. It would be useful to collect data further from implementation to determine whether user familiarity with CPOE reduces its impact on workflow over time given previously described learning curves with CPOE implementation.¹⁷

The retrospective cohort design encompasses potential sampling errors that limit the generalizability of the authors' findings.¹⁸ Gray et al.³ drew data from only July and August during the pre- and post-implementation phases of CPOE in one Canadian city. It is difficult to state whether this would be a representative sampling for a number of reasons.

First, the authors did not collect or report on potential confounding characteristics between the pre- and post-implementation time periods such as mean boarding times, hospital over-capacity statistics, population and ED utilization rates, and bed allocation resources.

Second, the choice of July and August for data extraction may have impacted outcomes, given that July 1 marks the beginning of residency for new trainees in Canada, and junior residents in the ED environment would be adjusting to CPOE in addition to new clinical responsibilities. The existence of the "July effect" (whereby medical trainees beginning new clinical roles

Table 1. Key results from Gray et al.'s CPOE study³

	Pre-CPOE (2013)	Post-CPOE (2014)	<i>p</i> -value
Median wait time (minutes [IQR])	78 (33-165)	83 (33-166)	0.036
Median length of stay (minutes [IQR])	254 (147-417)	264 (153-442)	0.001
Left without being seen (all patients)	1364 (7.2%)	1448 (8.1%)	0.002
Left without being seen (CTAS 3 patients)	706 (7.7%)	776 (8.6%)	0.025
Left without being seen (CTAS 4 patients)	575 (13.1%)	536 (15.4%)	0.004
Left without being seen (CTAS 5 patients)	42 (24.3%)	81 (42.0%)	<0.001
Admitted length of stay (minutes [IQR])	713 (443.5-1204.5)	776 (486-1260)	<0.001

affect medical error rates, patient outcomes, and departmental productivity) remains uncertain in the literature.¹⁹

Third, in the post-implementation group, a substantial proportion of patient data was excluded, mostly due to clerical and administrative data error. It is impossible to determine what impact this may have had on the results.

Finally, CPOE software programs can be highly variable in their design, implementation strategy, and user interfaces. Some CPOE programs may include components of clinical decision support and order set bundles to prompt evidence-based reminders during clinical encounters, whereas others may simply be a collection of individual medications and orders requiring manual entry. Without further detail on the specifics of the CPOE design and implementation in the authors' study, it is difficult to draw parallels between this study's results and CPOE use in other centres.

TAKE-TO-WORK POINTS

CPOE implementation may statistically impair ED flow metrics, although this is of uncertain clinical significance given the slight differences in flow metric results (5 and 10 minutes' difference in median WT and LOS, respectively) and the large interquartile ranges measured. The initial stages of implementing CPOE may lead to increased ED WTs, LOS, and more patients leaving without being seen by an emergency physician. The generalizability of Gray et al.'s study³ is uncertain because the impact of CPOE on ED work flow may be highly dependent on both software characteristics and implementation strategy. Further studies looking at the long-term outcomes of CPOE implementation are still needed.

METHODOLOGY OF THE SOCIAL MEDIA RESPONSE ANALYSIS

The social media discussion started with the launch of the blog post and podcast on July 5, 2016, and continued for 2 weeks until July 19, 2016. An invitation to comment on the article was included in the audio of the podcast, the text of the blog post, and on social media communications (Twitter and Facebook). Social media responses written in the SGEM blog's comment section, the SGEM Facebook page, and on Twitter (directed at @thesgem, @socmobem, or using the #SGEMHOP hashtag) between July 5 and July 19,

2016, were reviewed by the authorship team. KL compiled and reviewed all aforementioned social media commentary to identify tweets and posts related to the CPOE SGEM podcast and blog post. A thematic analysis was conducted using a qualitative framework approach as outlined below.

Framework approach for thematic analysis:

- Provisional classification: content from each of the various analysed social media platforms was classified as either promotional (i.e., containing only a link to the blog post with no further content) or commentary-based.
- Thematic framework development: each commentary-type item was evaluated individually to identify key issues, concepts, and themes raised.
- Indexing: commonly identified themes across all of the commentary-type items was compiled and coded with short phrases for ease of comparison and tracking.
- Charting: the thematic framework was organized into a comparison chart presented in Table 2.
- Mapping and interpretation: as soon as common thematic groupings were identified and a comparison chart was created, all authors participated in a consensus-based analysis to determine which comments were most representative of the general themes of the discussion.

Multiple metrics of dissemination were further tracked by the SGEM HOP team for analysis:

- Blog post page views were monitored using the Jet Pack plugin by Wordpress.com (available from <https://wordpress.org/plugins/jetpack/>).
- Facebook "reach" analytics were provided by Facebook and represented the number of users who saw the original SGEM Facebook post on their own newsfeeds.²⁰
- Twitter impressions (the number of users whose newsfeeds contained a tweet featuring the #SGEMHOP hashtag) were tracked using Symplur, a software program that monitors health care related twitter conversations.²¹ Tweets not containing the hashtag were not tracked by Symplur. The number of impressions was calculated by taking the number of tweets per twitter user using the #SGEMHOP hashtag and multiplying it by the number of followers that each participant had.

Table 2. Framework approach for thematic analysis – comparison chart

Initial themes	Initial categories	Refined categories	Final themes
Flawed external validity	Need to adjust for boarding and emergency inpatient changes to determine causality of relationship between CPOE and increased ED WT (EL). Learning curve during implementation extends beyond the first few months, so authors did not account for effects of familiarity with CPOE over long term (SD).	Lack of reporting or adjusting for ED overcrowding metrics. Lack of long-term follow-up to account for post-implementation flow.	Confounding variables unaccounted for.
Utility of order sets	Common order sets can be useful, but complex patients requiring manual entry slow things down (AG). Variable components of CPOE in different sites (Nanaimo v. Calgary v. Saskatchewan) (CW, NL, EL, SD). Order sets can be consistently designed across multiple sites in one city to improve standardization (NL).	Order set limitations and patient factors. Site-congruent order sets improve consistency.	Variability of CPOE software design.
CPOE implementation strategy	When used as part of a knowledge translation strategy, CPOE provides standardization, improved safety outcomes, and better communication among the health care team (EL). CPOE components can reduce practice variation and cut down costs overall if used effectively (KK). Implementation without ED physician input led to clinical inefficiency and impaired utility (CW).	Importance of implementation strategy. Integrated clinical decision support. ED physician input on implementation is key.	Collaborative implementation.

- The altmetric score is a proprietary, standardized tool that tracks the disseminative impact of research articles in social media forums (e.g., Facebook, Twitter) and on blogs, podcasts, and news outlets.²² The altmetrics of the featured article by Gray et al.³ (Figure 1) were compared to all other articles published in *CJEM*, all published research analysed by altmetrics, and the articles covered in the first three *CJEM-SGEM HOPs*.²³⁻²⁵

RESULTS OF THE SOCIAL MEDIA RESPONSE

Table 3 provides details on the social media reach of Gray et al.’s article³ during the *SGEM HOP* campaign. During the 2-week period following the podcast release, #SGEMHOP was used in 84 tweets by 43 individual users, representing 216,926 Twitter impressions. Thirty-four of these tweets were from the study’s authors or *CJEM* personnel prompting social media responses. Online conversation through Twitter and the *SGEM* blog remained active for 14 days following podcast release. The altmetrics score for Gray et al.’s article³ was 41, placing it 10th highest amongst all previous *CJEM* publications and within the top 5% of all published research. Prior articles featured in the *SGEM HOPs* series scored in the 46–71 range.²³⁻²⁵ The mean altmetrics score for *CJEM* publications is 5.9.²⁶

ONLINE DISCUSSION SUMMARY

The majority of the discourse was held directly on the *SGEM* blog with dissemination of the URL link through Twitter, Facebook, and Google +. The *SGEM* blog hosted a total of 17 comments from 9 discrete users, including direct feedback from the study’s primary author, Dr. Andrew Gray.² Clinicians across Canada engaged in conversation to share their experiences with CPOE implementation at their institutions. Consistently identified themes included the limited generalizability of the study’s results due to confounding factors, variability of CPOE software design across centres, and the importance of a collaborative implementation strategy when introducing CPOE to a department.

There were multiple confounding variables that could have influenced the study’s results. Dr. Eddy Lang, Academic Department Head for Emergency Medicine at the University of Calgary, cautioned that




Figure 1. Screen capture of the altmetrics data retrieved August 3, 2016, from <https://cambridgejournals.altmetric.com/details/9299925>.

Table 3. Aggregate analytic data of social media platform discussions following the SGEM blog posting

Social media analytic	Metric	Metric definition	Count
SGEM blog statistics	Number of page views	Number of times the Web page containing the post was viewed	338
	Number of blog comments	Comments made directly on the website in the blog comments section	17
	Average word count of comments	Mean number of words per comment on the blog	288
Symplur analytics	Number of tweets	Number of tweets containing the hashtag #SGEMHOP	84
	Number of Twitter participants	Number of unique Twitter users who participated with tweeting during 2 weeks around the event	43
	Twitter Impressions	The number of potential views of a tweet or a tweet containing a specific hashtag in users' Twitter streams, as calculated by the number of tweets per participant and multiplying it with the number of followers that participant has	216,926
Facebook statistics	Reach	The number of users whose newsfeeds featured the SGEM	8,387
	Likes	The number of "likes" on the SGEM post	2

the study's results needed to be considered within a systemic context of ED overcrowding: "CPOE may have an association with increased [length of stay] for admitted patients but without adjustment for boarding and [emergency inpatients] there is no way to know if any relationship exists." This was consistent with the discussion held on the SGEM podcast, where the authors' lack of reporting for important potential

confounding factors was addressed. Variables, such as boarding time and proportion of emergency inpatients, were not tracked between the pre- and post-implementation time periods.

Physicians across the country shared their experiences with CPOE implementation and highlighted the variability of CPOE software across different centres. Dr. Wurster felt that implementation of CPOE at his

centre was “nothing short of a disaster. [...] Our training was not adequate and we were given little input in how we would like the computer system to work for us. No one examined how using CPOE would affect our daily processes in the emergency department.” This was contrasted with Dr. Lalani’s experiences with CPOE in Calgary, where physician input is a critical component of CPOE development and maintenance: “I really value my colleagues [...] who have worked so hard to generate easily workable order sets that include things like ‘Well’s Score’ for VTE. I like that if I type ‘ED meningitis’ I get all the orders for this presentation (including CT and all four LP tubes pre-labelled), and I like that if I go to any of the four hospitals [in Calgary] that I work at, it’s the exact same process.”

Finally, the importance of a collaborative implementation strategy was summarized by Dr. Shawn Dowling, an emergency clinician-researcher with a funded role for maintenance and optimization of CPOE order sets in Calgary: “[CPOE’s] success and failure is dependent on the components (order sets), its fit (usability of the software), the engine (physician buy in/engagement) and its mechanics (the team responsible for maintaining and optimizing the CPOE environment). CPOE is most powerful when it’s part of an integrated knowledge translation strategy.”

LIMITATIONS OF SOCIAL MEDIA ANALYSIS

There are a number of inherent limitations to any study evaluating social media engagement. The utilization of social media data involves sampling bias, and we recognize that the online discussion summarized in this paper likely represents the subset of emergency physicians who are heavily engaged in free open access medical education (FOAMed) activities. As such, the opinions of the wider audience may not have been captured through the social media platforms analysed.

Furthermore, commonly used social media analytic scores, such as the number of Twitter impressions or the altmetrics score, are useful for quantifying article dissemination, but do not necessarily provide information on the quality of the discourse generated.²⁷⁻²⁹ While the blog post hosted informative discussion around Gray et al.’s publication,³ virtually all of the tweets containing the #SGEMHOP hashtag and various Facebook posts simply directed readers to the blog itself and did not directly add to the discussion.

We have summarized the social media dialogue in this paper; however, it remains important for readers to independently evaluate the primary literature and continue to critically appraise the social media feedback.

CONCLUSION

Gray et al.’s paper³ suggested that CPOE implementation impaired ED flow to both a statistically significant and clinically variable degree. The *SGEM*’s blog post summarized the podcast discussion for readers, highlighted key limitations of generalizing the original study’s results to CPOE at other centres, and allowed physician leaders from across the country to share their experiences with the research topic. Online distribution of Gray et al.’s article³ resulted in disseminative impact scores reaching the top 5% of all published research. Collaborative knowledge translation and online engagement have the potential to increase awareness of primary literature, harness the perspectives of academic experts to enrich critical appraisal, and bridge the gap between literature findings and pragmatic applications for clinical practice.

Keywords: medical order entry systems, emergency department, efficiency

Competing interests: None declared.

REFERENCES

1. Thoma B, Mohindra R, Artz JD, et al. CJEM and the changing landscape of medical education and knowledge translation. *CJEM* 2015;17(2):184-7.
2. Milne K. SGEM #159: Computer games – computer provider order entry (CPOE); 2016. Available at: <http://thesgem.com/2016/07/sgem159-computer-games-computer-provider-order-entry-cpoe> (accessed 27 July 2016).
3. Gray A, Fernandes C, Van Aarsen K, et al. The impact of computerized provider order entry on emergency department flow. *CJEM* 2016;18(4):264-9.
4. Kuperman GJ, Gibson RF. Computer physician order entry: benefits, costs, and issues. *Ann Intern Med* 2003;139:31-9.
5. Miller R, Waitman L, Chen S, et al. The anatomy of decision support during inpatient care provider order entry (CPOE): empirical observations from a decade of CPOE experience at Vanderbilt. *J Biomed Info* 2005;38(6):469-85.
6. Nuckols TK, Smith-Spangler C, Morton SC, et al. The effectiveness of computerized order entry at reducing preventable adverse drug events and medication errors in hospital settings: a systematic review and meta-analysis. *Syst Rev* 2014;3:56.

7. Blunkerfield JF, Rogers L, White J, et al. Prospective evaluation of the treatment of pain in the ED using computerized physician order entry. *Am J Emerg Med* 2012; 30(8):1613-6.
8. Netherton SJ, Lonergan K, Wang D, et al. Computerized physician order entry and decision support improves ED analgesic ordering for renal colic. *Am J Emerg Med* 2014; 32(9):958-61.
9. Yang JM, Park YS, Chung SP, et al. Implementation of a clinical pathway based on a computerized physician order entry system for ischemic stroke attenuates off-hour and weekend effects in the ED. *Am J Emerg Med* 2014; 32(8):884-9.
10. Cresswell K, Bates D, Williams R, et al. Evaluation of medium-term consequences of implementing commercial computerized physician order entry and clinical decision support prescribing systems in two "early adopter" hospitals. *J Am Med Inform Assoc* 2014;21(e2):e194-202.
11. Bastani A, Walch R, Todd B, et al. Computerized prescriber order entry decreases patient satisfaction and emergency physician productivity. *Am J Emerg Med* 2011;29(2): 207-11.
12. Shanafelt TD, Dyrbye LN, Sinsky C, et al. Relationship between clerical burden and characteristics of the electronic environment with physician burnout and professional satisfaction. *Mayo Clin Proc* 2016;91(7):836-48.
13. Schiff GD, Amato MG, Egulae T, et al. Computerised physician order entry-related medication errors: analysis of reported errors and vulnerability testing of current systems. *BMJ Qual Saf* 2015;24(4):264-71.
14. Spalding SC, Mayer PH, Ginde AA, et al. Impact of computerized physician order entry on ED patient length of stay. *Am J Emerg Med* 2011;29(2):207-11.
15. Brunette DD, Tersteeg J, Brown N, et al. Implementation of computerized physician order entry for critical patients in an academic emergency department is not associated with a change in mortality rate. *West J Emerg Med* 2013;14(2):114-20.
16. Georgiou A, Prgomet M, Paoloni R, et al. The effect of computerized provider order entry systems on clinical care and work processes in emergency departments: a systematic review of the quantitative literature. *Ann Emerg Med* 2013;61(6):644-53.
17. Abramson E, Patel V, Malhotra S, et al. Physician experiences transitioning between an older versus newer electronic health record for electronic prescribing. *Int J Health Info* 2012;81(8):539-48.
18. Kaji AH, Schriger D, Green S. Looking through the retrospectroscope: reducing bias in emergency medicine chart review studies. *Ann Emerg Med* 2014;64:292-8.
19. Young JQ, Ranji SR, Wachter RM, et al. July effect": impact of the academic year-end changeover on patient outcomes: a systematic review. *Ann Intern Med* 2011;155(5): 309-15.
20. Facebook. "How is reach defined for each of my Page posts?"; 2016. Available at: <https://www.facebook.com/help/241332825914969> (accessed 27 July 2016).
21. Symplur LLC. Symplur – connecting the dots in healthcare social media; 2016. Available at: www.symplur.com (accessed 27 July 2016).
22. Cambridge Journals. Altmetric scores; 2016. Available at: <https://cambridgejournals.altmetric.com/details/9299925#score> (accessed 26 July 2016).
23. Lockett-Gatopoulos S, Thoma B, Milne K, et al. SGEM Hot Off the Press: regional nerve blocks for hip and femoral neck fractures: a systematic review. *CJEM* 2016; 18(4):296-300.
24. Purdy E, Thoma B, Milne K, et al. SGEM Hot Off the Press: hypertonic saline in severe traumatic brain injury: a systematic review and meta-analysis of randomized controlled trials. *CJEM* 2016;18(5):379-84.
25. McKenna P, Thoma B, Milne K, et al. SGEM Hot Off the Press: ultrasound during critical care simulation: a randomized crossover study. *CJEM* 2017;19(1):50-4.
26. Cambridge Journals. Altmetric summary; 2016. Available at: <https://cambridgejournals.altmetric.com/details/9299925/twitter> (accessed 26 July 2016).
27. Eysenbach G. Can tweets predict citations? Metrics of social impact based on Twitter and correlation with traditional metrics of scientific impact. *J Med Internet Res* 2011;13(4): e123, doi:10.2196/jmir.2012.
28. Ringelhan S, Wollersheim J, Welpel I. I like, I cite? Do Facebook likes predict the impact of scientific work? *PLoS One* 2015;10(8):e0134389, doi:10.1371/journal.pone.0134389.
29. Fox CS, Gurary EB, Ryan J, et al. Randomized controlled trial of social media: effect of increased intensity of intervention. *J Am Heart Assoc* 2016;5(5):pii:e004088, doi:10.1161/JAHA.115.003088.