

Spiders (Arachnida: Araneae) from the Insect Limestone (Bembridge Marls, Late Eocene) of the Isle of Wight, southern England

Paul A. Selden^{1,2}

¹ Paleontological Institute and Department of Geology, University of Kansas, 1475 Jayhawk Boulevard, Lawrence, Kansas 66045, USA. Email: selden@ku.edu

² Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK.

ABSTRACT: The arachnids from the late Eocene Insect Limestone of the Isle of Wight are reviewed. Only spiders (Araneae) have been recorded, of which several dozen are known, mostly juveniles. Nearly all are araneomorphs; the putative mygalomorph *Eoatypus woodwardii* McCook, 1888 was removed from that infraorder by Selden (2001), but new specimens, yet to be described, could belong in this group. Among the adult specimens, which are thus identifiable to family level, most belong to the argyronetid water-spider *Vectaraneus yulei* Selden, 2001; some resemble araneids, and a small number can be referred to other families. New morphological information on the holotype of *Vectaraneus* is presented, and three other specimens are described in more detail here, in Segestriidae, Araneidae and Salticidae. Segestriids are nocturnal sedentary hunters living in tubular retreats in rock or bark crevices. Araneids are the archetypal weavers of orb webs. Salticids are the familiar jumping spiders; the one described here is an ant mimic. Their presence suggests a drier habitat was present somewhere in the source environment for the fossil biota, and they were probably inhabitants of the raised hammocks suggested for the Insect Bed palaeoenvironment.

KEY WORDS: Araneidae, Araneoidea, Argyronetidae, Dysderoidea, Salticidae, Salticoidea, Segestriidae



Cenozoic spiders are best known from the amber deposits of the Baltic region of Europe and the Dominican Republic, for example, but fossils preserved in strata with an aqueous sedimentary origin can also yield high-quality morphological information. The Insect Bed of the Isle of Wight has already produced a number of spider fossils which have been informative for studies on spider phylogeny and palaeoecology (Selden 2001, 2002). There are two formally described Insect Bed spiders: *Eoatypus woodwardii* McCook, 1888 and *Vectaraneus yulei* Selden, 2001. *Eoatypus* was described by McCook (1888) as a mygalomorph spider, but the restudy by Selden (2001) showed that this was unlikely and he placed it in Opisthothelae *incertae sedis*. *Vectaraneus* was placed by Selden (2001) in Araneomorphae: Cybaeidae, Argyronetinae, as part of the stem connecting the European Water spider *Argyroneta aquatica* (Clerck 1757) to the remainder of the family. (Note that the family name Argyronetidae has been shown to have priority over Cybaeidae by Grothendieck & Kraus (1994), although its usage has not been accepted by Platnick (2009)). Between Eocene *Vectaraneus* and Recent *Argyroneta* is another fossil spider described by Bertkau (1878) from the Miocene Brown Coal of Rott, Germany.

Here, the araneofauna from the Insect Bed is reviewed, and new specimens are described belonging to the families Segestriidae Simon, 1893, Araneidae Latreille, 1806 and Salticidae Blackwall, 1841. Segestriidae have a reasonable fossil record. They occur in Dominican, Baltic and Bitterfeld ambers (Wunderlich 2004a), and also Cretaceous ambers from Lebanon (Wunderlich & Milki 2004), New Jersey (Penney 2004) and possibly Siberia (Eskov & Wunderlich 1995). They are also known from some Cenozoic rock matrix deposits, such as Florissant, Colorado (Scudder 1890; Petrunkevitch 1922). Ara-

neids have a similar record, being known from most Cenozoic ambers (Wunderlich 2004b), and it is mostly juveniles which have been reported from Cretaceous ambers, e.g. Lebanon (Wunderlich 2004c), Canada (McAlpine & Martin 1969), Siberia (Eskov & Wunderlich 1995) and New Jersey (Penney 2004). Araneidae are also known from some Cenozoic rock matrix localities such as Aix-en-Provence (Gourret 1886), Florissant (Scudder 1890; Petrunkevitch 1922), Messel, Germany (Wunderlich 1986a), Öhningen, Switzerland (Heer, 1865, 1872, 1876) and Shanwang, China (Zhang *et al.* 1994). Wunderlich (1986b) suggested that the fossil described by Eskov (1984) as *Juraneus* in the extinct family Juraneidae from Transbaikalia, Siberia may actually belong in the Araneidae, which would extend the known range of the family back to the Jurassic; however, this synonymy has not yet been formally established (Penney 2004). Salticids have an interesting fossil record. Because they are the most diverse extant spider family, they would be expected to turn up frequently as fossils and, indeed, they are common in Cenozoic ambers (Wunderlich 2004d). They are also known from rock matrix deposits of the Cenozoic, such as Aix-en-Provence (Gourret 1886), Florissant, Colorado (Scudder 1890), Randecker Maar, Germany (Schawaller & Ono 1979), Willershausen, Germany (Schawaller 1982), British Columbia, Canada (Selden & Penney 2009) and Shanwang, China (Zhang *et al.* 1994). No salticid is known from the Mesozoic, which is surprising if they were indeed present then. Quite possibly, they originated or radiated in the Cenozoic, and their absence as fossils in Mesozoic strata is real.

The fossil spiders described most likely come from a variety of habitats which, together with the preponderance of juveniles, suggests that extraordinary events, rather than continuous attrition, led to the preservation of the spiders on the Insect Bed.

1. Locality and stratigraphy

Specimens from the Smith, Brodie and Hooley collections in the Natural History Museum are labelled 'Gurnard Bay' or 'Gurnet Bay' (an old name for Gurnard Bay). However, Smith collected specimens all the way from West Cowes to Newtown River on the northwest side of the Isle of Wight (Jarzembowski 1980). Most of the specimens probably came from Thorness Bay (Jarzembowski 1976). The specimens come from concretions or tabular bands of very fine-grained micrite, known as Insect Limestone. The unit in which these concretions/bands occur is known as the Insect Bed, which lies towards the base of the Bembridge Marls Member (Solent Group: Bouldnor Formation). The Bembridge Marls were considered to be early Oligocene by Gale *et al.* (2006), but later work by Hooker *et al.* (2007, 2009) place them as latest Eocene (Priabonian) in age.

2. Material and methods

Specimens studied for the INTAS project came from the Natural History Museum, London (NHMUK), the Sedgwick Museum of Earth Sciences, Cambridge (CAMSM), the Isle of Wight Museum of Geology, Sandown, Isle of Wight (MIWG) and the Booth Museum of Natural History, Brighton, Sussex (BMB). Specimens I.8442, I.8704 and In.17177, described in detail here, are in the Natural History Museum, London. Most of the NHMUK specimens belong to the A'Court Smith (purchased 1877, 1883), P. B. Brodie (purchased 1898) and R. W. Hooley (purchased 1924) collections; and specimens collected over the last two decades by Mr Andrew Yule have also been deposited in the NHMUK and MIWG, and thus made available for study. Several dozens (but <100) of arachnid specimens from the museums mentioned above have been studied; there are 53 currently in the Natural History Museum collections alone (see Appendix I). All are spiders; to date, no other arachnids have been identified in the Insect Bed collections. The vast majority of specimens are juveniles, several of which can be referred to *Vectaraneus* on account of their general habitus and features such as sternum shape. Others appear to be araneoids, but firm identification to family is generally impossible in immatures.

The most remarkable aspect of the Bembridge spiders is the extremely fine preservation of internal anatomical structure. Some organic matter is preserved, mainly thin laths of cuticle lining the external mould cavities of the chelicerae, and setae within the rock which can be seen when the specimens are observed under alcohol. Within the moulds of the chelicerae, labium and coxae, fine fibres of cream-coloured calcite can be seen. Presumably, these represent muscle fibres replaced by calcite during diagenesis, as described for the Bembridge Marls insects by McCobb *et al.* (1998). The holotype of *Vectaraneus yulei*, BMB 021960/1, is a good example of the preservation of the Insect Bed spiders (Selden 2001). Much of the opisthosoma of BMB 021960/1 is filled with cream-coloured calcite, which strikingly preserves the large tracheae as tubes running forward from the wide spiracle near the middle of the opisthosoma. The inner surfaces of these tubes bear reinforcements preserved in the same cream calcite. Book-lung lamellae are preserved in buff-coloured calcite, and the book-lung atria anterior to the lamellae are lined with tiny, buff, drusy calcite crystals. Before its removal to reveal the nature of the spinnerets, the posterior part of the opisthosoma of BMB 021960/1 was filled with calcite showing a spheroidal structure in places. This could represent silk glands or ova, or could be botryoidal mineral growth inside a cavity. Because of the preservation of internal anatomy, their three-dimensional nature, and their completeness (e.g. carapaces are intact, not

detached) most of the fossils probably represent dead animals rather than moults.

The specimens were photographed with a Leica digital camera, and drawn using a camera lucida attachment, on Leica MZ16 stereomicroscopes. Specimens were scanned on a Metris X-Tek HMX-ST scanner (Natural History Museum, London) with a tungsten reflection target at 200mA and 225kV, 0.17–1 second exposure times for 3142 projections, and a 1 mm copper filter. The 4MP (2000 × 2000) Perkin Elmer detector panel provided a voxel-size (resolution) of ~20 microns. Analysis, reconstruction, manual preparation and colour coding of the three-dimensional model from tomographic slices was performed using the custom SPIERS software suite (Sutton 2009, unpublished), implementing the methods described by Sutton *et al.* (2001, 2002). Visualisations are screen captures from the SPIERSview renderer. Drawings and photographs were prepared for publication using the Adobe Creative Suite software. All measurements are in mm.

3. Systematic palaeontology

Order Araneae Clerck, 1757

Superfamily Dysderoidea C. L. Koch, 1837

Family Segestriidae Simon, 1893

Remarks. Specimen NHMUK I.8442 can be assigned to this family on the following combination of characters: medium-sized; suboval sternum, projected between coxae (but not as dramatically as in dysderids); large, elongate palpal endites; elongate labium (if notched apically then only slightly); chelicerae not obviously fused basally; coxae subequal in length (in Dysderidae C. L. Koch, 1837 coxae 1 and 2 are nearly twice as long as coxae 3 and 4 (Almquist 2005)); legs 1 and 2 with spines (only legs 3 and 4 have spines in Dysderidae: Ubick (2005a)). The punctate ornament on all body parts is rather more reminiscent of dysderids than segestriids.

Genus *Segestria* Latreille, 1804

Remarks. There are three genera of Segestriidae: *Ariadna* Audouin, 1826, *Gippsicola* Hogg, 1900 and *Segestria* Latreille, 1804 (Platnick 2009). *Gippsicola* is known from a single species in Victoria, Australia. *Ariadna* and *Segestria* are cosmopolitan and each has numerous species. The former has relatively short legs, whilst those of the latter genus are longer (Ubick 2005b). The leg lengths of the fossil specimen are closer to those of *Segestria* than *Ariadna*, so the fossil is referred to that genus. There are insufficient additional features to place the species or to create a new species name for the fossil.

Segestria sp.

Fig. 1

Material. NHMUK I.8442 in the Natural History Museum, London, Brodie Collection; female or immature; Insect Bed of the Bembridge Marls Member of the Bouldnor Formation (Solent Group), latest Eocene (Priabonian), of Thorness Bay, Isle of Wight, England.

Description. Integument regularly punctate. Carapace not preserved. Sternum suboval, projected between coxae and labium; length 1.83, width 1.52. Labium twice as long as wide (length 0.84, width 0.42), if notched apically then not strongly, small projections laterally at mid-length. Pedipalps coxa length 1.13, broad posteriorly, narrowing to endite as long as labium. Pedipalps short: left femur length 0.95, right




Figure 1 *Segestria* sp., NHMUK I.8442, Insect Bed, Bembridge Marls Member, Bouldnor Formation (Solent Group), Thorness Bay, Isle of Wight: (A) photograph $\times 80$; (B) explanatory camera lucida drawing. Scale bar = 0.5 mm. Abbreviations: 1–4 = leg numbers; ch = chelicera; cx = coxa; fe = femur; l = labium; m = maxilla (endite of pedipalp coxa); mt = metatarsus; pa = patella; Pd = pedipalp; st = sternum; ta = tarsus; ti = tibia.

patella length 0.41. Right leg 1 podomere lengths: coxa 0.95; femur 2.47; tibia 2.53; metatarsus 1.95. Right leg 2 podomere lengths: coxa 1.05; tibia ≥ 1.75 ; metatarsus 1.73. Right leg 3 coxa length 0.91. Right leg 4 podomere lengths: coxa 0.89; femur 2.07; patella 0.81. All other podomeres not fully preserved to measure lengths. Spines present at least on pedipalp tibia and tarsus, leg 1 femur (distal prolateral), leg 1 tibia (distal prolateral), legs 1 and 2 metarsi (pro- a retrolateral from $1/3$ length to distal). Opisthosoma not preserved.

Superfamily Araneoidea Latreille, 1806

Remarks. Specimen NHMUK In.17177 is chosen as an example of one of the small, three-dimensionally-preserved immature spiders which are common in the Insect Bed. It can be referred to Araneoidea on the following characters. The legs are long and slender, particularly the first; the opisthosoma is subcircular and the spinnerets are grouped towards the centre of the venter. There are no noticeable specialities, so the spider is almost certainly an orbicularian (orb-web weaver). The lack of a cribellum (which would be present in immatures of Deinopoidea C. L. Koch, 1850, the sister-group of Araneoidea) places it in Araneoidea.

Family Araneidae Latreille, 1806

Remarks. Specimen NHMUK In.17177 can be placed in this family on account of the rebordered labium, sternum pointed posteriorly (it is truncated in Theridiosomatidae: Coddington (2005), Levi (2005), and short palpal endites.

Araneidae incertae sedis

Fig. 2

Material. NHMUK In.17177 in the Natural History Museum, London, Smith Collection; immature; Insect Bed of the Bembridge Marls Member of the Bouldnor Formation (Solent Group), latest Eocene (Priabonian), of Thorness Bay, Isle of Wight, England.

Description. Carapace not preserved. Sternum longer than wide (length 0.73, width 0.52), widest near anterior margin, procurved at labium, pointed posteriorly between coxae 4. Labium length 0.19, width 0.12, rebordered. Leg podomeres difficult to measure, but leg 1 at least 2.5 long (distal podomeres patella–tarsus ≥ 1.1). Podomere lengths: leg 1 coxa 0.31; leg 2 coxa 0.30; leg 3 coxa 0.28; leg 3 femur 0.72; leg 4 coxa 0.26. Opisthosoma longer than wide (length 1.22, width 0.99), spinnerets located just posterior to centre of venter; anterior spinneret (AS) and posterior spinneret (PS) similar in size, median spinneret (MS) very small; tiny colulus present; anal tubercle about same size as AS/PS.

Superfamily Dictynoidea Simon, 1874

Family Argyronetidae Thorell, 1870

Genus *Vectaraneus* Selden, 2001

Vectaraneus yulei Selden, 2001

Figs 3, 4

Material. BMB 021960 part and BMB 021961 counterpart (ventral and dorsal) respectively, in the Booth Museum of Natural History, Brighton, England, Jarzembowski Collec-




Figure 2 *Araneidae incertae sedis*, NHMUK In.17177, Insect Bed, Bembridge Marls Member, Bouldnor Formation (Solent Group), Thorness Bay, Isle of Wight. (A) photograph $\times 25$; (B) explanatory camera lucida drawing. Scale bar = 0.5 mm. Abbreviations: 1–4 = leg numbers; a t = anal tubercle; AS = anterior spinneret; ch = chelicera; col = colulus; cx = coxa; fe = femur; l = labium; m = maxilla (endite of pedipalp coxa); MS = median spinneret; op = opisthosoma; Pd = pedipalp; PS = posterior spinneret.

tion; adult male; Insect Bed of the Bembridge Marls Member of the Bouldnor Formation (Solent Group), latest Eocene (Priabonian), of Thorness Bay, Isle of Wight, England.

Remarks. The holotype specimen was described by Selden (2001) as questionably a mature female, on basis of lack of evidence of male characters. Since then, Timothy Chalk, in an undergraduate research project in the Department of Earth Science & Engineering, Imperial College, London, under the supervision of Mark Sutton, has produced some x-ray computer tomography (XMT) scans of the holotype which shows modified male palps (Figs 3B, 4), thus proving that the specimen is, in fact, an adult male. The elongate cymbium is typical of Argyronetidae (and also present in other, related families). The XMT scans (Figs 3, 4) also show some parts of the anterior legs and mouthparts which are not visible using light microscopy on the rock specimens. The long, tapering fang and large, bifid tooth on the distal end of the promargin of the chelicera, and the spatulate shape of the labium, are confirmed.

Additional description of holotype BMB 021960/1. Adult male. Adult male palp (Fig. 4) with elongate cymbium, length 0.92. Legs 1 and 2 thin, elongate; approximate (minimum) podomere lengths: leg 1 tibia 3.4; metatarsus 2.5; tarsus 1.2; leg 2 metatarsus 1.9; tarsus 0.8.

Superfamily Salticoidea Blackwall, 1841
Family Salticidae Blackwall, 1841

Remarks. Specimen NHMUK I.8704 shows characteristics of this unmistakable spider family: parallel-sided carapace with prominent anterior lateral eyes; long pedicel connecting

prosoma and opisthosoma; and relatively short legs. The only other extant family to which NHMUK I.8704 could be referred would be Corinnidae Karsch, 1880, but the prominent anterior lateral eyes and relatively short legs lend no support for this placement. Prominent ?anterior lateral eyes are also present in the extinct family Lagonomegopidae Eskov & Wunderlich, 1995; however, apart from this family being unknown outside of the Mesozoic (Penney 2005, 2006), lagonomegopids have longer legs and an inconspicuous pedicel, thus giving these spiders a quite different appearance to the fossil described here.

Salticidae is the most diverse spider family today, with 5202 species in 563 genera at the last count (Platnick 2009). Most are small and commonly brightly coloured, diurnal predators, using visual clues for prey capture and mating. Salticidae are jumping spiders which use the unique aranean hydrostatic muscle extensor system for saltation. Consequently, the prosoma of many salticids is high, to accommodate the pump required to produce a burst of haemolymph pressure to the rather short legs and initiate a jump. In many salticids, the carapace is not short and tall, but elongated, as in the fossil described here. Such salticids may be more commonly found running on the ground and many mimic ants and other ground-living biting insects. Lower systematics of salticids is not related to gross morphology, so it is not possible to determine to which subfamily or genus the fossil is related, hence it is referred to *Salticidae incertae sedis*.

Salticidae incertae sedis
Fig. 5




Figure 3 *Vectaraneus yulei* Selden, 2001, BMB 021960 (part) and 021961 (counterpart), Insect Bed, Bembridge Marls Member, Bouldnor Formation (Solent Group), Thorness Bay, Isle of Wight. Combined XMT scans of part and counterpart: (A) dorsal view; (B) ventral view; (C) lateral view. Note that the dorsal and ventral views are not from the same angle, hence legs appear in different orientations. X–Y in C shows approximate line of join of part and counterpart. Scale bars = 1 mm. Abbreviations: 1–4 = leg numbers; a t = anal tubercle; AS = anterior spinneret; car = carapace; ch = chelicera; cx = coxa; lab = labium; MS = median spinneret; mt = metatarsus; Pd = pedipalp; PS = posterior spinneret; st = sternum; ta = tarsus; ti = tibia.




Figure 4 *Vectaranus yulei* Selden, 2001, BMB 021960 (part) and 021961 (counterpart), Insect Bed, Bembridge Marls Member, Bouldnor Formation (Solent Group), Thorness Bay, Isle of Wight. Combined XMT scans of part and counterpart showing detail of male pedipalps. See Figure 3 for scale. Abbreviations: pa = patella; ti = tibia.

Material. NHMUK I.8704, Brodie Collection; sex and maturity not known; Insect Bed of the Bembridge Marls Member of the Bouldnor Formation (Solent Group), latest Eocene (Priabonian), of Thorness Bay, Isle of Wight, England.

Description. Carapace about twice as long as wide (length 1.36, width 0.72), subrectangular, anterior border straight, parallel-sided. Presumed anterior lateral eyes present at anterolateral corners of carapace. Legs relatively short, stout, approximately equal in length. Podomere lengths: leg 1 tarsus 0.82; leg 2 metatarsus 0.75; tarsus ≥ 0.49 ; leg 3 metatarsus 0.87; tarsus 0.80; leg 4 tibia 0.98; metatarsus 0.73; tarsus ≥ 0.41 . Opisthosoma connected to prosoma by long pedicel. Opisthosoma about twice as long as wide (length 1.95, width 1.09), wrinkled posteriorly.

4. Discussion

The spiders found in the Insect Bed come from a number of extant families, each with different habits and from a variety of habitats. *Vectaranus yulei* is an argyronetid water-spider, and is likely to have spent some of its time under water, as evidenced by the enlarged tracheal system. Its mode of life within the Insect Bed ecosystem is clearly linked with the aquatic habitat. Araneidae and other orb-web weavers could have made their prey-capture webs in reeds and other emergent aquatic vegetation; indeed, orb-weavers are common in such habitats today. Such spiders are generally fairly adept at avoiding falling into the water and, if they did, would then remain at the surface due to surface tension. This, together with the large number of juveniles in the collection, suggest that it was unusual events which led to their entombment in the lime mud, rather than chance natural attrition of the living population. Segestriids live in silken tubes in cracks in bark or under stones; 'trip-lines' radiating from the tube entrance provide information to the spider when possible prey passes close by. Segestriids generally respond actively to prey at night. Their presence in the Insect Bed suggests the presence of trees with creviced bark, rocky outcrops or stony debris in the vicinity of the centre of deposition, although it is possible that they could make tube webs in dead reed stems. Again, their presence in the thanatocoenosis is unlikely to have been a normal chance occurrence. Unlike segestriids, Salticidae are diurnal spiders which have shunned the prey-capture web in favour of active hunting. They could occur in any habitat, but prefer sunny sites where insects constantly land (e.g. flies) or are in abundance (e.g. ants). The salticid described here has the morphology of an active ground-dweller, and so is suggestive of an emerged dry habitat, but could also occur in a reed swamp, jumping from plant to plant. Catastrophic events are more likely to knock such an agile hunter into the water than normal mortality.





Figure 5 Salticidae *incertae sedis*, NHMUK I.8704, Insect Bed, Bembridge Marls Member, Bouldnor Formation (Solent Group), Thorness Bay, Isle of Wight: (A) photograph $\times 50$; (B) explanatory camera lucida drawing. Scale bar = 0.5 mm. Abbreviations: 1–4 = leg numbers; ALE = anterior lateral eye; car = carapace; mt = metatarsus; op = opisthosoma; ta = tarsus; ti = tibia.

In conclusion, the spider fossils in the Insect Bed suggest unusual preservation events; storms perhaps, which collected together specimens from a variety of disparate habitats.

5. Acknowledgements

This paper is a contribution to INTAS Project No. 03-51-4367, who are thanked for their support. I thank Paulyn Cartwright and Annalise Nawrocki (Ecology and Evolutionary Biology, University of Kansas) for their help with microscopy, and Timothy Chalk and Mark Sutton, Imperial College, London, for XMT scans of the holotype of *Vectaraneus*.

6. Appendix I. List of specimens labelled as Araneae in the collections of the Natural History Museum, London (NHMUK)

Acc. No.	Comments
I.470	six pieces in various states of preservation; some are <i>Vectaraneus</i>
I.8437	label over specimen
I.8438	<i>Vectaraneus</i> paratype
I.8439	3D, no abdomen
I.8440	<i>Vectaraneus</i> paratype
I.8441	Old, rubbed and poor preservation
I.8442	Dysderidae
I.8443	imm. <i>Vectaraneus</i>
I.8444	half a spider
I.8445	poor preservation
I.8448	3D linyphiid?
I.8449	large but poor preservation
I.8452	<i>Vectaraneus</i>
I.8454	<i>Vectaraneus</i> ?
I.8455	tiny specimen
I.8457	tiny 3D specimen
I.8458	3D specimen
I.8459	poor preservation
I.8462	spider?
I.8463	some brown cuticle preserved
I.8464	poor preservation
I.8529	counterpart of In.25368
I.8563	largish but poorly preserved spider on weathered surface
I.8579	poor preservation
I.8704	Salticidae
I.8787	spider, rather mashed
I.9742	tiny spider
In.17121	tiny spider
In.17139	tiny spider
In.17147	poor preservation
In.17149	tiny spider; linyphiid?
In.17152	poor preservation
In.17177	3D theridiid?
In.17206	poor preservation
In.17207	part and counterpart, poor preservation; salticid?
In.17323	possibly interesting
In.17354	small, 3D female linyphiid
In.17487	lateral preservation; mygalomorph?
In.24866	tiny, 3D, poor preservation
In.24954	mess of arthropods
In.25235	two specimens in mess of arthropods
In.25264	mess of arthropods
In.25368	many insects, large but poor spider, counterpart of I.8529
In.25731	3D spider, legs contracted; part and counterpart

In.25732	probably <i>Vectaraneus</i>
In.25733	3D
In.25734	tiny specimen
In.25735	tiny 3D salticid?
In.25736	not a spider
In.25737	tiny, 3D carapace
In.25738	tiny 3D specimen
In.61271	<i>Eoatypus woodwardi</i> holotype
In.77151	<i>Vectaraneus</i> paratype

7. References

- Almqvist, S. 2005. Swedish Araneae, part. 1: families Atypidae to Hahnidae (Linyphiidae excluded). *Insect Systematic and Evolution Supplement* **62**, 1–284.
- Audouin, V. 1826. Explication sommaire des planches d'arachnides de l'Égypte et de la Syrie. In Savigny, J-C (ed.) *Description de l'Égypte ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française* **1**(4), 99–186. Paris: C. L. F. Panckoucke. 339 pp.
- Bertkau, P. 1878. Einige Spinnen und ein Myriapode aus der Braunkohle von Rott. *Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande und Westfalens, Bonn* **35**, 346–60, pl. 5.
- Blackwall, J. 1841. The difference in the number of eyes with which spiders are provided proposed as the basis of their distribution into tribes; with descriptions of newly discovered species and the characters of a new family and three new genera of spiders. *Transactions of the Linnean Society of London* **18**, 601–70.
- Clerck, C. 1757. *Araneae suecici, descriptionibus et figuris oeneis illustrati, ad genera subalterna redacti speciebus ultra LX determinati*. Stockholm: L. Salvii. 154 pp, 6 pls. [In Swedish and Latin.]
- Coddington, J. A. 2005. Theridiosomatidae. In Ubick, D., Paquin, P., Cushing, P. E. & Roth, V. (eds) *Spiders of North America: an identification manual*, 244–45. College Park, MD: American Arachnological Society. v + 377 pp.
- Eskov, K. Y. 1984. A new fossil spider family from the Jurassic of Transbaikalia (Araneae, Chelicerata). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* **1984**, 645–53.
- Eskov & Wunderlich, J. 1995. On the spiders from Taimyr ambers, Siberia, with the description of a new family and with general notes on the spiders from the Cretaceous resins. *Beiträge zur Araneologie* **4**, 95–107.
- Gale, A. S., Huggett, J. M., Pälke, H., Laurie, E., Hailwood, E. A. & Hardenbol, J. 2006. Correlation of Eocene–Oligocene marine and continental records: orbital cyclicity, magnetostratigraphy and sequence stratigraphy of the Solent Group, Isle of Wight, UK. *Journal of the Geological Society, London* **163**, 401–15.
- Grothendieck, K. & Kraus, O. 1994. Die Wasserspinne *Argyroneta aquatica*: Verwandtschaft und Spezialisierung (Arachnida, Araneae, Agelenidae). *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg (NF)* **34**, 259–73.
- Gourret, P. 1886. Recherches sur les arachnides tertiaires d'Aix en Provence. *Recueil Zoologique Suisse* **4**, 431–96, pls XX–XXIII.
- Heer, O. 1865. *Die Urwelt der Schweiz*. Zurich: Friedrich Schulthess. 622 pp.
- Heer, O. 1872. *Le monde primitif de la Suisse*. Genève and Basle: H. Georg. 801 pp. [French translation of Heer 1865].
- Heer, O. 1876. *The primaevial world of Switzerland*. Volume 2. Heywood, J. (ed.). London: Longmans, Green & Co. 324 pp. [English translation of Heer 1865.]
- Hogg, H. R. 1900. A contribution to our knowledge of the spiders of Victoria; including some new species and genera. *Proceedings of the Royal Society of Victoria* **13**, 68–123.
- Hooker, J. J., Collinson, M. E., Grimes, S. T., Sille, N. P. & Matthey, D. P. 2007. Discussion on the Eocene–Oligocene boundary in the UK. *Journal of the Geological Society, London* **164**, 685–88.
- Hooker, J. J., Grimes, S. T., Matthey, D. P., Collinson, M. E. & Sheldon, N. D. 2009. Refined correlation of the UK Late Eocene–Early Oligocene Solent Group and timing of its climate history. *Geological Society of America Special Paper* **452**, 179–95.
- Jarzewowski, E. A. 1976. Report of Easter field meeting: the Lower Tertiaries of the Isle of Wight, 27–31.III.1976. *Tertiary Research* **1**, 11–16.
- Jarzewowski, E. A. 1980. Fossil insects from the Bembridge Marls, Palaeogene of the Isle of Wight, southern England. *Bulletin of the British Museum (Natural History) Geology* **33**, 237–93.
- Karsch, F. 1880. Arachnologische Blätter (Decas I). *Zeitschrift für die Gesamte Naturwissenschaft* **53**, 373–409.

- Koch, C. L. 1837. *Übersicht des Arachnidensystems*, Heft 1. Nürnberg. 39 pp.
- Koch, C. L. 1850. *Übersicht des Arachnidensystems*, Heft 5. Nürnberg. 77 pp.
- Latreille, P.-A. 1804. Tableau méthodique des Insectes. *Nouveau Dictionnaire d'Histoire Naturelle* **24**, 129–200.
- Latreille, P.-A. 1806. *Genera crustaceorum et insectorum, volume 1, Aranéides*, 82–127. Paris: A. Koenig.
- Levi, H. W. 2005. Araneidae. In Ubick, D., Paquin, P., Cushing, P. E. & Roth, V. (eds) *Spiders of North America: an identification manual*, 68–74. College Park, MD: American Arachnological Society. v + 377 pp.
- Mcalpine, J. F. & Martin, J. E. H. 1969. Canadian amber – a paleontological treasure chest. *Canadian Entomologist* **101**, 819–38.
- Mccobb, L. M. E., Duncan, I. J., Jarzembowski, E. A., Stankiewicz, B. A., Wills, M. A. & Briggs, D. E. G. 1998. Taphonomy of the insects from the Bembridge Marls (Late Eocene), Isle of Wight, England. *Geological Magazine* **135**, 553–63.
- Mccook, H. C. 1888. A new fossil spider, *Eoatypus woodwardii*. *Proceedings of the Academy of Natural Sciences of Philadelphia* **1888**, 200–02.
- Penney, D. 2004. New spiders in Upper Cretaceous amber from New Jersey in the American Museum of Natural History (Arthropoda: Araneae). *Palaeontology* **47**, 367–75.
- Penney, D. 2005. The fossil spider family Lagonomegopidae in Cretaceous ambers with descriptions of a new genus and species from Myanmar. *Journal of Arachnology* **33**, 439–44.
- Penney, D. 2006. The oldest lagonomegopid spider, a new species in Lower Cretaceous amber from Alava, Spain. *Geologica Acta* **4**, 377–82.
- Petrunkovitch, A. 1922. Tertiary spiders and opilionids of North America. *Transactions of the Connecticut Academy of Arts and Sciences* **25**, 211–79.
- Platnick, N. I. 2009. *The world spider catalog, version 9.5*. New York: American Museum of Natural History. (<http://research.amnh.org/entomology/spiders/catalog/index.html>)
- Schawaller, W. 1982. Zur fossilen Spinnenfauna des Pliozäns von Willershausen in Norddeutschland (Arachnida, Araneae). *Bericht der naturhistorischen Gesellschaft zu Hannover*, **125**, 89–95.
- Schawaller, W. & Ono, H. 1979. Fossile Spinnen aus miozänen Sedimenten des Randecker Maars in SW-Deutschland (Arachnida: Araneae). *Jahreshefte der Gesellschaft für Naturkunde in Württemberg* **134**, 131–41.
- Scudder, S. H. 1890. The Tertiary insects of North America. *Report of the United States Geological Survey of the Territories* **13**, 1–734.
- Selden, P. A. 2001. Eocene spiders from the Isle of Wight with preserved respiratory structures. *Palaeontology* **44**, 695–729.
- Selden, P. A. 2002. Missing links between *Argyroneta* and Cybaeidae revealed by fossil spiders. *Journal of Arachnology* **30**, 189–200.
- Selden, P. A. & Penney, D. 2009. A fossil spider (Araneae: Pisauridae) of Eocene age from Horsefly, British Columbia, Canada. In Kropf, C. & Horak, P. (eds) *Towards a natural history of arthropods and other organisms. In memoriam Konrad Thaler. Contributions to Natural History* **12**, 1269–82.
- Simon, E. 1874. *Les arachnides de France*. Volume 1, Paris: 1–272. Librairie encyclopédique de Roret.
- Simon, E. 1893. *Histoire naturelle des araignées*. Volume 1, part 2, 257–488. Paris: Librairie encyclopédique de Roret.
- Sutton, M. D., Briggs, D. E. G., Siveter, D. J. & Siveter, D. J. 2001. Methodologies for the visualization and reconstruction of three-dimensional fossils from the Silurian Herefordshire Lagerstätte. *Palaeontologia Electronica* **4**, 1–17. [http://palaeo-electronica.org/2001_1/s2/issue1_01.htm]
- Sutton, M. D., Briggs, D. E. G., Siveter, D. J., Siveter, D. J. & Orr, P. J. 2002. The arthropod *Offacolus kingi* (Chelicerata) from the Silurian of Herefordshire, England: computer based morphological reconstructions and phylogenetic affinities. *Proceedings of the Royal Society, London* **B269**, 1195–203.
- Thorell, T. 1870. On European spiders. Part 2. *Nova Acta Societas Scientiae Uppsalensis* (3) **7**, 109–242.
- Ubick, D. 2005a. Dysderidae. In Ubick, D., Paquin, P., Cushing, P. E. & Roth, V. (eds) *Spiders of North America: an identification manual*, 103. College Park, MD: American Arachnological Society. v + 377 pp.
- Ubick, D. 2005b. Segestriidae. In Ubick, D., Paquin, P., Cushing, P. E. & Roth, V. (eds) *Spiders of North America: an identification manual*, 219–10. College Park, MD: American Arachnological Society, v + 377 pp.
- Wunderlich, J. 1986a. Die ersten Spinnen aus dem Mittel-Eozän der Grube Messel. *Senckenbergiana lethaea* **67**, 171–76.
- Wunderlich, J. 1986b. *Spinnenfauna gestern und heute. Fossile Spinnen in Bernstein und ihre heute lebenden Verwandten*. Wiesbaden: Erich Bauer Verlag bei Quelle und Meyer. 283 pp.
- Wunderlich, J. 2004a. Fossil spiders (Araneae) of the superfamily Dysderoidea in Baltic and Dominican amber, with revised family diagnoses. *Beiträge zur Araneologie* **3**, 633–746.
- Wunderlich, J. 2004b. Fossil taxa of the family Araneidae (Araneae) inclusively Nephilinae in Baltic and Dominican amber, with the description of a new extinct subfamily and notes on selected extant taxa. *Beiträge zur Araneologie* **3**, 956–97.
- Wunderlich, J. 2004c. Report on spiders (Araneae) of the families Araneidae and Zygellidae in Lebanese amber. *Beiträge zur Araneologie* **3**, 1864–65.
- Wunderlich, J. 2004d. Fossil jumping spiders (Araneae: Salticidae) in Baltic and Dominican amber, with remarks on salticid subfamilies. *Beiträge zur Araneologie* **3**, 1761–819.
- Wunderlich, J. & Milki, R. 2004. Description of the extinct new subfamily Microsegestriinae (Araneae: Segestriidae) in Cretaceous Lebanese amber. *Beiträge zur Araneologie* **3**, 1867–73.
- Zhang, J., Sun, B. & Zhang, X. 1994. *Miocene insects and spiders from Shanwang, Shandong*. Beijing: Science Press. 298 pp. [In Chinese with English summary.]