

# The singularity spectrum of the inverse of cookie-cutters

JULIEN BARRAL<sup>†</sup> and STÉPHANE SEURET<sup>‡</sup>

<sup>†</sup> *INRIA Rocquencourt, 78153 Le Chesnay Cedex, France*  
(e-mail: julien.barral@inria.fr)

<sup>‡</sup> *Laboratoire d'Analyse et de Mathématiques Appliquées, Université Paris-Est,  
CNRS UMR 8050-61, avenue du Général de Gaulle, 94010 Créteil Cedex, France*  
(e-mail: seuret@univ-paris12.fr)

(Received 12 March 2008 and accepted in revised form 20 June 2008)

*Abstract.* Gibbs measures  $\mu$  on cookie-cutter sets are the archetype of multifractal measures on Cantor sets. We compute the singularity spectrum of the inverse measure of  $\mu$ . Such a measure is discrete (it is constituted only by Dirac masses), it satisfies a multifractal formalism and its  $L^q$ -spectrum possesses one point of non-differentiability. The results rely on heterogeneous ubiquity theorems.

## 1. Introduction

Gibbs measures  $\mu$  on cookie-cutter sets are the archetype of multifractal measures on Cantor sets. The singularity spectrum of such a measure  $\mu$  has been obtained in one of the pioneer papers on multifractal analysis [24]. Then, many papers have extended this study for Gibbs measures or Birkhoff averages on conformal repellers (see, for instance, [7, 16, 21, 22]). As noted in [19], a very natural object associated with a Gibbs measure on a cookie-cutter set is its inverse measure  $\nu$ , which belongs to the class of purely discrete measures. The multifractal nature of the measure  $\nu$  cannot be described only by the techniques developed for the study of measures generated by a multiplicative procedure and enjoying self-similarity properties. The aim of this work is to perform the multifractal analysis for the measure  $\nu$ , and to prove that  $\nu$  also obeys some multifractal formalism. It appears that the  $L^q$ -spectrum  $\tau_\nu(q)$  of  $\nu$  is analytic except at the point  $q$  equal to the Hausdorff dimension of the cookie-cutter set, where it is non-differentiable. This non-differentiability is reminiscent of the phase transition phenomenon occurring in the thermodynamic formalism setting. The multifractal analysis of the discrete measure  $\nu$  is explained by a subtle combination between the fine local structure of Gibbs measures and the distribution of the jump points of  $\nu$ . The results rely on the notion of *heterogeneous ubiquity* introduced in [2, 4].

Subsequently, taking the inverse of Gibbs measures on cookie-cutter sets provides us with a very natural way of generating new objects obeying the multifractal formalism, although they are of a very different nature than Gibbs measures. The relationship between the spectra of  $\mu$  and  $\nu$  will be given.

We start by recalling the definition of a cookie-cutter set.

For every  $n \geq 0$ , let  $\Sigma_n = \{0, 1\}^n$ , where  $\Sigma_0$  contains only the empty word  $\emptyset$ . Also, let  $\Sigma^* = \bigcup_{n \geq 0} \Sigma_n$  and let  $\Sigma = \{0, 1\}^{\mathbb{N}^*}$  be the set of infinite words on the alphabet  $\{0, 1\}$ . The concatenation operation from  $\Sigma^* \times (\Sigma^* \cup \Sigma)$  to  $(\Sigma^* \cup \Sigma)$  is denoted by  $\cdot$ .

The notation  $\bar{0}$  and  $\bar{1}$  is for the infinite words in  $\Sigma$  whose letters are respectively all equal to 0 and 1.

For  $w = w_1 w_2 \dots \in \Sigma$  and  $n \geq 1$ ,  $w|n := w_1 w_2 \dots w_n$  is the projection of  $w$  on  $\Sigma_n$  and  $w|\infty = w$ . For  $w \in \Sigma_n$ , the cylinder with root  $w$  is  $[w] = \{x \in \Sigma : x|n = w\}$ . Given two words of infinite length  $(w_1, w_2) \in \Sigma^2$ , one defines  $w_1 \wedge w_2$  as  $w_1|n_0$ , where  $n_0 = \sup\{n \geq 1 : w_1|n = w_2|n\}$ . We adopt the convention that  $\inf \emptyset = 0$  and  $w|0$  is the empty word  $\emptyset$ .

The length of any element  $w$  of  $\Sigma_n$  is equal to  $n$  and is denoted by  $|w|$ .

The set  $\Sigma$  is endowed with the shift operation denoted by  $\sigma : w = w_1 w_2 \dots \in \Sigma \mapsto \sigma(w) = w_2 w_3 \dots \in \Sigma$ .

A *cookie-cutter set* is defined as follows (see, for instance, [24]). Let  $U_0$  and  $U_1$  be two compact disjoint subintervals of  $[0, 1]$ . For  $i \in \{0, 1\}$  consider  $T_i : U_i \rightarrow [0, 1]$ , a  $C^{1+\gamma}$  ( $\gamma > 0$ ) diffeomorphism such that  $|T_i'| > 1$ , and denote by  $g_i = (T_i)^{-1}$  the inverse of  $T_i$ .

We denote by  $T$  the mapping from  $U_0 \cup U_1$  to  $[0, 1]$  whose restrictions to  $U_0$  and  $U_1$  are respectively  $T_0$  and  $T_1$ .

Then consider the subset  $X$  of  $[0, 1]$ , called the cookie-cutter set associated with  $T$ , given by

$$X = \bigcap_{n \geq 1} \bigcup_{w \in \Sigma_n} g_{w_1} \circ \dots \circ g_{w_n}([0, 1]).$$

By construction  $T^{-1}(X) = X$ , and  $X$  is the unique compact set satisfying this equation. The cookie-cutter set we work with is naturally associated with a dyadic tree, but up to technical modifications, the rest of the paper is easily adapted to cookie-cutter sets constructed with more than two contractions. Figure 1 illustrates the first steps of the construction of  $X$ .

For any finite word  $w \in \Sigma^*$ , we introduce the sets

$$X_w = g_{w_1} \circ \dots \circ g_{w_{|w|}}(X).$$

There is a natural identification between  $\Sigma$  and  $X$  via the homeomorphism

$$\begin{aligned} \pi : \Sigma &\rightarrow X \\ w &\mapsto \lim_{n \rightarrow \infty} g_{w|1} \circ \dots \circ g_{w|n}(0). \end{aligned}$$

The dynamical systems  $(\Sigma, \sigma)$  and  $(X, T)$  are thus topologically conjugate.

In order to define a Gibbs measure on the cookie-cutter set  $X$ , we eventually need a Hölder continuous function  $\varphi : X \rightarrow \mathbb{R}$ . As usual, for  $x \in X$  and  $n \geq 1$ ,  $S_n \varphi(x)$  stands for  $\sum_{k=0}^{n-1} \varphi(T^k(x))$ , the  $n$ th Birkhoff sum of  $\varphi$  at  $x$ .




FIGURE 1. Left: the mapping  $T : U_0 \cup U_1 \rightarrow [0, 1]$ ; right: first three steps of the construction of the cookie-cutter set.

We recall the definition of the topological pressure and Gibbs measure associated with  $\varphi$ .

*Definition 1.1.* The topological pressure  $P(\varphi)$  of  $\varphi$  is the unique real number such that for some constant  $C \geq 1$  we have

$$\text{for all } n \in \mathbb{N}, C^{-1} \exp(nP(\varphi)) \leq \sum_{w \in \Sigma_n} \sup_{x \in X_w} \exp(S_n \varphi(x)) \leq C \exp(nP(\varphi)). \quad (1.1)$$

The Gibbs measure  $\mu_\varphi$  on  $(X, T)$  associated with  $\varphi$  is the unique ergodic measure such that for some constant  $C > 0$

$$\text{for all } w \in \Sigma^*, \text{ for all } x \in X_w, C^{-1} \leq \frac{\mu_\varphi(X_w)}{\exp(S_{|w|} \varphi(x) - |w|P(\varphi))} \leq C. \quad (1.2)$$

This measure  $\mu_\varphi$  has a unique extension to a probability Borel measure on  $[0, 1]$  that we also denote by  $\mu_\varphi$ .

The existence of the topological pressure as well as the existence and uniqueness of  $\mu_\varphi$  are parts of the thermodynamic formalism theory. Proofs of these assertions can be found in [9]. We can assume that  $P(\varphi) = 0$ . Clearly the latter can always be achieved by replacing  $\varphi$  by  $\varphi - P(\varphi)$ , if necessary.

The next proposition states some classical results for the topological pressure (see [13, 24] for proofs and related references). Let us first introduce, for  $(q, t) \in \mathbb{R}^2$ , the mappings

$$\psi = -\log |T'| : X \rightarrow \mathbb{R} \quad \text{and} \quad \psi_{q,t} = q\psi - t\varphi : X \rightarrow \mathbb{R}. \quad (1.3)$$

**PROPOSITION 1.2.** *We recall the following properties.*

- (1) *The mapping  $(q, t) \mapsto P(q\psi - t\varphi)$  is strictly decreasing in  $q$  and strictly increasing in  $t$ .*
- (2) *For every  $q \in \mathbb{R}$ , there exists a unique  $\theta(q) \in \mathbb{R}$  such that  $P(\psi_{q,\theta(q)}) = P(q\psi - \theta(q)\varphi) = 0$ .*
- (3) *The mapping  $q \mapsto \theta(q)$  is analytic, montone and concave.*

Let us now introduce the inverse measure of  $\mu_\varphi$ , which is the focus of the rest of the paper. Let  $F_m$  denote the distribution function of a probability measure  $m$  on  $[0, 1]$ : for  $x \in [0, 1]$ ,  $F_m(x) = m([0, x])$ .

*Definition 1.3.* The inverse measure  $\nu$  of  $\mu_\varphi$  is the unique Borel probability measure on  $[0, 1]$  such that

$$\text{for all } x \in [0, 1], \quad F_\nu(x) = \sup\{t \in [0, 1] \mid F_{\mu_\varphi}(t) \leq x\}. \quad (1.4)$$

Investigations of inverse measures in terms of multifractals were first carried out in [19], where the authors point out the interest of such studies from the multifractal viewpoint.

Our aim is to study the local scaling properties of  $\nu$ . Let us recall how the local regularity of any positive Borel measure  $m$  on  $[0, 1]$  is described. For  $x \in [0, 1]$ , the pointwise Hölder exponent of  $m$  at  $x$  is given by

$$h_m(x) = \liminf_{r \rightarrow 0^+} \frac{\log m(B(x, r))}{\log(r)}.$$

Performing the multifractal analysis of  $m$  consists in computing the Hausdorff dimensions of the level sets of  $h_m$ , i.e. the sets

$$E_m(h) = \{x \in [0, 1] \mid h_m(x) = h\}, \quad h \geq 0.$$

We denote the singularity spectrum of  $m$  by  $d_m(h) = \dim E_m(h)$ , where  $\dim$  stands for the Hausdorff dimension (see, for instance, [13] for the definition of the dimension), and  $\dim \emptyset = -\infty$  by convention. The knowledge of the singularity spectrum provides us with information about the geometric distribution of the singularities of the measure  $m$ .

Recall that if  $g$  is a function from  $\mathbb{R}$  to  $\mathbb{R} \cup \{-\infty\}$ , then its Legendre transform  $g^*$  is defined for any  $\alpha \in \mathbb{R}$  by

$$g^*(\alpha) = \inf_{q \in \mathbb{R}} (\alpha q - g(q)) \in \mathbb{R} \cup \{-\infty\}. \quad (1.5)$$

In [24], the multifractal analysis of  $\mu_\varphi$  is performed. Recall that for any positive Borel measure  $\mu$  on  $[0, 1]$ , the  $L^q$ -spectrum  $\tau_\mu : \mathbb{R} \rightarrow \mathbb{R} \cup \{-\infty\}$  is given by

$$\tau_\mu(q) = \liminf_{r \rightarrow 0} \frac{\log \sup\{\sum_i \mu(B_i)^q\}}{\log(r)},$$

the supremum being taken over all families of disjoint closed intervals  $B_i$  of radius  $r$  with centers in  $\text{supp}(\mu)$  (see [8, 10, 20, 21, 23] for definitions and a study of multifractal formalisms).

With our notation, the result of [24] is the following.

**THEOREM 1.4.** *For every  $q \in \mathbb{R}$ ,  $\tau_{\mu_\varphi}(q) = -\theta^{-1}(-q)$ , and the measure  $\mu_\varphi$  satisfies the multifractal formalism, i.e. for every  $h \geq 0$ ,  $d_{\mu_\varphi}(h) = \tau_{\mu_\varphi}^*(h)$ .*

*The singularity spectrum of  $\mu_\varphi$  can thus be written as*

$$d_{\mu_\varphi}(h) = \inf\{qh + \theta^{-1}(-q) \mid q \in \mathbb{R}\}. \quad (1.6)$$

The main goal of this paper is to derive similar multifractal statements for the inverse measure  $\nu$  of  $\mu_\varphi$ . These main results are summarized in the following theorem.

**THEOREM 1.5.** *Let  $X$  be a cookie-cutter set, and let  $\mu_\varphi$  be a Gibbs measure on  $X$  associated with a Hölder continuous potential  $\varphi$ .*

*Let  $\nu$  be the inverse measure of  $\mu_\varphi$ .*

- (1) *The measure  $\nu$  is discrete (i.e. it can be written as an infinite sum of Dirac masses), and its singularity spectrum  $d_\nu$  satisfies:*
  - *for all  $h \in [0, \theta'(\dim X)]$ ,  $d_\nu(h) = h \cdot \dim X$ ;*
  - *for all  $h > \theta'(\dim X)$  such that  $\theta^*(h) > 0$ ,  $d_\nu(h) = \theta^*(h)$ ;**In addition,  $E_\nu(h) = \emptyset$  for all  $h > \theta'(\dim X)$  such that  $\theta^*(h) < 0$ .*
- (2) *The  $L^q$ -spectrum of  $\nu$  is given by*

$$\text{for every } q \in \mathbb{R}, \quad \tau_\nu(q) = \min(\theta(q), 0).$$

- (3) *The measure  $\nu$  obeys the multifractal formalism: for every  $h > 0$ , such that  $\tau_\nu^*(h) > 0$ , one has  $d_\nu(h) = \tau_\nu^*(h)$ .*

Let  $h_{\max} := \sup\{h \mid \tau_\nu^*(h) \geq 0\}$ . The attentive reader has noticed that Theorem 1.5 does not include the case  $\tau_\nu^*(h_{\max}) = 0$ , i.e. when the singularity spectrum touches zero at the right end of its decreasing part. We discuss this point in §4.5.

Note that there is an interesting relationship between the singularity spectrum of  $\mu_\varphi$  and the spectrum of  $\nu$ . This can be expressed in terms of their Legendre transforms:

$$\tau_\nu(q) = -(\tau_{\mu_\varphi})^{-1}(-q) \quad \text{when } q \leq \dim X. \tag{1.7}$$

The main difficulty here is that the measure  $\nu$  is constituted only by Dirac masses. Subsequently, the multifractal analysis of  $\nu$  is not entirely based on the study of auxiliary Gibbs measures as in the case of  $\mu_\varphi$  (see [24]). This measure  $\nu$  is an example of discrete measures whose multifractal analysis requires other mathematical tools, namely ubiquity results (see §3.3).

The singularity spectrum of  $\nu$  is composed of two parts: a linear part with slope  $\dim X$ , and a strictly concave part. The linear part is essentially due to the presence of Dirac masses, and the concave part follows from standard multifractal arguments involving auxiliary (diffuse) Gibbs measures. As claimed above, the  $L^q$ -spectrum of  $\nu$  possesses a point of non-differentiability as soon as the graph of the function  $q \mapsto \theta(q)$  does not have a horizontal tangent when it reaches zero (see Figure 2).

Other examples of infinite homogeneous [1, 14, 17] or heterogeneous [3, 5] sums of Dirac masses have been studied. Note that the measure  $\nu$  does not belong to the class of discrete measures considered in these papers, except when  $\mu_\varphi$  is the measure of maximal entropy associated with a system where the mappings  $g_0$  and  $g_1$  are affine maps with same contraction ratio. Moreover, compared with the measures appearing in [1, 3, 5, 14, 17], an additional important property is that  $\nu$  is naturally associated with a dynamical system.

2. *A reformulation of the problem: some remarks*

2.1. *Identifying the intensities of the Dirac masses of  $\nu$ .* We assume that  $\sup U_0 < \inf U_1$ . For  $w \in \Sigma^*$  (including  $w = \emptyset$ ) we introduce the real numbers

$$\begin{aligned} m_w^0 &= \min X_{w \cdot 0} = \pi(w \cdot \bar{0}) \quad \text{and} \quad M_w^0 = \max X_{w \cdot 0} = \pi(w \cdot 0 \cdot \bar{1}), \\ m_w^1 &= \min X_{w \cdot 1} = \pi(w \cdot 1 \cdot \bar{0}) \quad \text{and} \quad M_w^1 = \max X_{w \cdot 1} = \pi(w \cdot \bar{1}). \end{aligned}$$

Obviously, for every integer  $n \geq 1$ ,

$$X \subset \bigcup_{w \in \Sigma_n} [m_w^0, M_w^0] \cup [m_w^1, M_w^1].$$

Let us also introduce the mapping

$$\begin{aligned} F_\varphi : X &\rightarrow [0, 1] \\ t &\mapsto \mu_\varphi([0, t]). \end{aligned}$$

For every  $w \in \Sigma^*$ , let

$$x_w = F_\varphi(M_w^0) \quad (\text{this includes the case } x_\emptyset = F_\varphi(\max(X_{0,1}))). \tag{2.1}$$

We prove that  $\nu$  is a discrete measure whose Dirac masses are located at the family of points  $x_w$ , where  $w$  ranges in  $\Sigma^*$ . Figure 3 illustrates the construction of the inverse measure  $\nu$ .

PROPOSITION 2.1. *With the notation above, we have*

$$\nu = m_\emptyset^0 \cdot \delta_0 + \sum_{w \in \Sigma^*} (m_w^1 - M_w^0) \cdot \delta_{x_w} + (1 - M_\emptyset^1) \cdot \delta_1. \tag{2.2}$$

*Proof.* Denote the right-hand side of (2.2) by  $\rho$ .

For every  $w \in \Sigma^* \setminus \{\emptyset\}$ ,  $m_w^1 - M_w^0$  represents the length of the interval located between  $X_{w \cdot 0}$  and  $X_{w \cdot 1}$ . Recall that for every  $x \in [0, 1]$ ,  $\nu([0, x]) = \inf\{t \mid \mu_\varphi([0, t]) > x\}$ . Hence, if  $x < x_w < y$ , then  $\nu([0, y]) - \nu([0, x]) \geq m_w^1 - M_w^0$ . Since this holds for every  $(x, y)$  for which  $x < x_w < y$ , we deduce that  $\nu$  has a Dirac mass with weight greater than  $m_w^1 - M_w^0$  at  $x_w$ . The same arguments also work for the Dirac masses at zero,  $x_\emptyset$  and at one.

Subsequently,  $\nu - \rho$  is still a positive Borel measure on  $[0, 1]$ .

Moreover,  $\rho([0, 1]) = m_\emptyset^0 + \sum_{w \in \Sigma^*} (m_w^1 - M_w^0) + (1 - M_\emptyset^1) = \max X + 1 - M_\emptyset^1 = 1$ , and by construction  $\nu([0, 1])$  is also equal to one. We conclude that  $\nu = \rho$ .  $\square$

2.2. *A tractable reformulation of (2.2).* If  $w \in \Sigma^*$ , then we denote

$$I_w = F_\varphi(X_w) \setminus F_\varphi(\max X_w). \tag{2.3}$$

By construction,  $I_w$  is an interval, since the support of  $\mu_\varphi$  (the derivative of  $F_\varphi$ ) restricted to the interval  $[\min(X_w), \max(X_w)]$  is  $X_w$ , and  $\mu_\varphi$  is atomless. The families of intervals  $\mathcal{F}_n = \{I_w\}_{w \in \Sigma_n}$ ,  $n \geq 1$ , form a nested grid of  $[0, 1]$ , since  $\text{supp}(\mu_\varphi) \subset X$  and  $\bigcup_{w \in \Sigma_n} X_w = X$ .

Now we make full use of the structure of the construction in order to slightly transform  $\nu$  into a measure enjoying the same multifractal nature as  $\nu$ . This will make the calculations easier. For this, note that for  $w \in \Sigma^*$  we have

$$[M_\emptyset^0, m_\emptyset^1] = T^{|w|}([M_w^0, m_w^1]). \tag{2.4}$$

Here  $T^{|w|}$  is the  $|w|$ th iterate of  $T$ . Owing to the definition (1.3), for  $w \in \Sigma_n$  and  $x \in X_w$  we have  $S_n \psi(x) = -\log |(T^n)'(x)|$ .




FIGURE 2. Top left: a typical  $L^q$ -spectrum for a Gibbs measure  $\mu_\varphi$  on a cookie-cutter set  $X$ ; top right: the typical singularity spectrum of  $\mu_\varphi$ ; bottom left: a typical  $L^q$ -spectrum for the inverse measure  $\nu$ ; bottom right: the typical singularity spectrum of  $\nu$ . The two graphs on the left-hand side are related via the formula (1.7). The graphs on the right-hand side are the Legendre transforms of the graphs on the left-hand side.




FIGURE 3. Top left: the integral of a Gibbs measure  $\mu_\varphi$  associated with a cookie-cutter set; top right: the inverse of the integral of  $\mu_\varphi$ ; bottom: the discrete measure  $\nu$ . The first two graphs are symmetric with respect to the first diagonal.

We often use the standard bounded distortion principle, for which we refer the interested reader to [24] or to [13, Ch. 4, Propositions 4.1 and 4.2]. This principle yields a control, uniform in  $n$ , of the variations of the Birkhoff sums  $S_n\psi(x)$  inside each  $X_w$ ,  $w \in \Sigma_n$ . Specifically, there exists a constant  $C > 1$  such that, for every  $w \in \Sigma^*$ , for every  $(x, y) \in (X_w)^2$ ,  $C^{-1} \leq |X_w| \cdot \exp(-S_n\psi(x)) \leq C$  and

$$C^{-1}|T^{|w|}(x) - T^{|w|}(y)| \leq \frac{|x - y|}{|X_w|} \leq C|T^{|w|}(x) - T^{|w|}(y)|. \tag{2.5}$$

Let us introduce the mapping  $\tilde{\psi} : \Sigma^* \rightarrow \mathbb{R}$

$$\tilde{\psi}(w) = \sup_{x \in X_w} S_{|w|}\psi(x). \tag{2.6}$$

This yields

$$C^{-1} \leq |X_w| \cdot \exp(\tilde{\psi}(w)) \leq C. \tag{2.7}$$

Applying (2.5) to  $x = M_w^0$  and  $y = m_w^1$ , and using (2.4), we obtain that there is a constant  $C > 0$  such that, for all  $w \in \Sigma^*$ ,

$$C^{-1}(m_\emptyset^1 - M_\emptyset^0) \leq \frac{m_w^1 - M_w^0}{\exp(\tilde{\psi}(w))} \leq C(m_\emptyset^1 - M_\emptyset^0). \tag{2.8}$$

Since  $\psi$  is Hölder ( $T$  is supposed to belong to  $C^{1+\gamma}$ ), we can consider  $\mu_\psi$ , the Gibbs measure on  $X$  associated with  $\psi$ . Then, (1.2) holds for  $(\psi, \mu_\psi)$  instead of  $(\varphi, \mu_\varphi)$ . There exists another constant  $C > 0$  such that for all  $w \in \Sigma^*$ , for every  $x \in X_w$ ,

$$C^{-1} \leq \frac{\mu_\psi(X_w)}{\exp(S_{|w|}\psi(x) - |w|P(\psi))} \leq C.$$

This implies that, for all  $w \in \Sigma^*$  (using our definition for  $\tilde{\psi}(w)$ ),

$$C^{-1} \leq \frac{\exp(|w|P(\psi))\mu_\psi(X_w)}{\exp(\tilde{\psi}(w))} \leq C.$$

Combining (2.8) and the last estimate, we see that there is a constant  $C > 0$  such that, for every  $w \in \Sigma^*$ ,

$$C^{-1} \leq \frac{m_w^1 - M_w^0}{\exp(|w|P(\psi))\mu_\psi(X_w)} \leq C.$$

With  $\mu_\psi$  can be associated the unique Borel measure  $\mu_{\psi,\varphi}$  on  $[0, 1]$  such that  $\mu_{\psi,\varphi}(I_w) = \mu_\psi(X_w)$ , that is to say  $\mu_{\psi,\varphi} = \mu_\psi \circ F_\varphi^{-1}$ . The measure  $\mu_{\psi,\varphi}$  has its support equal to  $[0, 1]$ , while the support of  $\mu_\psi$  is equal to  $X$ . The last inequalities can thus be rewritten

$$C^{-1} \leq \frac{m_w^1 - M_w^0}{\exp(|w|P(\psi))\mu_{\psi,\varphi}(I_w)} \leq C. \tag{2.9}$$

Remembering (2.2), we conclude that the discrete measure  $\nu$  is equivalent to any of the following discrete measures:

$$\nu_1 = \sum_{w \in \Sigma^*} \exp(\tilde{\psi}(w)) \cdot \delta_{x_w} + m \cdot \delta_0 + (1 - M) \cdot \delta_1, \tag{2.10}$$

$$\nu_2 = \sum_{w \in \Sigma^*} \exp(|w|P(\psi))\mu_{\psi,\varphi}(I_w) \cdot \delta_{x_w} + m \cdot \delta_0 + (1 - M) \cdot \delta_1. \tag{2.11}$$



It appears that  $\nu_1$  and  $\nu_2$  are more convenient to work with than (2.2). We are going to perform the multifractal analysis of these measures, that we again denote by  $\nu$  for ease of notation.

3. Some definitions and tools for the multifractal analysis

If  $m$  is a positive Borel measure on  $\mathbb{R}$ , then the lower Hausdorff dimension of  $m$  is defined as

$$\dim(m) = \inf\{\dim E \mid m(E) > 0\}.$$

We refer the reader to [13] for more details and for the definition of Hausdorff measures and dimension.

3.1. Definitions of exponents, singularity spectrum and approximation degree.

Definition 3.1. For  $w \in \Sigma^*$ , we set

$$\begin{aligned} \lambda_w &= 2|I_w| (= 2\mu_\varphi(X_w)), \\ \chi(w) &= \log |I_w|, \\ \tilde{\alpha}(w) &= \frac{\tilde{\psi}(w)}{\chi(w)} \quad \text{when } |w| \geq 1. \end{aligned}$$

From (2.7), there exist  $0 < \alpha_0 \leq \alpha_1 < \infty$  such that

$$\text{for all } w \in \Sigma^*, \quad \alpha_0 \leq \tilde{\alpha}(w) \leq \alpha_1. \tag{3.1}$$

Definition 3.2. For  $x \in [0, 1)$  and  $n \geq 1$ ,  $w_n(x)$  stands for the unique element  $w$  of  $\Sigma_n$  such that  $x \in I_w$ . One then defines

$$\alpha_n(x) = \tilde{\alpha}(w_n(x)) = \frac{\tilde{\psi}(w_n(x))}{\log |I_{w_n(x)}|} \quad \text{and} \quad \alpha(x) = \liminf_{n \rightarrow \infty} \alpha_n(x).$$

Let us now focus on the family  $\{(x_w, \lambda_w)\}_{w \in \Sigma}$ . By construction (recall formula (2.1)), for every  $n \geq 1$ , we have  $[0, 1] \subset \bigcup_{w \in \Sigma_n} B(x_w, \lambda_w/2)$ . Hence, any real number  $x \in [0, 1]$  is covered by infinitely many intervals  $B(x_w, \lambda_w/2)$ . The family  $\{(x_w, \lambda_w)\}_{w \in \Sigma}$  is referred to as a *ubiquitous system*.

Given a ubiquitous system, two approximation degrees of any real number  $x \in [0, 1]$  by the family  $\{(x_w, \lambda_w)\}_{w \in \Sigma}$  can be defined as follows.

Definition 3.3. For  $x \in (0, 1) \setminus \{x_w \mid w \in \Sigma^*\}$ , the approximation degrees  $\xi_x$  and  $\tilde{\xi}_x$  of  $x$  by the ubiquitous system  $\{(x_w, \lambda_w)\}_{w \in \Sigma^*}$  are defined as

$$\xi_x = \limsup_{n \rightarrow \infty} \left( \sup_{w \in \Sigma_n} \frac{\log |x - x_w|}{\log \lambda_w} \right) \quad \text{and} \quad \tilde{\xi}_x = \limsup_{n \rightarrow \infty} \frac{\log |x - x_{w_n(x)}|}{\log \lambda_{w_n(x)}}.$$

Owing to the covering property of the ubiquitous system  $\{(x_w, \lambda_w)\}_{w \in \Sigma}$ , one obviously has  $\xi_x \geq \tilde{\xi}_x \geq 1$ , for every  $x \in [0, 1]$ .

3.2. *Auxiliary Gibbs measures.* Recall the definitions of  $\psi$ ,  $\psi_{q,t}$  and  $\theta$  given in Proposition 1.2.

*Definition 3.4.* For every  $q \in \mathbb{R}$ , we set  $\mu_q = \mu_{\psi_{q,\theta(q)},\varphi} := \mu_{\psi_{q,\theta(q)}} \circ F_\varphi^{-1}$ .

PROPOSITION 3.5. *For each  $q \in \mathbb{R}$ , the following hold.*

- (1) *The set  $\{x \in (0, 1) \mid \alpha(x) = \theta'(q)\}$  is of full  $\mu_q$ -measure.*
- (2) *We have  $\dim(\mu_q) \geq \theta^*(\theta'(q))$ .*

The second statement in the proposition implies that if a set  $E$  has a Hausdorff dimension less than  $\theta^*(\theta'(q))$ , then  $\mu_q(E) = 0$ . This is useful in Proposition 4.9, where we derive lower bounds for the Hausdorff dimensions of the level sets  $E_\nu(h)$ .

*Proof.* By construction of  $\theta(q)$ ,  $P(\psi_{q,\theta(q)}) = 0$ . Hence, there exists  $C_q > 0$  such that for all  $w \in \Sigma^*$ , for every  $x \in X_w$ ,

$$C_q^{-1} \leq \frac{\mu_{\psi_{q,\theta(q)}}(X_w)}{\exp(S_{|w|}\psi_{q,\theta(q)}(x))} = \frac{\mu_{\psi_{q,\theta(q)}}(X_w)}{\exp(qS_{|w|}\psi(x) - \theta(q)S_{|w|}\varphi(x))} \leq C_q.$$

Consequently, using (2.3), (2.6) and the fact that  $\mu_{\psi_{q,\theta(q)}}(X_w) = \mu_q(I_w)$ , it follows for  $C_q > 0$  some further constant that

$$C_q^{-1} \leq \frac{\mu_q(I_w)}{\exp(q\tilde{\psi}(w))|I_w|^{-\theta(q)}} \leq C_q, \quad \text{for all } w \in \Sigma^* \tag{3.2}$$

Let  $n \geq 1$  and  $\varepsilon > 0$ . We prove that  $\{x \in (0, 1) \mid \alpha(x) = \theta'(q)\}$  is of full  $\mu_q$ -measure. Let us set  $\alpha = \theta'(q)$ , and let  $\delta > 0$ . Let  $K_n$  be defined by  $K_n = \{x \in (0, 1) \mid \tilde{\psi}(w_n(x)) \geq (\alpha - \varepsilon) \log |I_{w_n(x)}|\}$ . We have

$$\begin{aligned} \mu_q(K_n) &= \mu_q(\{x \in (0, 1) : \exp(\delta\tilde{\psi}(w_n(x)))|I_{w_n(x)}|^{-\delta(\alpha-\varepsilon)} \geq 1\}) \\ &\leq \int_0^1 \exp(\delta\tilde{\psi}(w_n(x)))|I_{w_n(x)}|^{-\delta(\alpha-\varepsilon)} d\mu_q(x) \\ &= \sum_{w \in \Sigma_n} \exp(\delta\tilde{\psi}(w))|I_w|^{-\delta(\alpha-\varepsilon)} \mu_q(I_w) \\ &\leq C_q \sum_{w \in \Sigma_n} \exp((\delta + q)\tilde{\psi}(w))|I_w|^{-(\theta(q)+\delta(\alpha-\varepsilon))} \\ &\leq C'_q \sum_{w \in \Sigma_n} \sup_{x \in X_w} \exp((\delta + q)S_n\psi(x) - (\theta(q) + \delta(\alpha - \varepsilon))S_n\varphi(x)) \\ &\leq C''_q \exp(n\gamma_q), \end{aligned}$$

where  $C'_q, C''_q > 0$  are constants, and  $\gamma_q = P((q + \delta)\psi - (\theta(q) + \delta(\alpha - \varepsilon))\varphi)$ . Now, since the function  $\theta$  is right-differentiable at  $q$  and  $\alpha = \theta'(q)$ , for  $\delta$  small enough we have  $\theta(q) + \delta(\alpha - \varepsilon) < \theta(q + \delta)$ . Consequently, applying Proposition 1.2(1), since by Proposition 1.2(2)  $P((\delta + q)\psi - \theta(q + \delta)\varphi) = 0$ , we obtain  $\gamma_q < 0$ . Therefore, using the Borel–Cantelli lemma, it follows that  $\mu_q(\limsup_{n \rightarrow +\infty} K_n) = 0$ . Hence, for  $\mu_q$ -almost all  $x \in [0, 1]$ , we have that  $\tilde{\psi}(w_n(x)) < (\alpha - \varepsilon) \log |I_{w_n(x)}|$  for  $n$  large enough. This holds for any  $\varepsilon > 0$ , and thus we have that  $\mu_q$ -almost every  $x \in [0, 1]$  verifies  $\alpha(x) \geq \alpha$ .

The converse inequality is obtained using a similar argument, and this is left to the reader. This shows that  $\mu_q(\{x \mid \alpha(x) = \theta'(q)\}) = 1$ , and the first part of the proposition follows. □

Note that by the bounded distortion principle, the ratio  $|I_{w_n(x)}|/|I_{w_{n+1}(x)}|$  is bounded from below and above by constants independent of  $n \geq 1$  and  $x \in [0, 1]$ . To obtain the second part of the proposition, we only have to prove that

$$\mu_q\text{-almost everywhere, } \liminf_{n \rightarrow +\infty} \frac{\log \mu_q(I_{w_n(x)})}{\log |I_{w_n(x)}|} \geq \theta^*(\theta'(q)).$$

Indeed, in this case the *mass distribution principle* [13] allows us to conclude that  $\dim \mu_q \geq \theta^*(\theta(q))$ .

For this, note that (3.2) implies

$$\left| \frac{\log \mu_q(I_{w_n(x)})}{\log |I_{w_n(x)}|} - q \frac{\tilde{\psi}(w_n(x))}{\log |I_{w_n(x)}|} + \theta(q) \right| \leq \frac{\log C_q}{|\log(|I_{w_n(x)}|)|}.$$

For  $\mu_q$ -almost every  $x$ , we proved before that

$$\lim_{n \rightarrow +\infty} \frac{\tilde{\psi}(w_n(x))}{\log |I_{w_n(x)}|} = \alpha.$$

Since  $\alpha = \theta'(q)$ , we obtain that

$$\lim_{n \rightarrow +\infty} \frac{\log \mu_q(I_{w_n(x)})}{\log |I_{w_n(x)}|} = q\alpha - \theta(q) = q\theta'(q) - \theta(q).$$

By definition of the Legendre transform, we have that  $q\theta'(q) - \theta(q) = \theta^*(\theta'(q))$  and, hence, the result follows.

3.3. *An heterogeneous ubiquity theorem.* Let  $q_0 := \dim X$  denote the unique solution of  $\theta(q_0) = 0$ . Let  $\tilde{\varepsilon} = (\varepsilon_n)_{n \geq 1}$  be a sequence of positive numbers which tend to 0 for  $n$  tending to infinity. For  $\xi \geq 1$  we then introduce the limsup set

$$S(\xi, \tilde{\varepsilon}) = \bigcap_{N \geq 1} \bigcup_{n \geq N} \bigcup_{\substack{w \in \Sigma_n: \\ \theta'(q_0) - \varepsilon_n \leq \alpha(w) \leq \theta'(q_0) + \varepsilon_n}} B(x_w, \lambda_w^\xi). \tag{3.3}$$

Using the definition of  $\alpha(w)$  and the measure  $\mu_{q_0}$  (for which there exists a constant  $C_{q_0} > 0$  such that  $C_{q_0}^{-1}|X_w|^{1/q_0} \leq \mu_{q_0}(I_w) \leq C_{q_0}|X_w|^{1/q_0}$  for every  $w \in \Sigma^*$ ), one can rewrite  $S(\xi, \tilde{\varepsilon})$  as

$$S(\xi, \tilde{\varepsilon}) = \bigcap_{N \geq 1} \bigcup_{n \geq N} \bigcup_{\substack{w \in \Sigma_n: \\ |I_w|^{q_0(\theta'(q_0) + \varepsilon_n)} \leq \mu_{q_0}(B(x_w, \lambda_w)) \leq |I_w|^{q_0(\theta'(q_0) - \varepsilon_n)}}} B(x_w, \lambda_w^\xi), \tag{3.4}$$

where the sequence  $\tilde{\varepsilon}$  has been slightly modified to take into account the constant  $C_{q_0}$ .

Limsup sets of the form (3.4) arise in *ubiquity* theory. We present here a short version of ubiquity and heterogeneous ubiquity theorems, adapted to our context. Given  $(x_n)_{n \geq 1}$  a sequence of real numbers in  $[0, 1]$ , and  $(\lambda_n)_{n \geq 1}$  a sequence of positive real numbers tending to zero when  $n$  tends to infinity, the classical ubiquity results are concerned with the computations of limsup sets of the form

$$A(\xi) = \bigcap_{N \geq 1} \bigcup_{n \geq N} B(x_n, \lambda_n^\xi),$$

where  $\xi > 1$  is a contraction exponent. The classical ubiquity theorems, proved in [11, 18], state the following: as soon as the set  $A(1)$  is of full Lebesgue measure, we have  $\dim A(\xi) \geq 1/\xi$  for every  $\xi \geq 1$ .

Now suppose that  $\{(x_n, \lambda_n)\}_{n \geq 1} = \{(x_w, \lambda_w)\}_{w \in \Sigma^*}$ . Although the sets  $S(\xi, \tilde{\varepsilon})$  defined in (3.3) are very similar to the sets  $A(\xi)$ , the Hausdorff dimension of the sets  $S(\xi, \tilde{\varepsilon})$  cannot be reached by such theorems. This is due to the fact that the measure playing a key role is not the Lebesgue measure any more, but the monodimensional measure  $\mu_{q_0}$ .

The notion of heterogeneous ubiquity, introduced in [2–4], has been developed to estimate the Hausdorff dimension of the limsup sets  $S(\xi, \tilde{\varepsilon})$  (and other limsup sets of similar form). This is a crucial point in the following, since such limsup sets arise naturally when performing the multifractal analysis of the inverse measure  $\nu$  of  $\mu_\varphi$ . Proposition 3.6 has been obtained in [4, Theorem 2.2] (case  $\rho = 1$ ).

We now give the assumptions needed to obtain sharp lower bounds for the Hausdorff dimension of  $S(\xi, \tilde{\varepsilon})$ . In order to apply [4, Theorem 2.2], three properties are required on the measure  $\mu_{q_0}$  and the system  $\{(x_w, \lambda_w)\}_{w \in \Sigma^*}$ :

- The measure  $\mu_{q_0}$  is quasi-Bernoulli [10] with respect to the grid constituted by the intervals  $\{I_w\}_{w \in \Sigma^*}$ , i.e. there is a constant  $C > 1$  such that for every word  $w, w' \in \Sigma^*$ ,

$$\begin{cases} C^{-1} \mu_{q_0}(I_w) \mu_{q_0}(I_{w'}) \leq \mu_{q_0}(I_{ww'}) \leq C \mu_{q_0}(I_w) \mu_{q_0}(I_{w'}) \\ C^{-1} |I_w| \cdot |I_{w'}| \leq |I_{ww'}| \leq C |I_w| \cdot |I_{w'}|. \end{cases}$$

- The measure  $\mu_{q_0}$  is monofractal, in the sense that

$$\mu_{q_0}\text{-almost everywhere, } \lim_{n \rightarrow +\infty} \frac{\log \mu_{q_0}(I_{w_n(x)})}{\log |I_{w_n(x)}|} = \theta^*(\theta'(q_0)) = q_0 \theta'(q_0). \quad (3.5)$$

Note that since  $\theta(q_0) = 0$ , we have  $\theta^*(\theta'(q_0)) = q_0 \theta'(q_0)$ .

- We have

$$\mu_{q_0} \left( \limsup_{n \rightarrow +\infty} \{B(x_w, \lambda_w/2) \mid w \in \Sigma_n\} \right) = 1,$$

since, by construction, the system  $\{(x_w, \lambda_w)\}_{w \in \Sigma^*}$  enjoys the covering property:  $\limsup_{n \rightarrow +\infty} \{B(x_w, \lambda_w/2) \mid w \in \Sigma_n\} = [0, 1]$ .

Then, according to the definitions of [2, 4], the system of points  $\{(x_w, \lambda_w)\}_{w \in \Sigma^*}$  forms an heterogeneous ubiquitous system with respect to  $\mu_{q_0}$  and its almost sure Hölder exponent  $\theta^*(\theta'(q_0))$ .

In [4, Theorem 2.2] it is stated that, as soon as  $\mu_{q_0}(A(1)) = \|\mu_{q_0}\|$ , there exists  $\tilde{\varepsilon}$  such that  $\dim S(\xi, \tilde{\varepsilon}) \geq \dim(\mu_{q_0})/\xi$  for every  $\xi \geq 1$ . In our context, this yields the following proposition.

**PROPOSITION 3.6.** *There is a positive sequence  $\tilde{\varepsilon}$  converging to zero at  $\infty$  such that, for every  $\xi \geq 1$ , there exists a positive Borel measure  $m_\xi$  such that  $m_\xi(S(\xi, \tilde{\varepsilon})) > 0$  and  $\dim(m_\xi) \geq (q_0 \theta'(q_0))/\xi$ .*

More on heterogeneous ubiquity can be found in [4, 6, 12].

4. Proof of Theorem 1.5

4.1. A first result on the local regularity analysis of  $\nu$ . For the following proposition the reader might like to recall Definition 3.3.

PROPOSITION 4.1. Let  $x \in [0, 1]$ . If  $x \in \{x_w \mid w \in \Sigma^*\}$ , then  $h_\nu(x) = 0$ , otherwise

$$\frac{\alpha(x)}{\xi_x} \leq \frac{\alpha(x)}{\tilde{\xi}_x} \leq h_\nu(x) \leq \alpha(x).$$

Here, if  $\xi_x = +\infty$  then  $\alpha(x)/\xi(x) := 0$ . Likewise if  $\tilde{\xi}_x = +\infty$ , then  $\alpha(x)/\tilde{\xi}(x) := 0$ .

Proof. Let  $x \in [0, 1] \setminus \{x_w \mid w \in \Sigma^*\}$  and  $r > 0$ .

- Obviously,  $\xi_x \geq \tilde{\xi}_x$ , so the left inequality is trivial.
- Let  $n_{x,r} = \min\{n \mid \exists w \in \Sigma_n, x_w \in B(x, r)\}$ . Let  $w(x, r) = w_{n_{x,r}}(x)$  be the unique word of  $\Sigma_{n_{x,r}}$  such that  $x_{w(x,r)} \in B(x, r)$ . Indeed, if two such words  $w$  and  $w'$  exist (without loss of generality we can assume that  $x_w < x_{w'}$ ), then by construction there is another word  $w''$  such that  $|w''| < n_{x,r}$  and  $x_w < x_{w''} < x_{w'}$ . This contradicts the minimality of  $n_{x,r}$ . This implies that, using (2.10),  $\nu(B(x, r)) \geq \exp(\tilde{\psi}(w(x, r)))$ .

To find an upper bound for  $\nu(B(x, r))$ , we use the form (2.11) of  $\nu$ . Note that the minimality of  $n_{x,r}$  yields that we necessarily have

$$x_{w(x,r)} \in B(x, r) \subset I_{w(x,r)}. \tag{4.1}$$

Hence, for every  $n > n_{x,r}$ , we obtain

$$\sum_{v \in \Sigma_n: x_v \in B(x,r)} \exp(|v|P(\psi))\mu_{\psi,\varphi}(I_v) \leq \exp(nP(\psi))\mu_{\psi,\varphi}(I_{w(x,r)}).$$

Since  $P(\psi) < 0$ , we deduce combining (2.8) and (2.9) that

$$\begin{aligned} \nu(B(x, r)) &\leq \exp(\tilde{\psi}(w(x, r))) + \left( \sum_{n > n_{x,r}} \exp(nP(\psi)) \right) \mu_{\psi,\varphi}(I_{w(x,r)}) \\ &\leq C \exp(\tilde{\psi}(w(x, r))). \end{aligned}$$

Summarizing the above, we now have that there exists a constant  $C \geq 1$  such that

$$C^{-1} \exp(\tilde{\psi}(w(x, r))) \leq \nu(B(x, r)) \leq C \exp(\tilde{\psi}(w(x, r))). \tag{4.2}$$

Fix now  $\varepsilon > 0$ . By definition of  $\tilde{\xi}_x$ , for  $r$  small enough we have  $r \geq |x - x_{w(x,r)}| \geq (2|I_{w(x,r)}|)^{\tilde{\xi}_x + \varepsilon}$ . Moreover, again for  $r$  small enough,  $\exp(\tilde{\psi}(w(x, r))) \leq |I_{w(x,r)}|^{\alpha(x) - \varepsilon}$  by definition of  $\alpha(x)$ .

These estimates yield

$$\nu(B(x, r)) \leq C \exp(\tilde{\psi}(w(x, r))) \leq Cr^{(\alpha(x) - \varepsilon)/(\tilde{\xi}_x + \varepsilon)},$$

and, thus, by letting  $r$  tend to zero, it follows that  $h_\nu(x) \geq (\alpha(x) - \varepsilon)/(\tilde{\xi}_x + \varepsilon)$ .

- Finally, for the right inequality, let  $(n_j)_{j \geq 1}$  be an increasing sequence of integers such that  $\exp(\tilde{\psi}(w_{n_j}(x))) \geq |I_{w_{n_j}(x)}|^{\alpha(x) + \varepsilon}$  for all  $j \geq 1$ . Such a sequence exists by definition of  $\alpha(x)$ . By construction,  $x_{w_j} \in B(x, 2|I_{w_j}|)$  so that, using formula (2.10) for  $\nu$ , we obtain  $\nu(B(x, 2|I_{w_{n_j}}(x)|)) \geq |I_{w_{n_j}(x)}|^{\alpha(x) + \varepsilon}$  and  $h_\nu(x) \leq \alpha(x) + \varepsilon$ . Since the previous estimates hold for all  $\varepsilon > 0$ , we obtain the desired result.  $\square$

4.2. *The singularity sets of  $\nu$ .* In view of Proposition 4.1, it is natural to introduce the following property for real numbers of  $[0, 1]$ .

*Definition 4.2.* Let  $\alpha > 0, \xi \geq 1$  and  $\varepsilon > 0$ .

A real number  $x \in [0, 1]$  is said to satisfy the property  $\mathcal{P}(\alpha, \xi, \varepsilon)$  if there exists an increasing sequence of positive integers  $(n_j)_{j \geq 1}$  such that for every  $j \geq 1$ , there exists  $w \in \Sigma_{n_j}$  such that  $x \in B(x_w, \lambda_w^{\xi-\varepsilon})$  and  $\tilde{\alpha}(w) \in [\alpha - \varepsilon, \alpha + \varepsilon]$ .

We say that  $x$  is approximated at degree  $\xi - \varepsilon$  by the pair  $(x_w, \lambda_w)$  when  $x \in B(x_w, \lambda_w^{\xi-\varepsilon})$ . Hence, a real number  $x$  satisfies  $\mathcal{P}(\alpha, \xi, \varepsilon)$  when  $x$  is approximated at degree  $\xi - \varepsilon$  by some couples among the family  $\{(x_w, \lambda_w)\}_{w \in \Sigma^*}$ , those couples being selected according to the value of  $\tilde{\alpha}(w)$  (we impose that  $\tilde{\alpha}(w) \in [\alpha - \varepsilon, \alpha + \varepsilon]$ ).

*Definition 4.3.* For  $h > 0$  one sets

$$F(h) = \left\{ x \in (0, 1) \mid \left\{ \begin{array}{l} \forall \varepsilon > 0, \exists \alpha > 0, \exists \xi \geq 1 \text{ such that} \\ \alpha/\xi \leq h + \varepsilon \text{ and } \mathcal{P}(\alpha, \xi, \varepsilon) \text{ holds} \end{array} \right. \right\}.$$

We explore the relationship between the singularity sets  $E_\nu(h)$  and our sets  $F(h)$ .

PROPOSITION 4.4. *Let  $h > 0$ .*

- (1) *One has  $F(h) \subset \bigcup_{h' \leq h} E_\nu(h')$ .*
- (2) *One has  $E_\nu(h) \subset F(h)$ .*

*Proof.* (1) Let  $x \in F(h)$ . Then fix  $\varepsilon \in (0, 1/2)$ , as well as  $\alpha > 0$  and  $\xi \geq 1$  such that  $x$  satisfies  $\mathcal{P}(\alpha, \xi, \varepsilon)$ . Let  $(n_j)_{j \geq 1}$  be an increasing sequence of integers such that, for every  $j \geq 1$ , there exists  $w_j \in \Sigma_{n_j}$  such that  $x \in B(x_{w_j}, (\lambda_{w_j})^{\xi-\varepsilon})$  and  $\tilde{\alpha}(w_j) \in [\alpha - \varepsilon, \alpha + \varepsilon]$ . Recall that  $\tilde{\alpha}(w_j) = ((\tilde{\psi}(w_j))/(\log |I_{w_j}|))$ .

Taking  $r_j = (\lambda_{w_j})^{\xi-\varepsilon}$  for  $j \geq 1$ , using (2.10) we obtain that

$$\nu(B(x, r_j)) \geq \exp(\tilde{\psi}(w_j)) \geq (\lambda_{w_j})^{\tilde{\alpha}(w_j)} \geq r_j^{(\alpha+\varepsilon)/(\xi-\varepsilon)}.$$

This implies that

$$h_\nu(x) \leq \frac{\alpha + \varepsilon}{\xi - \varepsilon} \leq \frac{\alpha}{\xi} \frac{\xi}{\xi - \varepsilon} + 2\varepsilon \leq h + O(\varepsilon).$$

Consequently, letting  $\varepsilon$  tend to zero yields  $h_\nu(x) \leq h$ .

(2) Fix  $x \in E_\nu(h)$  and  $\varepsilon > 0$ . By definition there is a sequence  $(r_j)_{j \geq 1}$  of positive real numbers decreasing to zero such that for all  $j \geq 1$ ,  $\nu(B(x, r_j)) \geq (r_j)^{h+\varepsilon}$ . Then (4.2) yields

$$\exp(\tilde{\psi}(w(x, r_j))) \geq C^{-1}(r_j)^{h+\varepsilon},$$

which is equivalent to

$$|I_{w(x, r_j)}|^{\tilde{\alpha}(w(x, r_j))} \geq C^{-1}(r_j)^{h+\varepsilon}.$$

Thus, writing  $|x - x_{w(x, r_j)}| = (2|I_{w(x, r_j)}|)^{\xi_j} = (\lambda_{w(x, r_j)})^{\xi_j}$ , we obtain

$$|I_{w(x, r_j)}|^{\tilde{\alpha}(w(x, r_j))} \geq C^{-1}(2|I_{w(x, r_j)}|)^{\xi_j(h+\varepsilon)} \tag{4.3}$$

since  $|x - x_{w(x, r_j)}| \leq r_j$ .

Recall that  $B(x, r_j) \subset I_{w(x, r_j)}$  by (4.1). This implies that  $\xi_j \geq 1$ . Moreover,  $\limsup_{j \rightarrow \infty} \xi_j < \infty$ : if this limsup is  $+\infty$ , then combining (3.1) and part (1) of the current proposition, we would have  $h_\nu(x) = 0$ .

Consequently, there exists  $(\alpha, \xi) \in \mathbb{R}_+^* \times [1, \infty)$  and an increasing sequence of integers,  $(j_k)_{k \geq 1}$ , such that simultaneously:

$$\begin{aligned} |\alpha - \tilde{\alpha}(w(x, r_{j_k}))| &\leq \varepsilon, \\ |\xi - \xi_{j_k}| &\leq \varepsilon, \\ (\tilde{\alpha}(w(x, r_{j_k})), \xi_{j_k}) &\longrightarrow (\alpha, \xi) \text{ as } k \rightarrow +\infty. \end{aligned}$$

Note that (4.3) implies that  $\alpha/\xi \leq h + \varepsilon$ .

Since these properties hold for all  $\varepsilon > 0, x \in F(h)$ . □

**COROLLARY 4.5.** For every  $h > 0, E_\nu(h) = F(h) \setminus \bigcup_{h' < h} F(h')$ .

*Proof.* This follows directly when combining parts (1) and (2) of Proposition 4.4. □

**4.3. Upper bound for the dimensions of the singularity sets.** It is known that, for any positive Borel measure  $\nu$  with a bounded support, we always have the upper bound  $\dim E_\nu(h) \leq \tau_\nu^*(h)$  (see, for instance, [10]). Using the definition of the Legendre transform (1.5), in order to find an upper bound for  $\dim E_\nu(h)$ , we need a lower bound for  $\tau_\nu$ .

**PROPOSITION 4.6.** For every  $q \in \mathbb{R}$ , we have  $\tau_\nu(q) \geq \min(\theta(q), 0)$ .

This immediately yields the desired upper bound for the spectrum. Indeed, using the definition of the Legendre transform (1.5), we obtain that  $\tau_\nu^*(h) = h \cdot \dim X$  if  $h \in [0, \theta'(\dim X)]$ , and  $\tau_\nu^*(h) = \theta^*(h)$  if  $h > \theta'(\dim X)$ .

In the next section, we prove that  $\dim E_\nu(h) \geq \tau_\nu^*(h)$ , for every  $h \geq 0$ . By applying the inverse Legendre transform to the inequality claimed by Proposition 4.6, we obtain the equality  $\tau_\nu(q) = \min(\theta(q), 0)$ , as stated in Theorem 1.5.

*Proof.* Let  $r > 0$  and consider  $\mathcal{B} = \{B_i\}$ , a packing of  $[0, 1]$  by disjoint closed intervals  $B_i$  of radius  $r$ .

- First fix  $q < 0$ . Owing to the bounded distortion principle (which rules the size of the elements of the grid  $\{I_w\}_{w \in \Sigma^*}$ ), there exists a constant  $C > 0$  depending on  $\varphi$  only, such that each  $B_i$  contains an interval  $I_{w_i}$  satisfying  $|I_{w_i}| \geq r/C$ .

Using the formula (2.11) for  $\nu$ , we have that  $\nu(B_i) \geq \exp(|w_i|P(\psi))\mu_{\psi, \varphi}(I_{w_i})$  and since  $q < 0$ , we get  $\nu(B_i)^q \leq \exp(q|w_i|P(\psi))\mu_{\psi, \varphi}(I_{w_i})^q$ . Formula (3.2) yields

$$\nu(B_i)^q \leq C_q \mu_q(I_{w_i}) |I_{w_i}|^{\theta(q)}.$$

Finally, since  $\theta(q) < 0$  and  $|I_{w_i}| \geq r/C$ , we obtain

$$\nu(B_i)^q \leq C_q C^{-q} r^{\theta(q)} \mu_q(I_{w_i}).$$

Consequently,

$$\sum_{B_i \in \mathcal{B}} \nu(B_i)^q \leq C_q C^{-q} r^{\theta(q)}$$

independently of the choice of the  $r$ -packing  $\mathcal{B} = \{B_i\}$ . This yields  $\tau_\nu(q) \geq \theta(q)$ .

- Second fix  $q \in (0, \dim X)$ . Let  $M = \sup_{t \in I_0 \cup I_1} |T'(t)|$ . Let  $S_r = \{w \in \Sigma^* \mid 2r \leq |I_w| \leq 2r(M + 1)\}$ . We also set  $n_r = \max\{|w| : w \in S_r\}$  and  $n'_r = \min\{|w| : w \in S_r\}$ . Since the mapping  $T$  is  $C^1$  and  $|T'| > 1$ , we have  $n_r = O(|\log(r)|)$  and  $n'_r = O(|\log(r)|)$ .

For  $w \in S_r$ ,  $I_w$  meets at most  $M + 1$  balls  $B_i$  of the packing  $\mathcal{B}$ . Reciprocally, each  $B_i$  is included in the union of at most two intervals belonging to  $S_r$ , say  $I_w$  and  $I_{w'}$  (and possibly by the singleton  $\{1\}$ ).

The subadditivity of the application  $t \geq 0 \mapsto t^q$  yields  $v(B_i)^q \leq v(\{1\}) + v(I_w)^q + v(I_{w'})^q$  if  $1 \in B_i$ , and  $v(B_i)^q \leq v(I_w)^q + v(I_{w'})^q$  otherwise. Thus,

$$\begin{aligned} \sum_{B_i \in \mathcal{B}} v(B_i)^q &\leq v(\{1\}) + (M + 1) \sum_{n=n'_r}^{n_r} \sum_{w \in S_r \cap \Sigma_n} v(I_w)^q \\ &\leq v(\{1\}) + (M + 1) \sum_{n=0}^{n_r} \sum_{w \in \tilde{S}_r \cap \Sigma_n} v(I_w)^q, \end{aligned}$$

where  $\tilde{S}_r = \{w \in \Sigma^* : 2r \leq |I_w| \} \supset S_r$ . Since  $q \in (0, \dim X)$  and  $\dim X < 1$ , we have

$$\begin{aligned} v(I_w)^q &= \left( \sum_{\substack{n' < |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\}) + \sum_{\substack{n' \geq |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\}) \right)^q \\ &\leq \sum_{\substack{n' < |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\})^q + \left( \sum_{\substack{n' \geq |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\}) \right)^q \\ &\leq \sum_{\substack{n' < |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\})^q + \left( \sum_{\substack{n' \geq |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} \mu_{\psi, \varphi}(I_v) \exp(n' P(\psi)) \right)^q \\ &\leq \sum_{\substack{n' < |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\})^q + \mu_{\psi, \varphi}(I_w)^q \left( \sum_{n' \geq |w|} \exp(n' P(\psi)) \right)^q \\ &= \sum_{\substack{n' < |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\})^q + K_q \exp(q|w|P(\psi)) \mu_{\psi, \varphi}(I_w)^q \end{aligned}$$

where we used the formula (2.11) for  $v$  and  $K_q = (1 - \exp(qP(\psi)))^{-q}$ . Consequently,

$$\sum_{B_i \in \mathcal{B}} v(B_i)^q \leq v(\{1\}) + (M + 1)(A_1 + A_2),$$

where

$$A_1 = \sum_{n=0}^{n_r} \sum_{w \in \tilde{S}_r \cap \Sigma_n} \sum_{\substack{n' < |w| \\ v \in \Sigma_{n'} \\ x_v \in I_w}} v(\{x_v\})^q$$

and  $A_2 = \sum_{n=0}^{n_r-1} \sum_{w \in S_r \cap \Sigma_n} K_q \exp(q|w|P(\psi)) \mu_{\psi, \varphi}(I_w)^q.$



Obviously,

$$A_2 \leq K_q n_r \sum_{n=0}^{n_r} \sum_{w \in \tilde{S}_r \cap \Sigma_n} \exp(q|w|P(\psi)) \mu_{\psi, \varphi}(I_w)^q.$$

Then, a simple reordering of the terms in  $A_1$  yields

$$A_1 \leq \sum_{n=0}^{n_r-1} \sum_{v \in \Sigma_n} \left( \# \left\{ w \in \tilde{S}_r \cap \bigcup_{p=n+1}^{n_r} \Sigma_p \mid x_v \in I_w \right\} \right) \nu(\{x_v\})^q.$$

Each cardinality is, by construction, less than  $n_r$ . Hence,

$$A_1 \leq n_r \sum_{n=0}^{n_r-1} \sum_{v \in \tilde{S}_r \cap \Sigma_n} \nu(\{x_v\})^q \leq n_r \sum_{n=0}^{n_r-1} \sum_{v \in \tilde{S}_r \cap \Sigma_n} \exp(q|v|P(\psi)) \mu_{\psi, \varphi}(I_v)^q.$$

Finally, formula (3.2) yields

$$\begin{aligned} \sum_{B_i \in \mathcal{B}} \nu(B_i)^q &\leq \nu(\{1\}) + (M + 1)(K_q + 1)n_r \sum_{n=0}^{n_r} \sum_{w \in \tilde{S}_r \cap \Sigma_n} |I_w|^{\theta(q)} \mu_q(I_w) \\ &\leq \nu(\{1\}) + (M + 1)(K_q + 1)n_r \sum_{n=0}^{n_r} r^{\theta(q)} \|\mu_q\| \\ &\leq \nu(\{1\}) + (M + 1)(K_q + 1)(n_r + 1)^2 r^{\theta(q)}, \end{aligned}$$

where we used that  $r \leq |I_w|$  for  $w \in \tilde{S}_r$  and  $\theta(q) < 0$ . The estimate on  $n_r$  finally gives that

$$\sum_{B_i \in \mathcal{B}} \nu(B_i)^q \leq O(r^{\theta(q)} |\log(r)|^2).$$

The above upper bound is independent of the packing  $\mathcal{B}$ . This yields  $\tau_\nu(q) \geq \theta(q)$ .

- Finally, let  $q \geq \dim X$ . Since  $\nu$  is discrete, we necessarily have  $\tau_\nu(q) = 0$  for every  $q \geq 1$ . In addition, if  $q = \dim X$ , then  $\tau_\nu(q) \geq \theta(q) = 0$ . The concavity of  $\tau_\nu$  then implies that  $\tau_\nu(q) = 0$  for every  $q \geq \dim X$ . □

4.4. *Lower bounds for the dimensions of the singularity sets.* In view of Definition 4.2 and Corollary 4.5, in order to find lower bounds for the Hausdorff dimensions of the level sets  $E_\nu(h)$ , it is natural to introduce the following limsup sets: for every  $\alpha, \varepsilon > 0$  and  $\xi \geq 1$ , set

$$G(\alpha, \xi, \varepsilon) = \bigcap_{N \geq 1} \bigcup_{n \geq N} \bigcup_{w \in \Sigma_n: \alpha - \varepsilon \leq \tilde{\alpha}(w) \leq \alpha + \varepsilon} B(x_w, \lambda_w^\xi).$$

It is obvious that, if  $x \in F(h)$ , then for every  $\varepsilon > 0$ , there exists  $\alpha$  and  $\xi \geq 1$  such that  $\alpha + \xi \leq h + \varepsilon$  and  $x$  satisfies  $\mathcal{P}(\alpha, \xi, \varepsilon)$ . In particular,  $x$  belongs to  $G(\alpha, \xi, \varepsilon)$ . Our aim here is to control the Hausdorff dimension of the sets  $G(\alpha, \xi, \varepsilon)$  for the following reason: if the Hausdorff dimension of  $G(\alpha, \xi, \varepsilon)$  is proved to be strictly less than  $\tau_{\mu_q}^*(\tau'_\mu(q))$ , then  $G(\alpha, \xi, \varepsilon)$  is  $\mu_q$ -negligible.

PROPOSITION 4.7. *There exists  $C > 0$ , depending on  $\theta$  only, such that for  $\varepsilon > 0$  small enough, for all  $\alpha > 0$  and  $\xi \geq 1$ ,*

$$\dim G(\alpha, \xi, \varepsilon) \leq C\varepsilon + \frac{\max(\theta^*(\alpha - \varepsilon), \theta^*(\alpha + \varepsilon))}{\xi}.$$

When the right-hand side of the above inequality is negative, the set  $G(\alpha, \xi, \varepsilon)$  is empty.

*Proof.* Fix  $\alpha, \varepsilon > 0$ ,  $\xi \geq 1$ , and for  $N \geq 1$  let  $\delta_N = \sup_{w \in \Sigma_N} \lambda_w^\xi$ . For any  $s > 0$  and  $\delta > 0$ , we denote by  $\mathcal{H}_\delta^s$  the  $s$ -Hausdorff pre-measure computed using coverings by sets of diameter less than  $\delta$  [13]. By construction,

$$\mathcal{H}_{\delta_N}^s(G(\alpha, \xi, \varepsilon)) \leq \sum_{n \geq N} \sum_{\substack{w \in \Sigma_n: \\ \alpha - \varepsilon \leq \tilde{\alpha}(w) \leq \alpha + \varepsilon}} 2^s (\lambda_w)^{s\xi}.$$

We proceed as follows.

- Suppose that  $\alpha + \varepsilon \leq \theta'(0)$ . If  $q \geq 0$ , using that  $\tilde{\alpha}(w) = \tilde{\psi}(w)/\chi(w)$ ,  $\tilde{\psi}(w) < 0$  and  $\chi(w) = \log |I_w| < 0$ , we have for some constant  $C > 0$  that

$$\begin{aligned} \mathcal{H}_{\delta_N}^s(G(\alpha, \xi, \varepsilon)) &\leq C \sum_{n \geq N} \sum_{\substack{w \in \Sigma_n: \\ q\tilde{\psi}(w) \geq q(\alpha + \varepsilon)\chi(w)}} (\lambda_w)^{s\xi} \\ &\leq C \sum_{n \geq N} \sum_{w \in \Sigma_n} \exp(q\tilde{\psi}(w) - q(\alpha + \varepsilon)\chi(w)) \exp(s\xi\chi(w)). \end{aligned}$$

Now take  $s = (\eta + \theta^*(\alpha + \varepsilon))/\xi$  with  $\eta > 0$ . We deduce that

$$\mathcal{H}_{\delta_N}^s(G(\alpha, \xi, \varepsilon)) \leq C \sum_{n \geq N} \sum_{w \in \Sigma_n} \exp(q\tilde{\psi}(w) - (q(\alpha + \varepsilon) - \eta - \theta^*(\alpha + \varepsilon))\chi(w)). \tag{4.4}$$

We know that,  $n \geq N$  being fixed, the second sum in the previous double sum is comparable to

$$\exp(nP(\psi_{q,t_{q,\alpha,\varepsilon}})) \quad \text{where } t_{q,\alpha,\varepsilon} = q(\alpha + \varepsilon) - \eta - \theta^*(\alpha + \varepsilon).$$

Since  $\alpha + \varepsilon \leq \theta'(0)$ , we can choose  $q \geq 0$  such that  $\theta^*(\alpha + \varepsilon) = (\alpha + \varepsilon)q - \theta(q) - \gamma_q$ , with  $0 \leq \gamma_q \leq \eta/2$ . This implies that

$$P(\psi_{q,t_{q,\alpha,\varepsilon}}) \leq P(\psi_{q,(\theta(q) - \eta/2)}) < 0.$$

Using (4.4), this implies that  $\lim_{N \rightarrow \infty} \mathcal{H}_{\delta_N}^s(G(\alpha, \xi, \varepsilon)) = 0$ . Consequently,  $\dim G(\alpha, \xi, \varepsilon) \leq (\eta + \theta^*(\alpha + \varepsilon))/\xi$ . Since this holds for all  $\eta > 0$ , we have the desired conclusion.

- When  $\alpha - \varepsilon \geq \theta'(0)$ , the same approach with  $q < 0$  yields the desired estimate.
- When  $\alpha - \varepsilon \leq \theta'(0) \leq \alpha + \varepsilon$ , for  $\eta > 0$  and  $s = (\eta + \theta(0))/\xi$  we can write

$$\mathcal{H}_{\delta_N}^s(G(\alpha, \xi, \varepsilon)) \leq C \sum_{n \geq N} \sum_{w \in \Sigma_n} (\lambda_w)^{s\xi} \leq C \sum_{n \geq N} \exp(nP(-(\eta + \theta(0))\varphi)).$$

This last sum tends to 0 as  $N$  tends to infinity. Thus,  $\dim G(\alpha, \xi, \varepsilon) \leq \theta(0)/\xi$ .

Now, since  $\theta^*$  is differentiable near  $\theta'(0)$ , there exists a constant  $C > 0$ , depending only on  $\theta$ , such that  $\theta(0)/\xi \leq C\varepsilon + \max(\theta^*(\alpha - \varepsilon), \theta^*(\alpha + \varepsilon))/\xi$  when  $\varepsilon > 0$  is sufficiently small and  $\alpha - \varepsilon \leq \theta'(0) \leq \alpha + \varepsilon$ . The result follows.  $\square$

COROLLARY 4.8. For all  $h > 0$ ,  $\dim F(h) \leq h \cdot \dim X$ .

*Proof.* Let  $x \in F(h)$ , and  $\varepsilon > 0$ . There exists  $\alpha$  and  $\xi \geq 1$  such that  $\alpha + \xi \leq h + \varepsilon$  and  $x$  satisfies  $\mathcal{P}(\alpha, \xi, \varepsilon)$ . It follows that  $x$  belongs to the set  $G(\alpha, \xi, \varepsilon)$ , whose Hausdorff dimension is bounded from above by the last Proposition 4.7. This yields that for every  $\varepsilon > 0$

$$\dim F(h) \leq \sup_{\alpha, \xi: \alpha/\xi \leq h+\varepsilon} C\varepsilon + \frac{\max(\theta^*(\alpha - \varepsilon), \theta^*(\alpha + \varepsilon))}{\xi}.$$

Letting  $\varepsilon$  tend to zero, yields

$$\dim F(h) \leq \sup_{\alpha, \xi: \alpha/\xi \leq h} \frac{\theta^*(\alpha)}{\xi} \leq h \cdot \sup_{\alpha \geq 0} \frac{\theta^*(\alpha)}{\alpha} \leq h \cdot \dim X.$$

The last inequality follows from the fact that the measure  $\mu_\varphi$  and  $\mu_\psi$  have their support included in  $X$  and from the definition of the Legendre transform (1.5).  $\square$

PROPOSITION 4.9. We make the following propositions.

- (1) If  $h \geq \theta'(\dim X)$  and  $\theta^*(h) \geq 0$ , then  $\dim E_\nu(h) \geq \theta^*(h)$ .
- (2) If  $h \in [0, \theta'(\dim X)]$ , then  $\dim E_\nu(h) \geq h \cdot \dim X$ .

*Proof.* (1) If  $h \geq \theta'(\dim X)$  and  $\theta^*(h) \geq 0$ , then  $h \in \{\theta'(q) \mid q \geq \dim X\} \cup \{\lim_{q \rightarrow -\infty} \theta'(q)\}$ . Proposition 4.1 shows that  $E_\nu(h)$  contains the set

$$\begin{aligned} \tilde{E}(h) &= \left\{ x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h \text{ and } \tilde{\xi}(x) = 1 \right\} \\ &= \left\{ x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h \right\} \setminus \bigcup_{m \geq 1} \left\{ x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h \text{ and } \tilde{\xi}(x) > 1 + 1/m \right\}. \end{aligned}$$

Consider first the case where  $h = \theta'(q)$  for some  $q \geq \dim X$ .

Part (1) of Proposition 3.5 gives that  $\mu_q(\{x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h\}) = 1$ .

From Proposition 4.7, we easily deduce that for all  $m \geq 1$ ,

$$\dim \left\{ x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = \theta'(q) \text{ and } \tilde{\xi}(x) > 1 + 1/m \right\} < \theta^*(\theta'(q)) = \theta^*(h).$$

Then, Proposition 3.5(2) gives that the countable union of sets

$$\bigcup_{m \geq 1} \left\{ x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h \text{ and } \tilde{\xi}(x) > 1 + 1/m \right\}$$

is  $\mu_q$ -negligible. Hence,  $\mu_q(\tilde{E}(h)) = 1$ . Finally,  $\dim E_\nu(h) \geq \dim(\mu_q) \geq \theta^*(\theta'(q)) = \theta^*(h)$ .

It remains us to treat the case of the right endpoint of the singularity spectrum  $h_{\max} := \sup\{h \geq 0 \mid \tau_\nu^*(h) \geq 0\}$ . We have  $h_{\max} = \lim_{q \rightarrow -\infty} \theta'(q) < +\infty$ . In this case there is no natural Gibbs measure supported by the set  $\tilde{E}_\nu(h_{\max}) = \{x \mid \lim_{n \rightarrow +\infty} \alpha_n(x) = h_{\max}\}$ .

When  $\tau_\nu^*(h_{\max}) > 0$ , the approach used in [15] to achieve the multifractal analysis of Birkhoff averages associated with a continuous, but not necessarily Hölderian, potential on  $\Sigma$  allows us to construct a Cantor set included in  $\{x \mid \lim_{n \rightarrow +\infty} \alpha_n(x) = h_{\max}\}$  (usually called in this context a Moran set) and a measure  $\mu_{-\infty}$  supported by this Cantor

set, for which  $\dim(\mu_{-\infty}) \geq \theta^*(h_{\max})$ . The conclusion is then obtained as in the previous case.

We discuss the case  $\tau_v^*(h_{\max}) = 0$  briefly in §4.5.

(2) Let  $h \in (0, \theta'(\dim X)]$  and write  $h = \theta'(\dim X)/\xi$ , with  $\xi \geq 1$ . Consider the set  $S(\xi, \tilde{\varepsilon})$  of formula (3.3) with a suitable sequence  $\tilde{\varepsilon}$  such that Proposition 3.6 can be applied. This provides us with a positive Borel measure  $m_\xi$  such that the following hold:

- $m_\xi(S(\xi, \tilde{\varepsilon})) > 0$ ;
- $m_\xi(E) = 0$  as soon as  $\dim E < \dim(X)((\theta'(\dim(X)))/(\xi)) = h \cdot \dim X$ .

It follows from Proposition 4.4 that

$$S(\xi, \tilde{\varepsilon}) \setminus \bigcup_{0 < h' < h} F(h') \subset E_v(h).$$

By Corollary 4.8 we have  $\dim F(h') \leq h' \cdot \dim X < h \cdot \dim X$ , for all  $0 < h' < h$ . Hence,  $m_\xi(F(h')) = 0$  for all  $h' < h$ . Moreover, the set family  $(F(h'))_{0 < h' < h}$  is non-decreasing.

Summarizing the above, we now have that  $m_\xi(E_v(h)) > 0$ , and hence  $\dim E_v(h) \geq h \cdot \dim X$ . □

4.5. *The case  $\tau_v^*(h_{\max}) = 0$ .* When  $\tau_v^*(h_{\max}) = 0$ , by the same computations as in [15], it is possible to construct a Cantor set included in  $\{x \mid \lim_{n \rightarrow +\infty} \alpha_n(x) = h_{\max}\}$  and a non-trivial measure  $\mu_{-\infty}$  supported by this Cantor set. Obviously  $\dim(\mu_{-\infty}) = 0$ , but there is a gauge function  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  naturally associated with  $\mu_{-\infty}$ , such that for every  $\varepsilon > 0$ , we have that  $f(x) = o(x^\varepsilon)$ , for  $x$  tending to zero from above, and for every Borel set  $E$ ,

$$\mu_\infty(E) \leq f(|E|).$$

The difficulty lies in the fact that the argument of previous section do not apply here. Indeed, every set  $\{x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h_{\max} \text{ and } \tilde{\xi}(x) > 1 + 1/m\}$  has also Hausdorff dimension zero, and these sets are not negligible *a priori* from the  $\mu_{-\infty}$  viewpoint. Hence, we cannot conclude directly that  $\mu_{-\infty}(\{x \mid \lim_{n \rightarrow +\infty} \alpha_n(x) = h_{\max} \text{ and } \tilde{\xi}(x) = 1\}) = 1$ .

This can certainly be circumvented by at least two methods:

- one could modify the construction of the Cantor set included in  $\{x \mid \lim_{n \rightarrow +\infty} \alpha_n(x) = h_{\max}\}$  to ensure that this Cantor set is indeed included in  $\{x \mid \lim_{n \rightarrow +\infty} \alpha_n(x) = h_{\max} \text{ and } \tilde{\xi}(x) = 1\}$ ;
- one could replace the dimension argument (to neglect the sets  $\{x \mid \lim_{n \rightarrow \infty} \alpha_n(x) = h_{\max} \text{ and } \tilde{\xi}(x) > 1 + 1/m\}$ ) by an argument based on the study of generalized Hausdorff measure using gauge functions.

Both approaches, which are very technical, will certainly work. However, this would require some lengthy extra work, and therefore we do not go into further details here.

4.6. *The  $L^q$ -spectrum.* If  $q \geq \dim X$ , then  $\tau_v(q) = 0 \leq \theta(q)$ . This follows from the observations that  $\tau_v(\dim X) = \tau_v(1) = 0$  and that  $\tau_v$  is concave.

When  $q \leq \dim X$ , we proved that  $\tau_v^*(\theta'(q)) = \theta^*(\theta'(q))$ . Taking the inverse Legendre transform yields that we have  $\tau_v(q) = \theta(q)$ .

Summarizing the above, we have  $\tau_v(q) = \min(0, \theta(q))$  for every  $q \in \mathbb{R}$ .

*Acknowledgements.* The authors thank the anonymous referee for his valuable remarks and suggestions.

## REFERENCES

- [1] V. Aversa and C. Bandt. The multifractal spectrum of discrete measures. 18th Winter School on Abstract Analysis (Srn, 1990). *Acta Univ. Carolin. Math. Phys.* **31** (1990), 5–8.
- [2] J. Barral and S. Seuret. Sums of Dirac masses and conditioned ubiquity. *C. R. Math. Acad. Sci. Paris* **339**(11) (2004), 787–792.
- [3] J. Barral and S. Seuret. Combining multifractal additive and multiplicative chaos. *Comm. Math. Phys.* **257**(2) (2005), 473–497.
- [4] J. Barral and S. Seuret. Heterogeneous ubiquitous systems in  $R^d$  and Hausdorff dimensions. *Bull. Braz. Math. Soc.* **38**(3) (2007), 467–515.
- [5] J. Barral and S. Seuret. The multifractal nature of heterogeneous sums of Dirac masses. *Math. Proc. Cambridge Philos. Soc.* **144**(3) (2008), 707–727.
- [6] J. Barral and S. Seuret. Ubiquity and large intersections properties under digit frequencies constraints. *Math. Proc. Cambridge Philos. Soc.* **145** (2008), 527–548.
- [7] L. Barreira, B. Saussol and J. Schmeling. Higher-dimensional multifractal analysis. *J. Math. Pures Appl.* **81** (2002), 67–91.
- [8] F. Ben Nasr. Analyse multifractale de mesures. *C. R. Acad. Sci. Paris Série I. Math.* **319** (1994), 807–810.
- [9] R. Bowen. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms (Springer Lecture Notes in Mathematics, 470)*. Springer, New York, 1975.
- [10] G. Brown, G. Michon and J. Peyrière. On the multifractal analysis of measures. *J. Stat. Phys.* **66** (1992), 775–790.
- [11] M. M. Dodson, M. V. Melián, D. Pestana and S. L. Vélani. Patterson measure and Ubiquity. *Ann. Acad. Sci. Fenn. Ser. A I Math.* **20** (1995), 37–60.
- [12] A. Durand. Ubiquitous systems and metric number theory. *Adv. Math.* **218**(2) (2008), 368–394.
- [13] K. J. Falconer. *Techniques in Fractal Geometry*. Wiley, New York, 1997.
- [14] K. J. Falconer. Representation of families of sets by measures, dimension spectra and Diophantine approximation. *Math. Proc. Cambridge Philos. Soc.* **128** (2000), 111–121.
- [15] A.-H. Fan and D.-J. Feng. On the distribution of long-term average on the symbolic space. *J. Stat. Phys.* **99** (2000), 813–856.
- [16] D.-J. Feng, K. S. Lau and J. Wu. Ergodic limits on the conformal repellers. *Adv. Math.* **169**(1) (2002), 58–91.
- [17] S. Jaffard. Old friends revisited: the multifractal nature of some classical functions. *J. Fourier Anal. Appl.* **3** (1997), 1–22.
- [18] S. Jaffard. On lacunary wavelet series. *Ann. Appl. Probab.* **10**(1) (2000), 313–329.
- [19] B. Mandelbrot and R. Riedi. Exceptions to the multifractal formalism for discontinuous measures. *Math. Proc. Cambridge Philos. Soc.* **123** (1998), 133–157.
- [20] L. Olsen. A multifractal formalism. *Adv. Math.* **116** (1995), 82–196.
- [21] Y. Pesin. *Dimension Theory in Dynamical Systems: Contemporary Views and Applications (Chicago Lectures in Mathematics)*. The University of Chicago Press, Chicago, IL, 1997.
- [22] Y. Pesin and H. Weiss. The multifractal analysis of Gibbs measures: motivation, mathematical moudation, and examples. *Chaos* **7** (1997), 89–106.
- [23] J. Peyrière. A vectorial multifractal formalism. *Fractal Geometry and Applications (Proceedings of Symposia in Pure Mathematics, 72(2))*. Eds. M. L. Lapidus and M. van Frankenhuysen. American Mathematical Society, Providence, RI, 2004, pp. 217–230.
- [24] D. A. Rand. The singularity spectrum  $f(\alpha)$  for cookie-cutters. *Ergod. Th. & Dynam. Sys.* **9** (1989), 527–541.