

Short Communication

Interspecific hybridization between *Cajanus cajan* (L.) Millsp. and *C. lanceolatus* (WV Fitzg) van der Maesen

Sandhya Srikanth^{1,2*}, M. V. Rao² and Nalini Mallikarjuna¹

¹Grain Legumes Program, International Crops Research Institute for Semi-Arid tropics, Patancheru PO 502 324, Andhra Pradesh, India and ²Department of Plant Sciences, Bharathidasan University, Tiruchirappalli 620 024, Tamil Nadu, India

Received 10 April 2013; Accepted 1 August 2013 – First published online 28 August 2013

Abstract

Cultivated pigeonpea has a narrow genetic base. Wild relatives play an important role in the efforts to broaden its genetic base. In this report, we present a successful wide-cross between the cultivated pigeonpea and *Cajanus lanceolatus*, a wild relative from the secondary gene pool, native to Australia, with desirable traits such as frost and drought resistance. A range of F₁ progeny were obtained and the resultant F₁ hybrid plants set mature pods and seeds. The hybrids had intermediate morphology, sharing the traits of both the parents. All the F₁ hybrids flowered profusely. Some of the hybrids were completely male sterile and some were partially fertile with pollen fertility ranging from 35 to 50 %. Meiotic analysis of the fertile F₁ hybrids revealed a high degree of meiotic chromosome pairing between the two parental genomes. Meiotic analysis of the sterile F₁ hybrids revealed that the breakdown of microsporogenesis occurred at the post-meiotic stage after the formation of tetrads. Fertile plants formed regular bivalents with normal disjunction, except for occasional asynchrony at meiotic II division.

Keywords: bruchid; *Cajanus lanceolatus*; gene pools; meiosis; pigeonpea; pollen fertility; sterility

Introduction

Pigeonpea [*Cajanus cajan* (L.) Millsp.] is a multi-purpose grain legume grown by resource-poor farmers in the semi-arid tropics and subtropics. The crop has narrow genetic diversity and is susceptible to a range of diseases and pests such as pod borer [*Helicoverpa armigera* (Hub.)], pod fly [*Melanagromyza obtusa* (Malloch)] and bruchid [*Callosobruchus chinensis* (F.)]. High levels of

resistance to many of these pests and diseases are low to moderate in the cultivated germplasm (Sharma, 2005), but the wild relatives of pigeonpea have shown high levels of resistance to many of the constraints (Green *et al.*, 2006; Sujana *et al.*, 2008; Sharma *et al.*, 2009). The utilization of wild species from the secondary gene pool is important as they are closely related, leading to normal chromosome recombination. This helps in the transfer of useful genes/traits to the cultivated pigeonpea (Mallikarjuna *et al.*, 2011a, b, c). *Cajanus lanceolatus*, a native of northern Australia, is a wild relative from the secondary gene pool. Until now, *C. lanceolatus* had not been successfully crossed; for instance, a previous study

*Corresponding author. E-mail: sandhya.thudumu@gmail.com

(Sateesh Kumar, 1985) has reported that F_1 hybrids died during the vegetative stage. The present paper reports the successful crosses between the cultivated pigeonpea and *C. lanceolatus*.

Experimental

Cajanus lanceolatus (ICP 15639) and *C. cajan* (ICPL 85010) plants were grown and maintained in a glass-house. Crosses were made using *C. cajan* as the female parent and *C. lanceolatus* as the pollen donor. Pollinations were carried out soon after emasculations in the morning before 10 a.m. Out of 86 pollinations, 20 pods were obtained. The pods were harvested 40–45 d after pollination. For cytological analysis of meiocytes, immature flower buds from F_1 hybrids were fixed in Carnoy's II solution (acetic acid–chloroform–ethanol, 1:3:6) for 24 h at 4°C and transferred to Carnoy's I solution (acetic acid:ethanol, 1:3). Meiocytes were squashed and stained in 4% acetocarmine and well-spread meiotic preparations were taken for analysis and photographed.

Results and discussion

Pod formation was 23% when *C. cajan* was crossed with *C. lanceolatus*. More than half of the seeds were normal with the exception of few semi-shrunken seeds (34 %). Of the 35 morphologically normal seeds, 14 germinated to produce hybrid plants under *in vivo* germination conditions. The plants initially grew slowly, but, later on, normal growth was observed. Morphologically, the hybrid plants had excessive growth compared with both the parents. The F_1 hybrids were screened for morphological traits such as plant height, branching pattern, flower size and shape, pod shape and size, and seed colour. Hybrids were tall, measuring 325 cm (P_{10} - F_1) to 380 cm (P_{13} - F_1) in height, resembling the male parent *C. lanceolatus* with a height of 285 cm compared with the female parent *C. cajan* with a height of 185 cm (Fig. 1(a)). All the hybrids flowered at 98 to 160 d from the date of germination. Sateesh Kumar (1985) reported that F_1 hybrids died during the vegetative stage. It is possible that the authors of this study failed to notice that hybrids inherited the long-duration trait of the male




Fig. 1. Morphological observations and meiotic analysis of the F_1 hybrids derived from the cross *Cajanus cajan* (ICPL 85010) \times *Cajanus lanceolatus* (ICP 15639). (a) Comparison of the hybrids (middle) with the cultivar (female, left) and wild (male, right) parents. (b) Metaphase I (fertile plant F_1 - P_6) showing two rod and nine ring bivalents. (c) Metaphase I (sterile plant F_1 - P_7) showing four univalents and nine bivalents. (d) Anaphase I (fertile plant F_1 - P_6) showing normal disjunction of chromosomes. (e) Anaphase I (sterile plant F_1 - P_7) showing five laggards. (f) Fertile and sterile pollens. (g) Unseparated and empty pollen grains in the sterile anther.

Table 1. Meiotic studies of the hybrids derived from the cross *Cajanus cajan* (ICPL 85010) × *Cajanus lanceolatus* (ICP 15639)

| Hybrid plant no. | Metaphase I | | | | Anaphase I (%) | | Pollen fertility (%) |
|------------------|-------------|--------------|------------|--------------|----------------|----|----------------------|
| | Univalents | Bivalents | Trivalents | Tetravalents | ND | AD | |
| P ₁ | 1.4 (1–4) | 9.5 (8–11) | 0.2 (0–1) | 0 (0) | 50 | 50 | Sterile |
| P ₂ | 0.7 (1–4) | 10.2 (7–11) | 0.1 (0–1) | 0.15 (1–2) | 65 | 35 | 37.8 |
| P ₃ | 1.15 (1–4) | 10.1 (8–11) | 0.15 (0–1) | 0.05 (1–1) | 45 | 55 | 24.3 |
| P ₄ | 1.05 (1–5) | 10.1 (7–11) | 0.25 (0–1) | 0 (0) | 60 | 40 | Sterile |
| P ₅ | 0.7 (1–4) | 10.25 (8–11) | 0.2 (0–1) | 0.05 (0–1) | 65 | 35 | 40.5 |
| P ₆ | 0.35 (1–4) | 10.2 (7–11) | 0.2 (0–1) | 0 (0) | 65 | 35 | 40.7 |
| P ₇ | 1 (1–4) | 10.05 (8–11) | 0.1 (0–1) | 0.15 (1–2) | 25 | 75 | Sterile |
| P ₈ | 0.35 (1–3) | 10.2 (8–11) | 0.05 (0–1) | 0.3 (1–2) | 65 | 35 | 56 |
| P ₉ | 0.8 (1–4) | 10.3 (9–11) | 0.2 (0–1) | 0 (0) | 55 | 45 | 36 |
| P ₁₀ | 1.35 (1–4) | 9.75 (7–11) | 0.25 (0–1) | 0.1 (0–1) | 20 | 80 | Sterile |
| P ₁₁ | 0.6 (1–2) | 10.2 (7–11) | 0.2 (0–1) | 0.1 (0–1) | 50 | 50 | 45 |
| P ₁₂ | 1.1 (1–4) | 9.25 (8–11) | 0.3 (0–1) | 0.15 (0–1) | 35 | 65 | Sterile |
| P ₁₃ | 0.65 (1–4) | 9.9 (7–11) | 0.15 (0–1) | 0.3 (0–2) | 50 | 50 | 50 |
| P ₁₄ | 0.75 (1–4) | 9.15 (7–11) | 0.15 (0–1) | 0.35 (0–2) | 45 | 55 | 48 |

ND, normal distribution; AD, abnormal distribution at anaphase I.

parent, and did not maintain the hybrid plants until they reached the flowering stage. Alternatively, it is possible that the genotypes of the female cultivars used in their study, in combination with *C. lanceolatus*, were not genetically successful.

The meiotic analysis of pollen mother cells of the F₁ hybrids exhibited a regular formation of 11 bivalents that were predominantly rings. It is clear from Table 1 that the number of bivalents ranged from 11 in the anther from the fertile plant to 7 in the sterile F₁ plant (Fig. 1(b) and (c)). Univalents were also found in many cells, and the average number of univalents per cell varied from 1 to 5 in the sterile F₁ plant. Meanwhile, trivalents and tetravalents appeared at a lower frequency, ranging from 0 to 2. Normal bivalent formation in the majority of the pollen mother cells is an indication that there is good recombination between the parental genomes. Meiotic anaphase I showed 50–70% of the pollen mother cells with normal disjunction and remaining 30–50% with abnormal disjunction of chromosomes (Fig. 1(e)). At the tetrad stage, 100% normal tetrads were observed in all the hybrids except in P₇ in which 6% of the tetrads contained micronuclei. Pollen fertility was found to vary between 35 and 50 % in the fertile hybrids (Fig. 1(f)). In some of the F₁ hybrids (P₁, P₄, P₇, P₁₀ and P₁₂), total male sterility was observed in all the anthers having 100% sterile pollen grains, a result of unseparated tetrads (Fig. 1(g)). An important observation made was that male sterility was a post-meiotic process. The development of tetrads was normal, but none of them formed pollen grains. Instead, they grouped together and the tetrads did not separate into individual pollen grains. Such sources may be useful in the development of cytoplasmic male sterile systems in pigeonpea; as

such, a phenomenon was observed in the A₇ cytoplasmic male sterile (CMS) system derived from *Cajanus platycarpus* (Mallikarjuna *et al.*, 2012). Pigeonpea crossed with different wild *Cajanus* species has been reported to have given rise to different cytoplasmic male sterile systems (Saxena *et al.*, 2010; Mallikarjuna *et al.*, 2011a, b, c). Hence it is worth exploring if a CMS system can be developed from this cross, as complete male sterility was observed in the F₁ hybrids.

The cross between the cultivated pigeonpea and *C. lanceolatus* generated two categories of progenies. The first category is the fertile progeny with good recombination between the parental genomes, leading to fertile plants, good material for broadening the narrow genetic base of pigeonpea and traits of interest. The second progeny category is the CMS lines, i.e. F₁ hybrids with 100 % male sterility which can be used to develop another CMS source, distinct from the currently available A₅ CMS system (Mallikarjuna and Saxena, 2005), which was derived from the cross between cv. ICPL 85010 and *Cajanus acutifolius*, and developed on cultivated pigeonpea cytoplasm. CMS is developed as a result of the interaction between the cytoplasmic genome of the female parent and the nuclear genome of the pollen parent (Saxena *et al.*, 2010). It is envisaged that the gametic recombination between cv ICPL 85010 and *C. lanceolatus* may have given rise to fertile and sterile hybrid plants.

References

- Green PWC, Sharma HC, Stevenson PC and Simmonds MSJ (2006) Susceptibility of pigeonpea and some of its wild relatives to predation by *Helicoverpa armigera*: implications for

- breeding resistant cultivars. *Australian Journal of Agricultural Research* 57: 831–836.
- Mallikarjuna N and Saxena KB (2005) A new cytoplasmic male sterility system derived from cultivated pigeonpea cytoplasm. *Euphytica* 142: 143–148.
- Mallikarjuna N, Saxena KB and Jadhav DR (2011a) *Cajanus*. In: Kole C (ed.) *Wild Crop Relatives: Genomic and Breeding Resources*. Berlin Heidelberg: Springer Berlin Heidelberg, pp. 21–33.
- Mallikarjuna N, Senapathy S, Jadhav DR, Saxena KB, Sharma HC, Upadhyaya HD, Rathore A and Varshney RK (2011b) Progress in the utilization of *Cajanus platycarpus* (Benth.) Maesen in pigeonpea improvement. *Plant Breeding* 130: 507–514.
- Mallikarjuna N, Jadhav DR, Sandhya Srikanth and Saxena KB (2011c) *Cajanus platycarpus* (Benth.) Maesen as the donor of new pigeonpea cytoplasmic male sterile (CMS) system. *Euphytica* 182: 65–71.
- Mallikarjuna N, Jadhav DR, Saxena KB and Srivastava RK (2012) Cytoplasmic male sterile systems in pigeonpea with special reference to A7 CMS. *Electronic Journal of Plant Breeding* 3: 983–986.
- Sateesh Kumar P (1985) Crossability, genome relationships and inheritance studies in intergeneric hybrids of pigeonpea. PhD Thesis, University of Hyderabad, Hyderabad.
- Saxena KB, Sultana R, Mallikarjuna N, Saxena RK, Kumar RV, Sawargaonkar SL and Varshney RK (2010) Male-sterility systems in pigeonpea and their role in enhancing yield. *Plant Breeding* 129: 125–134.
- Sharma HC (2005) *Heliothis/Helicoverpa Management: Emerging Trends and Strategies for Future Research*. New Delhi: Oxford and India Book House Publishing Co. Pvt. Ltd.
- Sharma HC, Sujana G and Manohar Rao D (2009) Morphological and chemical components of resistance to pod borer, *Helicoverpa armigera* in wild relatives of pigeonpea. *Arthropod–Plant Interactions* 3: 151–161.
- Sujana G, Sharma HC and Manohar Rao D (2008) Antixenosis and antibiosis components of resistance to pod borer, *Helicoverpa armigera* in wild relatives of pigeonpea. *International Journal of Tropical Insect Science* 28: 191–200.