

# Equine piroplasmosis: an insight into global exposure of equids from 1990 to 2019 by systematic review and meta-analysis

## Review

**Cite this article:** Onyiche ThankGE, Taioe MO, Molefe NI, Biu AA, Luka J, Omeh IJ, Yokoyama N, Thekisoe O (2020). Equine piroplasmosis: an insight into global exposure of equids from 1990 to 2019 by systematic review and meta-analysis. *Parasitology* **147**, 1411–1424. <https://doi.org/10.1017/S0031182020001407>

Received: 4 March 2020

Revised: 21 June 2020

Accepted: 20 July 2020

First published online: 3 August 2020


### Key words:

Apicomplexa; donkeys; equine piroplasmosis; horses; meta-analysis

### Author for correspondence:

ThankGod E. Onyiche,

E-mail: [et.onyiche@unimaid.edu.ng](mailto:et.onyiche@unimaid.edu.ng)

ThankGod E. Onyiche<sup>1,2,3</sup> , Moeti O. Taioe<sup>4</sup>, Nthatsi I. Molefe<sup>1</sup>, Abdullahi A. Biu<sup>2</sup>, Joshua Luka<sup>2</sup>, Isaac J. Omeh<sup>5</sup>, Naoaki Yokoyama<sup>3</sup> and Oriel Thekisoe<sup>1</sup>

<sup>1</sup>Unit for Environmental Sciences and Management, North-West University, Potchefstroom Campus, Private Bag X6001, Potchefstroom 2520, South Africa; <sup>2</sup>Department of Veterinary Parasitology and Entomology, University of Maiduguri, P. M. B. 1069, Maiduguri 600230, Nigeria; <sup>3</sup>National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido 080-8555, Japan; <sup>4</sup>National Zoological Gardens of South Africa, South African National Biodiversity Institute, PO Box 754, Pretoria 0001, South Africa and <sup>5</sup>Department of Veterinary Physiology and Biochemistry, University of Maiduguri, P. M. B. 1069, Maiduguri 600230, Nigeria

## Abstract

Equine piroplasmosis (EP) is a tick-borne disease of economic importance, relevant in the international movement of equids. The causative agents are at least two apicomplexan protozoan parasites *Babesia caballi* and *Theileria equi*. To date, there is no study that estimates global and regional exposure of equids to EP. We therefore conducted a systematic review and meta-analysis to estimate the pooled prevalence and heterogeneity of EP using random-effects model. Six electronic databases were searched for publications on EP and assessed according to Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines. A total of 66 eligible studies published between 1990 and 2019 and representing 24 041 equids were included. The overall pooled prevalence estimates (PPEs) of *B. caballi* was 22.3% (95% CI 21.7–22.8), while the overall PPE for *T. equi* was 29.4% (95% CI 28.7–30.0). The overall pooled prevalence due to co-infection with both parasites was 11.8% (95% CI 11.32–12.32). Also, subgroup analysis according to sex, age, diagnostic technique, equid species, region and publication years showed a substantial degree of heterogeneity across studies computed for both *B. caballi* and *T. equi* infections in equids. Awareness of the current status of EP globally will alert the relevant authorities and stakeholders where necessary on the need for better preventive and control strategies against the disease.

## Introduction

Equine piroplasmosis (EP) is a disease of equids caused by at least two haemoparasites, *Babesia caballi* and *Theileria equi* transmitted by Ixodid ticks. Clinical manifestations of the disease in infected animals include lethargy, anorexia, fever, jaundice, haemolysis and petechia haemorrhages in mucous membranes (Scoles and Ueti, 2015).

EP is responsible for economic losses in the equid industry. *Babesia caballi* and *T. equi* are in the group of apicomplexan parasites collectively called piroplasms (Levine, 1985). *Babesia caballi* lack a pre-erythrocytic cycle and *T. equi* have no documented transovarial transmission (Homer *et al.*, 2000). *Babesia caballi* is considered less virulent than *T. equi* because the latter acute phase infects leucocytes before erythrocytes and the infection is long-lasting (Ramsay *et al.*, 2013). Surviving animals remain chronic carriers with low levels of parasitaemia and serve as reservoirs for ticks (Wise *et al.*, 2013; Scoles and Ueti, 2015).

EP is widespread in subtropical and tropical regions of the world (Uilenberg, 2006). It is endemic in several parts of Africa, Asia, America and Europe where competent tick vectors are present (Rothschild, 2013; Onyiche *et al.*, 2019). International movement of chronically infected animals has played some role in the epidemiology of this disease necessitating proper screening of animals prior to movement (Ayala-Valdovinos *et al.*, 2014).

Due to non-specific clinical signs associated with EP, diagnosis is often challenging. Furthermore, the sensitivity of different diagnostic tests such as microscopy, serology and PCR is another issue in diagnosis (Mans *et al.*, 2015). These issues have led to many different types of epidemiological prevalence studies from different areas. This is also complicated by several factors such as presence and abundance of competent vectors, management practices, host activity and effectiveness of control programmes for ticks (reviewed by Onyiche *et al.*, 2019).

To date, there has been no systematic review to ascertain the current global status of EP. Therefore, we conducted a systematic review and meta-analysis to determine the global exposure and evaluated risk factors potentially associated with their occurrence.

## Materials and methods

### Search strategy and selection criteria

The study was carried out in accordance with the methodology recommended by the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) (Fig. 1) (Moher *et al.*, 2015). We searched the primary literature of published articles from 1 January 1990 to 25 February 2019 in English databases of Scopus, Science Direct, PubMed, Web of Science, Embase and Springer link. Keywords used for the systemic search were 'Equine Piroplasmosis', 'Prevalence', 'Seroprevalence' 'Babesia caballi', 'Theileria equi', 'Tick-borne', 'Equids', 'Horses', 'Donkeys', 'Equines' and 'Mules'. Keywords were used individually and in combination with 'OR' and/or 'AND' operators. The titles of the articles were scanned, and relevant articles were downloaded. In addition, the reference list of the searched articles was also screened for relevant studies.

### Inclusion and exclusion criteria


After a review of titles and abstracts, the selected studies were screened further by a detailed review of the full text. Articles that were included in the study had to fulfil all the following criteria: (1) original research articles without geographical limitation (global); (2) the full texts were available; (3) the publication was in English; (4) conducted between 1 January 1990 and 25 February 2019; (5) study design was cross-sectional/prevalence study; (6) the diagnostic method was clearly stated; (7) the geographical location of the study was clearly stated; (8) the species of equid was clearly stated; (9) the number of positive cases and sample size were provided; (10) the species of the piroplasms was clearly identified; (11) the study screened for both *B. caballi* and *T. equi*; and (12) the sample size was at least a minimum of 50 equids. Any study that did not fulfil the criteria stated was excluded. Eligibility and inclusion as well as data extraction were carried out by two trained investigators working independently. At the end of the search and screening, the investigators met and compared findings. No attempt was made to contact the authors of the original manuscripts for any additional information or retrieval of unpublished studies.

### Subgroup analyses

We performed several subgroup analyses to study the independent effects of infection of *B. caballi* and *T. equi* on several risk factors including age, sex, publication years, diagnostic methods, species of equid and continent and/or region. In the estimation of the overall pooled prevalence, we used the individual infection rates reported for all the eligible studies that were arrived at using either microscopy, molecular or serological technique. Where more than one technique was used, we used the data for serological technique (IFAT/ELISA/ICT) ahead of molecular test and microscopy. Due to a small number of sample size for mules, this was excluded from the subgroup analysis. We combined data from both North and South America as a single subgroup called the Americas.

### Data extraction and analysis

From the eligible studies, data extracted included first author surname; publication year; sample size; number of positives; country of study; diagnostic method; the species of piroplasms and age; sex and species of equids. Data collected were entered into spreadsheets. Graph-Pad Prism version 5.0 was used for preliminary analysis. Meta-analysis was conducted with Comprehensive Meta-Analysis Version 3.0. For each of the eligible studies, the



**Fig. 1.** Flow diagram showing the selection process of eligible studies according to the PRISMA guidelines.

prevalence was calculated as a percentage by expressing positive cases to sample size. Pooled prevalence and their 95% confidence interval (CI) were determined using MedCalc® statistical software. The prevalence's estimates as well as the *P* value and 95% CI were obtained using a random-effects model (Hedges and Vevea, 1998). Cochran's heterogeneity (*Q*) within studies as well as percentage variation in prevalence ( $I^2$ ) was evaluated using the Cochran's *Q*-test. Heterogeneity was described as low, moderate or high depending on if  $I^2$  was  $\leq 25$ , 50 or  $\geq 75\%$ , respectively (Higgins and Thompson, 2002). Publication bias was evaluated using the Egger's regression intercept (Egger *et al.*, 1997). The effect size and corresponding CI for each subgroup was calculated and expressed on forest plots. Furthermore, to determine the source of heterogeneity within subgroups, e.g. sex, age, geographical regions, diagnostic technique and years of study, meta-regression analyses were performed.

## Results

### Search results and eligible studies

Following a search on the six databases, about 2054 relevant published materials were identified and retrieved. Following the review of their titles, abstracts and duplicates, a total of 1901 studies were excluded. The remaining 153 studies were further screened for eligibility. Studies ( $n = 87$ ) were excluded for failure to identify the parasite to species level ( $n = 18$ ); unidentified locations ( $n = 6$ ); inconsistent data ( $n = 25$ ); studies with sample size below 50 ( $n = 8$ ); non-availability of full text ( $n = 23$ ); and incomplete information on the number of samples tested ( $n = 7$ ). A total of 66 studies (Table 1) were eligible and subsequently used for the meta-analysis (Fig. 1).

The study parameters included the species of piroplasm; single or mixed infection, equine sex; equine species; equine age; diagnostic technique employed, region(s) of the world and year of publication. All eligible studies were conducted between 1 January 1990 and 25 February 2019. All studies included peer-

**Table 1.** List and characteristics of 66 eligible studies included in the meta-analysis

Study reference	Country	Host type	Sample size	<i>B. caballi</i> No. of positives; prevalence %	<i>T. equi</i> No. of positives; prevalence %	Mixed infection No. of positives; prevalence %
Gizachew <i>et al.</i> (2013)	Ethiopia	Donkey	395	52; 13.2	220; 55.7	39; 9.9
Gummow <i>et al.</i> (1996)	South Africa	Horses	176	58; 32.9	95; 53.9	–
Hawkins <i>et al.</i> (2015)	Kenya	Donkey	71	0; 0	67; 30.9	–
Mahmoud <i>et al.</i> (2016)	Egypt	Horse/donkey	139	31; 22.3	37; 26.6	–
Motloang <i>et al.</i> (2008)	South Africa	Horses	138	51; 36.9	97; 70.3	–
Oduori <i>et al.</i> (2015)	Kenya	Donkeys	314	0; 0	255; 81.2	–
Ros-García <i>et al.</i> (2013)	Tunisia	Horses	104	1; 0.9	11; 10.6	1; 0.9
Salim <i>et al.</i> (2008)	Sudan	Horses	158	7; 4.4	80; 50.6	–
Salim <i>et al.</i> (2013)	Sudan	Horses/donkey	308	0; 0	121; 39.3	–
Sanusi <i>et al.</i> (2014)	Nigeria	Horses	400	60; 15.0	8; 2.0	4; 1.0
Turaki <i>et al.</i> (2014)	Nigeria	Horses	240	6; 2.5	94; 39.2	–
Xu <i>et al.</i> (2003)	China	Horses	111	36; 32.4	38; 34.2	–
Wang <i>et al.</i> (2014)	China	Horses	1990	1018; 51.2	229; 11.5	–
Sumbria <i>et al.</i> (2016)	India	Horses/donkeys/ mules	180	113; 62.8	133; 73.9	–
Ybañez <i>et al.</i> (2018)	Philippines	Horses	105	2; 1.9	23; 21.9	–
Seo <i>et al.</i> (2011)	Korea	Horses	184	0; 0	2; 1.1	–
Rüegg <i>et al.</i> (2007)	Mongolia	Horses	499	328; 65.7	393; 78.8	–
Nugraha <i>et al.</i> (2018)	Indonesia	Horses	235	15; 6.4	5; 2.1	–
Sevinc <i>et al.</i> (2008)	Turkey	Horses	481	4; 0.8	78; 16.2	7; 1.5
Munkhjargal <i>et al.</i> (2013)	Mongolia	Horses	250	129; 51.6	49; 19.6	26; 10.4
Kizilarlan <i>et al.</i> (2015)	Turkey	Horses	203	4; 1.9	6; 2.9	–
Karatepe <i>et al.</i> (2009)	Turkey	Horses	125	12; 9.6	16; 12.8	5; 4.0
Kamyinkird <i>et al.</i> (2016)	Thailand	Horses/mules	240	12; 5.0	21.0; 8.8	0; 0.0
Ikadai <i>et al.</i> (2002)	Japan	Horses	2019	109; 5.4	44; 2.2	–
Hussain <i>et al.</i> (2014)	India	Horses/donkey/ mules	430	93; 21.6	177; 41.2	44; 10.2
Akkan <i>et al.</i> (2003)	Turkey	Horses	110	5; 4.6	71; 64.6	1; 0.9
Al-Obaidi <i>et al.</i> (2016)	Malaysia	Horses	306	193; 63.07	157; 51.3	105; 34.3
Acici <i>et al.</i> (2008)	Turkey	Horses/donkey/ mules	153	53; 34.6	33; 21.6	8; 5.2
Guven <i>et al.</i> (2017)	Turkey	Horses	125	0; 0	11; 8.8	–
Cruz-Flores <i>et al.</i> (2010)	Philippines	Horses	104	45; 43.3	6; 5.8	12; 11.5
Boldbaatar <i>et al.</i> (2005)	Mongolia	Horses	254	102; 40.2	185; 72.8	78; 30.7
Chahan <i>et al.</i> (2006)	China	Donkey	93	36; 38.7	9; 9.7	2; 2.2
Avarzed <i>et al.</i> (1997)	Mongolia	Horses	110	93; 84.5	97; 88.18	–
Sigg <i>et al.</i> (2010)	Switzerland	Horses	689	10; 1.5	30; 4.4	10; 1.5
Piantedosi <i>et al.</i> (2014)	Italy	Donkey	203	72; 35.5	90; 44.3	46; 22.7
Moretti <i>et al.</i> (2010)	Italy	Horses	412	74; 17.9	51; 12.4	157; 38.1
Kouam <i>et al.</i> (2010)	Greece	Horses/mules	544	12; 2.2	60; 11.0	9.0; 1.7
Guidi <i>et al.</i> (2015)	France	Horses	443	57; 12.9	257; 58.01	36; 8.1
Grandi <i>et al.</i> (2011)	Italy	Horses	294	1; 0.3	24; 8.2	0; 0.0
García-Bocanegra <i>et al.</i> (2013)	Spain	Horses/donkey/ mules	537	61; 11.4	270; 50.3	45; 8.4

(Continued)

Table 1. (Continued.)

Study reference	Country	Host type	Sample size	<i>B. caballi</i> No. of positives; prevalence %	<i>T. equi</i> No. of positives; prevalence %	Mixed infection No. of positives; prevalence %
Gallusová et al. (2014)	Romania	Horses	178	8; 4.5	69; 38.8	–
Cortés et al. (2017)	Spain	Horses	3100	643; 20.7	1381; 44.6	398; 12.8
Camino et al. (2018)	Spain	Horses	536	25; 4.7	117; 21.8	11; 2.1
Camacho et al. (2005)	Spain	Horses	60	17; 28.3	24; 40.0	12; 20
Butler et al. (2012)	Netherland	Horses	300	9; 3.0	5; 1.7	3; 1.0
Bartolome et al. (2016)	Italy	Horses	673	60; 8.9	268; 39.8	–
Abedi et al. (2014)	Iran	Horses	100	2; 2.0	48; 48.0	3, 3.0
Abedi et al. (2015)	Iran	Donkey	106	0; 0	54; 50.9	–
Abutarbush et al. (2012)	Jordan	Horses	253	0; 0	37; 14.6	–
Alanazi et al. (2012)	Saudi Arabia	Horses	241	18; 7.5	25; 10.4	7; 2.9
Jaffer et al. (2010)	United Arab Emirates	Horses	105	11; 10.5	35; 33.3	13; 12.4
Kakekhani et al. (2017)	Iran	Horses	186	0; 0	1; 0.5	–
Qablan et al. (2013)	Jordan	Horses/donkey	288	21; 7.3	54; 18.8	–
Malekifard et al. (2014)	Iran	Horses	240	14; 5.8	26; 10.8	4; 1.67
Posada-Guzmán et al. (2015)	Costa Rica	Horses	285	90; 31.6	115; 40.4	81; 28.4
Cantú-Martínez et al. (2012)	Mexico	Horses	248	41; 16.5	85; 34.3	27; 10.9
Díaz-Sánchez et al. (2018)	Cuba	Horses	100	25; 25.0	73; 73.0	20; 20.0
Asgarali et al. (2007)	Trinidad	Horses	93	64; 68.8	31; 33.3	18; 19.4
Vieira et al. (2013)	Brazil	Horses	198	137; 69.2	155; 78.3	99; 50.0
Vieira et al. (2018)	Brazil	Horses	90	5; 5.6	17; 18.9	2; 10.5
Rosales et al. (2013)	Venezuela	Horses	694	161; 23.2	97; 13.9	90; 12.9
Mujica et al. (2011)	Venezuela	Horses	360	254; 70.6	181; 50.3	128; 35.6
Machado et al. (2012)	Brazil	Donkey	88	82; 93.2	65; 73.9	59; 67.0
Kerber et al. (2009)	Brazil	Horses	582	405; 69.6	155; 26.6	–
Heuchert et al. (1999)	Brazil	Horses	740	505; 68.2	211; 28.5	–
Heim et al. (2007)	Brazil	Horses	487	443; 90.9	404; 82.9	–

reviewed journal articles and no attempt was made to check dissertations or thesis. Studies were from Africa ( $n = 11$ ), Asia ( $n = 20$ ), Europe ( $n = 14$ ), the Middle East ( $n = 8$ ) and Americas ( $n = 13$ ) (Table 1).

### Pooled prevalence estimates

An overall pooled prevalence estimate (PPE) due to EP caused by *B. caballi* was 22.3% (95% CI 21.7–22.8) from the 66 eligible studies that reported 5348 cases in over 24 041 equids screened (Table 2). Individual point estimates were determined for studies reporting the occurrence of *B. caballi* (Fig. 2). Furthermore, a significant difference between study heterogeneity was observed ( $Q = 8531.7$ ,  $I^2 = 99.9$ , 95% CI 99.2–99.3,  $P < 0.0001$ ). The overall PPE due to *T. equi* was 29.4% (95% CI 28.7–30.0) from the 66 eligible studies with 7074 cases in 24 041 equids screened (Table 3). Individual point estimates were determined for the 66 eligible studies with regards to infection with *T. equi* (Fig. 3). Finally, 43 studies reported mixed infection with an overall PPE of 11.8% (95% CI 11.3–12.3) of the 16 250 equid samples.

### According to region

The Americas had the highest prevalence of 47.9% (95% CI 45.8–49.9%,  $Q = 1583.3$ ,  $I^2 = 99.3$ ,  $P < 0.0001$ ) while the lowest prevalence was in the Middle East (4.8%; 95% CI 3.7–5.8%,  $Q = 92.1$ ,  $I^2 = 92.4$ ,  $P < 0.0001$ ) (Table 2 and Fig. 4). Although Asia region had the highest number of eligible studies examined within the period ( $n = 22$ ) as well as the largest number of animals ( $n = 8307$ ; 2124 cases), the prevalence was (25.6%; 24.5–26.6%). Similarly, the prevalence due to *T. equi* was highest in the Americas (46.5%; 95% CI 44.5–48.6%,  $Q = 1131.5$ ,  $I^2 = 99.0$ ,  $P < 0.0001$ ) compared to the Middle East (18.1%; 95% CI 16.1–20.2%,  $Q = 211.7$ ,  $I^2 = 96.7%$ ,  $P < 0.0001$ ) and Asia (18.1%, 95% CI 17.2–18.9,  $Q = 2379.4$ ,  $I^2 = 99.1%$ ,  $P < 0.0001$ ) (Table 3).

### According to sex

Infection due to both piroplasms was slightly higher in males. For infection due to *B. caballi*, the PPE in male equids was 5.5% (95% CI 4.9–6.0%,  $Q = 813.7$ ,  $I^2 = 98.4$ ,  $P < 0.0001$ ) compared with 4.5% (95% CI 4.0–4.9%,  $Q = 549.7$ ,  $I^2 = 97.6$ ,  $P < 0.0001$ ) in females

**Table 2.** Pooled prevalence and risk factors associated with *Babesia caballi* infection in equines 1990–2019

Risk factors	Number of studies	Pooled prevalence estimates			Measure of heterogeneity		
		Sample size	No of positives	Prevalence 95% CI (%)	Q	I <sup>2</sup> (95% CI)	Q-P
Overall							
<i>B. caballi</i>	66	24 041	5348	22.3 (21.7–22.8)	8531.7	99.9 (99.2–99.3)	0.0001
Region							
Africa	11	2443	266	10.9 (9.90–12.1)	416.5	97.6 (96.8–98.2)	0.0001
Asia	22	8307	2124	25.6 (24.5–26.6)	2915.7	99.3 (99.2–99.4)	0.0001
Middle East	8	1519	73	4.8 (3.7–5.8)	92.1	92.4 (87.4–95.4)	0.0001
Europe	13	7807	987	12.6 (11.9–13.4)	827.7	98.6 (98.2–98.9)	0.0001
Americas	12	3965	1898	47.9 (45.8–49.9)	1583.3	99.3 (99.2–99.4)	0.0001
Sex							
Male	14	6952	383	5.5 (4.9–6.0)	813.7	98.4 (97.9–98.7)	0.0001
Female	14	6952	312	4.5 (4.0–4.9)	549.7	97.6 (96.9–98.2)	0.0001
Age							
<5	11	2489	342	13.7 (12.4–15.1)	554.7	98.2 (97.6–98.6)	0.0001
>5	11	2489	283	11.4 (10.1–12.6)	420.9	97.6 (96.8–98.2)	0.0001
Diagnostic technique							
Microscopy	23	5129	212	4.1 (3.6–4.7)	459.8	95.2 (93.9–96.3)	0.0001
ELISA	26	11 006	3014	27.4 (26.5–28.3)	4779.3	99.5 (99.4–99.5)	0.0001
IFAT	26	10 230	2344	22.9 (22.0–23.8)	2849.9	99.1 (99.0–99.2)	0.0001
PCR	30	6143	515	8.4 (7.7–9.1)	1039.1	97.2 (96.6–97.7)	0.0001
Species							
Horses	56	21 358	4870	22.8 (21.2–23.4)	77 805	99.3 (99.2–99.4)	0.0001
Donkey	11	2148	333	15.5 (14.4–17.7)	807.4	98.9 (98.6–99.1)	0.0001
Years of study							
1990–1999	3	1026	277	26.9 (23.9–30.0)	64.7	96.9 (93.7–98.5)	0.0001
2000–2009	15	5363	1589	29.6 (28.2–31.0)	3236.6	99.6 (99.5–99.6)	0.0001
2010–2019	48	17 652	3482	19.7 (19.1–20.4)	4976.3	99.1 (98.9–99.2)	0.0001

ELISA, enzyme linked immunosorbent assay; IFAT, Immunofluorescence Antibody Test; PCR, polymerase chain reaction.

(Table 2). A similar observation was also noted in respect to infection with *T. equi*, males had a prevalence of 16.9% (95% CI 15.5–18.5%,  $Q = 1003.3$ ,  $I^2 = 98.2$ ) compared with 16.4% in females (95% CI 15.6–17.3%,  $Q = 682.7$ ,  $I^2 = 97.4$ ,  $P < 0.0001$ ) (Table 3 and Fig. 5).

#### According to age

For *T. equi* infections, the prevalence was slightly higher for those <5 years (16.9%; 95% CI 15.5–18.5%,  $Q = 397.2$ ,  $I^2 = 97.5$ ,  $P < 0.0001$ ) compared with those >5 years (16.4%; 95% CI 14.9–17.9%,  $Q = 229.9$ ,  $I^2 = 95.7$ ,  $P < 0.0001$ ) (Table 3 and Fig. 6). A similar observation was noted in infection due to *B. caballi*, with prevalence higher in those animals <5 years (13.7%; 95% CI 12.4–15.1%,  $Q = 554.7$ ,  $I^2 = 98.2$ ,  $P < 0.0001$ ) compared with equids >5 years (11.4%; 95% CI 10.1–12.6%,  $Q = 420.9$ ,  $I^2 = 97.6$ ,  $P < 0.0001$ ) (Table 2, S1-Supplementary file).

#### According to diagnostic technique

The PPE for different *B. caballi* diagnostic methods indicated that ELISA tests were associated with the highest exposure (27.4%;

95% CI 26.5–28.3%,  $Q = 4779.3$ ,  $I^2 = 99.5$ ,  $P < 0.0001$ ) (Table 2, Fig. 7), followed by IFAT (22.9%; 95% CI 22.0–23.8%,  $Q = 2849.9$ ,  $I^2 = 99.1$ ,  $P < 0.0001$ ), PCR (8.4%; 95% CI 7.7–9.1%) and microscopy (4.1%; 95% CI 3.6–4.7%,  $Q = 459.8$ ,  $I^2 = 95.2$ ,  $P < 0.0001$ ) (Table 2). The PPE for *T. equi* using different diagnostic methods indicates that IFAT technique was associated with the highest exposure (41.1%; 95% CI 39.9–42.3%,  $Q = 3167.4$ ,  $I^2 = 99.2$ ,  $P < 0.0001$ ) (Table 3, Fig. 8), followed by PCR (31.6%; 95% CI 30.2–32.9%,  $Q = 2610.6$ ,  $I^2 = 98.9$ ,  $P < 0.0001$ ), ELISA (21.9%; 95% CI 21.0–22.7%,  $Q = 3037.3$ ,  $I^2 = 99.2$ ,  $P < 0.0001$ ) and microscopy (8.1%; 95% CI 7.4–8.9%,  $Q = 863.0$ ,  $I^2 = 97.5$ ,  $P < 0.0001$ ) (Table 3).

#### According to equid species

Infection due to *B. caballi* was higher in horses (*Equus caballus*) (22.8%; 95% CI 21.2–23.4%,  $Q = 77 805$ ,  $I^2 = 99.3$ ,  $P < 0.0001$ ) as compared to donkeys (*Equus asinus*) (15.5%, 95% CI 14.4–17.7%,  $Q = 807.4$ ,  $I^2 = 98.9$ ,  $P < 0.0001$ ) (Table 2, S2-Supplementary file). Infection due to *T. equi* was higher in donkeys (50.9%; 95% CI 48.1–53.9%,  $Q = 379.1$ ,  $I^2 = 97.4$ ,  $P < 0.0001$ ) as compared to horses (27.8%; 95% CI 27.1–28.5%,  $Q = 7478.8$ ,  $I^2 = 99.3$ ,  $P < 0.0001$ ) (Table 3).






Fig. 2. Forest plot of the prevalence estimates of *Babesia caballi* in equids globally from 1990 and 2019. Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.

According to years of study

The time span of 1990–1999 had a higher PPE of *B. caballi* (26.9%;  $n = 3$ ,  $Q = 64.7$ ,  $I^2 = 96.9$ ,  $P < 0.0001$ ) as compared to the period of 2010–2019 (19.7%;  $n = 46$ ,  $Q = 4976.3$ ,  $I^2 = 99.1$ ,  $P < 0.0001$ ) (Table 2; S3-Supplementary file). Similarly, the time span of 1990–1999 had a higher *T. equi* PPE (59.7%;  $Q = 202.1$ ,  $I^2 = 99.0$ ,  $P < 0.0001$ ) as compared to the period of 2010–2019 (29.3%;  $Q = 4399.2$ ,  $I^2 = 98.9$ ,  $P < 0.0001$ ) (Table 3).

Heterogeneity and publication bias

Results from our study showed strong heterogeneity between the selected studies which were largely influenced by the number of studies and diagnostic techniques. No publication bias was observed in subgroup analysis due to diagnostic technique, age, sex and region. Major publication bias was observed only in the overall PPEs due to *B. caballi* [Egger’s intercept ( $B_0$ ) = -4.79,  $P = 0.003$ ] and *T. equi* [Egger’s intercept ( $B_0$ ) = -3.67,  $P < 0.05$ ].

Discussion

It is evident that EP is widespread and endemic in various regions of the world. The PPE for *T. equi* infection was 29.4%. This estimate is relatively similar to the prevalence of 30.9% reported in Kenya (Hawkins et al., 2015) and 26.6% in Brazil and Egypt (Kerber et al., 2009; Mahmoud et al., 2016). Higher prevalences have been reported in several studies across different regions of the world (Gummow et al., 1996; Garcia-Bocanegra et al., 2013; Sumbria et al., 2016; Diaz-Sánchez et al., 2018; Onyiche et al., 2020). The PPE due to *B. caballi* was 22.3%, lower than that of *T. equi*. Generally, the prevalence of *T. equi* has been found to be higher than that of *B. caballi* likely due to the fact that *T. equi*-infected animals remain infected for life (Rüegg et al., 2007). Another possible reason for the differences in the PPE between the two pathogens could be due to differences in vector distribution (Salim et al., 2008). Mixed infection of the two piroplasmids has been reported in different studies and is unconnected with the presence of the tick vectors responsible for the transmission of both pathogens within the same geographical area infesting their host.

**Table 3.** Pooled prevalence and risk factors associated with *Theileria equi* infection in equines 1990–2019

Risk factors	Number of studies	Pooled prevalence estimates			Measure of heterogeneity		
		Sample size	No of positives	Prevalence 95% CI (%)	Q	I <sup>2</sup> 95% CI	Q-P
Overall							
<i>T. equi</i>	66	24 041	7074	29.4 (28.7–30.0)	8265.4	99.2 (99.2–99.3)	0.0001
Region							
Africa	11	2443	1085	44.4 (41.9–46.9)	1583.3	99.3 (99.2–99.4)	0.0001
Asia	22	8307	1503	18.1 (17.2–18.9)	2379.4	99.1 (98.9–99.2)	0.0001
Middle East	8	1519	275	18.1 (16.1–20.2)	211.7	96.7 (95.1–97.8)	0.0001
Europe	13	7807	2366	30.3 (29.2–31.5)	1786.2	99.3 (99.2–99.4)	0.0001
Americas	12	3965	1845	46.5 (44.5–48.6)	1131.5	99.0 (98.8–99.2)	0.0001
Sex							
Male	19	8449	1435	16.9 (16.2–17.8)	1003.3	98.2 (97.8–98.5)	0.0001
Female	19	8449	1391	16.5 (15.6–17.3)	682.7	97.4 (96.7–97.9)	0.0001
Age							
<5	11	2489	423	16.9 (15.5–18.5)	397.2	97.5 (96.6–98.1)	0.0001
>5	11	2489	409	16.4 (14.9–17.9)	229.9	95.7 (93.8–96.9)	0.0001
Diagnostic technique							
Microscopy	23	5129	418	8.1 (7.4–8.9)	863.0	97.5 (96.9–97.9)	0.0001
ELISA	26	11 006	2406	21.9 (21.0–22.7)	3037.3	99.2 (99.1–99.3)	0.0001
IFAT	26	10 230	4209	41.1 (39.942.3)	3167.4	99.2 (99.1–99.3)	0.0001
PCR	30	6143	1940	31.6 (30.2–32.9)	2610.6	98.9 (98.7–99.0)	0.0001
Species							
Horses	56	21 358	5932	27.8 (27.1–28.5)	7478.8	99.3 (99.2–99.3)	0.0001
Donkey	11	2148	1095	50.9 (48.1–53.9)	379.1	97.4 (96.4–98.1)	0.0001
Years of study							
1990–1999	3	1026	614	59.8 (55.3–64.4)	202.1	99.0 (98.4–99.4)	0.0001
2000–2009	15	5363	1573	29.3 (27.9–30.7)	2885.5	99.5 (99.4–99.6)	0.0001
2010–2019	48	15 809	4635	29.3 (28.5–30.1)	4399.2	98.9 (98.9–99.1)	0.0001

ELISA, enzyme linked immunosorbent assay; IFAT, Immunofluorescence Antibody Test; PCR, polymerase chain reaction.

Diagnosis of EP can be achieved by either the use of direct or indirect methods (Abedi *et al.*, 2015). The gold standard for piroplasm's diagnosis is microscopy but poor sensitivity during low parasitaemia limits its use (Böse *et al.*, 1995). Microscopy and PCR techniques are considered direct methods as they indicate active infection and serological assays are considered indirect as they detect the presence of antibodies which is an indicator of exposure rather than an indication of infection status (Abedi *et al.*, 2015). We observed that the IFAT method detected higher exposure to *T. equi*, and ELISA detected higher exposure of *B. caballi*. Infection with *B. caballi* is transient and best detected during the acute phase of the infection due to low parasitaemia associated with it. Competitive ELISA (cELISA) based on *rap-1* demonstrated higher exposure to antibodies of *B. caballi* as observed in the Venezuelan isolates (Rosales *et al.*, 2013). Nonetheless, the *rap-1* region is believed to be highly polymorphic as demonstrated in some epidemiological studies with no positive samples detected using cELISA in Egypt, South Africa and Israel (Bhoora *et al.*, 2010; Rapoport *et al.*, 2014; Mahmoud *et al.*, 2016). Due to variation in the *rap-1* gene between geographically diverse isolates with differences in their amino acid sequences, this has led to inconsistency in the

commercial *rap-1* cELISA assays for the detection of *B. caballi* strain (Idoko *et al.*, 2020). Therefore, the commercial cELISA for *B. caballi* is problematic and can lead to a high number of false negatives hence leading to lack of positive samples in some region of the world (Bhoora *et al.*, 2010).

On the other hand, infection with *T. equi* is often lifelong and exposed equids seroconvert after a brief period of infection, usually within 14–16 days. According to the OIE (2005), horses deemed for export must have a negative result to EP when screened using either IFAT or ELISA techniques which remain to be recommended diagnostic methods based on the OIE manual for diagnostic tests and vaccines for terrestrial manual. Therefore, it is not surprising that serological techniques (IFAT and ELISA) were the most efficient in determining the exposure of equids to EP.

Furthermore, microscopy was associated with low prevalence for both pathogens. In several studies, piroplasms were not detected in blood smears but were detected using other techniques such as PCR and serology on same samples that were initially negative (Abutarbush *et al.*, 2012; Munkhjargal *et al.*, 2013). However, the OIE diagnostic manual recommends that microscopic examination to be used in some situations (OIE,




Fig. 3. Forest plot of the prevalence estimates of *Theileria equi* in equids globally from 1990 and 2019. Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.






Fig. 4. Forest plot of the prevalence estimates due to equine piroplasm in the Americas. Prevalence due to *T. equi* in the Americas is illustrated in (A) while estimates due to *B. caballi* are shown in (B). Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.





**Fig. 5.** Forest plot of the prevalence estimates of *Theileria equi* in male equids (A) compared with females (B) from 1990 and 2019. Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.


2018). The method continues to be applied in resource-poor countries despite its disadvantages of poor sensitivity during low parasitaemia.

The Americas had the highest PPE for both *B. caballi* and *T. equi* infection while the Middle East had the least estimates for both pathogens. The difference in prevalences among geographical regions may be due to the sensitivity of the various diagnostic tests that have been used in the different epidemiological studies; abundance and occurrence of competent tick vectors; husbandry system; activity of the equids; effectiveness of the control measures instituted at the farm and national levels (Kouam *et al.*, 2010). In some parts of Africa, the prevalence of EP caused by *T. equi* was high (Motloang *et al.*, 2008; Hawkins *et al.*, 2015; Oduori *et al.*, 2015). Also, few epidemiological studies have been conducted in the continent despite a handful of equestrian sports and traditional local festival where the use of horses is common. It is therefore expedient that more testing be conducted which is necessary before the institution of treatment and control.


The PPE for both *B. caballi* and *T. equi* indicates that these parasites are more prevalent in males as compared to females.

Individual studies have reported contrasting observations (Sigg *et al.*, 2010; Abedi *et al.*, 2014). Nevertheless, the difference between sexes has not been significant in majority of the individual studies. However, males may have higher tick exposure and immune-suppression due to stress arising from strenuous physical activities (Vieira *et al.*, 2013). This may consequently lead to higher infection rates in males. Furthermore, younger equids (<5 years) had a slightly higher PPE for both pathogens compared to the older ones (>5 years). Generally, young horses may reside longer in the fields and consequently, more exposure to tick vectors which increases their likelihood of infection with tick-borne pathogens as compared to adults.

A majority of EP studies focused on horses. The high interest in research-related studies on horses compared to other equids could be due to their high economic value compared with donkeys and other equids (Onyiche *et al.*, 2019). *Theileria equi* PPE was higher in the donkeys as compared to horses. Donkeys are asymptomatic carriers of piroplasmids with low parasitaemia and positive antibody titres throughout their lifetime (Balkaya *et al.*, 2010). However, PPE was higher in the infection of horses with *B. caballi*.




**Fig. 6.** Forest plot of the prevalence estimates of *Theileria equi* in equids <5 years old (A) compared with those above 5 years old (B) from 1990 and 2019. Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.



**Fig. 7.** Forest plot of the prevalence estimates of *Babesia caballi* using ELISA as a diagnostic technique in equids from 1990 and 2019. Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.

In the period spanning 2010–2019, the global PPE due to *T. equi* has remained stable at 29.3% down from the earlier 59.8% during the period 1990–1999. Similarly, the global PPE due to

*B. caballi* has decreased from 26.9% between the period spanning 1990 and 1999 to 19.7% covering the period 2010–2019. The decrease in the prevalence could be attributed to a better



**Fig. 8.** Forest plot of the prevalence estimates of *Theileria equi* using IFAT as a diagnostic technique in equids from 1990 and 2019. Note: The squares show the individual point estimate. The diamond at the base indicates the pooled estimates from the total studies.

understanding of the epidemiology of the parasite and more efficient control of the vectors. Furthermore, the testing of equids before their export as recommended by the OIE may have further helped to decrease the burden of the disease and help in the curtailment of the spread of the disease between different regions of the world. Additionally, we speculate that the decrease in EP over the time period could also be attributed to differences in the diagnostic techniques over the years.

We have attempted to present a systematic review and meta-analysis of exposure of equids to EP to gain more insight on the global epidemiology of the disease. Due to the pooling of data, we acknowledge that this will lead to significant heterogeneity as a result of the differences in the characteristics among the identified studies despite the use of random-effect model. Some of the limitations include paucity of data which varies across region, publication bias, uneven distribution of prevalence across countries and low sample size in some studies. Therefore, results must be interpreted with caution as apparent prevalence may vary from the actual estimates. Nevertheless, we believe that our report is very close to true estimates of the global exposure of equids to agents of EP.

## Conclusion

To the best of our knowledge, this study represents the first systematic review and meta-analysis on the global exposure of equids to agents of EP to better understand the distribution of the disease across the world in the last three decades. All eligible studies incorporated in this systematic review were cross-sectional, further studies incorporating case-control and cohort studies will be required to expand our knowledge horizon on the risk factors and exposures to this disease. Lastly, they are urgent needs for discovering candidate antigens for improved diagnostic tools for the control of equine babesiosis most especially in Africa

and the Middle East. Therefore, further studies to fill in this knowledge gap are expedient.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182020001407>

**Acknowledgements.** The first author was financially supported by the North West University (NWU) Post graduate student bursary. We thank Cornelia Silaghi, Ana Vasic and Cristian Raileanu for their helpful suggestions on the manuscript.

**Financial support.** This research work did not receive any specific grant from funding agencies. The first author was financially supported by the North West University (NWU) Post graduate student bursary. The Unit for Environmental Sciences and Management (NWU) also financially supported the first author's scientific travel to Japan through an international travel award.

**Conflict of interest.** No conflict of interest exists among the authors.

**Ethical standards.** Not applicable.

## References

- Abedi V, Razmi G, Seifi H and Naghibi A (2014) Molecular and serological detection of *Theileria equi* and *Babesia caballi* infection in horses and ixodid ticks in Iran. *Ticks and Tick-Borne Diseases* **5**, 239–244.
- Abedi V, Razmi G, Seifi H and Naghibi A (2015) Molecular detection of equine piroplasms in donkeys (*Equus asinus*) in North Khorasan province, Iran. *Iranian Journal of Veterinary Research* **16**, 202.
- Abutarbush SM, Alqawasmeh DM, Mukbel RM and Al-Majali AM (2012) *Equine babesiosis*: seroprevalence, risk factors and comparison of different diagnostic methods in Jordan. *Transboundary and Emerging Disease* **59**, 72–78.
- Acici M, Umur S, Guvenc T, Arslan HH and Kurt M (2008) Seroprevalence of equine babesiosis in the Black Sea region of Turkey. *Parasitology International* **57**, 198–200.



- Akkan HA, Karaca M, Tutuncu M, Deger S, Keles I and Agaoglu Z (2003) Serologic and microscopic studies on babesiosis in horses in the eastern border of Turkey. *Journal of Equine Veterinary Science* 5, 181–183.
- Al-Obaidi QT, Arshad MM, Al-Sultan II, Azlinda AB and Mohd-Azam KGK (2016) Comparison between microscopic examination and competitive ELISA for diagnosis of equine piroplasmosis in Kelantan. *Malaysia. Malaysia Journal of Veterinary Research* 7, 23–29.
- Alanazi AD, Alyousif MS and Hassieb MM (2012) Seroprevalence study on *Theileria equi* and *Babesia caballi* antibodies in horses from central province of Saudi Arabia. *Journal of Parasitology* 98, 1015–1017.
- Asgarali Z, Coombs DK, Mohammed F, Campbell MD and Caesar E (2007) A serological study of *Babesia caballi* and *Theileria equi* in thoroughbreds in Trinidad. *Veterinary Parasitology* 144, 167–171.
- Avarzed A, De Waal DT, Igarashi I, Saito A, Oyamada T, Toyoda Y and Suzuki N (1997) Prevalence of equine piroplasmosis in Central Mongolia. *Onderstepoort Journal of Veterinary Research* 64, 141–145.
- Ayala-Valdovinos MA, Lemus-Flores C, Galindo-García J, Sánchez-Chipres D, Duifhuis-Rivera T, Anguiano-Estrella R, Bañuelos-Pineda J and Rodríguez-Carpena JG (2014) Nested PCR detection of *Theileria equi* infection and frequency in horses imported into Mexico. *Journal of Animal Veterinary Advances* 13, 859–863.
- Balkaya I, Utuk AE and Piskin FC (2010) Prevalence of *Theileria equi* and *Babesia caballi* in donkeys from Eastern Turkey in winter season. *Pakistan Veterinary Journal* 30, 245–246.
- Bartolome LED, Roberto N, Vincenzo V, Francesca I, Antonella C, Luca A and Teresa SM (2016) *Babesia caballi* and *Theileria equi* infections in horses in Central-Southern Italy: Sero-molecular survey and associated risk factors. *Ticks and Tick-Borne Diseases* 7, 462–469.
- Bhoora R, Quan M, Zweygarth E, Guthrie AJ, Prinsloo SA and Collins NE (2010) Sequence heterogeneity in the gene encoding the rhoptry-associated protein-1 (RAP-1) of *Babesia caballi* isolates from South Africa. *Veterinary Parasitology* 169, 279–288.
- Boldbaatar D, Xuan X, Battsetseg B, Igarashi I, Battur B, Batsukh Z and Fujisaki K (2005) Epidemiological study of equine piroplasmosis in Mongolia. *Veterinary Parasitology* 127, 29–32.
- Böse R, Jorgensen WK, Dalgliesh RJ, Friedhoff KT and De Vos AJ (1995) Current state and future trends in the diagnosis of babesiosis. *Veterinary Parasitology* 57, 61–74.
- Butler CM, van Oldruitenborgh-Oosterbaan MMS, Stout TA, van der Kolk JH, Van den Wollenberg L, Nielen M and Houwers DJ (2012) Prevalence of the causative agents of equine piroplasmosis in the South West of the Netherlands and the identification of two autochthonous clinical *Theileria equi* infections. *The Veterinary Journal* 193, 381–385.
- Camacho AT, Guitian FJ, Pallas E, Gestal JJ, Olmeda AS, Habela MA and Spielman A (2005) *Theileria (Babesia) equi* and *Babesia caballi* infections in horses in Galicia, Spain. *Tropical Animal Health Production* 37, 293–302.
- Camino E, de la Cruz ML, Dominguez L, Carvajal KA, Fores P, de Juan L and Cruz-Lopez F (2018) Epidemiological situation of the exposure to agents causing equine piroplasmosis in Spanish purebred horses in Spain: seroprevalence and associated risk factors. *Journal of Equine Veterinary Science* 67, 81–86.
- Cantú-Martínez MA, Segura-Correa JC, Silva-Páez ML, Avalos-Ramírez R and Wagner GG (2012) Prevalence of antibodies to *Theileria equi* and *Babesia caballi* in horses from northeastern Mexico. *Journal of Parasitology* 98, 869–871.
- Chahan B, Zhang S, Seo J Y, Nakamura C, Zhang G, Bannai H and Kabeya H (2006) Seroepidemiological evidence for the possible presence of *Babesia (Theileria) equi* and *Babesia caballi* infections in donkeys in western Xinjiang, China. *Journal of Veterinary Medical Science* 68, 753–755.
- Cortés MGM, Fernández-García JL and Martínez-Estélez MÁH (2017) Seroprevalence of *Theileria equi* and *Babesia caballi* in horses in Spain. *Parasite* 24, 14.
- Cruz-Flores MJ, Bata M, Co B, Claveria FG, Verdida R, Xuan X and Igarashi I (2010) Immunochromatographic assay of *Babesia caballi* and *Babesia equi* Laveran 1901 (*Theileria equi* Mehlhorn and Schein, 1998) (Phylum Apicomplexa) infection in Philippine horses correlated with parasite detection in blood smears. *Veterinarski Arhiv* 80, 715–722.
- Díaz-Sánchez AA, Pires MS, Estrada CY, Cañizares EV, del Castillo Domínguez SL, Cabezas-Cruz A and Corona-González B (2018) First molecular evidence of *Babesia caballi* and *Theileria equi* infections in horses in Cuba. *Parasitology Research* 117, 3109–3118.
- Egger M, Smith GD, Schneider M and Minder C (1997) Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315, 629–634.
- Gallusová M, Qablan MA, D'Amico G, Oborník M, Petrželková KJ, Mihalca AD and Modrý D (2014) Piroplasms in feral and domestic equines in rural areas of the Danube Delta, Romania, with survey of dogs as a possible reservoir. *Veterinary Parasitology* 206, 287–292.
- García-Bocanegra I, Arenas-Montes A, Hernández E, Adaszek Ł, Carbonero A, Almeria S and Arenas A (2013) Seroprevalence and risk factors associated with *Babesia caballi* and *Theileria equi* infection in equids. *The Veterinary Journal* 195, 172–178.
- Gizachew A, Schuster RK, Joseph S, Wernery R, Georgy NA, Elizabeth SK and Wernery U (2013) Piroplasmosis in donkeys – a hematological and serological study in Central Ethiopia. *Journal of Equine Veterinary Science* 33, 18–21.
- Grandi G, Molinari G, Tittarelli M, Sasseria D and Kramer LH (2011) Prevalence of *Theileria equi* and *Babesia caballi* infection in horses from northern Italy. *Vector-Borne and Zoonotic Diseases* 11, 955–956.
- Guidi E, Pradier S, Lebert I and Leblond A (2015) Piroplasmosis in an endemic area: analysis of the risk factors and their implications in the control of theileriosis and babesiosis in horses. *Parasitology Research* 114, 71–83.
- Gummow B, De Wet CS and De Waal DT (1996) A sero-epidemiological survey of equine piroplasmosis in the northern and eastern Cape Provinces of South Africa. *Journal of the South African Veterinary Association* 67, 204–208.
- Güven E, Avcioglu H, Deniz A, Balkaya İ, Abay U, Yavuz Ş and Akyüz M (2017) Prevalence and molecular characterization of *Theileria equi* and *Babesia caballi* in jereed horses in Erzurum, Turkey. *Acta Parasitologica* 62, 207–213.
- Hawkins E, Kock R, McKeever D, Gakuya F, Musyoki C, Chege SM and Skilton RA (2015) Prevalence of *Theileria equi* and *Babesia caballi* as well as the identification of associated ticks in sympatric Grevy's zebras (*Equus grevyi*) and donkeys (*Equus africanus asinus*) in northern Kenya. *Journal of Wildlife Diseases* 51, 137–147.
- Hedges LV and Vevea JL (1998) Fixed-and random-effects models in meta-analysis. *Psychological Methods* 3, 486.
- Heim A, Passos LM, Ribeiro MF, Costa-Júnior LM, Bastos CV, Cabral DD and Pfister K (2007) Detection and molecular characterization of *Babesia caballi* and *Theileria equi* isolates from endemic areas of Brazil. *Parasitology Research* 102, 63–68.
- Heuchert CMS, de Giulli Jr V, De Athaide DF, Böse R and Friedhoff KT (1999) Seroepidemiological studies on *Babesia equi* and *Babesia caballi* infections in Brazil. *Veterinary Parasitology* 85, 1–11.
- Higgins JP and Thompson SG (2002) Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* 21, 1539–1558.
- Homer MJ, Aguilar-Delfin I, Telford SR, Krause PJ and Persing DH (2000) Babesiosis. *Clinical Microbiology Reviews* 13, 451–469.
- Hussain MH, Saqib M, Raza F, Muhammad G, Asi MN, Mansoor MK and Jabbar A (2014) Seroprevalence of *Babesia caballi* and *Theileria equi* in five draught equine populated metropolises of Punjab, Pakistan. *Veterinary Parasitology* 202, 248–256.
- Idoko SI, Tirosch-Levy S, Leszkowicz Mazuz M, Mohammed AB, Sikiti Garba B, Wesley ND and Steinman A (2020) Genetic characterization of piroplasms in donkeys and horses from Nigeria. *Animals* 10, 324.
- Ikadai H, Nagai A, Xuan X, Igarashi I, Kamio T, Tsuji N and Fujisaki K (2002) Seroepidemiologic studies on *Babesia caballi* and *Babesia equi* infections in Japan. *Journal of Veterinary Medical Science* 64, 325–328.
- Jaffer O, Abdishakur F, Hakimuddin F, Riya A, Wernery U and Schuster RK (2010) A comparative study of serological tests and PCR for the diagnosis of equine piroplasmosis. *Parasitology Research* 106, 709–713.
- Kakekhani S, Rahbari S, Madani R and Bokaei S (2017) Molecular and microscopic detection of *Theileria equi* and *Babesia caballi* in horses in Kurdistan Province, Iran. *Archives of Razi Institute* 72, 51–55.
- Kamyngkird K, Yangtara S, Desquesnes M, Cao S, Moumouni A, Jittapalpong S and Igarashi I (2016) Seroprevalence of *Babesia caballi* and *Theileria equi* in horses and mules from Northern Thailand. *The Journal of Protozoology Research* 24, 11–17.
- Karatepe B, Karatepe M, Çakmak A, Karaer Z and Ergün G (2009) Investigation of seroprevalence of *Theileria equi* and *Babesia caballi* in



- horses in Nigde province, Turkey. *Tropical Animal Health and Production* **41**, 109–113.
- Kerber CE, Labruna MB, Ferreira F, De Waal DT, Knowles DP and Gennari SM** (2009) Prevalence of equine piroplasmosis and its association with tick infestation in the State of São Paulo, Brazil. *Revista Brasileira de Parasitologia Veterinária* **18**, 1–8.
- Kizilarlan F, Yildirim A, Duzlu O, Inci A, Onder Z and Ciloglu A** (2015) Molecular detection and characterization of *Theileria equi* and *Babesia caballi* in horses (*Equus ferus caballus*) in Turkey. *Journal of Equine Veterinary Science* **35**, 830–835.
- Kouam MK, Kantzoura V, Gajadhar AA, Theis JH, Papadopoulos E and Theodoropoulos G** (2010) Seroprevalence of equine piroplasms and host-related factors associated with infection in Greece. *Veterinary Parasitology* **169**, 273–278.
- Levine ND** (1985) Apicomplexa: the piroplasms. In Soulsby EJL (ed.), *Veterinary Protozoology*. Ames, IA: Iowa State University Press, pp. 291–328.
- Machado RZ, Toledo CZP, Teixeira MCA, André MR, Freschi CR and Sampaio H** (2012) Molecular and serological detection of *Theileria equi* and *Babesia caballi* in donkeys (*Equus asinus*) in Brazil. *Veterinary Parasitology* **186**, 461–465.
- Mahmoud MS, El-Ezz NTA, Abdel-Shafy S, Nassar SA, El Namaky AH, Khalil WK and Suarez CE** (2016) Assessment of *Theileria equi* and *Babesia caballi* infections in equine populations in Egypt by molecular, serological and hematological approaches. *Parasites and Vectors* **9**, 260.
- Malekifard F, Tavassoli M, Yakhchali M and Darvishzadeh R** (2014) Detection of *Theileria equi* and *Babesia caballi* using microscopic and molecular methods in horses in suburb of Urmia, Iran. *Veterinary Research Forum* **5**, 129–134.
- Mans BJ, Pienaar R and Latif AA** (2015) A review of *Theileria* diagnostics and epidemiology. *International Journal for Parasitology: Parasites and Wildlife* **4**, 104–118.
- Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M and Stewart LA** (2015) Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) statement. *Systematic Reviews* **4**, 1.
- Moretti A, Mangili V, Salvatori R, Maresca C, Scoccia E, Torina A and Pietrobelli M** (2010) Prevalence and diagnosis of *Babesia* and *Theileria* infections in horses in Italy: a preliminary study. *The Veterinary Journal* **184**, 346–350.
- Motloang MY, Thekisoe OMM, Alhassan A, Bakheit M, Motheo MP, Masangane FES and Mbatia PA** (2008) Prevalence of *Theileria equi* and *Babesia caballi* infections in horses belonging to resource-poor farmers in the north-eastern Free State Province, South Africa. *Onderstepoort Journal of Veterinary Research* **75**, 141–146.
- Mujica FF, Perrone T, Forlano M, Coronado A, Meléndez RD, Barrios N and Granda F** (2011) Serological prevalence of *Babesia caballi* and *Theileria equi* in horses of Lara State, Venezuela. *Veterinary Parasitology* **178**, 180–183.
- Munkhjargal T, Sivakumar T, Battsetseg B, Nyamjargal T, Aboulaila M, Purevtseren B and Igarashi I** (2013) Prevalence and genetic diversity of equine piroplasms in Tov province, Mongolia. *Infection, Genetics and Evolution* **16**, 178–185.
- Nugraha AB, Cahyaningsih U, Amrozi A, Ridwan Y, Agungpriyono S, Taher DM and Sivakumar T** (2018) Serological and molecular prevalence of equine piroplasmosis in Western Java, Indonesia. *Veterinary Parasitology: Regional Studies and Reports* **14**, 1–6.
- Oduori DO, Onyango SC, Kimari JN and MacLeod ET** (2015) A field survey for the seroprevalence of *Theileria equi* and *Babesia caballi* in donkeys from Nuu Division, Kenya. *Ticks and Tick-Borne Diseases* **6**, 683–688.
- OIE** (2005) Manual of Diagnostic Tests and Vaccines for Terrestrial Animals, ch. 2.5.6 Equine Piroplasmosis. Available at [http://www.oie.int/eng/normes/mmanual/A\\_00084.htm](http://www.oie.int/eng/normes/mmanual/A_00084.htm) (accessed 25 January 2011).
- OIE** (2018) Manual of Diagnostic Tests and Vaccines for Terrestrial Animals, ch. 3.5.8 Equine Piroplasmosis. Available at <https://www.oie.int/standard-setting/terrestrial-manual/access-online/> (accessed 29 April 2020).
- Onyiche TE, Suganuma K, Igarashi I, Yokoyama N, Xuan X and Thekisoe O** (2019) A review on equine piroplasmosis: epidemiology, vector ecology, risk factors, host immunity, diagnosis and control. *International Journal of Environmental Research and Public Health* **16**, 1736.
- Onyiche T, Taioe M, Ogo N, Sivakumar T, Biu A, Mbaya A, Xuan X, Yokoyama N and Thekisoe O** (2020) Molecular evidence of *Babesia caballi* and *Theileria equi* in equines and ticks in Nigeria: prevalence and risk factors analysis. *Parasitology*, 1–35. doi:10.1017/S0031182020000992.
- Piantedosi D, D'Alessio N, Di Loria A, Di Prisco F, Mariani U, Neola B and Veneziano V** (2014) Seroprevalence and risk factors associated with *Babesia caballi* and *Theileria equi* infections in donkeys from Southern Italy. *The Veterinary Journal* **202**, 578–582.
- Posada-Guzmán MF, Dolz G, Romero-Zúñiga JJ and Jiménez-Rocha AE** (2015) Detection of *Babesia caballi* and *Theileria equi* in blood from equines from four indigenous communities in Costa Rica. *Veterinary Medicine International*, <http://dx.doi.org/10.1155/2015/236278>.
- Qablan MA, Obornik M, Petrželková KJ, Sloboda M, Shudiefat MF, Hořín P and Modrý D** (2013) Infections by *Babesia caballi* and *Theileria equi* in Jordanian equids: epidemiology and genetic diversity. *Parasitology* **140**, 1096–1103.
- Ramsay JD, Ueti MW, Johnson WC, Scoles GA, Knowles DP and Mealey RH** (2013) Lymphocytes and macrophages are infected by *Theileria equi*, but T cells and B cells are not required to establish infection in vivo. *PLoS ONE* **8**, e76996.
- Rapoport A, Aharonson-Raz K, Berlin D, Tal S, Gottlieb Y, Klement E and Steinman A** (2014) Molecular characterization of the *Babesia caballi* rap-1 gene and epidemiological survey in horses in Israel. *Infection, Genetics and Evolution* **23**, 115–120.
- Ros-García A, M'ghirbi Y, Hurtado A and Bouattour A** (2013) Prevalence and genetic diversity of piroplasm species in horses and ticks from Tunisia. *Infection, Genetics and Evolution* **17**, 33–37.
- Rosales R, Rangel-Rivas A, Escalona A, Jordan LS, Gonzatti MI, Aso PM and Mijares A** (2013) Detection of *Theileria equi* and *Babesia caballi* infections in Venezuelan horses using Competitive-Inhibition ELISA and PCR. *Veterinary Parasitology* **196**, 37–43.
- Rothschild CM** (2013) Equine piroplasmosis. *Journal of Equine Veterinary Science* **33**, 497–508.
- Rüegg SR, Torgerson P, Deplazes P and Mathis A** (2007) Age-dependent dynamics of *Theileria equi* and *Babesia caballi* infections in southwest Mongolia based on IFAT and/or PCR prevalence data from domestic horses and ticks. *Parasitology* **134**, 939–947.
- Salim BOM, Hassan SM, Bakheit MA, Alhassan A, Igarashi I, Karanis P and Abdelrahman MB** (2008) Diagnosis of *Babesia caballi* and *Theileria equi* infections in horses in Sudan using ELISA and PCR. *Parasitology Research* **103**, 1145.
- Salim B, Bakheit MA, Kamau J and Sugimoto C** (2013) Current status of equine piroplasmosis in the Sudan. *Infection, Genetics and Evolution* **16**, 191–199.
- Sanusi M, Ahmed IA, Tahir I, Mai HM, Kalla DJU and Shuaibu I** (2014) Survey of equine piroplasmosis in the Savanna areas, Bauchi state, North-eastern Nigeria. *Ippologia* **25**, 3–8.
- Scoles GA and Ueti MW** (2015) Vector ecology of equine piroplasmosis. *Annual Review of Entomology* **60**, 561–580.
- Seo MG, Yun SH, Choi SK, Cho GJ, Park YS, Kwon OD and Kwon YS** (2011) Seroprevalence of equine piroplasms in the Republic of Korea. *Veterinary Parasitology* **179**, 224–226.
- Sevinc F, Maden M, Kumas C, Sevinc M and Ekici OD** (2008) A comparative study on the prevalence of *Theileria equi* and *Babesia caballi* infections in horse sub-populations in Turkey. *Veterinary Parasitology* **156**, 173–177.
- Sigg L, Gerber V, Gottstein B, Doherr MG and Frey CF** (2010) Seroprevalence of *Babesia caballi* and *Theileria equi* in the Swiss horse population. *Parasitology International* **59**, 313–317.
- Sumbria D, Singla LD and Sharma A** (2016) *Theileria equi* and *Babesia caballi* infection of equids in Punjab, India: a serological and molecular survey. *Tropical Animal Health and Production* **48**, 45–52.
- Turaki UA, Kumsha HA, Biu AA and Bokko PB** (2014) Prevalence of piroplasmosis amongst local horses in Northeastern Nigeria. *Journal of Agricultural and Veterinary Science* **4**, 27.
- Uilenberg G** (2006) *Babesia* – a historical overview. *Veterinary Parasitology* **138**, 3–10.
- Vieira TS, Vieira RF, Finger MA, Nascimento DA, Sicupira PM, Dutra LH and Vidotto O** (2013) Seroepidemiological survey of *Theileria equi* and *Babesia caballi* in horses from a rural and from urban areas of Paraná State, southern Brazil. *Ticks and Tick-Borne Diseases* **4**, 537–541.
- Vieira MIB, Costa MM, de Oliveira MT, Gonçalves LR, André MR and Machado RZ** (2018) Serological detection and molecular characterization of piroplasmids in equids in Brazil. *Acta Tropica* **179**, 81–87.
- Wang M, Guo W, Igarashi I, Xuan X, Wang X, Xiang W and Jia H** (2014) Epidemiological investigation of equine piroplasmosis in China by enzyme-

- linked immunosorbent assays. *Journal of Veterinary Medical Science* **76**, 549–552.
- Wise LN, Kappmeyer LS, Mealey RH and Knowles DP (2013) Review of equine piroplasmosis. *Journal of Veterinary Internal Medicine* **27**, 1334–1346.
- Xu Y, Zhang S, Huang X, Bayin C, Xuan X, Igarashi I and Mikami T (2003) Seroepidemiologic studies on *Babesia equi* and *Babesia caballi* infections in horses in Jilin province of China. *Journal of Veterinary Medical Science* **65**, 1015–1017.
- Ybañez AP, Ybañez RHD, Talle MG, Arreglo RMT, Geens MJC, Villas JGI and Liu M (2018) Serological and molecular detection of *Theileria equi* and *Babesia caballi* in Philippine horses. *Ticks and Tick-Borne Diseases* **9**, 1125–1128.