

effectiveness in mania (Ferrier, 1998). After talking to an advocate at a day centre who told her that it was an 'experimental' treatment, the patient became anxious and asked to discontinue treatment. Her mania subsequently worsened.

While junior doctors sometimes see patients with a nurse or other doctor on the ward, and are thus fairly open in their interactions, advocates usually see patients alone either on or off the ward (sometimes in day centres). There is thus a feeling that advocates are giving advice in private which others may not be aware of.

The title of Thomas & Bracken's article was "Putting ethics into practice". I am concerned that the ethics of some advocacy movements are not those of doctors but that they may be using their access to vulnerable people (psychiatric patients), to promote their own anti-medical establishment political agenda. It would be mistaken to 'dismiss' them as being 'anti-psychiatry' as Thomas & Bracken state. To dismiss them would be to ignore their destructive ideology-driven power. Local advocates have told ward patients that nurses are unable to fight back if attacked, a tacit encouragement of violence against staff.

My experience of advocacy has suggested that while the concept is a good one, in practice there are problems.

Reference

FERRIER, I. N. (1998) Lamotrigine and gabapentin. Alternative in the treatment of bipolar disorder. *Neuropsychobiology*, **38**, 192-197.

DAMIAN GAMBLE, *Senior House Officer in Psychiatry, Department of Psychiatry, 1st Floor, Out-Patients Department, Royal London Hospital, Whitechapel Road, London E1 1BB*

Guidelines for the police and psychiatric services

Sir: Your special article 'Police training for the management of dangerous patients' (*Psychiatric Bulletin* January 1999, **23**, 46-48) drew attention to an important aspect of the relationship between the police and psychiatric services.

Another area of police practice impinging on psychiatry needs to be highlighted, as it is a cause for concern. This is the widespread practice of the police bringing people from the community to be 'assessed' at psychiatric hospitals, without placing them on Section 136 of the Mental Health Act 1983.

The usual scenario is that the police attempt to arrest an individual, but on becoming aware of a

psychiatric history, decide to divert the person to hospital.

This practice is worrying for a number of reasons: it results in a large number of patients being escorted/detained by the police outside of the protection of Section 136 of the Act. It may encourage people to escape justice by hiding behind a psychiatric label. Also the person may be coerced unwillingly to go to hospital, which subsequently may leave trusts open to litigation by patients claiming they were taken to hospital and detained on wards illegally.

It appears imperative that this practice is monitored on a local basis, and guidelines drawn up between the police and psychiatric services regarding best practice in this grey area.

RAJEN SHAH, *Specialist Registrar in Adult Psychiatry, Medical Centre, Farnham Road Hospital, Farnham Road, Guildford, Surrey GU2 5LX*

Healthy debate about the fragmentation of services

Sir: I am pleased to see that my article (*Psychiatric Bulletin*, January 1999, **23**, 31-33) has generated questions and debate from medical colleagues, National Health Service Management and the media. They all had specific and valid questions and concerns on the feasibility and difficulties of single gender wards.

The most frequent question was about problems which may arise in all male wards once single gender wards are established universally in a district. Most colleagues fear that single gender male wards may become extremely difficult to deal with and may turn into unofficial intensive care units.

The concern around fragmentation of services and difficulties in coordination has been brought forward by a senior practitioner with a special interest in services. I have to admit that the coordination of continuity of care in our specialist service has taken an enormous effort. It has been through personal endeavour, universal good will and collaboration that the difficulties in working with four sectors and six adult general psychiatry consultants have been, only partly, overcome. Being served by several different community teams, organising Care Programme Approach meetings, keyworking systems, out-patient follow-up by each one of them has been a daunting task. We still are in the process of reorganising the follow-up system.

Questions around length of stay in hospital have come from different sources including Department of Health officials. The problems of aggregating difficult to treat female patients with complex needs has not been overcome in our