

A comparison between mitochondrial DNA and the ribosomal internal transcribed regions in prospecting for cryptic species of platyhelminth parasites

R. VILAS*, C. D. CRISCIONE and M. S. BLOUIN

Department of Zoology, Oregon State University, 3029 Cordley Hall, Corvallis, Oregon 97331, USA

(Received 25 March 2005; revised 12 May and 31 May 2005; accepted 31 May 2005)

SUMMARY

We examined the relative merits of mitochondrial DNA loci and ribosomal DNA internal transcribed spacers for their use in prospecting for cryptic species of platyhelminth parasites. Sequence divergence at ITS1 and ITS2 was compared with divergence at 2 mtDNA loci (NADH dehydrogenase-1 and cytochrome *c* oxidase I) between closely related species of trematodes and cestodes. Both spacers accumulated substitutions substantially more slowly than mtDNA, which clearly shows a higher level of divergence among species relative to intra-specific variation. Besides a slow rate of substitution, other caveats that may be encountered when using ITS sequences as a prospecting marker are discussed. In particular, we note recent studies that suggest the existence of substantial levels of intra-individual variation in ITS sequences of flatworms. Because it is likely that closely related species share this phenomenon, it may confound the detection of cryptic species, especially if small sample sizes are studied. Although potential limitations of mtDNA are also recognized, the higher rate of evolution and smaller effective population size of this marker increases the probability of detecting diagnostic characters between cryptic species.

Key words: cytochrome *c* oxidase I, NADH dehydrogenase-1, rDNA internal transcribed spacer, platyhelminth parasites, cryptic species, molecular prospecting.

INTRODUCTION

The application of DNA methods to studies on the systematics and population genetics of platyhelminth parasites has become widespread in recent years. These molecular studies have led to the chance discovery of many genetically distinct but morphologically very similar species (Jousson, Bartoli and Pawlowski, 2000; Macnish *et al.* 2002; Criscione and Blouin, 2004; Xiao *et al.* 2005). What is not commonly done is to use a DNA marker to deliberately search for evidence of cryptic species (e.g., Jousson and Bartoli, 2000; Haukisalmi *et al.* 2001; Hu *et al.* 2005; Wu *et al.* 2005), an approach sometimes referred to as molecular prospecting (Blouin, 2002; Criscione, Poulin and Blouin, 2005). Molecular prospecting is usually initiated when one suspects the presence of cryptic species or when one needs to be sure they are dealing with a single species or population. For example, if a species occurs in multiple hosts or habitats it may be prudent to check for evidence of cryptic species. If one uses DNA sequence data, then cryptic species might be indicated if one finds unusually large genetic distances between certain individuals (distances greater than typically

found within species of that taxon), or if one finds evidence of reciprocally monophyletic groups in large population samples. Note that prospecting is simply a first pass to look for evidence that species may be present. Given such evidence (e.g., a few individuals having unusually large genetic distance), the next step would be to formally test the hypothesis of absence of gene flow or lineage exclusivity by collecting more samples and applying multiple markers. This second step can be called species delimitation. In that sense prospecting is a form of exploratory data analysis that generates hypotheses, and delimitation is the next step of hypothesis testing. Finally, we of course recognize that using an arbitrary genetic distance yardstick is not a sound way to delimit species. But it *is* a rational approach in prospecting.

An ideal molecular marker for prospecting would be easy to obtain, provide unambiguous data, and have a rapid evolutionary rate. Mitochondrial DNA and ITS are commonly used on platyhelminths for phylogenetics and species identification (distinguishing between previously identified, but morphologically similar, species; Morgan and Blair, 1995, 1998; León-Règagnon, Brooks and Pérez-Ponce de León, 1999; Tkach, Pawlowski and Sharpilo, 2000; Kostadinova *et al.* 2003; Scholz *et al.* 2004). Given that cryptic species of platyhelminth parasites are probably common, it is useful to review the relative merits of mtDNA and ITS for their use

* Corresponding author: Department of Zoology, Oregon State University, 3029 Cordley Hall, Corvallis, Oregon 97331, USA. Tel: +541 737 4360. Fax: +541 737 0501. E-mail: vilasr@science.oregonstate.edu

Table 1. Species and GenBank accessions corresponding to the sequences used in this study

Species	ITS1	ITS2	ND1	COI
TREMATODA				
<i>Deropeges aspina</i> A	AY269443	–	AY269445– AY269466	–
<i>Deropeges aspina</i> B	AY269444	–	AY269511– AY2969532	–
<i>Echinostoma caproni</i>	U58098	U58098	AF025837	–
<i>Echinostoma paraensei</i>	U58100	U58100	AF025834	–
<i>Echinostoma revolutum</i>	AY168930	AY168930	AY168933	–
<i>Echinostoma robustum</i>	U58102	U58102	AF025832	–
<i>Echinostoma trivolvis</i>	U58097	U58097	AF025831	–
<i>Fasciola hepatica</i>	–	AJ557568	NC002546	–
<i>Fasciola gigantica</i>	–	AJ557569	AF286338	–
<i>Glypthelmins brownorumae</i>	–	AY278062	–	AY278055
<i>Glypthelmins facioi</i>	–	AY278060	–	AY278053
<i>Glypthelmins tuxtlaensis</i>	–	AY278061	–	AY278054
<i>Lecithochirium rufoviride</i> *	Unpubl.	–	Unpubl.	–
<i>Lecithochirium furcolabiatum</i> A	Unpubl.	–	Unpubl.	–
<i>Lecithochirium furcolabiatum</i> B	Unpubl.	–	Unpubl.	–
<i>Paragonimus hokuoensis</i>	–	AY618756	–	AY618837
<i>Paragonimus miyazaki</i>	–	AY618757	–	U97215
<i>Paragonimus skrjabini</i>	–	AY618752	–	U97216
<i>Paragonimus westermani</i> (Jap.)	AF040942	U96907	–	U97208
<i>Paragonimus westermani</i> (Mal.)	AF071426	U96909	–	U97211
<i>Paragonimus westermani</i> (Phil.)	AF040943	U96910	–	U97213
<i>Paragonimus westermani</i> (Thail.)	AB094992	–	–	U97212
<i>Schistosoma bovis</i>	–	AF146035	–	AY157212
<i>Schistosoma edwardiense</i>	AY197344	AY197344	–	AY197347
<i>Schistosoma haematobium</i>	Z21716	Z21716	–	AY157209
<i>Schistosoma hippopotami</i>	AY197343	AY197343	–	AY197346
<i>Schistosoma intercalatum</i>	Z21717	Z21717	–	AY157208
<i>Schistosoma japonicum</i>	–	U22167	NC002544	U82264
<i>Schistosoma malayensis</i>	–	U82398	AF295106	U82262
<i>Schistosoma mansoni</i>	AF029309	U22168	–	NC002545
<i>Schistosoma matheei</i>	Z21718	Z21718	–	AY157211
<i>Schistosoma mekongi</i>	U89871	U22169	NC002529	U82263
<i>Schistosoma rodhaini</i>	AF531312	AF531312	–	AY157202
CESTODA				
<i>Echinococcus equinus</i>	AJ237773	–	AF346403	AF346403
<i>Echinococcus granulosus</i> G1	AJ237777	–	AF297617	AF297617
<i>Echinococcus granulosus</i> G7	AJ237821	–	AJ237638	AF458876
<i>Echinococcus granulosus</i> G10	AY185199	–	AF525297	AF525457
<i>Echinococcus multilocularis</i>	AJ237778	–	NC000928	NC000928
<i>Echinococcus ortleppi</i>	AJ237774	–	AJ237636	M84665
<i>Paranoplocephala alternata</i>	AY299557	–	–	AY181502
<i>Paranoplocephala arctica</i>	AY752661	–	–	AY181509
<i>Paranoplocephala fellmani</i>	AY752655	–	–	AY586612
<i>Paranoplocephala kalelai</i>	AY752660	–	–	AY189959
<i>Paranoplocephala macrocephala</i>	AY752658	–	–	AY181518
<i>Paranoplocephala primordialis</i>	AY752662	–	–	AY568218
<i>Paranoplocephala serrata</i>	AF314414	–	–	AY568220
<i>Taenia asiatica</i>	AY606272	–	NC004826	NC004826
<i>Taenia saginata</i>	AY392045	–	AJ239106	AB107242

* Unpublished data for *Lecithochirium* species.

in the molecular prospecting in this group of organisms. Little is known about the relative divergence rates of nuclear versus mtDNA in flatworms (Morgan and Blair, 1998), so here we first establish the relative levels of intra-versus inter-specific variation typical of the two types of markers in flatworms. Then we review data on molecular evolution of each marker and discuss caveats that may be

encountered when using either mtDNA and ITS as a prospecting marker in platyhelminths.

MATERIALS AND METHODS

We obtained DNA sequences from GenBank for pairs of congeners for which both mitochondrial sequence data and ITS data were available (Table 1).

When available, previous phylogenetic information was used to ensure that comparisons were among closely related or sister species. Our goal is not to use this information to identify a minimum genetic distance for defining species, but to compare the relative rates of evolution of the two markers, and evaluate which is more useful in molecular prospecting for cryptic species. Because certain studies suggest that ITS1 appears to accumulate substitution at higher rate than ITS2 (Morgan and Blair, 1995; van Herwerden, Blair and Agatsuma, 1999; Tkach *et al.* 2000), we included data from both loci separately. Although the mitochondrial genome contains different functional units (genes), the whole molecule is a single genealogical unit due to a lack of recombination. However, the rate of substitution among mitochondrial genes is variable. Thus, we restricted the data set to species pairs for which nicotinamide adenine dinucleotide dehydrogenase subunit 1 (ND1) and cytochrome *c* oxidase subunit 1 (COI) sequences were available. These are the most commonly used mitochondrial markers in molecular systematics of flatworms. According to the identification given in Kostadinova *et al.* (2003), the *Echinostoma robustum* sequences used in this study were those provided by Morgan and Blair (1998) as a German isolate of *Echinostoma revolutum*. Several strains of *Echinococcus granulosus* that could represent different species were included in the comparative analyses. We followed the recent proposition of species status for 2 'strains' (G4 and G5) of *Echinococcus granulosus*, which were named *E. equinus* and *E. ortleppi*, respectively (see Thompson and McManus, 2002). We also compared specimens identified as *Paragonimus westermani* from different localities that probably constitute cryptic species (Blair *et al.* 1997). Recently, the reduction of *Paragonimus skrjabini* and *P. miyazakii* to subspecific status has been suggested (Blair *et al.* 2005); however, we retain the original designation. Species of *Lecithochirium* show both repetitive elements within ITS1 and length variation for this spacer within a single individual (unpublished observation). We assumed that intra-individual ITS1 length variation found within *Lecithochirium* species was due to variation in the number of repeats and not owing to the presence of paralogues in other rDNA loci. Because of the size of the repetitive region, only the 5' end was compared among species. Similarly, a repeat region present in ITS1 of some species of *Schistosoma* was not included in the comparisons between *S. mekongi* and *S. malayensis*, and species of the *S. haematobium* group. Repeats in the ITS were not observed for other species in our data set. The mtDNA sequences aligned unambiguously with no gaps. However, because ITS is a non-coding sequence, frequent insertions or deletions (indels) were present in the comparison of different taxa, making the alignment more complicated than in the case of




Fig. 1. Cytochrome *c* oxidase I uncorrected *p* distance versus ribosomal DNA internal transcribed spacer1 uncorrected *p* distance for each species pair. The diagonal line indicates where equal distances for both markers would lie.

mtDNA genes. In order to be consistent we aligned each species pair using Clustal W, applying the same algorithm in all cases. The percentage of sequence difference was computed as the proportion of nucleotide sites at which 2 sequences are different (uncorrected *p*). No correction is necessary for multiple changes because the sequences of closely related species exhibit little divergence. Thus, uncorrected *p* is an estimate of the accumulated number of nucleotide substitutions per site. Because ND1 sequences of multiple individuals were available for *Lecithochirium* and *Deropegeus* species, we computed average distances between sibling species in each case. MEGA 2.0 (Kumar *et al.* 2001) was used to make all calculations.

RESULTS AND DISCUSSION

Comparisons of genetic distances from pairs of congeners for both mtDNA and ITS sequences clearly demonstrated that mtDNA sequences of platyhelminths accumulate nucleotide substitutions at a much higher rate than ITS (Fig. 1). It is notable that several pairs of congeners showed up to 10% divergence at the mtDNA, but about 1% at the ITS (Table 2). Species pairs of platyhelminths that have approximately 1% divergence between ITS sequences are relatively frequent (Després *et al.* 1992; Kane and Rollinson, 1994; Anderson and Barker, 1998; Criscione and Blouin, 2004). This degree of variation in the ITS was also typical of the variation within and among individuals of the same species (e.g., van Herwerden *et al.* 1999; Macnish *et al.* 2002). In contrast, large population samples from

Table 2. Mitochondrial and internal transcribed spacer distances for each species pair

	ITS1	ITS2	ND1	COI
TREMATODA				
Deropegus				
<i>D. aspina</i> A- <i>D. aspina</i> B	0.018	–	0.101*	–
Echinostoma				
<i>E. caproni</i> - <i>E. paraensei</i>	0.022	0.019	0.150	–
<i>E. caproni</i> - <i>E. robustum</i>	0.010	0.026	0.147	–
<i>E. caproni</i> - <i>E. trivolvis</i>	0.022	0.024	0.150	–
<i>E. caproni</i> - <i>E. revolutum</i>	0.032	0.038	0.166	–
<i>E. paraensei</i> - <i>E. robustum</i>	0.015	0.007	0.161	–
<i>E. paraensei</i> - <i>E. revolutum</i>	0.035	0.019	0.162	–
<i>E. paraensei</i> - <i>E. trivolvis</i>	0.017	0.005	0.138	–
<i>E. revolutum</i> - <i>E. trivolvis</i>	0.044	0.024	0.130	–
<i>E. revolutum</i> - <i>E. robustum</i>	0.024	0.026	0.119	–
<i>E. trivolvis</i> - <i>E. robustum</i>	0.021	0.124	0.130	–
Fasciola				
<i>F. hepatica</i> - <i>F. gigantica</i>	–	0.014	0.103	–
Glyphelmins				
<i>G. brownorumae</i> - <i>G. facioi</i>	–	0.010	–	0.126
<i>G. brownorumae</i> - <i>G. tuxtlasensis</i>	–	0.014	–	0.119
<i>G. facioi</i> - <i>G. tuxtlasensis</i>	–	0.007	–	0.119
Lecithochirium				
<i>L. rufoviride</i> - <i>L. furcolabiatum</i> A	0.026	–	0.118*	–
<i>L. rufoviride</i> - <i>L. furcolabiatum</i> B	0.026	–	0.121*	–
<i>L. furcolabiatum</i> A- <i>L. furcolab.</i> B	0.026	–	0.076*	–
Paragonimus				
<i>P. skrjabini</i> - <i>P. miyazaki</i>	–	0.003	–	0.097
<i>P. skrjabini</i> - <i>P. hokuoensis</i>	–	0.003	–	0.063
<i>P. hokuoensis</i> - <i>miyazaki</i>	–	0.006	–	0.097
<i>P. westermani</i> Malaysia-Japan	0.036	0.017	–	0.107
<i>P. westermani</i> Malaysia-Philippines	0.021	0.003	–	0.087
<i>P. westermani</i> Malaysia-Thailand	0.015	–	–	0.090
<i>P. westermani</i> Japan-Philippines	0.028	0.020	–	0.110
<i>P. westermani</i> Japan-Thailand	0.035	–	–	0.128
<i>P. westermani</i> Philippines-Thailand	0.019	–	–	0.067
Schistosoma				
<i>S. mekongi</i> - <i>S. malayensis</i>	0.046	0.015	0.108	0.094
<i>S. mekongi</i> - <i>S. japonicum</i>	–	0.060	0.347	0.158
<i>S. malayensis</i> - <i>S. japonicum</i>	–	0.080	0.354	0.164
<i>S. haematobium</i> - <i>S. intercalatum</i>	0.000	0.013	–	0.116
<i>S. haematobium</i> - <i>S. mattheei</i>	0.066	0.029	–	0.162
<i>S. intercalatum</i> - <i>S. mattheei</i>	0.066	0.016	–	0.142
<i>S. bovis</i> - <i>S. mattheei</i>	–	0.016	–	0.138
<i>S. bovis</i> - <i>S. intercalatum</i>	–	0.000	–	0.065
<i>S. bovis</i> - <i>S. haematobium</i>	–	0.013	–	0.113
<i>S. hippopotami</i> - <i>S. edwardiense</i>	0.048	0.019	–	0.214
<i>S. rodhaini</i> - <i>S. mansoni</i>	0.011	0.010	–	0.133
CESTODA				
Echinococcus				
<i>E. granulosus</i> G1- <i>E. granulosus</i> G7	0.050	–	0.155	0.098
<i>E. granulosus</i> G10- <i>E. equinus</i>	0.033	–	0.129	0.082
<i>E. granulosus</i> G10- <i>E. ortleppi</i>	0.009	–	0.068	0.043
<i>E. granulosus</i> G10- <i>E. multilocularis</i>	0.054	–	0.158	0.100
<i>E. equinus</i> - <i>E. ortleppi</i>	0.038	–	0.122	0.093
<i>E. equinus</i> - <i>E. multilocularis</i>	0.035	–	0.145	0.091
<i>E. ortleppi</i> - <i>E. multilocularis</i>	0.059	–	0.157	0.093
Paranoplocephala				
<i>P. arctica</i> - <i>P. alternata</i>	0.005	–	–	0.052
<i>P. arctica</i> - <i>P. serrata</i>	0.034	–	–	0.128
<i>P. serrata</i> - <i>P. alternata</i>	0.059	–	–	0.127
<i>P. macrocephala</i> - <i>P. kalelai</i>	0.031	–	–	0.084
<i>P. primordialialis</i> - <i>P. fellmani</i>	0.041	–	–	0.167
Taenia				
<i>T. saginata</i> - <i>T. asiatica</i>	0.024	–	0.053	0.047

* Average distances between sibling species.

Table 3. ND1 average and maximum percentage of sequence differences observed between individuals within populations of trematodes

(*N* is the number of individuals studied.)

Species	Average	Maximum	<i>N</i>
<i>Deropegus aspina</i> A*	0.5	1.6	66
<i>Deropegus aspina</i> B*	1.1	2.0	89
<i>Lecithochirium furcolabiatum</i>	1.3	2.2	19
<i>Lecithochirium fusiforme</i>	1.1	2.3	27
<i>Lecithochirium musculus</i>	0.03	0.3	19
<i>Lecithochirium rufoviride</i>	0.5	1.6	22
<i>Nanophyetus salmicola</i> *	0.8	2.2	91
<i>Plagioporus shawi</i> *	0.8	2.2	52

* Data over 3 or 4 geographical populations.

species of trematodes show maximum intraspecific divergences ranging from 0.3 to 2.2% at mtDNA (Table 3). These values do not overlap with divergences between species (Table 2). Thus, for example, if one has a sample of individuals that are putatively from a single population, then the presence of individuals differing by more than say, 5% at mtDNA should raise a red flag and motivate further investigation.

In accordance with previous data on flatworms (Gasser, Zhu, and McManus, 1999; Obwaller *et al.* 2004), our results suggest that COI exhibits less divergence between pairs than ND1 (Table 2). Nevertheless, even COI accumulates substitutions substantially more quickly than ITS1 (Fig. 1), which is considered more variable than ITS2. Interestingly, comparisons between ITS1 and ITS2 did not reveal a consistent pattern in terms of which region had a higher nucleotide substitution rate. This inconsistent pattern is in contrast to previous studies in flatworms that suggest ITS1 is more variable than ITS2 (Luton, Walker and Blair, 1992; Morgan and Blair, 1995; van Herwerden *et al.* 1999; Tkach *et al.* 2000; Galazzo *et al.* 2002).

A prospecting marker should exhibit relatively low intra-specific variation and a high level of divergence between closely related species. Moreover, it is important that the marker lacks intra-individual variation because one may not be able to determine which variant should be compared among samples. Low intra-individual and intra-specific variation for ITS markers is traditionally assumed due to the action of molecular turnover mechanisms (Dover, 1982). However, the efficiency of concerted evolution depends on whether rDNA variants are on nonhomologous chromosomes (Polanco, González and Dover, 1998; Parkin and Butlin, 2004), on the speciation rate (Vogler and DeSalle, 1994; Harris and Crandall, 2000), and on the frequency of chiasmata (the sites of crossing over), which may be highly variable in trematodes (van Herwerden *et al.* 1999). Under circumstances of slow concerted evolution,

processes such as hybridization, incomplete lineage sorting or retention of an ancestral polymorphism could cause the sharing of different ITS types among species of flatworms, and could therefore confound a prospecting study by using this marker (e.g., Rollinson *et al.* 1990; Huang *et al.* 2004). Hybrids having mtDNA and ITS sequences from different species were recently reported for *Schistosoma* (Morgan *et al.* 2003). These potential problems are generally recognized for mtDNA (Ballard and Whitlock, 2004). However, the importance of hybridization and lineage sorting for ITS probably has been overlooked because of the assumption of concerted evolution. There are several observations that indicate concerted evolution may not be efficient in platyhelminths. For example, 2 distinct ITS1 types were found in a single isolate of *Echinococcus granulosus* (see Bowles, Blair and McManus, 1995). Intra-individual variation in ITS of trematodes and cestodes may result from a variable copy number of tandem repeats within the spacer (van Herwerden, Blair and Agatsuma, 1998, 1999; Luo *et al.* 2002). Such repetitive elements are common in ITS of flukes (Luton *et al.* 1992; Kane and Rollinson, 1994; Kane *et al.* 1996; van Herwerden *et al.* 1998; León-Regañon *et al.* 1999; van Herwerden *et al.* 1999; Jousson and Bartoli, 2000; Bell and Sommerville, 2002; Dvorák *et al.* 2002). Indeed, the evolution by internal repetition in the ITS may be a general process in a variety of taxa (von der Schulenburg *et al.* 2001).

Intra-individual variation for mtDNA may be explained by heteroplasmy or the presence of pseudogenes. Heteroplasmy has been described for flatworms (Pena *et al.* 1995), and can result from somatic mutations, introgression or bi-parental inheritance (Avice, 2000). The rapid sorting of mtDNA molecules in cell lineages should maintain identical or very similar sequences within individuals (Avice, 2000). Thus, heteroplasmy by somatic mutations should not present a problem for molecular prospecting. However, mtDNA may introgress faster than nuclear genes (Ballard and Whitlock, 2004; Chan and Levin, 2005), and both introgression and bi-parental inheritance can lead to divergent mtDNA lineages within individuals. Bi-parental inheritance was reported for *Schistosoma mansoni* (Janotti-Passos *et al.* 2001, but see Bieberich and Minchella, 2001 for an alternative explanation). Although more work is needed to determine if mtDNA heteroplasmy is common among parasitic flatworms, the typically maternal inheritance of animal mtDNA suggests that the influence of heteroplasmy in molecular prospecting for cryptic species is probably of little importance. Large divergence between samples may also be falsely concluded if a comparison is made between a pseudogene and a functional gene. Both mitochondrial and ITS pseudogenes have been found in parasitic

helminths (Morgan and Blair, 1998; Hugall, Stanton and Moritz, 1999; van Herwerden, Blair and Agatsuma, 2000; Obwaller *et al.* 2004). Because frame-shift mutations or premature stop codons are common in pseudogenes, protein-coding genes of the mtDNA can be error checked by translating the nucleotide sequence (Benasson *et al.* 2001). However, detection of ITS pseudogenes might be more difficult because they are not protein-coding genes.

Undetected paralogy, incomplete lineage sorting and introgressive hybridization could mislead a prospecting study using either ITS or mitochondrial sequences. However, a rapid rate of evolution is a key requirement for a prospecting marker, and the results obtained here indicate mtDNA of flukes evolves much faster than ITS. Furthermore, mtDNA generally has a smaller effective population size than nuclear genes. Given that small populations reach a state of reciprocal monophyly faster than large populations, the probability of detecting diagnostic characters between cryptic species is increased by the use of mtDNA. Thus, despite limitations, mtDNA is preferable to ITS to prospect for cryptic species of parasitic plathyhelminths owing to its higher sensitivity to reveal species with recent ancestry.

R. V. was the recipient of a PD/MEC Fellowship from the European Union and the Spanish Ministry of Education and Science. C. D. C. is supported by a US Environmental Protection Agency STAR Fellowship.

REFERENCES

- Anderson, G. R. and Barker, S. C.** (1998). Inference of phylogeny and taxonomy within the didymozoidae (Digenea) from the second internal transcribed spacer ITS2 of ribosomal DNA. *Systematic Parasitology* **41**, 87–94.
- Awise, J. C.** (2000). *Phylogeography: the History and Formation of Species*. Harvard University Press, Cambridge, MA, USA.
- Ballard, J. W. O. and Whitlock, M. C.** (2004). The incomplete natural history of mitochondria. *Molecular Ecology* **13**, 729–744.
- Bell, A. S. and Sommerville, C.** (2002). Molecular evidence for a synonymy of two species of *Apatemon* Szidat, 1928, *A. gracilis* (Rudolphi, 1819) and *A. anuligerum* (von Nordman, 1832) (Digenea: Strigeidae) parasitic as metacercariae in British fishes. *Journal of Helminthology* **76**, 193–198.
- Benasson, D., Zhang, D. X., Hartl, D. L. and Hewitt, G. M.** (2001). Mitochondrial pseudogenes: Evolution's misplaced witnesses. *Trends in Ecology and Evolution* **16**, 314–321.
- Bieberich, A. A. and Minchella, D. J.** (2001). Mitochondrial inheritance in *Schistosoma mansoni*: mitochondrial variable number tandem repeat mutation produces noise on top of the signal. *Journal of Parasitology* **87**, 1011–1015.
- Blair, D., Agatsuma, T., Watanobe, T., Okamoto, M. and Ito, A.** (1997). Geographical genetic structure within the human lung fluke, *Paragonimus westermani*, detected from DNA sequences. *Parasitology* **115**, 411–417.
- Blair, D., Chang, Z., Chen, M., Cui, A., Wu, B., Agatsuma, T., Iwagami, M., Corlis, D., Fu, C. and Zhan, X.** (2005). *Paragonimus skrjabini* Chen, 1959 (Digenea: Paragonimidae) and related species in eastern Asia: a combined molecular and morphological approach to identification and taxonomy. *Systematic Parasitology* **60**, 1–21.
- Blouin, M. S.** (2002). Molecular prospecting for cryptic species of nematodes: mitochondrial DNA versus internal transcribed spacer. *International Journal for Parasitology* **32**, 527–531.
- Bowles, J., Blair, D. and McManus, D. P.** (1995). A molecular phylogeny of the genus *Echinococcus*. *Parasitology* **110**, 317–328.
- Chan, K. M. A. and Levin, S. A.** (2005). Leaky prezygotic isolation and porous genomes: rapid introgression of maternally inherited DNA. *Evolution* **59**, 720–729.
- Criscione, C. D. and Blouin, M. S.** (2004). Life cycles shape parasite evolution: comparative population genetics of salmon trematodes. *Evolution* **58**, 198–202.
- Criscione, C. D., Poulin, R. and Blouin, M. S.** (2005). Molecular ecology of parasites: elucidating ecological and microevolutionary processes. *Molecular Ecology* **14**, 2247–2257.
- Després, L., Imbert-Establet, D., Combes, C. and Bonhomme, F.** (1992). Molecular evidence linking hominid evolution to recent radiation of Schistosomes (Platyhelminthes: Trematoda). *Molecular Phylogenetics and Evolution* **1**, 295–304.
- Dover, G. A.** (1982). Molecular drive: a cohesive mode of species evolution. *Nature, London* **299**, 111–117.
- Dvorák, J., Vanáčová, S., Hampl, V., Flegr, J. and Horák, P.** (2002). Comparison of European *Trichobilharzia* species based on ITS1 and ITS2 sequences. *Parasitology* **124**, 307–313.
- Galazzo, D. E., Dayanandan, S., Marcogliese, D. J. and McLaughlin, J. D.** (2002). Molecular systematics of some North American species of *Diplostomum* (Digenea) based on rDNA-sequence data and comparisons with European congeners. *Canadian Journal of Zoology* **80**, 2207–2217.
- Gasser, R. B., Zhu, X. and McManus, D. P.** (1999). NADH dehydrogenase subunit 1 and cytochrome c oxidase subunit I sequences compared for members of the genus *Taenia* (Cestoda). *International Journal for Parasitology* **29**, 1965–1970.
- Harris, D. J. and Crandall, K. A.** (2000). Intragenomic variation within ITS1 and ITS2 of freshwater crayfishes (Decapoda: Cambaridae): implications for phylogenetic and microsatellite studies. *Molecular Biology and Evolution* **17**, 284–291.
- Haukisalmi, V., Wickström, L. M., Hantula, J. and Henttonen, H.** (2001). Taxonomy, genetic differentiation and Holarctic biogeography of *Paranoplocephala* spp. (Cestoda: Anoplocephalidae) in collared lemmings (*Dicrostonyx*: Arvicolinae). *Biological Journal of the Linnean Society* **74**, 171–196.
- Hu, M., Gasser, R. B., Chilton, N. B. and Beveridge, I.** (2005). Genetic variation in the mitochondrial cytochrome c oxidase subunit 1 within three species of

- Progamotaenia* (Cestoda: Anoplocephalidae) from macropodid marsupials. *Parasitology* **130**, 117–129.
- Huang, W. Y., He, B., Wang, C. R. and Zhu, X. Q.** (2004). Characterization of *Fasciola* species from Mailand China by ITS-2 ribosomal DNA sequence. *Veterinary Parasitology* **120**, 75–83.
- Hugall, A., Stanton, J. and Moritz, C.** (1999). Reticulate evolution and the origins of ribosomal internal transcribed spacer diversity in apomictic *Meloidogyne*. *Molecular Biology and Evolution* **16**, 157–164.
- Janotti-Passos, L. K., Souza, C. P., Parra, J. C. and Simpson, A. J. G.** (2001). Biparental mitochondrial DNA inheritance in the parasitic trematode *Schistosoma mansoni*. *Journal of Parasitology* **87**, 79–82.
- Jousson, O. and Bartoli, P.** (2000). The life cycle of *Opecoeloides columbellae* (Pagenstecher, 1863) n.comb. (Digenea, Opecoelidae): evidence from molecules and morphology. *International Journal for Parasitology* **30**, 747–760.
- Jousson, O., Bartoli, P. and Pawlowki, J.** (2000). Cryptic speciation among intestinal parasites (Trematoda: Digenea) infecting sympatric host fishes (Sparidae). *Journal of Evolutionary Biology* **13**, 773–775.
- Kane, R. A. and Rollinson, D.** (1994). Repetitive sequences in the ribosomal DNA internal transcribed spacer of *Schistosoma haematobium*, *Schistosoma intercalatum* and *Schistosoma mattheei*. *Molecular and Biochemical Parasitology* **63**, 153–156.
- Kane, R. A., Ridgers, I. L., Johnston, D. A. and Rollinson, D.** (1996). Repetitive sequences within the first internal transcribed spacer of ribosomal DNA in schistosomes contain a Chi-like site. *Molecular and Biochemical Parasitology* **75**, 265–269.
- Kostadinova, A., Herniou, E. A., Barrett, J. and Littlewood, D. T.** (2003). Phylogenetic relationships of *Echinostoma rudolphi*, 1809 (Digenea: Echinostomatidae) and related genera re-assessed via DNA and morphological analysis. *Systematic Parasitology* **54**, 159–176.
- Kumar, S., Tamura, K., Jakobsen, I. B. and Nei, M.** (2001). *MEGA2: Molecular Evolutionary Genetics Analysis software*, Arizona State University, Tempe, Arizona, USA.
- León-Régagnon, V., Brooks, D. R. and Pérez-Ponce De León, G.** (1999). Differentiation of Mexican species of *Haematoloechus* loos, 1899 (Digenea: Plagiorchiiformes): molecular and morphological evidence. *Journal of Parasitology* **85**, 935–946.
- Luo, H., Nie, P., Zhang, Y. A., Wang, G. T. and Yao, W. J.** (2002). Molecular variation of *Bothriocephalus acheilognathi* Yamaguti, 1934 (Cestoda: Pseudophyllidae) in different fish host species based on ITS rDNA sequences. *Systematic Parasitology* **52**, 159–166.
- Luton, K., Walker, D. and Blair, D.** (1992). Comparisons of ribosomal internal transcribed spacers from two congeneric species of flukes (Platyhelminthes: Trematoda: Digenea). *Molecular and Biochemical Parasitology* **56**, 323–327.
- Macnish, M. G., Morgan-Ryan, U. M., Monis, P. T., Behnke, J. M. and Thompson, R. C. A.** (2002). A molecular phylogeny of nuclear and mitochondrial sequences in *Hymenolepis nana* (Cestoda) supports the existence of a cryptic species. *Parasitology* **125**, 567–575.
- Morgan, J. A. T. and Blair, D.** (1995). Nuclear rDNA ITS sequence variation in the trematode genus *Echinostoma*: an aid to establishing relationships within the 37-collar spine group. *Parasitology* **111**, 609–615.
- Morgan, J. A. T. and Blair, D.** (1998). Relative merits of nuclear ribosomal internal transcribed spacer and mitochondrial CO1 and ND1 genes for distinguishing among *Echinostoma* species (Trematoda). *Parasitology* **116**, 289–297.
- Morgan, J. A. T., DeJong, J., Lwambo, J. S., Mungai, B. N., Mkoji, G. M. and Loker, E. S.** (2003). First report of a natural hybrid between *Schistosoma mansoni* and *S. rodhaini*. *Journal of Parasitology* **89**, 416–418.
- Obwaller, A., Schneider, R., Walochnik, J., Gollackner, B., Deutz, A., Janitschke, K., Aspöck, H. and Auer, H.** (2004). *Echinococcus granulosus* strain differentiation based on sequence heterogeneity in mitochondrial genes of cytochrome c oxidase-1 and NADH dehydrogenase-1. *Parasitology* **128**, 569–575.
- Parkin, E. J. and Butlin, R. K.** (2004). Within and between-individual sequence variation among ITS1 copies in the meadow grasshopper *Chorthippus parallelus* indicates frequent intrachromosomal gene conversion. *Molecular Biology and Evolution* **21**, 1595–1601.
- Polanco, C., González, A. I. and Dover, G. A.** (1998). Multigene family of ribosomal DNA in *Drosophila melanogaster* reveals contrasting patterns of homogenization for IGS and ITS spacer regions. A possible mechanism to resolve this paradox. *Genetics* **149**, 243–256.
- Pena, H. B., de Souza, C. P., Simpson, A. J. G. and Pena, S. D. J.** (1995). Intracellular promiscuity in *Schistosoma mansoni*: Nuclear transcribed nuclear sequences are part of a mitochondrial minisatellite region. *Proceedings of the National Academy of Sciences, USA* **92**, 915–919.
- Rollinson, D., Walker, T. K., Knowles, R. J. and Simpson, A. J. G.** (1990). Identification of schistosome hybrids and larval parasites using rRNA probes. *Systematic Parasitology* **15**, 65–73.
- Scholz, T., Skeriková, A., Shimazu, T. and Grygier, M. J.** (2004). A taxonomic study of species of *Bothriocephalus* Rudolphi, 1808 (Cestoda: Pseudophyllidae) from eels in Japan: morphological and molecular evidence for the occurrence of *B. claviceps* (Goeze, 1782) and confirmation of the validity of *B. japonicus* Yamaguti, 1934. *Systematic Parasitology* **57**, 87–96.
- Thompson, R. C. A. and McManus, D. P.** (2002). Towards a taxonomic revision of the genus *Echinococcus*. *Trends in Parasitology* **18**, 452–457.
- Tkach, V. V., Pawlowski, J. and Sharpilo, V. P.** (2000). Molecular and morphological differentiation between species of the *Plagiorchis vespertilionis* group (Digenea, Plagiorchiidae) occurring in European bats, with a re-description of *P. vespertilionis* (Müller, 1780). *Systematic Parasitology* **47**, 9–22.
- van Herwerden, L., Blair, D. and Agatsuma, T.** (1998). Intra- and inter-specific variation in nuclear ribosomal internal transcribed spacer 1 of the *Schistosoma japonicum* species complex. *Parasitology* **116**, 311–317.
- van Herwerden, L., Blair, D. and Agatsuma, T.** (1999). Intra- and interindividual variation in ITS1 of *Paragonimus westermani* (Trematoda: Digenea) and

- related species: implications for phylogenetic studies. *Molecular Phylogenetics and Evolution* **12**, 67–73.
- van Herwerden, L., Blair, D. and Agatsuma, T.** (2000). Multiple lineages of the mitochondrial gene NADH dehydrogenase subunit 1 (ND1) in parasitic helminthes: implications for molecular evolutionary studies of facultatively anaerobic eukaryotes. *Journal of Molecular Evolution* **51**, 339–352.
- Vogler, A. P. and DeSalle, R.** (1994). Evolution and phylogenetic information content of the ITS-1 region in the tiger beetle *Cincindela dorsalis*. *Molecular Biology and Evolution* **11**, 393–405.
- von der Schulenburg, J. H., Hancock, J. M., Pagnamenta, A., Slogett, J. J., Majerus, M. E. and Hurst, G. D.** (2001). Extreme length and length variation in the first ribosomal internal transcribed spacer of ladybird beetles (Coleoptera: Coccinellidae). *Molecular Biology and Evolution* **18**, 648–660.
- Wu, X. Y., Chilton, N. B., Zhu, X. Q., Xie, M. Q. and Li, A. X.** (2005). Molecular and morphological evidence indicates that *Pseudorhabdosynochus lantauensis* (Monogenea: Diplectanidae) represents two species. *Parasitology* **130**, 669–677.
- Xiao, N., Qiu, J., Nakao, M., Li, T., Yang, W., Chen, X., Schantz, P. M., Craig, P. S. and Ito, A.** (2005). *Echinococcus shiquicus* n.sp., a taeniid cestode from Tibetan fox and plateau pika in China. *International Journal for Parasitology* **35**, 693–701.