

Bounds for equilibrium states on amenable group subshifts

C. EVANS HEDGES 

Mathematics, University of Denver, Denver, Colorado, USA
(e-mail: evans.hedges@du.edu)

(Received 26 September 2024 and accepted in revised form 13 November 2024)

Abstract. We prove a result on equilibrium measures for potentials with summable variation on arbitrary subshifts over a countable amenable group. For finite configurations v and w , if v is always replaceable by w , we obtain a bound on the measure of v depending on the measure of w and a cocycle induced by the potential. We then use this result to show that under this replaceability condition, we can obtain bounds on the Lebesgue–Radon–Nikodym derivative $d(\mu_\phi \circ \xi)/d\mu_\phi$ for certain holonomies ξ that generate the homoclinic (Gibbs) relation. As corollaries, we obtain extensions of results by Meyerovitch [Gibbs and equilibrium measures for some families of subshifts. *Ergod. Th. & Dynam. Sys.* **33**(3) (2013), 934–953], and García-Ramos and Pavlov [Extender sets and measures of maximal entropy for subshifts. *J. Lond. Math. Soc. (2)* **100**(3) (2019), 1013–1033] to the countable amenable group subshift setting. Our methods rely on the exact tiling result for countable amenable groups by Downarowicz, Huczek, and Zhang [Tilings of amenable groups. *J. Reine Angew. Math.* **2019**(747) (2019), 277–298] and an adapted proof technique from García-Ramos and Pavlov.

Key words: equilibrium state, thermodynamic formalism, Gibbs

2020 Mathematics Subject Classification: 37D35 (Primary); 37B10, 37C85 (Secondary)

1. Introduction

This paper is concerned with equilibrium states on subshifts over a countable amenable group. In particular, for an arbitrary subshift, given an equilibrium state μ for a potential with summable variation, we prove Gibbs-like bounds on the measures of finite configurations under a replicability condition. We use this result to prove a novel conformal Gibbs-like bound on the Lebesgue–Radon–Nikodym (LRN) derivative $d(\mu \circ \xi)/d\mu$ for a certain class of Borel isomorphisms ξ . Our results generalize results of Meyerovitch [17] and García-Ramos and Pavlov [11] and correct an error in the latter paper.

Let $X \subset \mathcal{A}^G$ be a subshift (that is, a closed and shift-invariant subset) over a countable amenable group G with a finite alphabet \mathcal{A} . The dynamics of the system will be induced

by the left-translation map $\{\sigma_g\}_{g \in G}$ where X is required to be compact and σ -invariant. A potential is a continuous, real valued function $\phi : X \rightarrow \mathbb{R}$. Equilibrium states are σ -invariant Borel probability measures maximizing the pressure of μ with respect to ϕ : $P_\phi(\mu) = h(\mu) + \int \phi d\mu$, where $h(\mu)$ is the classical Kolmogorov–Sinai entropy of μ . Although we make no further assumptions on our subshift, we will require that the potential under consideration have summable variation, which will be defined in §3.2.

There is an extensive history in statistical physics and dynamical systems relating the global property of being an equilibrium state to local properties depending on the potential. Thermodynamic formalism, at its core, concerns itself with relating these global and local phenomena. Foundational results in this area were obtained by Dobrushin in [9], and Lanford and Ruelle in [15], who considered well-behaved \mathbb{Z}^d -subshifts coupled with a sufficiently regular potential. In this setting, they were able to show that an invariant measure is an equilibrium state if and only if it can be locally characterized by the Gibbs property.

We say a measure μ is Gibbs for ϕ if it satisfies a conditional probability condition. For a finite $F \Subset G$, we define the F -language of X , $L_F(X) = \{w \in \mathcal{A}^F : \text{there exists } x \in X : x_F = w\}$, to be the set of all F -shape configurations that are legal in X . For $w \in L_F(X)$, we can define the extender set of w as in [14, 20], $E_X(w) = \{\eta \in \mathcal{A}^{F^c} : w\eta \in X\}$, to be the collection of all background configurations for w such that $w\eta \in X$. We say that μ is Gibbs for ϕ if for any configuration $w \in L_F(X)$, and almost every background configuration $\eta \in E_X(w)$, we have

$$\mu(w||\eta) = \frac{\exp(\phi(w\eta))}{\sum_{v \in L_F(X)} \exp(\phi(v\eta)) \cdot 1_X(v\eta)}.$$

The Dobrušin theorem and Lanford–Ruelle theorem relating Gibbs measures and equilibrium states have been extended to the countable amenable group subshift setting in [18, 23], where it was shown that for sufficiently regular subshifts and potentials with summable variation, an invariant measure is an equilibrium state if and only if it is Gibbs for the potential.

In the case where $\phi = 0$, equilibrium states correspond to measures of maximal entropy (MMEs). Parry showed in [22] that for \mathbb{Z} -subshifts of finite type, the MME is unique (and, in fact, by application of the Lanford–Ruelle theorem, it is Gibbs for $\phi = 0$). In [4], Bowen showed that for an expansive \mathbb{Z} -action on a compact metric space satisfying the specification property and a potential with summable variation, there exists a unique equilibrium state. However, in the \mathbb{Z}^d setting with $d \geq 2$, Burton and Steif in [6] were able to construct strongly irreducible subshifts of finite type with non-unique MMEs. Additionally, in the \mathbb{Z} setting, there are trivial examples of subshifts with positive entropy and a unique MME such that the MME is not Gibbs for $\phi = 0$. Take, for example, the product of a full shift with the orbit closure of the point $0^\infty 10^\infty$, whose unique MME is the product of the unique MME on the full shift with δ_{0^∞} .

While neither uniqueness nor the Gibbs property may be attainable for MMEs for a general subshift, García-Ramos and Pavlov proved in [11] that for arbitrary \mathbb{Z}^d -subshifts, and any MME, one can obtain bounds on the measures of finite configurations under a replaceability condition. For a subshift $X \subset \mathcal{A}^{\mathbb{Z}^d}$ and $w, v \in L_F(X)$, we say v is

replaceable by w if $E_X(v) \subset E_X(w)$. García-Ramos and Pavlov showed that for any MME μ on a \mathbb{Z}^d -subshift, if v is replaceable by w , then $\mu([v]) \leq \mu([w])$.

The context considered in this paper will combine that of Meyerovitch in [17] and of García-Ramos and Pavlov in [11]. We make no assumptions on the subshift under consideration, and we require that the potential $\phi \in SV(X)$ has summable variation. Our results make use of a class of Borel isomorphisms: for finite configurations $v, w \in L_F(X)$, define $\xi_{v,w}$ pointwise to swap v and w in the F location whenever legal in X . Note here that we do not require $\xi_{v,w}$ to be continuous and, in general, it is not (see §2.2 for a precise definition and further discussion). Our first result can now be stated.

THEOREM 1.1. *Let G be a countable amenable group and X be a G -subshift. Let $\phi \in SV(X)$, μ_ϕ an equilibrium state for ϕ , $F \in G$ and $v, w \in L_F(X)$. If $E_X(v) \subset E_X(w)$, then*

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \sup_{x \in [v]} \exp\left(\sum_{g \in G} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x)))\right).$$

We note here that this inequality holds whenever the conclusion of the Lanford–Ruelle theorem also holds (Theorem 2.10 below), or even more generally when the conclusion of Theorem 1.3 holds (as noted in Observation 4.4). In particular, the equation immediately holds for all subshifts of finite type and potentials with summable variation, the novelty here is that we require no assumptions on the structure of the subshift X . It is in this general case where we must discuss extender sets, as in [11].

In general, this supremum may be hard to compute. However, an immediate corollary in the locally constant case allows us to easily compute this bound when v and w agree on a sufficient boundary. First, for finite $H \in G$, we call ϕ an H -potential when if $x_H = y_H$, then $\phi(x) = \phi(y)$. In particular, this means for any $v \in L_F(X)$ with $H \subset F$, we can write $\phi(v)$ unambiguously to mean $\phi(x)$ for any $x \in X$ such that $x_F = v$ since for all $x, y \in X$ with $x_F = y_F = v$, $\phi(x) = \phi(y)$. We also denote $H^\pm = H \cup H^{-1}$.

COROLLARY 1.2. *Let $H, F \in G$, $v, w \in L_F(X)$, and ϕ be an H -potential. Suppose that $E_X(v) \subset E_X(w)$ and for all $g \in F^c H^\pm \cap F$, $v_g = w_g$. Then, for any equilibrium state μ_ϕ for ϕ ,*

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \exp\left(\sum_{g \in F \setminus F^c H^{-1}} \phi(\sigma_g(v)) - \phi(\sigma_g(\xi_{v,w}(w)))\right).$$

As another immediate corollary, by letting $\phi = 0$, we extend [11, Theorem 4.4] by García-Ramos and Pavlov to arbitrary countable amenable groups. Although the theorem in [11] is stated for countable, finitely generated, torsion-free, amenable groups, due to an unfortunate error, their proof technique only applies immediately to the case where $G = \mathbb{Z}^d$. The error is the false assertion that for any torsion-free, finitely generated, countable amenable group $G = \langle g_1, \dots, g_k \rangle$, the subgroup generated by $\langle g_1^n, \dots, g_k^n \rangle$ has finite index in G . This is known to be false and can be shown not to hold in a variety of examples, including the Lamplighter group.

In addition to the classical definition of Gibbs in the sense of Dobrušin, Lanford, and Ruelle, another fruitful approach has been to consider a measure that is conformal Gibbs for a potential. It was shown by Borsato and MacDonald in [3] that for subshifts over a countable group and any potential ϕ , a measure is Gibbs for ϕ if and only if it is conformal Gibbs for ϕ (see §4 for a precise definition). As a consequence, this means that μ_ϕ is Gibbs for ϕ if and only if, for every Borel isomorphism of the form $\xi_{v,w}$ and for μ_ϕ -almost every (a.e.) $x \in X$,

$$\frac{d(\mu_\phi \circ \xi_{v,w})}{d\mu}(x) = \exp\left(\sum_{g \in G} \phi(\sigma_g(\xi_{v,w}(x))) - \phi(\sigma_g(x))\right).$$

Since, in general, equilibrium states are not necessarily Gibbs, this equality cannot always hold. In the general subshift setting of this paper, little can be said of this LRN derivative.

In [17], Meyerovitch showed that for a general \mathbb{Z}^d subshift X and potential ϕ with d -summable variation, if μ_ϕ is an equilibrium state for ϕ and if $E_X(v) = E_X(w)$, then $d(\mu_\phi \circ \xi_{v,w})/d\mu$ satisfies the equation above. In the language of Meyerovitch, μ_ϕ must be $(\mathfrak{T}_X^0, \psi_\phi)$ -conformal.

In §4, we will use Theorem 1.1 to obtain the following bound on this LRN, showing a conformal Gibbs-like result.

THEOREM 1.3. *Let $F \subseteq G$, $v, w \in L_F(X)$, $\phi \in SV(X)$, and μ_ϕ be an equilibrium state for ϕ . If $E_X(v) \subset E_X(w)$, then $\mu_\phi \circ \xi_{v,w}$ is absolutely continuous with respect to μ_ϕ when restricted to $[w]$ and for μ_ϕ -a.e. $x \in [w]$,*

$$\frac{d(\mu_\phi \circ \xi_{v,w})}{d\mu_\phi}(x) \leq \exp\left(\sum_{g \in G} \phi(\sigma_g(\xi_{v,w}(x))) - \phi(\sigma_g(x))\right).$$

We have become aware that Corollary 1.4 has been proven in even greater generality in the sofic group setting in [2]. However, we recover the fact in the countable amenable group setting as an easy corollary of Theorem 1.3.

COROLLARY 1.4. *Let $X \subset \mathcal{A}^G$ be a subshift over a countable amenable group G , let $\phi \in SV(X)$ be a potential with summable variation, and let μ_ϕ be an equilibrium state for ϕ . Then, μ_ϕ is $(\mathfrak{T}_X^0, \psi_\phi)$ -conformal.*

The structure of this paper is as follows. We begin with §2 on the relevant preliminaries, discussing countable amenable groups and their relevant properties. We then formally introduce subshifts over a countable amenable group, and discuss their thermodynamic formalism. Finally, we discuss equilibrium measures, the Gibbs property, and their relationship.

In §3, we prove Theorem 1.1, beginning with a lemma using Downarowicz, Huczek, and Zhang's exact tiling result from [10] to generate a sufficiently sparse almost partition of a given group G . After some preliminary lemmas in the subshift setting, we prove Theorem 1.1 and conclude Corollary 1.2 in the locally constant case.

Finally, §4 is concerned with the conformal Gibbs perspective where we formally introduce the concept and relevant definitions. We then prove Theorem 1.3, relate it to the

results of Meyerovitch, and conclude by extending [17, Theorem 3.1 and Corollary 3.2] to the countable amenable subshift setting.

2. Preliminaries

2.1. *Countable amenable groups.* Let G be a countable group and denote the identity of G by e . We use the notation $K \Subset G$ to indicate that K is a finite subset of G . A *Følner sequence* for G is a collection of finite subsets $\{F_n\}$ of G such that $G = \bigcup_{n \in \mathbb{N}} F_n$ and for all $K \Subset G$, $\lim_{n \rightarrow \infty} |KF_n \Delta F_n|/|F_n| = 0$. A countable group G is called *amenable* if there exists a Følner sequence in G .

For a given Følner sequence $\{F_n\}$, we say the sequence is *tempered* if there exists some $C > 0$ such that for all $n > 0$, $|\bigcup_{k < n} F_k^{-1} F_n| \leq C|F_n|$. For any Følner sequence, there exists a subsequence that is tempered, see [16, Proposition 1.5] for a proof of this fact. A deeper discussion of Følner sequences and their relevant properties can be found in [8, Ch. 4], but for our purposes, we note that for any amenable group G , there exists a tempered Følner sequence that can be taken such that $n!$ divides $|F_n|$ for every $n \in \mathbb{N}$.

Similar in spirit, we can define a relative almost-invariance: for any finite $F, T \Subset G$ and $\epsilon > 0$, we say T is *right (K, ϵ) -invariant* if $|TK \Delta T|/|T| < \epsilon$. We can equivalently say G is amenable if for every finite $K \Subset G$ and every $\epsilon > 0$, there exists some finite $T \Subset G$ that is right (K, ϵ) -invariant.

In our setting, we will also be interested in a sense of sparseness of sets, for this, we define the following.

Definition 2.1. For any $S \subset G, F \Subset G$, we say S is *left F -sparse* if for all distinct $s, s' \in S$,

$$sF \cap s'F = \emptyset.$$

Note here, by a trivial application of the definition, we know for any $F \subset H \Subset G$ and $S \subset G$, if S is left H sparse, then S is left F sparse. We will now define the right (respectively left) H -interior and H -boundary of F for $F, H \Subset G$.

Definition 2.2. The *right H -interior* of F , denoted by $\text{Int}_H(F)$, is defined by

$$\text{Int}_H(F) = \bigcap_{h \in H} Fh = \{f \in F : \text{for all } h \in H, fh \in F\}.$$

The *right H -boundary* of F , denoted $\partial_H(F)$, is all elements of F not in the H -interior. Precisely,

$$\partial_H(F) = F \setminus \text{Int}_H(F) = \{f \in F : \text{there exists } h \in H \text{ such that } fh \notin F\}.$$

The left H -interior/boundary of F is defined similarly.

The proof of our main theorem will also require taking advantage of a result by Downarowicz, Huczek, and Zhang regarding exact tilings of G from [10]. For a countable amenable group G , a *finite tiling* $\mathcal{T} = \{T_i : 1 \leq i \leq k\}$ is a collection of tiles such that $\bigcup_{i \leq k} T_i = G$ and for each $T_i \in \mathcal{T}$, there exists a finite shape $S_i \Subset G$, and a collection of centers $C(S_i) \subset G$ such that $T_i = S_i C(S_i)$. In particular, a tiling \mathcal{T} is called *exact* if for

distinct $c_1, c_2 \in C(S_i)$, $S_i c_1 \cap S_i c_2 = \emptyset$. We also note it can be assumed that $e \in S$ for every shape. We can now state their result.

THEOREM 2.3. [10, Theorem 4.3] *Given any infinite, countable amenable group G , any $\epsilon > 0$, and any finite $K \Subset G$, there exists an exact tiling \mathcal{T} where each shape is right (K, ϵ) -invariant.*

Their result in fact included statements regarding the entropy of the tiling space which we have omitted since they are not necessary for the purposes of this paper.

2.2. G -subshifts. Let G be a countable amenable group. Let \mathcal{A} be a finite set, called the alphabet, endowed with the discrete topology. Our configuration space is the set of functions $x : G \rightarrow \mathcal{A}$, which we denote \mathcal{A}^G , endowed with the product topology. For any $H \subset G$ and $x \in \mathcal{A}^G$, we denote the restriction of x to H by x_H .

We define a G -action of homeomorphisms on \mathcal{A}^G by the *left-translation map*: for all $g \in G$ and $x \in \mathcal{A}^G$, we define pointwise $\sigma_g(x)_h = x_{gh}$. The set \mathcal{A}^G and the collection of left-translations $(\sigma_g)_{g \in G}$ together form a topological dynamical system (\mathcal{A}^G, σ) which we call the full G -shift on \mathcal{A} . A G -subshift is a subset $X \subset \mathcal{A}^G$ that is closed in the product topology and is σ -invariant (that is, for all $g \in G$, $\sigma_g(X) \subset X$). When G is clear, we will refer to a G -subshift as just a subshift.

For any finite $F \Subset G$ and any $w \in \mathcal{A}^F$, we call w a *configuration* of shape F . For a subshift X and a finite configuration w of shape $F \Subset G$, we say w is in the *language of X* (or w is legal in X) if there exists some $x \in X$ such that $x_F = w$. We call $L_F(X) = \{x_F : x \in X\}$ the F -*language* of X , and $L(X) = \bigcup_{F \Subset G} L_F(X)$ the language of X .

For any $F \Subset G$ and $w \in \mathcal{A}^F$, we define the *cylinder set* of w as follows:

$$[w] = \{x \in \mathcal{A}^G : x_F = w\}.$$

In particular, the cylinder sets form a basis for the product topology on \mathcal{A}^G . Cylinder sets are frequently taken intersected with a subshift, which will be clear by context.

Finally, we note here that the product topology on \mathcal{A}^G is metrizable, and for any Følner sequence $\{F_n\}$, the metric

$$d(x, y) = 2^{-\min\{n \in \mathbb{N} : x_{F_n} = y_{F_n}\}}$$

induces the product topology. This metric serves primarily to establish that G -subshifts are expansive.

For any disjoint $H, K \subset G$ and any $x \in \mathcal{A}^H$, $y \in \mathcal{A}^K$, we define the *concatenation* of x and y by $xy \in \mathcal{A}^{H \cup K}$ such that $(xy)_H = x$ and $(xy)_K = y$. We also define for any $v \in L_F(X)$, the *extender set* of v , $E_X(v)$, to be the collection of all background configurations for which v is legal. Specifically, $E_X(v) = \{\eta \in \mathcal{A}^{G \setminus F} : v\eta \in X\}$. For any $v, w \in L_F(X)$, we say v is *replaceable* by w if $E_X(v) \subset E_X(w)$.

We will now define the following functions that will be useful in the proof of our main results and essential to our discussion of the conformal Gibbs corollaries.

Definition 2.4. For any $F \Subset G$, $v, w \in L_F(X)$, we define $\xi_{v,w} : X \rightarrow X$ through the following cases.

- For $x \in [v]$, if the concatenation $wx_{Fc} \in X$, then $\xi_{v,w}(x) = wx_{Fc}$; otherwise, $\xi_{v,w}(x) = x$.
- For $x \in [w]$, similarly define $\xi_{v,w}(x)$.
- Otherwise, let $\xi_{v,w}(x) = x$.

Note this is exactly switching v and w in the F -location whenever the resulting point is still in X . It will be noted in §4 that these functions are in fact Borel isomorphisms that generate the homoclinic (Gibbs) relation.

For a given subshift $X \subset \mathcal{A}^G$, we will denote the collection of σ -invariant probability measures on X by $\mathcal{M}_\sigma(X)$. The existence of such measures is guaranteed by the fact that G is amenable. For a given $\mu \in \mathcal{M}_\sigma(X)$, for any collection of legal finite configurations $W \subset L(X)$, we will use $\mu(W)$ to mean $\mu(\bigcup_{w \in W} [w])$.

We say $\mu \in \mathcal{M}_\sigma(X)$ is *ergodic* if for all measurable $A \subset X$, for all $g \in G$, if $\mu(A \Delta \sigma_g^{-1}A) = 0$, then $\mu(A) \in \{0, 1\}$. Here, $\mathcal{M}_\sigma(X)$ is convex and compact under the weak-* topology, in fact, it forms a Choquet simplex whose extreme points are exactly the ergodic measures.

2.3. Thermodynamic formalism. The theory of thermodynamic formalism bridges the gap between microscopic and macroscopic descriptions of systems with many interacting particles, extending concepts of statistical mechanics to symbolic dynamics. Gibbs measures are central to this framework, enabling the analysis of global statistical properties derived from local interactions in a broad range of dynamical systems. This section delves into the thermodynamic formalism for countable amenable group actions on finite alphabet subshifts, examining topological pressure and its connection to statistical physics, the construction of partition functions, and the characterization of equilibrium states.

To begin, we must define topological pressure of a given potential over our subshift X . We first let $\phi : X \rightarrow \mathbb{R}$ be a real valued, continuous function which we will call a *potential*. We will denote the set of all potentials on a subshift X by $C(X)$. For any $F \Subset G$, we define the *F-Birkhoff sum* of ϕ : $\phi_F = \sum_{g \in F} \phi \circ \sigma_g$. Finally, for any open cover \mathcal{U} of X , any $F \Subset G$, we define the open cover $\mathcal{U}^F = \bigvee_{f \in F} \sigma_f^{-1}\mathcal{U} = \{\bigcap_{f \in F} \sigma_f^{-1}(U_f) : U_f \in \mathcal{U}\}$. We use this notation to define the following partition function.

Definition 2.5. We define a *partition function* for any $F \Subset G$ and any open cover \mathcal{U} of X :

$$Z_F(\phi, \mathcal{U}) = \inf \left\{ \sum_{u \in \mathcal{U}'} \exp\left(\sup_{x \in u} \phi_F(x)\right) : \mathcal{U}' \text{ is a subcover of } \mathcal{U}^F \right\}.$$

We can define the topological pressure of ϕ with respect to a given open cover \mathcal{U} to be

$$P(\phi, \mathcal{U}) = \lim_{n \rightarrow \infty} |F_n|^{-1} \log Z_{F_n}(\phi, \mathcal{U})$$

for any Følner sequence $\{F_n\}$. This limit is guaranteed to exist and does not depend on the choice of Følner sequence, further discussion of which can be found in [5] and [12]. We can now define the following.

Definition 2.6. The topological pressure of ϕ is then

$$P_{\text{top}}(\phi) = \sup_{\mathcal{U}} P(\phi, \mathcal{U}).$$

Since, in the subshift setting, the dynamical system (X, σ) is expansive, the above supremum is attained at an open cover \mathcal{U} of diameter less than or equal to the expansiveness constant. In particular, we can compute topological pressure by

$$P_{\text{top}}(\phi) = \lim_{n \rightarrow \infty} |F_n|^{-1} \log \sum_{w \in L_{F_n}(X)} \exp\left(\sup_{x \in [w]} \phi_{F_n}(x)\right).$$

In addition to the topological pressure, for a given invariant measure $\mu \in \mathcal{M}_\sigma(X)$, we can define the pressure of μ with respect to $\phi \in C(X)$ as follows:

$$P_\phi(\mu) = h(\mu) + \int \phi \, d\mu,$$

where $h(\mu)$ is the Kolmogorov–Sinai entropy of the invariant probability measure μ . As shown by Ollagnier and Pinchon [19], in the countable amenable subshift setting, the variational principle holds. In particular, for all $\phi \in C(X)$,

$$P_{\text{top}}(\phi) = \sup_{\mu \in \mathcal{M}_\sigma(X)} P_\phi(\mu).$$

In statistical physics, equilibrium states correspond with probability measures on the state space that minimize the Gibbs free energy of the system. Up to a multiplicative constant, the free energy of an invariant measure corresponds with the negative pressure, and so we similarly define an equilibrium state as follows.

Definition 2.7. We say $\mu \in \mathcal{M}_\sigma$ is an equilibrium state for ϕ if it attains the variational principle supremum, that is,

$$P_{\text{top}}(\phi) = h(\mu) + \int \phi \, d\mu.$$

When h is upper semicontinuous (as in the case for expansive dynamical systems), we know the collection of equilibrium states for a given ϕ is non-empty, compact under the weak- $*$ topology, and convex, where the extreme points are exactly the ergodic equilibrium states.

2.4. Equilibrium measures and the Gibbs property. A rich theory has developed around relating equilibrium states and measures with the Gibbs property (for some definition of Gibbs property). In the classical results of Dobrušin, Lanford, and Ruelle, the Gibbs property can be defined in terms of conditional probabilities as follows.

Definition 2.8. For a subshift $X \subset \mathcal{A}^{\mathbb{Z}^d}$ and a potential $\phi \in C(X)$, we say a measure μ is Gibbs for ϕ if for all $F \Subset \mathbb{Z}^d$, for all $w \in L_F(X)$, and for μ -a.e. $\eta \in \mathcal{A}^{\mathbb{Z}^d \setminus F}$ such that $w\eta \in X$,

$$\mu(w||\eta) = \frac{\exp(\phi(w\eta))}{\sum_{v \in L_F(X)} \exp(\phi(v\eta)) \cdot 1_X(v\eta)}.$$

In other words, we say a measure is Gibbs for ϕ when the probability of a local configuration, w , given a background configuration, η , can be computed in the typical way one computes Gibbs measures in the finite case. The results of Dobrušin combined with those of Lanford and Ruelle show equivalence to the Gibbs property and being an equilibrium state under certain assumptions on subshift.

The following theorem involves a technical condition called property (D), which is described in [9].

THEOREM 2.9. (Dobrushin [9]) *Let $X \subset \mathcal{A}^{\mathbb{Z}^d}$ be a subshift satisfying property (D) and $\phi \in C(X)$ be a potential with d -summable variation. If $\mu \in \mathcal{M}_\sigma(X)$ is an invariant probability measure that is Gibbs for ϕ , then μ is an equilibrium state for ϕ .*

THEOREM 2.10. (Lanford and Ruelle [15]) *Let $X \subset \mathcal{A}^{\mathbb{Z}^d}$ be a subshift of finite type and $\phi \in C(X)$ be a potential with d -summable variation. If μ is an equilibrium state for ϕ , then μ is Gibbs for ϕ .*

Subshifts of finite type, SFTs, are an important and well-studied class of subshifts that are not defined here. Analogous results have since been shown in the countable amenable subshift setting [18, 23]. Note that these statements rely on relatively strong assumptions on both the subshift and the potential. When no assumptions are made of the potential, little can be said about the equilibrium state. In fact, by upper semicontinuity of the entropy map, it can be shown that for any subshift $X \subset \mathcal{A}^G$ and any ergodic state $\mu \in \mathcal{M}_\sigma(X)$, there exists a potential $\phi \in C(X)$ for which μ is the unique equilibrium state [13]. It is therefore quite natural to retain some regularity assumptions on the class of potentials under consideration.

3. Proof of Theorem 1.1

For our results, we will impose a natural regularity assumption on our potential ϕ , but impose no restriction on the subshift $X \subset \mathcal{A}^G$. Our proof of Theorem 1.1 will be adapted from the proof technique of García-Ramos and Pavlov in [11], which involves replacing v with w along a sufficiently sparse and regular grid in G . Lemma 3.1 allows us to construct a partition of G from which we may choose the appropriate grid. We will then prove a few technical lemmas, and finally we will show our main result and conclude with a few notable corollaries.

3.1. Sufficiently sparse almost partitions. The following lemma allows us to generate a finite ϵ -almost partition of G where each part is sufficiently sparse relative to some fixed $F \in G$.

LEMMA 3.1. *For any $F \in G$, $\epsilon > 0$, and Følner sequence $\{F_n\}$, there exists a finite collection $\mathcal{P} = \{P_i : 1 \leq i \leq N\}$ of pairwise disjoint, left F -sparse subsets of G and a subsequence $\{F_{n_k}\}$ such that*

$$\liminf_{k \rightarrow \infty} \frac{|\bigcup_{P \in \mathcal{P}} P \cap F_{n_k}|}{|F_{n_k}|} \geq 1 - \epsilon.$$

In lieu of a complete and technical proof, we provide an outline of how this result can be shown. The result can be viewed as a weakened reformulation of a quasitiling result of Ornstein and Weiss in [21], or an application of the exact tiling results in [10]. In essence, one can construct a quasitiling (or an exact tiling) of G using a finite collection of shapes that are sufficiently F -invariant. The F -invariance ensures that we can consider only the F -interiors of these shapes and maintain a $(1 - \epsilon)$ -covering of G . We then let $P \in \mathcal{P}$ represent all the shifts of a particular location of an element in a specified shape. This allows us to ensure that \mathcal{P} is left F -sparse since we have restricted to the F -interiors of the relevant shapes.

3.2. *Subshift lemmas.* We begin by describing our regularity constraints imposed on the potential ϕ .

Definition 3.2. The F -variation of ϕ for any $F \subset G$ is defined as

$$\text{Var}_F(\phi) = \sup\{\phi(x) - \phi(y) : x, y \in X \text{ and } x_F = y_F\}.$$

Definition 3.3. We say $\{E_n\}$ is an *exhaustive sequence* in G if $E_1 \subset E_2 \subset \dots$ and $G = \bigcup_{n \in \mathbb{N}} E_n$.

Definition 3.4. We say ϕ has *summable variation according to the exhaustive sequence* $\{E_n\}$ if

$$\sum_{n \in \mathbb{N}} |E_{n+1}^{-1} \setminus E_n^{-1}| \cdot \text{Var}_{E_n}(\phi) < \infty.$$

Additionally, we will say $\phi \in C(X)$ has *summable variation* if ϕ has summable variation according to some exhaustive sequence.

For a fixed exhaustive sequence $\{E_n\}$, we define $SV_{\{E_n\}}(X)$ to be the collection of all potentials with summable variation according to $\{E_n\}$. We also let $SV(X) = \bigcup_{\{E_n\}} SV_{\{E_n\}}(X)$ denote the collection of all potentials with summable variation according to some exhaustive sequence.

It is worth noting here that when $G = \mathbb{Z}^d$, we can take $E_n = \{k \in \mathbb{Z}^d : |k_i| \leq n\}$ and summable variation according to this sequence corresponds to d -summable variation in the typical sense.

Definition 3.5. For an exhaustive sequence $\{E_n\}$, we define the *summable variation norm* on $SV_{\{E_n\}}(X)$ by

$$\|\phi\|_{SV_{\{E_n\}}} = 2|E_1| \cdot \|\phi\|_\infty + \sum_{n \in \mathbb{N}} |E_{n+1}^{-1} \setminus E_n^{-1}| \cdot \text{Var}_{E_n}(\phi).$$

OBSERVATION 3.6. *If ϕ has summable variation according to some exhaustive sequence $\{E_n\}$, then for any $F \Subset G$ such that $e \in F$, ϕ has summable variation according to exhaustive sequence $\{E_n F\}$.*

Proof. First, let $H \Subset G$ and note $H F^{-1}$ is also a finite set. We therefore have some $N \in \mathbb{N}$ such that for all $n \geq N$, $H F^{-1} \subset E_n$ and so $H \subset E_n F$. It immediately follows that $\{E_n F\}$ is an exhaustive sequence.

Since $e \in F$ by assumption, we have $E_n \subset E_n F$. We now see that

$$\sum_{n \in \mathbb{N}} |E_{n+1} F \setminus E_n F| \text{Var}_{E_n F}(\phi) \leq |F| \sum_{n \in \mathbb{N}} |E_{n+1} \setminus E_n| \text{Var}_{E_n}(\phi) < \infty. \quad \square$$

We now restate relevant definitions and lemmas from García-Ramos and Pavlov in [11].

Definition 3.7. For $v, u \in L(X)$, we define $O_v(u) = \{g \in G : \sigma_g(u)_F = v\}$.

In particular, $O_v(u)$ represents the indices where v occurs in u and can be read as ‘occurrences of v in u ’. We will now define a replacement function that for a given finite configuration u , replaces occurrences of v with w in specified locations.

Definition 3.8. Let $F, H \subseteq G$ and fix any $v, w \in L_F(X)$ and $u \in L_H(X)$. Let $S \subset O_v(u)$ be an F -sparse set of occurrences of v in u . We now define $R_u^{v \rightarrow w}(S) = u'$ as follows:

- for $s \in S, f \in F$, let $u'_{sf} = w_f$; and
- for all other $g \in H \setminus SF$, let $u'_g = u_g$.

Since S is an F -sparse subset of $O_v(u)$, we know u' is well defined and uniquely determined. Note here that u' is exactly the configuration obtained by replacing v with w in the S -locations.

We remind the reader that for any finite configuration $v \in L_F(X)$, the extender set of v is defined as $E_X(v) = \{\eta \in \mathcal{A}^{F^c} : v\eta \in X\}$. Note here, whenever $E_X(v) \subset E_X(w)$, then $R_u^{v \rightarrow w}(S) \in L(X)$ for any left F -sparse set $S \subset O_v(u)$.

LEMMA 3.9. [11, Lemma 4.2] For any $F, v, w \in L_F(X)$ with $v \neq w$, and left F -sparse set $T \subset O_v(u)$, $R_u^{v \rightarrow w}$ is injective on subsets of T .

LEMMA 3.10. [11, Lemma 4.3] For any F and $v, w \in L_F(X)$, any left F -sparse set $T \subset O_v(u)$, any u' , and any $m \leq |T \cap O_w(u')|$,

$$|\{(u, S) : S \text{ is left } F\text{-sparse, } |S| = m, S \subseteq T, u' = R_u^{v \rightarrow w}(S)\}| \leq \binom{|T \cap O_w(u')|}{m}.$$

The following lemma will be useful for computing topological pressure.

LEMMA 3.11. Let μ_ϕ be an ergodic equilibrium measure for any $\phi \in C(X)$. For any tempered Følner sequence $\{F_n\}$, if $S_n \subset L_{F_n}(X)$ such that $\mu_\phi(S_n) \rightarrow 1$, then

$$P_{\text{top}}(\phi) = \lim_{n \rightarrow \infty} \frac{1}{|F_n|} \log \left(\sum_{w \in S_n} \sup_{x \in [w]} \exp(\phi_{F_n}(x)) \right).$$

Proof. Let $\epsilon > 0$ and note by definition of topological pressure above, we have

$$\limsup_{n \rightarrow \infty} \frac{1}{|F_n|} \log \left(\sum_{w \in S_n} \sup_{x \in [w]} \exp(\phi_{F_n}(x)) \right) \leq P_{\text{top}}(\phi).$$

For every n , define

$$T_n = \left\{ w \in L_{F_n}(X) : \mu_\phi([w]) < \exp \left(|F_n| \left(\sup_{x \in [w]} \phi_{F_n}(x) - (P_{\text{top}}(\phi) - \epsilon) \right) \right) \right\}.$$

Note since F_n is a tempered Følner sequence and μ_ϕ is ergodic, we can apply the pointwise ergodic theorem and the Shannon–Macmillan–Breiman theorem proven in [12, 21] to see for μ_ϕ -a.e. $x \in X$,

$$P_{\text{top}}(\phi) = h(\mu_\phi) + \int \phi \, d\mu_\phi = \lim_{n \rightarrow \infty} |F_n|^{-1}(\phi_{F_n}(x) - \log \mu_\phi([x_{F_n}])).$$

By the definition of T_n , it therefore follows that $\mu_\phi(\bigcup_{N \in \mathbb{N}} \bigcap_{n \geq N} T_n) = 1$, and so $\mu_\phi(T_n) \rightarrow 1$. Further, $\mu(S_n \cap T_n) \rightarrow 1$ and by definition of T_n ,

$$\sum_{w \in S_n \cap T_n} \sup_{x \in [w]} \exp(\phi_{F_n}(x)) \geq \mu_\phi(S_n \cap T_n) \exp(|F_n|(P_{\text{top}}(\phi) - \epsilon)).$$

By taking sufficiently large n , we therefore have

$$\sum_{w \in S_n} \sup_{x \in [w]} \exp(\phi_{F_n}(x)) \geq 0.5 \exp(|F_n|(P_{\text{top}}(\phi) - \epsilon)).$$

Since $\epsilon > 0$ was arbitrary, we are finished. □

We remind the reader that for $v, w \in L_F(X)$, $\xi_{v,w} : X \rightarrow X$ is the map that swaps v and w in the F location whenever legal.

OBSERVATION 3.12. *If ϕ has summable variation according to the exhaustive sequence $\{E_n\}$, then for any $F \in G$, $v, w \in L_F(X)$, and any $x \in [v]$,*

$$\sum_{g \in G} |\phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x)))| \leq |F| \cdot \|\phi\|_{SV_{\{E_n\}}}.$$

We note the proof of Observation 3.12 is standard and the statement appears nearly identically as [1, Proposition 3.1]. The final lemma in this section is a Stirling approximation that will be necessary to compute a lower bound on pressure in the proof of Theorem 1.1.

LEMMA 3.13. *For $b, a \in \mathbb{Q}_+$, and any sequence $n_k \in \mathbb{N}$ such that for all k , $k!$ divides n_k , let $D \in \mathbb{R}$ satisfying $\log(a/b) > D$. Then, for sufficiently small $c \in \mathbb{Q}_+$, we have*

$$\lim_{k \rightarrow \infty} n_k^{-1} \left(\log \binom{an_k}{cn_k} - \log \binom{bn_k}{cn_k} \right) > cD.$$

Proof. We define for all $c \in (0, \min\{a, b\}) \cap \mathbb{Q}$,

$$f(c) = \lim_{k \rightarrow \infty} n_k^{-1} \left(\log \binom{an_k}{cn_k} - \log \binom{bn_k}{cn_k} \right).$$

First note, we have for all $k \in \mathbb{N}$,

$$\left(\log \binom{an_k}{cn_k} - \log \binom{bn_k}{cn_k} \right) = \log(an_k)! + \log((b - c)n_k)! - \log(bn_k)! - \log((a - c)n_k)!.$$

If we divide by n_k and take the limit as k goes to infinity, Stirling’s approximation implies

$$f(c) = a \log \frac{a}{a - c} + b \log \frac{b - c}{b} + c \log \frac{a - c}{b - c}.$$

We now examine

$$f'(c) = \frac{a}{a-c} - \frac{b}{b-c} + \log \frac{a-c}{b-c} + c \frac{b-a}{(a-c)(c-b)}.$$

Note here $f'(0) = \log(a/b) > D$. Since $f(0) = 0$, it immediately follows that for sufficiently small $c \in \mathbb{Q}_+$, we know $f(c) > cD$. □

3.3. Proof of Theorem 1.1.

THEOREM 3.1. *Let G be a countable amenable group and X be a G -subshift. Let $\phi \in SV(X)$, μ_ϕ an equilibrium state for ϕ , $F \subseteq G$ and $v, w \in L_F(X)$. If $E_X(v) \subset E_X(w)$, then*

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \sup_{x \in [v]} \exp \left(\sum_{g \in G} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x))) \right).$$

First, we let X, ϕ, F, v and w be as in the statement of the theorem. By ergodic decomposition, it is sufficient to show the desired result for ergodic equilibrium states, so we let μ_ϕ be an ergodic equilibrium state for ϕ . Fix $C = \sup_{x \in [v]} \sum_{g \in G} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x)))$ and note by Observation 3.12, we know $-\infty < C < \infty$. We now suppose for a contradiction that $\mu_\phi([v]) > \mu_\phi([w]) \cdot e^C$.

We now take $\delta \in (0, \frac{4}{3} \mu_\phi([v]))$ satisfying

$$e^{-C} \cdot \frac{\mu_\phi([v]) - 5\delta/4}{\mu_\phi([w]) + 2\delta} > 1.$$

Note here, this is attainable for all sufficiently small δ since we know

$$f(\delta) = e^{-C} \cdot \frac{\mu_\phi([v]) - 5\delta/4}{\mu_\phi([w]) + 2\delta}$$

is continuous and, by assumption, $f(0) > 1$.

Let F_n be a Følner sequence such that for each n , $n!$ divides $|F_n|$ and $F_n^{-1} = F_n$. It is noted by Xu and Zheng in [24] that we can assume F_n is tempered by passing to a subsequence. Re-index this sequence by n , and thus since F_n is tempered, we know it satisfies the requirements for the lemmas above.

We now define $S_n \subset L_{F_n}(X)$ as follows:

$$S_n := \{u \in L_{F_n}(X) : |O_v(u)| \geq |F_n|(\mu_\phi([v]) - \delta) \text{ and } |O_w(u)| \leq |F_n|(\mu_\phi([w]) + \delta)\}.$$

We note here since μ_ϕ is ergodic, $\mu_\phi(S_n) \rightarrow 1$ and we can therefore apply Lemma 3.11 to see

$$P_{\text{top}}(\phi) = \lim_{n \rightarrow \infty} |F_n|^{-1} \log \sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)).$$

We now apply Lemma 3.1 with respect to F and $\epsilon = \delta/8$, re-index our Følner sequence according to the lemma, and let $\mathcal{P} = \{P_i : 1 \leq i \leq N\}$ be our collection of pairwise

disjoint, left F -sparse sets. We also pass to the tail of the sequence of Følner sets ensuring that for each n , we have

$$\frac{|\bigcup_{i \leq N} P_i \cap F_n|}{|F_n|} \geq 1 - \frac{\delta}{4}.$$

For each $1 \leq i \leq N$, we now define

$$S_n^i = \left\{ u \in S_n : |O_v(u) \cap P_i| \geq \frac{\mu_\phi([v]) - 5\delta/4}{N} |F_n| \right\}.$$

LEMMA 3.2. For each $n \in \mathbb{N}$, $S_n = \bigcup_{1 \leq i \leq N} S_n^i$.

Proof. Suppose there exists some $u \in S_n \setminus \bigcup_{1 \leq i \leq N} S_n^i$. We therefore know for all $P \in \mathcal{P}$, $|O_v(u) \cap P| < (\mu_\phi([v]) - 5\delta/4)/N |F_n|$. Let $Q = F_n \setminus \bigcup_{i \leq N} P_i$ and note that

$$|O_v(u)| = \sum_{i=1}^N |O_v(u) \cap P_i| + |O_v(u) \cap Q| < (\mu_\phi([v]) - 5\delta/4) |F_n| + |O_v(u) \cap Q|.$$

However, since $u \in S_n$, we know $(\mu_\phi([v]) - \delta) \cdot |F_n| < |O_v(u)|$. It follows that

$$(\mu_\phi([v]) - \delta) \cdot |F_n| < (\mu_\phi([v]) - 5\delta/4) |F_n| + |O_v(u) \cap Q|.$$

By our construction of \mathcal{P} and since we took a sufficiently long tail for F_n , we know that $|O_v(u) \cap Q| \leq |Q \cap F_n| \leq (\delta/4) |F_n|$. We therefore have $\mu_\phi([v]) - \delta < \mu_\phi([v]) - \delta$, arriving at a contradiction, and we can conclude that $S_n = \bigcup_{1 \leq i \leq N} S_n^i$. \square

LEMMA 3.3. There exists some fixed $1 \leq i \leq N$ and a subsequence (n_k) for which we have

$$P_{\text{top}}(\phi) = \lim_{k \rightarrow \infty} |F_{n_k}|^{-1} \log \sum_{u \in S_{n_k}^i} \sup_{x \in [u]} \exp(\phi_{F_{n_k}}(x)).$$

Proof. First notice, since $S_n = \bigcup_{i \leq N} S_n^i$,

$$\sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \leq \sum_{i=1}^N \sum_{u \in S_n^i} \sup_{x \in [u]} \exp(\phi_{F_n}(x)),$$

and we therefore have for each n , some $1 \leq i_n \leq N$ such that

$$\frac{1}{N} \sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \leq \sum_{u \in S_n^{i_n}} \sup_{x \in [u]} \exp(\phi_{F_n}(x)).$$

We now use the pigeonhole principle to find a subsequence n_k such that $i_{n_k} = i$ is constant. Note, by our construction, we have

$$\sum_{u \in S_{n_k}^i} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \geq N^{-1} \sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)).$$

We now examine

$$\begin{aligned}
 P_{\text{top}}(\phi) &= \lim_{n \rightarrow \infty} |F_n|^{-1} \sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \\
 &\geq \lim_{n \rightarrow \infty} |F_n|^{-1} \log \sum_{u \in S'_{n_k}} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \\
 &\geq \lim_{n \rightarrow \infty} |F_n|^{-1} \log N^{-1} \sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \\
 &= \lim_{n \rightarrow \infty} |F_n|^{-1} \log N^{-1} + \lim_{n \rightarrow \infty} |F_n|^{-1} \sum_{u \in S_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) = P_{\text{top}}(\phi). \quad \square
 \end{aligned}$$

We now apply Lemma 3.3 and fix the resulting $1 \leq i \leq N$ and re-index along the resulting subsequence (n_k) . Denote $S'_k = S'_{n_k}$ and fix $P = P_i$. We can therefore compute topological pressure by restricting to the sequence of collections of finite configurations S'_n . We will now make replacements of v with w in every $u \in S'_n$ at locations in P with a small frequency to increase the pressure of the dynamical system to arrive at our contradiction.

We let $a, b \in \mathbb{Q}_+$ such that $a \leq (\mu_\phi([v]) - 5\delta/4)/N$, $b \geq (\mu_\phi([w]) + 2\delta)/N$, and $\log(a/b) > C$. Note, such $a, b \in \mathbb{Q}_+$ must exist since $\log(\mu_\phi([v]) - 5\delta/4)/(\mu_\phi([w]) + 2\delta) > C$ by assumption. We now let $\epsilon \in \mathbb{Q}_+$ be sufficiently small to satisfy Lemma 3.13 with respect to a, b and $D = C$.

Define for each $u \in S'_n$,

$$A_u = \{R_u^{v \rightarrow w}(S) : S \subset O_v(u) \cap P \text{ and } |S| = \epsilon \cdot |F_n|\}.$$

Since $\epsilon \in \mathbb{Q}$ and $n!$ divides $|F_n|$, for sufficiently large n , $\epsilon \cdot |F_n| \in \mathbb{N}$ and A_u is well defined. We restrict our consideration for all n large enough that this definition makes sense. Additionally, since $E_X(v) \subset E_X(w)$, we have $A_u \subset L(X)$.

We can now define for each $n \in \mathbb{N}$, $L_n = \bigcup_{u \in S'_n} A_u$.

LEMMA 3.4. For each $n \in \mathbb{N}$, $\sum_{u \in L_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x))$ is bounded below by

$$\frac{(\sum_{u \in S'_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x) - (\epsilon \cdot |F_n|/N)C)) \binom{\lceil (\mu_\phi([v]) - 5\delta/4) |F_n|/N \rceil}{\epsilon \cdot |F_n|}}{\binom{\lfloor (\mu_\phi([w]) + 2\delta) |F_n|/N \rfloor}{\epsilon \cdot |F_n|}}. \tag{1}$$

Proof. Note here, for each $u \in S'_n$,

$$|A_u| = \binom{|O_v(u) \cap P|}{\epsilon \cdot |F_n|} \geq \binom{\lceil (\mu_\phi([v]) - 5\delta/4) |F_n|/N \rceil}{\epsilon \cdot |F_n|}.$$

However, for every $u' \in \bigcup_{u \in S'_n} A_u$, we know u' came from some $u \in S_n$ by replacing the $\epsilon \cdot |F_n|/N$ v terms with w . We can therefore get a bound on the following:

$$|O_w(u') \cap P| \leq |O_w(u) \cap P| + \frac{\epsilon \cdot |F_n|}{N} < \frac{(\mu_\phi([w]) + \delta) |F_n|}{N} + \frac{\delta |F_n|}{|FF^{-1}| \cdot N}.$$

Since $|FF^{-1}| \geq 1$, we therefore have

$$|O_w(u') \cap P| < \frac{(\mu_\phi([w]) + 2\delta) |F_n|}{N}.$$

We therefore know for any fixed $u' \in A_u$, there are at most

$$\left(\frac{\lfloor (\mu_\phi([w]) + 2\delta) |F_n| / N \rfloor}{\epsilon \cdot |F_n|} \right)$$

u terms for which $u' \in A_u$.

It follows that $\sum_{u \in L_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x))$ is bounded below by

$$\frac{\left(\sum_{u \in S'_n} \sum_{u' \in A_u} \sup_{x \in [u']} \exp(\phi_{F_n}(x)) \right)^{\lceil (\mu_\phi([v]) - 5\delta/4) |F_n| / N \rceil}}{\left(\frac{\lfloor (\mu_\phi([w]) + 2\delta) |F_n| / N \rfloor}{\epsilon \cdot |F_n|} \right)}. \tag{2}$$

We now bound for any fixed $u \in S'_n$ and each $u' \in A_u$, $\sup_{x \in [u']} \phi_{F_n}(x)$. First, note $u' = R_u^{v \rightarrow w}(S)$ for some $S \subset F_n$. Since $E_X(v) \subset E_X(w)$ and by our choice of S , we know $E_X(u) \subset E_X(u')$. We therefore know $\xi_{u,u'}$ (as defined above) is injective and non-identity on $[u]$. In particular, we are replacing our v terms located at S with w terms just as we did in the construction of u' . Note here, $\xi_{u,u'}([u]) \subset [u']$ and thus, we have

$$\sup_{x \in [u']} \phi_{F_n}(x) \geq \sup_{x \in \xi_{u,u'}([u])} \phi_{F_n}(x) = \sup_{x \in [u]} \phi_{F_n}(\xi_{u,u'}(x)).$$

We now fix any $x \in [u]$. Note that since S is finite and left F -sparse, we may make the v to w replacements sequentially. Each of these replacements takes the form $\sigma_{g^{-1}}(\xi_{v,w}(\sigma_g(x)))$ for some $g \in G$. Note here, for each replacement of this kind, by definition of C , we have

$$\phi_{F_n}(\sigma_{g^{-1}}(\xi_{v,w}(\sigma_g(x)))) \geq \phi_{F_n}(x) - C.$$

Since we will make $|S|$ of these replacements, it follows that

$$\phi_{F_n}(\xi_{u,u'}(x)) \geq \phi_{F_n}(x) - |S|C.$$

We therefore know that

$$\sum_{u \in S'_n} \sum_{u' \in A_u} \sup_{x \in [u']} \exp(\phi_{F_n}(x)) \geq \sum_{u \in S'_n} |A_u| \sup_{x \in [u]} \exp(\phi_{F_n}(x) - |S|C).$$

Since $|A_u| \geq 1$, we therefore have

$$\sum_{u \in S'_n} \sum_{u' \in A_u} \sup_{x \in [u']} \exp(\phi_{F_n}(x)) \geq \sum_{u \in S'_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x) - |S|C). \tag{3}$$

Combining equations (2) and (3), we arrive at our desired result that $\sum_{u \in L_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x))$ is bounded below by equation (1). \square

We will conclude the proof of Theorem 1.1 by computing a lower bound on topological pressure and arrive at a contradiction. Since $L_n \subset L_{F_n}(X)$, it must be the case that

$$P_{\text{top}}(\phi) \geq \limsup_{n \rightarrow \infty} |F_n|^{-1} \log \left(\sum_{u \in L_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \right).$$

By application of Lemma 3.4, we can see that $P_{\text{top}}(\phi)$ is bounded below by

$$\liminf_{n \rightarrow \infty} |F_n|^{-1} \log \frac{(\sum_{u \in S'_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x) - (\epsilon \cdot |F_n|/N)C)) \binom{\lceil (\mu_\phi([v]) - 5\delta/4) |F_n|/N \rceil}{\epsilon \cdot |F_n|}}{\binom{\lfloor (\mu_\phi([w]) + 2\delta) |F_n|/N \rfloor}{\epsilon \cdot |F_n|}},$$

which is exactly equal to

$$\begin{aligned} & \liminf_{n \rightarrow \infty} |F_n|^{-1} \log \left(\sum_{u \in S'_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \right) - \epsilon \cdot C \\ & + |F_n|^{-1} \log \binom{\lceil (\mu_\phi([v]) - 5\delta/4) |F_n|/N \rceil}{\epsilon \cdot |F_n|} - |F_n|^{-1} \log \binom{\lfloor (\mu_\phi([w]) + 2\delta) |F_n|/N \rfloor}{\epsilon \cdot |F_n|}. \end{aligned} \tag{4}$$

We now note by our choice of a , b , and ϵ ,

$$\begin{aligned} & \lim_{n \rightarrow \infty} |F_n|^{-1} \log \binom{\lceil (\mu_\phi([v]) - 5\delta/4) |F_n|/N \rceil}{\epsilon \cdot |F_n|} - |F_n|^{-1} \log \binom{\lfloor (\mu_\phi([w]) + 2\delta) |F_n|/N \rfloor}{\epsilon \cdot |F_n|} \\ & \geq \lim_{n \rightarrow \infty} |F_n|^{-1} \log \binom{a|F_n|}{\epsilon \cdot |F_n|} - |F_n|^{-1} \log \binom{b|F_n|}{\epsilon \cdot |F_n|} > \epsilon C. \end{aligned}$$

It therefore follows that equation (4) is strictly greater than

$$\liminf_{n \rightarrow \infty} |F_n|^{-1} \log \left(\sum_{u \in S'_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \right),$$

and since equation (4) is a lower bound for pressure, we therefore know

$$P_{\text{top}}(\phi) > \liminf_{n \rightarrow \infty} |F_n|^{-1} \log \left(\sum_{u \in S'_n} \sup_{x \in [u]} \exp(\phi_{F_n}(x)) \right).$$

This contradicts Lemma 3.3. In particular, this means our assumption that $\mu_\phi([v]) > \mu_\phi([w]) \cdot e^C$ was incorrect, arriving at our desired result that if $E_X(v) \subset E_X(w)$, then

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \sup_{x \in [v]} \exp \left(\sum_{g \in G} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x))) \right).$$

3.4. *Locally constant corollaries.* We will now explore the case where ϕ is locally constant.

The following result shows that when the finite configurations v and w agree on a sufficiently chosen boundary, we may compute bounds for their relative measure using only information from the configurations themselves. Further, when they have equal extender sets, we have a closed form formula depending only on ϕ , v , and w to compute the ratio of their measures.

We will say that ϕ is an H -potential for some $H \in G$ if for all $x, y \in X$ such that $x_H = y_H$, then $\phi(x) = \phi(y)$, that is, $\text{Var}_H(\phi) = 0$. Additionally, for any $H \subset G$, we will denote $H^\pm = H \cup H^{-1}$.

COROLLARY 3.5. *Let $H, F \in G$, $v, w \in L_F(X)$, and ϕ be an H -potential. Suppose that $E_X(v) \subset E_X(w)$ and for all $g \in F^c H^\pm \cap F$, $v_g = w_g$ (that is, v and w agree on their H -boundary). Then, for any equilibrium state μ_ϕ for ϕ ,*

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \exp\left(\sum_{g \in F \setminus F^c H^{-1}} \phi(\sigma_g(v)) - \phi(\sigma_g(\xi_{v,w}(w)))\right).$$

Proof. We first note, for all $x \in [v]$ and for any $g \in F \cap F^c H^{-1}$, by assumption, we have $x_{gH} = \xi_{v,w}(x)_{gH}$. Since ϕ is an H -potential, it immediately follows that for all $g \in F \cap F^c H^{-1}$,

$$\phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x))) = 0.$$

We now consider $g \in F^c$. If $gH \cap F = \emptyset$, then $x_{gH} = \xi_{v,w}(x)_{gH}$ by definition of $\xi_{v,w}$. We now consider $gH \cap F \neq \emptyset$. However, by assumption, since $g \in F^c$, we know for all $f \in gH \cap F$, $v_f = w_f$ and again we can conclude $x_{gH} = \xi_{v,w}(x)_{gH}$. Thus, we know for all $g \in F^c$,

$$\phi \circ \sigma_g(x) - \phi \circ \sigma_g(\xi_{v,w}(x)) = 0.$$

Since ϕ is locally constant, it has summable variation, and we can apply Theorem 1.1 to see

$$\mu_\phi([v]) \leq \mu_\phi([w]) \sup_{x \in [v]} \exp\left(\sum_{g \in F \setminus F^c H^{-1}} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x)))\right).$$

We let $x, y \in [v]$, $g \in F \setminus F^c H^\pm$, and $h \in H$. By construction, we know $g \notin F^c H^{-1}$ and, thus, $gH \cap F^c = \emptyset$. Since ϕ is an H -potential, it follows that $\phi \circ \sigma_g(x) = \phi \circ \sigma_g(y)$, and similarly for $\xi_{v,w}(x)$ and $\xi_{v,w}(y)$. We can therefore conclude that $\sum_{g \in F \setminus F^c H^\pm} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x)))$ does not depend on the choice of $x \in [v]$ and can be computed by only looking at the cylinder set, and we have arrived at our desired result. \square

As an immediate corollary, we have the following.

COROLLARY 3.6. *Let $H, F \in G$, $v, w \in L_F(X)$, and ϕ be an H potential. Suppose that $E_X(v) = E_X(w)$ and for all $g \in F^c H^\pm \cap F$, $v_g = w_g$ (that is, v and w agree on their H -boundary). Then, for any equilibrium state μ_ϕ for ϕ ,*

$$\frac{\mu_\phi([v])}{\exp\left(\sum_{g \in F \setminus F^c H^{-1}} \phi \circ \sigma_g(v)\right)} = \frac{\mu_\phi([w])}{\exp\left(\sum_{g \in F \setminus F^c H^{-1}} \phi \circ \sigma_g(w)\right)}.$$

We remind the reader that in the case of $\phi = 0$, equilibrium states correspond with measures of maximal entropy. In this case, as a corollary of Theorem 1.1, we extend the results of García-Ramos and Pavlov in [11] to the countable amenable subshift setting.

COROLLARY 3.7. *Let X be a G -subshift, $F \in G$, $v, w \in L_F(X)$, and μ a measure of maximal entropy. If $E_X(v) \subset E_X(w)$, then $\mu([v]) \leq \mu([w])$.*

4. Conformal Gibbs results

4.1. *Homoclinic relation and conformal Gibbs.* As previously mentioned, a measure μ is Gibbs for a potential ϕ (in the sense of the theorem of Dobrušin and that of Lanford–Ruelle) if and only if it is conformal Gibbs for ϕ . This section will explore our conformal Gibbs-like result, for which we will follow the definition found in [3].

First, we define the *homoclinic relation*, also known as the *Gibbs relation*, as $\mathfrak{T}_X \subset X \times X$ such that $(x, y) \in \mathfrak{T}_X$ if and only if $x_{F^c} = y_{F^c}$ for some $F \in G$. We now say a Borel isomorphism $f : X \rightarrow X$ is a *holonomy* of \mathfrak{T}_X if for all $x \in X$, $(x, f(x)) \in \mathfrak{T}_X$. It is known (see [3]) that there exists a countable group Γ of holonomies of X such that $\mathfrak{T}_X = \{(x, \gamma(x)) : x \in X, \gamma \in \Gamma\}$. In fact, by a trivial extension of [3, Lemma 1], we can see immediately that $\Gamma = \langle \xi_{v,w} : v, w \in L_F(X), F \in G \rangle$ is a countable group of holonomies such that $\mathfrak{T}_X = \{(x, \gamma(x)) : x \in X, \gamma \in \Gamma\}$.

For any measurable $A \subset X$, we define $\mathfrak{T}_X(A) = \bigcup_{x \in A} \{y \in X : (x, y) \in \mathfrak{T}_X\}$. For a Borel measure μ on X , we say μ is *\mathfrak{T}_X -non-singular* if for all null sets $A \subset X$, $\mu(\mathfrak{T}_X(A)) = 0$. Note for any holonomy $f : X \rightarrow X$, if $A \subset X$ is a null set and μ is \mathfrak{T}_X -non-singular, then $\mu \circ f(A) \leq \mu(\mathfrak{T}_X(A)) = 0$ and therefore $\mu \circ f \ll \mu$.

We define a *cocycle on \mathfrak{T}_X* to be any measurable $\psi : \mathfrak{T}_X \rightarrow \mathbb{R}$ such that for all $x, y, z \in X$ with $(x, y), (y, z) \in \mathfrak{T}_X$, we have $\psi(x, y) + \psi(y, z) = \psi(x, z)$. We can now define conformality following [3, Definition 1].

Definition 4.1. Let μ be a \mathfrak{T}_X -non-singular Borel probability measure on X and let $\psi : \mathfrak{T}_X \rightarrow \mathbb{R}$ be a cocycle. We say μ is *(ψ, \mathfrak{T}_X) -conformal* if for any holonomy $f : X \rightarrow X$, we have

$$\frac{d\mu \circ f}{d\mu}(x) = \exp(\psi(x, f(x))).$$

Since we are concerned with Gibbs measures, in general, for a potential $\phi \in SV(X)$, we define the following cocycle:

$$\psi_\phi(x, y) = \sum_{g \in G} \phi \circ \sigma_g(y) - \phi \circ \sigma_g(x)$$

and we will say that a measure μ is *conformal Gibbs* for ϕ if it is $(\psi_\phi, \mathfrak{T}_X)$ -conformal.

Connecting this concept further to the classical notion of Gibbs measures, it was shown in [3] that a measure is conformal Gibbs for ϕ if and only if it is Gibbs for ϕ in the sense of Definition 2.8.

In addition to the homoclinic relation, in [17], Meyerovitch defined a subrelation as follows. First, let

$$\mathcal{F}(X) = \{f \in \text{Homeo}(X) : \text{there exists } F \in G \text{ such that for all } x \in X, f(x)_{F^c} = x_{F^c}\}.$$

In particular, $\mathcal{F}(X)$ is the group of all background preserving homeomorphisms. We then define $\mathfrak{T}_X^0 = \{(x, f(x)) : x \in X, f \in \mathcal{F}(X)\}$. Clearly, $\mathfrak{T}_X^0 \subset \mathfrak{T}_X$; however, in general, these are not necessarily equal.

Notice that when X is an SFT, then each $\xi_{v,w}$ is in fact a homeomorphism and thus $\xi_{v,w} \in \mathcal{F}(X)$. In this case, $\mathfrak{T}_X^0 = \mathfrak{T}_X$. Additionally, even if X is not an SFT, if

$E_X(v) = E_X(w)$, $\xi_{v,w}$ is again a homeomorphism. However, it need not be the case that $\xi_{v,w}$ is continuous in general.

Example 4.2. Take $X \subset \{0, 1\}^{\mathbb{Z}}$ to the orbit closure of $x = 0^\infty 10^\infty$ (which is sometimes referred to as the sunny-side-up subshift). We define $\xi_{1,0}$ to swap a 1 and 0 in the identity location, and note here that $E_X(1) \subsetneq E_X(0)$. For each $n \geq 0$, define $x_n \in X$ such that $(x_n)_n = 1$. Then, for all $n \geq 1$, $\xi_{1,0}(x_n) = x_n$. However, we have $x_n \rightarrow 0^\infty$ and $\xi_{1,0}(0^\infty) = x_0$. It immediately follows that

$$\lim_{n \rightarrow \infty} \xi_{1,0}(x_n) = 0^\infty \neq x_0 = \xi_{1,0}(\lim_{n \rightarrow \infty} x_n)$$

and we can conclude $\xi_{1,0}$ is not continuous.

In the case of $G = \mathbb{Z}^d$ and $\phi \in C(X)$ has d -summable variation, Meyerovitch proved the following result.

THEOREM 4.3. (Meyerovitch [17]) *Let $X \subset \mathcal{A}^{\mathbb{Z}^d}$ be a subshift, $\phi \in C(X)$ a potential with d -summable variation, and μ_ϕ be an equilibrium state for ϕ that is \mathfrak{T}_X -non-singular. Then, μ_ϕ is $(\psi_\phi, \mathfrak{T}_X^0)$ -conformal.*

In particular, μ_ϕ is conformal Gibbs with respect to the sub-relation \mathfrak{T}_X^0 . What this means in the terminology of our paper is for $G = \mathbb{Z}^d$ and ϕ with d -summable variation, if $E_X(v) = E_X(w)$, then for μ_ϕ -a.e. $x \in X$, $(d(\mu \circ \xi_{v,w})/d\mu)(x) = \exp(\psi_\phi(x, \xi_{v,w}(x)))$. We will use Theorem 1.1 to extend this result both to the general amenable group setting as well as to provide an inequality on the derivative when $E_X(v) \subset E_X(w)$.

4.2. Proof of conformal Gibbs result. In this section, we will prove the following theorem.

THEOREM 4.4. *Let $F \Subset G$, $v, w \in L_F(X)$, $\phi \in SV(X)$, and μ_ϕ be an equilibrium state for ϕ . If $E_X(v) \subset E_X(w)$, then $\mu_\phi \circ \xi_{v,w}$ is absolutely continuous with respect to μ_ϕ when restricted to $[w]$ and for μ_ϕ -a.e. $x \in [w]$,*

$$\frac{d(\mu \circ \xi_{v,w})}{d\mu}(x) \leq \exp(\psi_\phi(x, \xi_{v,w}(x))).$$

For the remainder of this section, we fix some $F \Subset G$, $v, w \in L_F(X)$, $\phi \in SV(X)$, and μ_ϕ an equilibrium state for ϕ .

OBSERVATION 4.5. *If $E_X(v) \subset E_X(w)$, then $\mu_\phi \circ \xi_{v,w} \ll \mu_\phi$ when restricted to $[w]$.*

Proof. Let $A \subset [w]$ be a null set. Note here

$$\xi_{v,w}(A) = \{x \in A : \xi_{v,w}(x) = x\} \sqcup \{\xi_{v,w}(x) : x \in A \text{ and } \xi_{v,w}(x) \in [v]\}.$$

Since $\{x \in A : \xi_{v,w}(x) = x\} \subset A$, it must be a null set, and thus it is sufficient to assume without loss of generality that $\xi_{v,w}(A) \cap A = \emptyset$.

Since we know μ_ϕ is outer regular, we can let $U_n = \bigcup_{i \leq N_n} [w_{n,i}]$ be an outer approximation by cylinder sets such that $\mu_\phi(A) = \lim_{n \rightarrow \infty} \mu_\phi(U_n)$. By taking sufficiently

large n , we can assume without loss of generality that $[w_{n,i}] \subset [w]$ for all n, i . Note for each n, i , we have $E_X(\xi_{v,w}(w_{n,i})) \subset E_X(w_{n,i})$. We now apply Theorem 1.1 with respect to $\phi, \mu_\phi, v_{n,i}$, and $w_{n,i}$ to see

$$\begin{aligned} \mu_\phi(\xi_{v,w}(A)) &= \lim_{n \rightarrow \infty} \sum_{i \leq N_n} \mu_\phi(\xi_{v,w}([w_{n,i}])) \\ &\leq \lim_{n \rightarrow \infty} \sum_{i \leq N_n} \mu_\phi([w_{n,i}]) \sup_{x \in [v]} \exp(f(x)) = 0. \end{aligned} \quad \square$$

In particular, this means we can discuss the LRN derivative when restricted to $[w]$. We now define for each $n \in \mathbb{N}$, $\phi_n : X \rightarrow \mathbb{R}$ to be an E_n potential approximating ϕ from above. In particular, for all $u \in L_{E_n}(X)$, for any $x \in [u]$, let $\phi_n(x) = \sup_{y \in [u]} \phi(y)$.

OBSERVATION 4.6. ϕ_n converges to ϕ with respect to the summable variation norm induced by $\{E_n\}$.

Proof. First note, for any $k, n \in \mathbb{N}$, we have

$$\text{Var}_{E_k}(\phi_n - \phi) \leq 2\|\phi\|_\infty \quad \text{and} \quad \text{Var}_{E_k}(\phi_n - \phi) \leq 2\text{Var}_{E_k}(\phi).$$

Let $\epsilon > 0$ and note since ϕ has summable variation according to $\{E_n\}$, by definition, there exists some $M \in \mathbb{N}$ such that $\sum_{k=M+1}^\infty |E_{k+1} \setminus E_k| \text{Var}_{E_k}(\phi) < \epsilon$. We fix this M . We now note by continuity of ϕ and since $\{E_n\}$ is an exhaustive sequence, we know there exists some $N \in \mathbb{N}$ such that for all $n \geq N$,

$$\|\phi_N - \phi\|_\infty < \epsilon \left(2|E_1| + \sum_{k=1}^M |E_{k+1} \setminus E_k| \right)^{-1}.$$

We now let $n \geq N$ and we examine

$$\begin{aligned} \|\phi_N - \phi\|_{SV_{\{E_n\}}} &= 2|E_1| \cdot \|\phi_n - \phi\|_\infty + \sum_{k=1}^M |E_{k+1} \setminus E_k| \text{Var}_{E_k}(\phi_n - \phi) \\ &\quad + \sum_{k=M+1}^\infty |E_{k+1} \setminus E_k| \text{Var}_{E_k}(\phi_n - \phi) \\ &\leq \|\phi_n - \phi\|_\infty \left(2|E_1| + \sum_{k=1}^M |E_{k+1} \setminus E_k| \right) + 2 \sum_{k=M+1}^\infty |E_{k+1} \setminus E_k| \text{Var}_{E_k}(\phi). \end{aligned}$$

By our choice of N and M , we know for all $n \geq N$, $\|\phi_N - \phi\|_{SV_{\{E_n\}}} < 3\epsilon$. Since ϵ was arbitrary, we can conclude the proof of the observation. \square

OBSERVATION 4.7. If $E_X(v) \subset E_X(w)$, then the function $f(x) = \psi_\phi(x, \xi_{v,w}(x))$ is continuous on $[v]$.

Proof. Using our locally constant approximations ϕ_n , we define for each $n \in \mathbb{N}$,

$$f_n(x) = \psi_{\phi_n}(x, \xi_{v,w}(x)) = \sum_{g \in G} \phi_n(\sigma_g(\xi_{v,w}(x))) - \phi_n(\sigma_g(x)).$$

Since ϕ_n is locally constant, we know the infinite sum is in fact a finite sum and so f_n is continuous on $[v]$. We now claim that $f_n \rightarrow f$ uniformly on $[v]$. By Observation 3.12, we know for all $x \in [v]$,

$$|f_n(x) - f(x)| \leq |F| \cdot \|\phi_n - \phi\|_{SV_{\{E_n\}}}.$$

Since $\phi_n \rightarrow \phi$ with respect to the summable variation norm by Observation 4.6, we can conclude that $f_n \rightarrow f$ uniformly on $[v]$. It immediately follows that f is continuous on $[v]$. □

We can combine the above observations with Theorem 1.1 to prove the desired result.

THEOREM 4.3. *Let $F \subseteq G$, $v, w \in L_F(X)$, $\phi \in SV(X)$, and μ_ϕ be an equilibrium state for ϕ . If $E_X(v) \subset E_X(w)$, then for μ_ϕ -a.e. $x \in [w]$,*

$$\frac{d\mu \circ \xi_{v,w}}{d\mu}(x) \leq \exp(\psi_\phi(x, \xi_{v,w}(x))).$$

Proof. First, we fix $F \subseteq G$, $v, w \in L_F(X)$, $\phi \in C(X)$, and μ_ϕ as in the theorem. We again define for all $x \in X$,

$$f(x) = \psi_\phi(x, \xi_{v,w}(x)) = \sum_{g \in G} \phi(\sigma_g(\xi_{v,w}(x))) - \phi(\sigma_g(x)),$$

which we know to be continuous on $[v]$ by Observation 4.7.

We fix any Følner sequence $\{F_n\}$ and assume without loss of generality that $F \subset F_n$ for all n . Fix any $x \in [w]$. We define $w_n = x_{F_n}$ and $v_n = \xi_{v,w}(x)_{F_n}$. Note here $\bigcap_{n \in \mathbb{N}} [w_n] = x$, $\bigcap_{n \in \mathbb{N}} [v_n] = \xi_{v,w}(x)$. Since $F \subset F_n$, we know $E_X(v_n) \subset E_X(w_n)$. We can directly apply Theorem 1.1 to see for each $n \in \mathbb{N}$,

$$\mu_\phi(v_n) \leq \mu_\phi(w_n) \sup_{y \in [v_n]} \exp(-f(y)).$$

Since by Observation 4.7, we know f is continuous on $[v]$ and thus there exists some $y_n \in [v_n]$ such that $\sup_{y \in [v_n]} \exp(-f(y)) = \exp(-f(y_n))$. We therefore have

$$\frac{\mu_\phi(v_n)}{\mu_\phi(w_n)} \leq \exp(-f(y_n)).$$

Since $y_n \in [v_n]$ and $\bigcap_{n \in \mathbb{N}} [v_n] = \{\xi_{v,w}(x)\}$, we know $y_n \rightarrow \xi_{v,w}(x)$. We can now take limits to see

$$\frac{d\mu_\phi \circ \xi_{v,w}}{d\mu_\phi}(x) = \lim_{n \rightarrow \infty} \frac{\mu_\phi(v_n)}{\mu_\phi(w_n)} \leq \lim_{n \rightarrow \infty} \exp(-f(y_n)) = \exp(f(x)). \quad \square$$

Finally, we note that whenever μ_ϕ satisfies the equations in Theorem 1.3, it is trivial to show μ_ϕ must also satisfy the equations in Theorem 1.1.

OBSERVATION 4.4. *Let X be a subshift and $u, v \in L_F(X)$ such that $E_X(v) \subset E_X(w)$. Let $\phi \in C(X)$ and μ_ϕ satisfy for μ_ϕ -a.e. $x \in [w]$,*

$$\frac{d(\mu \circ \xi_{v,w})}{d\mu}(x) \leq \exp(\psi_\phi(x, \xi_{v,w}(x))).$$

Then,

$$\mu_\phi([v]) \leq \mu_\phi([w]) \cdot \sup_{x \in [v]} \exp\left(\sum_{g \in G} \phi(\sigma_g(x)) - \phi(\sigma_g(\xi_{v,w}(x)))\right).$$

Proof. First, we notice that since $E_X(v) \subset E_X(w)$, then $[v] \subseteq \xi_{v,w}([w])$. Further, since $\xi_{v,w}$ is an involution, we know

$$\mu_\phi([v]) = \mu_\phi(\xi_{v,w}(\xi_{v,w}([v]))) = \int_{x \in \xi_{v,w}([v])} \frac{d(\mu_\phi \circ \xi_{v,w})}{d\mu_\phi}(x) d\mu_\phi.$$

Since by assumption, for μ_ϕ -a.e. $x \in [w]$,

$$\frac{d(\mu \circ \xi_{v,w})}{d\mu}(x) \leq \exp(\psi_\phi(x, \xi_{v,w}(x))),$$

we know that

$$\int_{x \in \xi_{v,w}([v])} \frac{d(\mu_\phi \circ \xi_{v,w})}{d\mu_\phi}(x) d\mu_\phi \leq \int_{x \in \xi_{v,w}([v])} \exp(\psi_\phi(\xi_{v,w}(x), \xi_{v,w}(\xi_{v,w}(x)))) d\mu_\phi.$$

It follows immediately that

$$\mu_\phi([v]) \leq \mu_\phi(\xi_{v,w}([v])) \cdot \sup_{y \in [v]} \exp(\psi_\phi(\xi_{v,w}(x), x)).$$

Since $\xi_{v,w}([v]) \subset [w]$, we arrive at our desired conclusion. □

We will now note that as a corollary of Theorem 1.3, we can extend [17, Theorem 3.1 and Corollary 3.2] to the countable amenable group subshift setting. We have become aware that [2, Theorem B] extends this further to the countable sofic group subshift setting.

COROLLARY 4.5. *Let $X \subset \mathcal{A}^G$ be a subshift over a countable amenable group G , let $\phi \in SV(X)$ be a potential with summable variation, and let μ_ϕ be an equilibrium state for ϕ . Then, μ_ϕ is $(\mathfrak{T}_X^0, \psi_\phi)$ -conformal.*

Proof. First note, as observed in [17], the collection $\{\xi_{v,w} : E_X(v) = E_X(w)\}$ generates $\mathcal{F}(X)$. By application of the results in [17], it is therefore sufficient to show that for any $v, w \in L_F(X)$ with $E_X(v) = E_X(w)$, we have for all $x \in X$,

$$\frac{d\mu \circ \xi_{v,w}}{d\mu}(x) = \exp(\psi_\phi(x, \xi_{v,w}(x))).$$

Let $v, w \in L_F(X)$ such that $E_X(v) = E_X(w)$. Let $x \in [w]$ and for all $n \in \mathbb{N}$, define $v_n, w_n \in L_{F_n}(X)$ as in the proof of Theorem 1.3. By application of Theorem 1.1, we can see that

$$\inf_{y \in [v_n]} \exp(-f(y)) \leq \frac{\mu_\phi([v_n])}{\mu_\phi([w_n])} \leq \sup_{y \in [v_n]} \exp(-f(y)).$$

By taking limits, we can conclude that for all $x \in [w]$,

$$\frac{d\mu \circ \xi_{v,w}}{d\mu}(x) = \exp(\psi_\phi(x, \xi_{v,w}(x))).$$

The proof is similar for $x \in [v]$ and for all $x \in X \setminus ([v] \cup [w])$, we know

$$\frac{d\mu \circ \xi_{v,w}}{d\mu}(x) = \exp(\psi_\phi(x, \xi_{v,w}(x))) = 1.$$

Again, by following the techniques in [17], it follows that μ_ϕ is (Σ_X^0, ψ_ϕ) -conformal. \square

Acknowledgements. This work was conducted during the author's PhD studies under the supervision of Dr Ronnie Pavlov. The author would like to thank Dr Pavlov for all of his assistance throughout the research and writing process. We would also like to thank the referee for their helpful report.

REFERENCES

- [1] S. Barbieri, R. Gómez, B. Marcus, T. Meyerovitch and S. Taati. Gibbsian representations of continuous specifications: the theorems of Kozlov and Sullivan revisited. *Comm. Math. Phys.* **382** (2021), 1111–1164.
- [2] S. Barbieri and T. Meyerovitch. The Lanford–Ruelle theorem for actions of sofic groups. *Trans. Amer. Math. Soc.* **376**(02) (2023), 1299–1342.
- [3] L. Borsato and S. MacDonald. Conformal measures and the Dobrušin–Lanford–Ruelle equations. *Proc. Amer. Math. Soc.* **149**(10) (2021), 4355–4369.
- [4] R. Bowen. Some systems with unique equilibrium states. *Math. Systems Theory* **8**(3) (1974/75), 193–202.
- [5] A. I. Bufetov. Pressure and equilibrium measures for actions of amenable groups on the space of configurations. *Sb. Math.* **202**(3) (2011), 341.
- [6] R. Burton and J. E. Steif. Non-uniqueness of measures of maximal entropy for subshifts of finite type. *Ergod. Th. & Dynam. Sys.* **14**(2) (1994), 213–235.
- [7] R. Burton and J. E. Steif. New results on measures of maximal entropy. *Israel J. Math.* **89**(1–3) (1995), 275–300.
- [8] T. Ceccherini-Silberstein, M. Coornaert, T. Ceccherini-Silberstein and M. Coornaert. Amenable groups. *Cellular Automata and Groups*. Springer, Berlin, 2010, pp. 77–110.
- [9] R. L. Dobrushin. Gibbsian random fields for lattice systems with pairwise interactions. *Funct. Anal. Appl.* **2**(4) (1968), 292–301.
- [10] T. Downarowicz, D. Huczek and G. Zhang. Tilings of amenable groups. *J. Reine Angew. Math.* **2019**(747) (2019), 277–298.
- [11] F. García-Ramos and R. Pavlov. Extender sets and measures of maximal entropy for subshifts. *J. Lond. Math. Soc. (2)* **100**(3) (2019), 1013–1033.
- [12] B. M. Gurevich and A. A. Tempelman. A Breiman type theorem for Gibbs measures. *J. Dyn. Control Syst.* **13**(3) (2007), 363–371.
- [13] O. Jenkinson. Every ergodic measure is uniquely maximizing. *Discrete Contin. Dyn. Syst.* **16**(2) (2006), 383.
- [14] S. Kass and K. Madden. A sufficient condition for non-soficness of higher-dimensional subshifts. *Proc. Amer. Math. Soc.* **141**(11) (2013), 3803–3816.
- [15] O. E. Lanford III and D. Ruelle. Observables at infinity and states with short range correlations in statistical mechanics. *Comm. Math. Phys.* **13**(3) (1969), 194–215.
- [16] E. Lindenstrauss. Pointwise theorems for amenable groups. *Invent. Math.* **146**(2) (2001), 259–295.
- [17] T. Meyerovitch. Gibbs and equilibrium measures for some families of subshifts. *Ergod. Th. & Dynam. Sys.* **33**(3) (2013), 934–953.
- [18] J. M. Ollagnier and D. Pinchon. Mesures de gibbs invariantes et mesures d'équilibre. *Z. Wahrsch. Verw. Gebiete* **55** (1981), 11–23.
- [19] J. M. Ollagnier and D. Pinchon. The variational principle. *Studia Math.* **72** (1982), 151–159.
- [20] N. Ormes and R. Pavlov. Extender sets and multidimensional subshifts. *Ergod. Th. & Dynam. Sys.* **36**(3) (2016), 908–923.

- [21] D. S. Ornstein and B. Weiss. Entropy and isomorphism theorems for actions of amenable groups. *J. Anal. Math.* **48**(1) (1987), 1–141.
- [22] W. Parry. Symbolic dynamics and transformations of the unit interval. *Trans. Amer. Math. Soc.* **122**(2) (1966), 368–378.
- [23] A. A. Tempelman. Specific characteristics and variational principle for homogeneous random fields. *Z. Wahrsch. Verw. Gebiete* **65**(3) (1984), 341–365.
- [24] L. Xu and L. Zheng. Weak mean equicontinuity for a countable discrete amenable group action. *J. Dynam. Differential Equations* **36** (2022), 1–14.