

THREE CHARACTERIZATIONS OF STRICT COHERENCE ON INFINITE-VALUED EVENTS

TOMMASO FLAMINIO

IIIA - CSIC, Campus de la Universidad Autònoma de Barcelona

Abstract. This article builds on a recent paper coauthored by the present author, H. Hosni and F. Montagna. It is meant to contribute to the logical foundations of probability theory on many-valued events and, specifically, to a deeper understanding of the notion of strict coherence. In particular, we will make use of geometrical, measure-theoretical and logical methods to provide three characterizations of strict coherence on formulas of infinite-valued Łukasiewicz logic.

§1. Introduction and motivation. In a collection of seminal contributions starting with [5] and culminating in [6], de Finetti grounded subjective probability theory on an ideal betting game between two players, a bookmaker and a gambler, who wager money on the occurrence of certain events e_1, \dots, e_k . For each event e_i , gambler's payoffs are 1 in case e_i occurs, and 0 otherwise. The *probability* of an event e_i is defined, by de Finetti, as the *fair selling price* fixed by the bookmaker for it.

Conforming to a standard notation, bookmaker's prices for the events e_1, \dots, e_k will be referred to as *betting odds* and an assignment $\beta : \{e_1, \dots, e_k\} \rightarrow [0, 1]$ of betting odds $\beta(e_i) = \beta_i$ will be called a *book*.

De Finetti had no particular inclination towards identifying events in a precise logical ground [9]. However, in order for his main result to be stated in precise mathematical terms, they will be understood, for the moment, as elements of a finitely generated free boolean algebra and hence coded by boolean formulas. Now, de Finetti's result reads as follows: let us fix finitely many events e_1, \dots, e_k and a book β on them. A gambler must choose *stakes* $\sigma_1, \dots, \sigma_k \in \mathbb{R}$, one for each event, and pay to the bookmaker the amount $\sigma_i \cdot \beta_i$ for each e_i . When a (classical propositional) valuation w determines e_i , the gambler gains σ_i if $w(e_i) = 1$ and 0 otherwise. The book β is said to be *coherent* if there is no choice of stakes $\sigma_1, \dots, \sigma_k \in \mathbb{R}$ such that for every valuation w

$$\sum_{i=1}^k \sigma_i \cdot \beta_i - \sum_{i=1}^k \sigma_i \cdot w(e_i) = \sum_{i=1}^k \sigma_i (\beta_i - w(e_i)) < 0. \quad (1)$$

The left-hand side of (1) captures the bookmaker's payoff, or *balance, relative to the book β under the valuation w* .

Note that a stake σ_i may be negative. Following tradition, money transfers are so oriented that "positive" means "gambler-to-bookmaker." Therefore, if $\sigma_i < 0$, the bookmaker is

Received: June 21, 2018.

2010 *Mathematics Subject Classification*: Primary 03B48, Secondary 60A05, 06D35, 06F20, 52B11.

Key words and phrases: de Finetti's coherence, strict coherence, faithful states, MV-algebras, Łukasiewicz logic.

forced to swap his role with the gambler: he has to pay $-\sigma_i \cdot \beta(e_i)$ to the gambler in hopes of winning $-\sigma_i$ in case e_i occurs.

De Finetti's Dutch-Book theorem characterizes coherent books as follows: a book β on events e_1, \dots, e_k pertaining to a boolean algebra \mathbf{A} is coherent iff it extends to a finitely additive probability P of \mathbf{A} , [5].

Along with the assumptions which regulate de Finetti's coherence criterion, condition (1) above effectively forces the bookmaker to set *fair prices* for gambling on events e_1, \dots, e_k . In other words, upon regarding each event e_i as a $\{0, 1\}$ -valued random variable, de Finetti's Dutch-Book theorem amounts to saying that coherent assessments are those with null expectation. For, if the bookmaker publishes a book with positive expectation (for him) a logically infallible gambler will choose negative stakes and inflict a *sure loss* on him, that is to say, a sure financial loss whatever the outcome of events.

Although coherence guards the bookmaker against the possibility of sure loss, at the same time it may bar him from making a profit. To illustrate the idea, consider an event e which is neither noncontradictory nor sure and the coherent book $\beta(e) = 0$. If the gambler bets 1 on e , then her balance is as follows: she pays $1 \cdot 0 = 0$ and gets back 0 if $w(e) = 0$ and 1 if $w(e) = 1$. Hence, the bookmaker never wins and possibly loses.

This rather odd feature of coherence was questioned in the mid 1950's first by Shimony [32] and then by Kemeny [16]. These authors studied a refinement of de Finetti's coherence that nowadays goes under the name of *strict coherence* (see [11]). Intuitively, a choice of prices is strictly coherent if every possibility of loss, for the bookmaker, is paired by a possibility of gain. Precisely, a book β is *strictly coherent* if, for each choice of stakes $\sigma_1, \dots, \sigma_k \in \mathbb{R}$, the existence of a valuation w such that $\sum_{i=1}^k \sigma_i(\beta_i - w(e_i)) < 0$ implies the existence of another one w' for which $\sum_{i=1}^k \sigma_i(\beta_i - w'(e_i)) > 0$.

Interest in the condition of strict coherence was prompted by Carnap's analysis of what he called "regular" probability functions in [1] (see also [31, Chap. 10]) and which we will term as *Carnap probabilities*. Those functions arise from the axiomatization of finitely additive probabilities by strengthening the usual normalization axiom in the right-to-left direction: 1 (respectively, 0) is assigned only to tautologies (respectively, contradictions). In other words, a probability function P is Carnap, if it is normalized, finitely additive and it satisfies $P(e) \neq 0$ for every noncontradictory event e . In [11] the authors characterized strictly coherent books on boolean events in terms of their extendability to Carnap probabilities.¹

Several authors proposed generalizations of de Finetti's coherence criterion and his Dutch-Book theorem to events not pertaining to boolean logic. Paris in [30] extended the classical Dutch-Book theorem to several nonclassical propositional logics including the modal logics K, T, S4, S5 and certain paraconsistent logics as well. In [34], Weatherston considered the case of events pertaining to intuitionistic logic and in [26], Mundici extended de Finetti's criterion to the case of infinite-valued Łukasiewicz logic and MV-algebras [3, 27].

In the MV-algebraic realm valuations are $[0, 1]$ -valued and hence they correspond to homomorphisms into the *standard MV-algebra* defined on the unit interval $[0, 1]$. De Finetti's coherence criterion immediately translates to the MV-setting with no extra

¹ Carnap probabilities are the same as Carnap-regular probabilities of [11]. In the present article we adopt this simplified notation in order to avoid any misleading interpretation of the adjective "regular" which has indeed different meanings if referred to probability functions or to Borel measures which will be discussed in §4.

conditions and the main result of [26] (see also [19]) is a de Finetti-like theorem which characterizes coherent books on Łukasiewicz events as those which are extendible to states, i.e., $[0, 1]$ -valued normalized and finitely additive maps of an MV-algebra.

From the perspective of reasoning about uncertainty, the interest in Łukasiewicz events is twofold: on the one hand these events capture properties of the world which are better described as *gradual* rather than *yes-or-no*; on the other hand, they also mimic bounded random variables. Indeed, any Łukasiewicz event e may be regarded as a $[0, 1]$ -valued continuous function f_e on a compact Hausdorff space (see [3, Theorem 9.1.5] and §2) and any state of e coincides with the expected value of f_e ([17, 29], [10, Remark 2.8] and §4). Therefore, up to renormalization, Mundici's generalization of de Finetti's theorem [26, Theorem 2.1] implies the Dutch-Book theorem for books on bounded continuous random variables.

For events pertaining to the restricted class of *finite-dimensional* MV-algebras, in [11, Theorem 6.4] the authors proved a de Finetti-like theorem for strictly coherent books in terms of their extendability to *faithful* states, i.e., states which satisfy $s(a) \neq 0$ for all $a \neq 0$. Nevertheless, extending [11, Theorem 6.4] to more general classes of MV-algebras is delicate because, as a consequence of seminal results by Mundici [25, Prop. 3.2], Kelley [15], and Gaifman [13], an MV-algebra may not have a faithful state.

In this article we will investigate strictly coherent books on Łukasiewicz events, i.e., elements of a finitely generated free MV-algebra. Our results sensibly extend the results of [11]. In particular, we will provide three characterizations of strict coherence by adopting geometrical, measure-theoretical and logical methods. In more details:

Geometrical approach: the functional representation of n -generated free MV-algebras in terms of n -variable, piecewise-linear continuous functions (see [3, Theorem 9.1.5] and [23]) implies that the set of all coherent books on a finite set Φ of Łukasiewicz events forms a convex polyhedron \mathcal{D}_Φ of \mathbb{R}^k . The main result of §3 shows that strictly coherent books on Φ form a subset of \mathbb{R}^k which coincides with the relative interior of \mathcal{D}_Φ .

Measure-theoretical approach: faithful states are the MV-algebraic analog of Carnap probabilities on a boolean algebra. In §4 we will first give an integral representation theorem for faithful states on finitely generated free MV-algebras and then we will characterize strictly coherent books on Łukasiewicz events as those which extend to a faithful state.

Logical approach: the relation among free MV-algebras, rational polyhedra and deducibility in propositional Łukasiewicz logic, will enable us to characterize the notions of coherence and strict coherence within propositional Łukasiewicz logic (see §5). In our opinion this result is interesting because it shows that propositional Łukasiewicz logic is capable to capture foundational aspects of probability theory on infinite-valued events.

In the next section we will introduce necessary preliminaries about MV-algebras and rational polyhedra.

§2. Preliminaries. The algebraic framework of this article is that of MV-algebras (see [3, 27]), i.e., the Lindenbaum algebras of Łukasiewicz infinite-valued logic [3, Definition 4.3.1]. A typical example of an MV-algebra is the *standard algebra* $[0, 1]_{MV} = ([0, 1], \oplus, \neg, 0)$ where $x \oplus y = \min\{1, x + y\}$ and $\neg x = 1 - x$. Further operations, together with their standard interpretation, are defined in $[0, 1]_{MV}$ as follows: $x \odot y = \neg(\neg x \oplus \neg y) = \max\{0, x + y - 1\}$, $x \rightarrow y = \neg x \oplus y = \min\{1, 1 - x + y\}$, $x \wedge y = x \odot (x \rightarrow y) = \min\{x, y\}$, $x \vee y = \neg(\neg x \wedge \neg y) = \max\{x, y\}$, $1 = \neg 0$. This structure generates the class of MV-algebras both as a variety and as quasi-variety [2].

Another relevant example of an MV-algebra is given by the free n -generated MV-algebra \mathbf{F}_n . By a standard universal algebraic argument, \mathbf{F}_n is the MV-algebra of functions $f : [0, 1]^n \rightarrow [0, 1]$ generated by the projection maps [3, Prop. 3.1.4] and whose operations $\odot, \oplus, \rightarrow, \wedge, \vee$ and \neg are defined via the pointwise application of those in $[0, 1]_{MV}$. By McNaughton theorem, up to isomorphism, \mathbf{F}_n coincides with the MV-algebra of n -variable *McNaughton functions*: maps from $[0, 1]^n$ to $[0, 1]$ which are continuous, piecewise linear, with finitely many pieces, and such that each piece has integer coefficients (cf. [3, Theorem 9.1.5] and [23]). For each $f \in \mathbf{F}_n$, the *oneset* of f is $\{x \in [0, 1]^n \mid f(x) = 1\}$ and the *zeroset* of f is $\{x \in [0, 1]^n \mid f(x) = 0\}$.

The free n -generated MV-algebra is, up to isomorphism, the Lindenbaum algebra of Łukasiewicz logic \mathbf{L} in a language with n propositional variables and $[0, 1]$ -valuations of \mathbf{L} are exactly the homomorphisms of \mathbf{F}_n to $[0, 1]_{MV}$. Furthermore, every $x \in [0, 1]^n$ determines the homomorphism $h_x : f \in \mathbf{F}_n \mapsto f(x) \in [0, 1]_{MV}$.

PROPOSITION 2.1 ([26, Lemma 3.1]). *For each finite n , homomorphisms of \mathbf{F}_n to $[0, 1]_{MV}$, $[0, 1]$ -valued valuations of Łukasiewicz logic on n variables and points of the n -cube $[0, 1]^n$ are in one-one correspondence.*

For every closed subset C of $[0, 1]^n$, let I_C be the subset of \mathbf{F}_n of those functions whose zeroset contains C . Then, I_C is an *ideal* of \mathbf{F}_n and the quotient \mathbf{F}_n/I_C is the MV-algebra whose universe coincides with the set given by the restrictions to C of the functions of \mathbf{F}_n (see [3, Prop. 3.4.5]). In particular, when C has k elements, the quotient MV-algebra \mathbf{F}_n/I_C is isomorphic to the product algebra $[0, 1]_{MV}^k$, [4]. The finite powers of $[0, 1]_{MV}$ — called *locally weakly finite* MV-algebras in [4]— are called, in this article, *finite-dimensional*.

2.1. Rational polyhedra, regular complexes and McNaughton functions. In this section we will prepare the necessary results concerning rational and regular complexes (see [8]) and their relation with finitely generated free MV-algebras. We invite the reader to consult [18, 27, 28] for background.

Let $k = 1, 2, \dots$. By a (*rational*) convex polyhedron (or (*rational*) *polytope*) of \mathbb{R}^k we mean the convex hull of finitely many points of \mathbb{R}^k (\mathbb{Q}^k respectively); a (*rational*) polyhedron is a finite union of (*rational*) convex polyhedra. Given any polytope \mathcal{P} , we respectively denote by $\text{ext } \mathcal{P}$, $\text{ri } \mathcal{P}$, $\text{rb } \mathcal{P}$ the set of its extremal points, its relative interior and its relative boundary. Since each polytope \mathcal{P} is closed, $\mathcal{P} = \text{ri } \mathcal{P} \cup \text{rb } \mathcal{P}$. Furthermore, for all vectors $x, y \in \mathbb{R}^k$, we denote $x \cdot y$ their scalar product and by $|x|$ the norm of x .

LEMMA 2.2. *For each polytope \mathcal{P} of \mathbb{R}^k , the following conditions hold:*

- (1) *For every $e \in \text{rb } \mathcal{P}$, there exists $\sigma \in \mathbb{R}^k$ such that, for all $\gamma \in \mathcal{P}$, $\sigma \cdot e \leq \sigma \cdot \gamma$;*
- (2) *Let $\beta \in \text{ri } \mathcal{P}$. Then, there exists $\sigma \in \mathbb{R}^k$ such that the sets $\mathcal{P}_\sigma^+ = \{\gamma \in \mathcal{P} \mid \gamma \cdot \sigma < \beta \cdot \sigma\}$ and $\mathcal{P}_\sigma^- = \{\gamma \in \mathcal{P} \mid \gamma \cdot \sigma > \beta \cdot \sigma\}$ are nonempty;*
- (3) *Let $\beta \in \text{ri } \mathcal{P}$. Then there exist $\sigma \in \mathbb{R}^k$, $e_1, e_2 \in \text{ext } \mathcal{P}$ such that $e_1 \in \mathcal{P}_\sigma^+$ and $e_2 \in \mathcal{P}_\sigma^-$;*
- (4) *$\gamma \in \text{ri } \mathcal{P}$ iff there exists a map $\lambda : \text{ext } \mathcal{P} \rightarrow [0, 1]$ such that $\sum_{e \in \text{ext } \mathcal{P}} \lambda(e) = 1$, $\lambda(e) > 0$ for all $e \in \text{ext } \mathcal{P}$ and $\gamma = \sum_{e \in \text{ext } \mathcal{P}} \lambda(e) \cdot e$.*

Before proving the lemma, recall that any hyperplane H of \mathbb{R}^k separates the space in two half spaces denoted H^+ and H^- . The above Claims (2) and (3) state that, if β is a point in the relative interior of a polytope \mathcal{P} , then there exists a hyperplane H passing through

β such that, respectively: both $H^+ \cap \mathcal{P}$ and $H^- \cap \mathcal{P}$ are nonempty; each $H^+ \cap \mathcal{P}$ and $H^- \cap \mathcal{P}$ contains an extremal point of \mathcal{P} .

Proof. (1) is the well-known supporting hyperplane theorem, see [22, Theorem 14].

(2) Let $\beta \in \text{ri } \mathcal{P}$. Let Σ be a sphere of radius r and centered at β and contained in $\text{ri } \mathcal{P}$. The existence of Σ is ensured by definition of relative interior [8, Chap. I, Definition 1.8]. Let σ be a vector of origin β . Suppose σ is not orthogonal to the affine hull of \mathcal{P} and also $0 < |\sigma| < r$. Trivially, $(\sigma - \beta) \cdot \sigma < \beta \cdot \sigma < (\sigma + \beta) \cdot \sigma$. Upon noting that $\sigma - \beta, \sigma + \beta \in \Sigma$, our claim is settled.

(3) By way of contradiction, assume that for no $e \in \text{ext } \mathcal{P}, e \cdot \sigma < \beta \cdot \sigma$. Equivalently, for all $e \in \text{ext } \mathcal{P}$,

$$e \cdot \sigma \geq \beta \cdot \sigma. \tag{2}$$

Since \mathcal{P}^+ is nonempty, in view of (2), let $\tau \in \mathcal{P} \cap \mathcal{P}^+$, i.e.,

$$\tau \cdot \sigma < \beta \cdot \sigma. \tag{3}$$

If $\tau \in \text{ext } \mathcal{P}$ the claim is settled. Assume that $\tau \notin \text{ext } \mathcal{P} = \{e_1, \dots, e_l\}$. Then there are $\lambda_1, \dots, \lambda_l \in [0, 1]$ such that $\sum_i \lambda_i = 1$ and $\tau = \sum_i \lambda_i \cdot e_i$. From (2) it follows that $e_i \cdot \sigma \geq \beta \cdot \sigma$, and hence $\sum \lambda_i e_i \cdot \sigma \geq \beta \cdot \sigma$, that is, $\tau \cdot \sigma \geq \beta \cdot \sigma$, which contradicts (3).

(4) See [11, Lemma 6.1(1)]. □

Let $k = 1, 2, \dots$ and let $x = \langle n_1/d_1, \dots, n_k/d_k \rangle$ be a rational vector in \mathbb{R}^k with n_i and d_i relatively prime for all $i = 1, \dots, k$. Denote by $\text{den}(x)$ the least common multiple of d_1, \dots, d_k . The *homogeneous correspondent* of x is the vector

$$\left\langle \frac{n_1}{d_1} \cdot \text{den}(x), \dots, \frac{n_k}{d_k} \cdot \text{den}(x), \text{den}(x) \right\rangle \in \mathbb{Z}^{k+1}.$$

Let $k = 1, 2, \dots$ and $m = 0, 1, \dots, k$. A rational m -simplex $\text{co}(x_1, \dots, x_m) \subseteq \mathbb{R}^k$ is said to be *regular* if the set of the homogeneous correspondents of x_1, \dots, x_m is part of a basis of the abelian group \mathbb{Z}^{m+1} [8, Chap. V, Definition 1.10]. A *regular complex* Δ is a simplicial complex all of whose simplexes are regular². Unless otherwise specified, all regular simplexes in this article are over the n -cube $[0, 1]^n$ (i.e., their faces constitute a *unimodular triangulations* of $[0, 1]^n$, in the terminology of [26]). Thus, we will say that Δ is a *regular complex of $[0, 1]^n$* without danger of confusion. From the regularity of Δ it follows that Δ is rational. We will denote by $V(\Delta)$ the finite set of rational vertices of Δ , i.e., the union of the set of the vertices of the simplexes in Δ .

Let Φ be a finite subset of the free n -generated MV-algebra \mathbf{F}_n . Up to isomorphism, we can (and we will, throughout this article) think of Φ as a finite set of n -variable McNaughton functions. Furthermore, if not otherwise specified, we will assume that Φ has k elements, denoted as f_1, \dots, f_k . Following [26, Sec. 3], for any Φ , there exists a regular complex Δ of $[0, 1]^n$ which *linearizes* Φ in the sense that each f_i is linear over each simplex of Δ .

EXAMPLE 2.3. *Let us fix $n = 2$ and consider $\Phi = \{x, y, x \oplus y\}$. Consider the regular complexes Δ_1 and Δ_2 of Figure 1 and whose vertices are $v_1^1 = \langle 0, 0 \rangle, v_2^1 = \langle 0, 1 \rangle, v_3^1 =$*

² Recall that a *simplicial complex* Δ is a nonempty finite set of simplexes such that: the face of each simplex in Δ belongs to Δ and for each pair of simplexes $T_1, T_2 \in \Delta$ their intersection is either empty, or it coincides with a common face of T_1 and T_2 .




Fig. 1. The two regular complexes Δ_1 (on the left) and Δ_2 (on the right) of the square $[0, 1]^2$. Every function x, y and $x \oplus y$ is linear over each simplex of Δ_1 and Δ_2 .

$\langle 1, 0 \rangle, v_4^1 = \langle 1, 1 \rangle, v_5^1 = \langle 1/2, 1/2 \rangle$ for Δ_1 and $v_1^2 = \langle 0, 0 \rangle, v_2^2 = \langle 0, 1 \rangle, v_3^2 = \langle 1, 0 \rangle, v_4^2 = \langle 1, 1 \rangle, v_5^2 = \langle 1/2, 1/2 \rangle, v_6^2 = \langle 1/3, 1/3 \rangle, v_7^2 = \langle 1/2, 0 \rangle, v_8^2 = \langle 0, 1/2 \rangle$ for Δ_2 . Then Δ_1 is union of four maximal simplexes, while Δ_2 is union of eight simplexes, see Figure 1

Both Δ_1 and Δ_2 linearize Φ . Indeed, for each regular simplex T of Δ_1 and each simplex T' of Δ_2 , the restriction of each function $f \in \Phi$ to T and T' is linear.

Let Δ be a regular complex of $[0, 1]^n$, and let v_i be one of its vertices. The *normalized Schauder hat at v_i (over Δ)* is the uniquely determined continuous function $\hat{h}_i : [0, 1]^n \rightarrow [0, 1]$ which is linear over each simplex of Δ and which attains the value 1 at v_i and 0 at all other vertices of Δ . The regularity of Δ ensures that each linear piece of each \hat{h}_i has integer coefficients and hence $\hat{h}_i \in \mathbf{F}_n$. By definition of normalized Schauder hat, Δ linearizes each \hat{h}_i . Furthermore, the following result holds:

LEMMA 2.4. Let Φ be a finite subset of \mathbf{F}_n , let Δ be a regular complex linearizing Φ and let v_1, \dots, v_t be the vertices of Δ . Then:

- (1) For each $i \neq j, \hat{h}_i \odot \hat{h}_j = 0$;
- (2) $\bigoplus_{i=1}^t \hat{h}_i = 1$;
- (3) For each $f \in \Phi, f = \bigoplus_{i=1}^t f(v_i) \cdot \hat{h}_i$;
- (4) Let $v_{i_1}, \dots, v_{i_l} \in V(\Delta)$, and let $\mathcal{P} = \text{co}(v_{i_1}, \dots, v_{i_l})$. Then the function $p = \bigoplus_{v_j \in \Delta \cap \mathcal{P}} \hat{h}_j$ is a McNaughton function whose onese is \mathcal{P} .

Proof. (1), (2), and (3) have been proved in [26, Lemma 3.4(ii), (iii), (iv), and (v)]. Let next prove (4). First of all, $p = \bigoplus_{v \in \Delta \cap \mathcal{P}} \hat{h}_v$ is a McNaughton function by definition. Further, for every vertex $v \in \Delta \cap \mathcal{P}, p(v) = 1$ by definition of normalized Schauder hat. If $x \in \mathcal{P} \setminus V(\Delta)$, let Σ be a simplex of Δ which contains x . The claim follows, since each \hat{h}_j is linear on Σ . □

§3. A geometric characterization of strict coherence. By Proposition 2.1, if $\Phi = \{f_1, \dots, f_k\}$ is a finite subset of \mathbf{F}_n and β a book on Φ , we can rephrase the definitions of coherence and strict coherence for β as follows:

- (1) β is coherent if for every $\sigma \in \mathbb{R}^k$, there exists $x \in [0, 1]^n$ such that $\sigma \cdot \langle f_1(x), \dots, f_k(x) \rangle \geq 0$.
- (2) β is strictly coherent if for every $\sigma \in \mathbb{R}^k$, the existence of $x \in [0, 1]^n$ such that $\sigma \cdot \langle f_1(x), \dots, f_k(x) \rangle < 0$, implies the existence of another $x' \in [0, 1]^n$ such that $\sigma \cdot \langle f_1(x'), \dots, f_k(x') \rangle > 0$.

Notice that a book β is coherent and not strictly coherent iff for any vector $\sigma \in \mathbb{R}^k$, one has that for all $x \in [0, 1]^n$, $\sigma \cdot \langle f_1(x), \dots, f_k(x) \rangle \leq 0$ and for some $x' \in [0, 1]^n$, $\sigma \cdot \langle f_1(x'), \dots, f_k(x') \rangle = 0$.

Throughout we will adopt the following notation:

$$\mathcal{D}_\Phi = \{\beta : \Phi \rightarrow [0, 1] \mid \beta \text{ is coherent}\}.$$

For any $X \subseteq [0, 1]^n$, $\mathcal{C}_\Phi(X)$ will denote the topological closure of the convex hull of all points of \mathbb{R}^k of the form $\langle f_1(x), \dots, f_k(x) \rangle$ for $\varphi_i \in \Phi$ and $x \in X$. In symbols,

$$\mathcal{C}_\Phi(X) = \text{cl co}\{\langle f_1(x), \dots, f_k(x) \rangle \mid \varphi_i \in \Phi, x \in X\}.$$

Whenever X is finite, $\mathcal{C}_\Phi(X) = \text{co}\{\langle f_1(x), \dots, f_k(x) \rangle \mid \varphi_i \in \Phi, x \in X\}$, which is a convex polytope. For the sake of readability, we will write \mathcal{C}_Φ instead of $\mathcal{C}_\Phi([0, 1]^n)$.

For every finite Φ , Mundici’s extension of de Finetti’s theorem to Łukasiewicz logic (see [26, Theorem 2.1]) shows that $\mathcal{D}_\Phi = \mathcal{C}_\Phi$.

LEMMA 3.1 ([26, Corollary 5.4]). *For any book $\beta : \Phi \rightarrow [0, 1]$ the following conditions are equivalent:*

- (1) β is coherent;
- (2) There exists a finite $X \subset [0, 1]^n$ such that, for each $f_i \in \Phi$, $\beta(f_i) \in \mathcal{C}_\Phi(X)$;
- (3) There exists a finite $X \subset [0, 1]^n$ with $|X| \leq n + 1$ such that, for each $f_i \in \Phi$, $\beta(f_i) \in \mathcal{C}_\Phi(X)$;
- (4) For each regular complex Δ of $[0, 1]^n$ linearizing Φ , $\beta(f_i) \in \mathcal{C}_\Phi(V(\Delta))$.

The next corollary is an immediate consequence of Lemma 3.1.

COROLLARY 3.2. *Let Φ be a finite set of Łukasiewicz formulas. Then, for every regular complex Δ which linearizes Φ ,*

$$\mathcal{D}_\Phi = \mathcal{C}_\Phi(V(\Delta)).$$

Thus, the coherence of a book β on Φ does not depend on the particular regular complex Δ chosen to linearize Φ . Moreover, since for each Δ , $V(\Delta)$ is finite, $\mathcal{C}_\Phi(V(\Delta))$ is a polytope coinciding with \mathcal{D}_Φ . Therefore, by the Krein–Milman theorem [8, Theorem 1.2], $\mathcal{C}_\Phi(V(\Delta))$ is the convex hull of the set of extremal points of \mathcal{D}_Φ , i.e., for every Δ ,

$$\mathcal{C}_\Phi(V(\Delta)) = \text{co ext } \mathcal{D}_\Phi.$$

The following example clarifies the claim made in Corollary 3.2.

EXAMPLE 3.3. *Let $\Phi = \{x, y, x \oplus y\}$ together with the regular complexes Δ_1 and Δ_2 of Example 2.3.*

Each of the five vertices v_i^1 of Δ_1 determines a point

$$p_i = \langle f_1(v_i^1), f_2(v_i^1), f_3(v_i^1) \rangle \in \mathbb{R}^3$$

(where $f_1(x, y) = x, f_2(x, y) = y$ and $f_3(x, y) = x \oplus y$) and

$$\mathcal{D}_\Phi = \mathcal{C}_\Phi(\{v_1^1, \dots, v_5^1\}) = \text{co}(\{p_1, \dots, p_5\}).$$

In particular: $p_1 = \langle 0, 0, 0 \rangle, p_2 = \langle 0, 1, 1 \rangle, p_3 = \langle 1, 0, 1 \rangle, p_4 = \langle 1, 1, 1 \rangle$ and $p_5 = \langle 1/2, 1/2, 1 \rangle$.

Similarly, for Δ_2 ,

$$\mathcal{D}_\Phi = \mathcal{C}_\Phi(\{v_1^2, \dots, v_8^2\}) = \text{co}(\{q_1, \dots, q_8\}),$$

where $q_1 = p_1 = \langle 0, 0, 0 \rangle$, $q_2 = p_2 = \langle 0, 1, 1 \rangle$, $q_3 = p_3 = \langle 1, 0, 1 \rangle$, $q_4 = p_4 = \langle 1, 1, 1 \rangle$, $q_5 = p_5 = \langle 1/2, 1/2, 1 \rangle$, $q_6 = \langle 1/3, 1/3, 2/3 \rangle$, $q_7 = \langle 1/2, 0, 1/2 \rangle$ and $q_8 = \langle 0, 1/2, 1/2 \rangle$.

Since both Δ_1 and Δ_2 linearize Φ ,

$$\mathcal{D}_\Phi = \mathcal{C}_\Phi(V(\Delta_1)) = \mathcal{C}_\Phi(V(\Delta_2))$$

(see Figure 2) and

$$\text{ext } \mathcal{C}_\Phi(V(\Delta_1)) = \text{ext } \mathcal{C}_\Phi(V(\Delta_2)) = \text{ext } \mathcal{D}_\Phi = \{p_1, p_2, p_3, p_4\}.$$

Let us write

$$\mathcal{K}_\Phi = \{\beta : \Phi \rightarrow [0, 1] \mid \beta \text{ is strictly coherent}\}.$$

The following theorem, which is the main result of this section, provides us with a geometric characterization of strict coherence for books on formulas of Łukasiewicz logic.

THEOREM 3.4. *Let Φ be a finite subset of \mathbf{F}_n . Then*

$$\mathcal{K}_\Phi = \text{ri } \mathcal{D}_\Phi.$$

Proof. Since $\mathcal{D}_\Phi = \mathcal{C}_\Phi$, we will prove the equivalent claim: $\mathcal{K}_\Phi = \text{ri } \mathcal{C}_\Phi$.

Trivially, $\mathcal{K}_\Phi \subseteq \mathcal{C}_\Phi$. Let us show that $\mathcal{K}_\Phi \subseteq \text{ri } \mathcal{C}_\Phi$. Since \mathcal{C}_Φ is closed,

$$\mathcal{C}_\Phi = \text{ri } \mathcal{C}_\Phi \cup \text{rb } \mathcal{C}_\Phi \text{ and } \text{ri } \mathcal{C}_\Phi \cap \text{rb } \mathcal{C}_\Phi = \emptyset.$$

Assume (absurdum hypothesis) that $\beta \in \mathcal{K}_\Phi \cap \text{rb } \mathcal{C}_\Phi$. By Lemma 2.2 (1) there exists $\sigma \in \mathbb{R}^k$ such that for all $\gamma \in \mathcal{C}_\Phi$,

$$\sigma \cdot \beta \leq \sigma \cdot \gamma.$$

Thus, for all $x \in [0, 1]^n$,

$$\sigma \cdot \beta \leq \sigma \cdot \langle f_1(x), \dots, f_k(x) \rangle \leq 0.$$

Therefore, β is coherent but not strictly coherent. This contradicts our hypothesis. Thus, $\mathcal{K}_\Phi \subseteq \text{ri } \mathcal{C}_\Phi$.

In order to prove the converse inclusion, assume that $\beta \in \text{ri } \mathcal{C}_\Phi$ and let $\sigma \in \mathbb{R}^k$ satisfy Lemma 2.2(2). Then

$$(\mathcal{C}_\Phi)_\sigma^+ = \{\gamma \in \mathcal{C}_\Phi \mid \gamma \cdot \sigma < \beta \cdot \sigma\}$$




Fig. 2. The convex polytope \mathcal{D}_Φ (in two perspectives) for $\Phi = \{x, y, x \oplus y\}$, and its extremal points p_1, p_2, p_3 and p_4 .

and

$$(\mathcal{C}_\Phi)_\sigma^- = \{\gamma \in \mathcal{C}_\Phi \mid \gamma \cdot \sigma > \beta \cdot \sigma\},$$

are nonempty.

Moreover, by Lemma 2.2(3), both $(\mathcal{C}_\Phi)_\sigma^+$ and $(\mathcal{C}_\Phi)_\sigma^-$ contain an extremal point of \mathcal{C}_Φ . Therefore, there are $x, x' \in [0, 1]^n$ such that $(f_1(x), \dots, f_k(x)) \cdot \sigma < \beta \cdot \sigma$ and $(f_1(x'), \dots, f_k(x')) \cdot \sigma > \beta \cdot \sigma$, that is, β is strictly coherent. \square

COROLLARY 3.5. *Let Φ and β be as above. Then the following conditions are equivalent:*

- (1) β is strictly coherent;
- (2) For each regular complex Δ which linearizes Φ , there is $\lambda : V(\Delta) \rightarrow [0, 1]$ such that $\sum_{v_i \in V(\Delta)} \lambda(v_i) = 1$, for all $v_i \in V(\Delta)$, $\lambda(v_i) > 0$ and $\beta(f_j) = \sum_{v_i \in V(\Delta)} \lambda(v_i) \cdot f_j(v_i)$;
- (3) There exists a map $\lambda : \text{ext } \mathcal{D}_\Phi \rightarrow [0, 1]$ such that $\sum_{e \in \text{ext } \mathcal{D}_\Phi} \lambda(e) = 1$, for all $e \in \text{ext } \mathcal{D}_\Phi$, $\lambda(e) > 0$ and $\beta(\varphi_i) = \sum_{e \in \text{ext } \mathcal{D}_\Phi} \lambda(e) \cdot f_i(e)$.

Furthermore, the set $\mathcal{K}_\Phi^{\mathbb{Q}}$ of rational-valued strictly coherent books on Φ is decidable.

Proof. The equivalence between (1), (2), and (3) follows from Theorem 3.4, Corollary 3.2 and Lemma 2.2(4). To conclude the proof we will prove the decidability of the set $\mathcal{K}_\Phi^{\mathbb{Q}}$. To this purpose, given Φ , the problem of determining a regular complex Δ which linearizes all McNaughton functions $f_i \in \Phi$, is computable by a Turing machine (see [26, Theorem 7.1, Claim 3]). Therefore, by (2), $\mathcal{K}_\Phi^{\mathbb{Q}}$ is decidable iff the following *bounded mixed integer programming* problem (see [14]) with unknowns $\lambda(v_i)$ for all $v_i \in V(\Delta)$, has a solution in $[0, 1] \cap \mathbb{Q}$:

$$(S_{\mathcal{K}_\Phi}) = \begin{cases} \lambda(v_i) > 0, \\ \sum_{v_i} \lambda(v_i) = 1, \\ \sum_{v_i} \lambda(v_i) \cdot f_\varphi(v_i) = \beta(\varphi). \end{cases}$$

Thus the decidability of $\mathcal{K}_\Phi^{\mathbb{Q}}$ follows from [14, Prop. 2]. \square

§4. Strict coherence, infinite-valued events and faithful states. Generalizing de Finetti’s theorem, a book on Łukasiewicz events is *coherent* iff it can be extended to a state in the sense of the following definition.

DEFINITION 4.1 ([25]). *A state of an MV-algebra \mathbf{A} is a map $s : A \rightarrow [0, 1]$ satisfying the following conditions:*

- (s1) *Normalization:* $s(1) = 1$,
- (s2) *Additivity:* $s(a \oplus b) = s(a) + s(b)$, for all $a, b \in A$ such that $a \odot b = 0$.

A state s is said to be faithful if $s(a) \neq 0$ for all $a \neq 0$.

Kroupa and Panti independently proved that for every state s of an MV-algebra \mathbf{A} there exists a unique regular Borel, and hence σ -additive, probability measure μ_s on the space of maximal ideals with the hull-kernel topology of \mathbf{A} such that s is the integral with respect to μ_s (see [17], [29], and [27, Sec. 10]). In particular, for n -generated free MV-algebras, the Kroupa–Panti theorem shows that for every state s of \mathbf{F}_n there exists a unique regular Borel probability measure μ_s on $[0, 1]^n$ such that for each $f \in \mathbf{F}_n$,

$$s(f) = \int_{[0,1]^n} f \, d\mu_s. \tag{4}$$

The correspondence between states of \mathbf{F}_n and regular Borel probability measures on $[0, 1]^n$ is one-one.

The next result, which to the best of our knowledge is new, represents faithful states of \mathbf{F}_n in a similar manner. Following [33], we say that a regular Borel measure μ of $[0, 1]^n$ is *strictly positive* if for every nonempty open $O \subseteq [0, 1]^n$, $\mu(O) > 0$.

PROPOSITION 4.2. *For any state s of \mathbf{F}_n the following conditions are equivalent:*

- (1) s is a faithful state;
- (2) There exists a unique strictly positive, regular probability Borel measure μ_s such that for every $f \in \mathbf{F}_n$,

$$s(f) = \int_{[0,1]^n} f \, d\mu_s.$$

The correspondence between faithful states of \mathbf{F}_n and strictly positive, regular probability Borel measures of $[0, 1]^n$ is one-one.

Proof. For every state s of \mathbf{F}_n , let μ_s be the unique regular probability Borel measure of $[0, 1]^n$ such that for every $f \in \mathbf{F}_n$, $s(f) = \int_{[0,1]^n} f \, d\mu_s$ as in the Kroupa–Panti theorem.

(1) \Rightarrow (2). Assume that O is a nonempty open set in the product topology of $[0, 1]^n$, and $\mu_s(O) = 0$. Let K_O be any nonempty compact subset of O and assume, without loss of generality, that K_O is a rational polyhedron. By [27, Corollary 2.10], there exists $f \in \mathbf{F}_n$ such that K_O is the onese of f . Since K_O is contained in O , by [18, Lemma 2.2(i)] there exists $n \in \mathbb{N}$ such that, for all $x \in [0, 1]^n$, the n -fold \odot -product $f^n = f \odot \dots \odot f$, satisfies $f^n(x) = 1$ if $x \in K_O$ and $f^n(x) = 0$ for all $x \notin O$. Therefore, $f^n \neq 0$ and $s(f^n) = \int_{[0,1]^n} f^n \, d\mu_s = 0$ whence s is not faithful.

(2) \Rightarrow (1). Assume that s is not faithful and in particular, let $f \in \mathbf{F}_n$ be such that $f \neq 0$ and $s(f) = 0$. Since f is continuous and not constantly 0, its support $\text{supp}(f) = \{x \in [0, 1]^n \mid f(x) > 0\}$ is nonempty and open. Thus,

$$0 = s(f) = \int_{[0,1]^n} f \, d\mu_s = \int_{\text{supp}(f)} f \, d\mu_s,$$

whence $\mu_s(\text{supp}(f)) = 0$. □

REMARK 4.3. *From Proposition 4.2 it follows that if $\mathbf{A} = \mathbf{F}_n/I_C$ is a finite-dimensional MV-algebra, there is a one-one correspondence between faithful states of \mathbf{A} , strictly positive distributions on the points c_1, \dots, c_t of C , and points in the relative interior of the simplex $\Sigma_\Delta = \{(\lambda_1, \dots, \lambda_t) \in \mathbb{R}^t \mid \sum_{i=1}^t \lambda_i = 1\}$ (see [11, Remark 6.3]). Every finitely generated free boolean algebra \mathbf{A} is, in particular, a finite-dimensional MV-algebra and every faithful state s of \mathbf{A} is a Carnap probability (recall §1). Therefore, Proposition 4.2 specializes on boolean events as follows: for every $n = 1, 2, \dots$, a finitely additive probability P of the n -generated free boolean algebra \mathbf{A} is Carnap iff there exists a unique strictly positive distribution μ_P on $[0, 1]^n$ such that for every $f \in \mathbf{A}$, $P(f) = \sum_{x \in [0,1]^n} \mu_P(x) \cdot f(x)$.*

In [11], the authors characterized strictly coherent books on finite subsets of a finite-dimensional MV-algebra \mathbf{A} (recall §2) as those books that can be extended to a faithful state of \mathbf{A} . In this section, we will provide two measure-theoretical characterizations of

strict coherence for books on \mathbf{F}_n . The first one (Theorem 4.6) involves states satisfying a *local* version of faithfulness which depends both on Φ and on the fixed regular complex linearizing its functions; the second one (Theorem 4.8) is given in terms of faithful states of \mathbf{F}_n .

DEFINITION 4.4. *Let Φ be a finite subset of \mathbf{F}_n and let Δ be a regular complex which linearizes Φ . Then s is said to be Δ -faithful provided that $s(\hat{h}_v) > 0$ for all $v \in V(\Delta)$.*

The next lemma collects some useful properties of states and Δ -faithful states.

LEMMA 4.5. *For each regular complex Δ of $[0, 1]^n$ with vertices $V(\Delta) = \{v_1, \dots, v_t\}$, the following conditions hold:*

- (1) *For each map $\lambda : V(\Delta) \rightarrow [0, 1]$ such that $\sum_{v \in V(\Delta)} \lambda(v) = 1$, the map $s_\lambda : \mathbf{F}_n \rightarrow [0, 1]$ defined as*

$$s_\lambda(f) = \sum_{v \in V(\Delta)} f(v) \cdot \lambda(v). \tag{5}$$

is a state of \mathbf{F}_n .

- (2) *The set of Δ -faithful states of \mathbf{F}_n is in one-one correspondence with the set of faithful states of $\mathbf{F}_n/I_{V(\Delta)}$ and hence is in one-one correspondence with the relative interior of the simplex*

$$\Sigma_\Delta = \left\{ \langle \lambda_1, \dots, \lambda_t \rangle \in \mathbb{R}^t \mid \sum_{i=1}^t \lambda_i = 1 \right\}.$$

Proof. (1). Every state of \mathbf{F}_n belongs to the closure of the convex hull of the homomorphisms of \mathbf{F}_n to $[0, 1]_{MV}$ (see [25, Theorem 2.5] and [12, Theorem 4.1.1]). Thus the claim follows immediately from Proposition 2.1

(2). The claim easily follows from Proposition 4.2, Remark 4.3 and the definition of Δ -faithfulness. □

The next theorem yields a characterization of strictly coherent books in terms of Δ -faithful states.

THEOREM 4.6. *Let Φ be a finite subset of \mathbf{F}_n and let β be a book on Φ . Then the following conditions are equivalent:*

- (1) *β is strictly coherent;*
- (2) *There exists a regular complex Δ which linearizes Φ and a Δ -faithful state s which extends β ;*
- (3) *For every regular complex Δ which linearizes Φ , there exists a Δ -faithful state s of \mathbf{F}_n which extends β .*

Proof. (1) \Rightarrow (3). Let β be strictly coherent. From Corollary 3.5(3), for every regular complex Δ linearizing Φ , there exists a map $\lambda : V(\Delta) \rightarrow [0, 1]$ such that $\sum_{v \in V(\Delta)} \lambda(v) = 1$, $\lambda(v) > 0$ for all $v \in V(\Delta)$ and for every $f_j \in \Phi$,

$$\beta(f_j) = \sum_{v \in V(\Delta)} f_j(v) \cdot \lambda(v). \tag{6}$$

Let s_λ be the state of \mathbf{F}_n defined in (5).

First of all notice that, directly from (5) and (6), s_λ extends β . Furthermore, for every vertex $v \in V(\Delta)$, the normalized Schauder hat \hat{h}_v takes value 1 on v and 0 on any $v' \neq v$.

Thus, $s_\lambda(\hat{h}_v) = \sum_{v' \in V(\Delta)} \hat{h}_v(v') \cdot \lambda(v') = \lambda(v)$ and hence $s_\lambda(\hat{h}_v) > 0$. Therefore s_λ is a Δ -faithful state of \mathbf{F}_n which extends β .

(3) \Rightarrow (1). Now assume that (3) holds and define $\lambda : V(\Delta) \rightarrow [0, 1]$ by

$$\lambda(v) = s(\hat{h}_v).$$

From Lemma 2.4,

$$\sum_{v \in V(\Delta)} \lambda(v) = \sum_{v \in V(\Delta)} s(\hat{h}_v) = s\left(\bigoplus_{v \in V(\Delta)} \hat{h}_v\right) = s(1) = 1.$$

Since s is Δ -faithful, $\lambda(v) > 0$ for each $v \in V(\Delta)$. Further, for all $f_i \in \Phi$, $\beta(f_i) = s(f_i) = \sum_{v \in V(\Delta)} \lambda_v \cdot f_i(v)$. Thus, β is strictly coherent by Corollary 3.5 ((2) \Rightarrow (1)).

Finally, (3) \Rightarrow (2) is trivial and (2) \Rightarrow (3) follows from (1) \Leftrightarrow (3) above, Corollary 3.2 and Theorem 3.4 □

Lemma 4.5(2) shows that for each regular complex Δ which linearizes Φ , Δ -faithful states of \mathbf{F}_n are in one-one correspondence with faithful states of $\mathbf{F}_n/I_{V(\Delta)}$. In particular, for every Δ -faithful state s of \mathbf{F}_n , let s_Δ be the unique faithful state of $\mathbf{F}_n/I_{V(\Delta)}$ such that: for every $f \in \mathbf{F}_n$, let f_Δ to be the restriction of f to $V(\Delta)$ and

$$s_\Delta(f_\Delta) = \sum_{v \in V(\Delta)} s(\hat{h}_v) \cdot f_\Delta(v).$$

Thus, if $f \in \Phi$, $s_\Delta(f_\Delta) = s(f)$. We then have:

COROLLARY 4.7. *Let Φ be a finite subset of \mathbf{F}_n and let β be a book on Φ . Then the following conditions are equivalent:*

- (1) β is strictly coherent;
- (2) There exists a regular complex Δ which linearizes Φ and a faithful state s_Δ of $\mathbf{F}_n/I_{V(\Delta)}$ such that, for all $f \in \Phi$, $\beta(f) = s_\Delta(f_\Delta)$;
- (3) For every regular complex Δ which linearizes Φ , there exists a faithful state s_Δ of $\mathbf{F}_n/I_{V(\Delta)}$ such that, for all $f \in \Phi$, $\beta(f) = s_\Delta(f_\Delta)$.

The following construction is used in the next result which characterizes strictly coherent books in terms of faithful states: let β be a strictly coherent book on a finite subset Φ of \mathbf{F}_n , fix an enumeration g_1, g_2, \dots of $\mathbf{F}_n \setminus \{\Phi, 0, 1\}$ and consider the following inductive construction:

- (S₁) Put $\Phi_1 = \Phi \cup \{g_1\}$. Each regular complex Δ_1 linearizing Φ_1 also linearizes Φ . Since β is strictly coherent, Theorem 4.6 yields a Δ_1 -faithful state s_1 which extends β . It follows that the extended book $\beta_1 = \beta \cup \{g_1 \mapsto s_1(g_1)\}$ is strictly coherent because s_1 extends it. Further, $0 < s_1(g_1) < 1$.
- (S₂) Consider $\Phi_2 = \Phi_1 \cup \{g_2\}$ and fix Δ_2 that linearizes Φ_2 and a Δ_2 -faithful state s_2 which extends β_1 . Again, $0 < s_2(g_2) < 1$ and $\beta_2 = \beta_1 \cup \{g_2 \mapsto s_2(g_2)\}$ is strictly coherent by Theorem 4.6. Further, $s_2(e) = s_1(e)$ for all $e \in \Phi_1$.
- (S_{*i*+1}) At step $i + 1$, arguing by induction, construct a regular complex Δ_{i+1} which linearizes $\Phi_i \cup \{g_{i+1}\} = \Phi \cup \{g_1, \dots, g_{i+1}\}$, a state s_{i+1} of \mathbf{F}_n which is Δ_{i+1} -faithful and a strictly coherent book $\beta_{i+1} = \beta_i \cup \{g_{i+1} \mapsto s_{i+1}(g_{i+1})\}$.

For each n , the state s_n agrees with s_{n-1} over Φ_{n-1} . Thus, for all n_0 and for all $n > n_0$, $s_n(g_{n_0})$ always attains the same value. In particular, for all $n, m \in \mathbb{N}$, $s_n(f) = s_m(f)$ for all $f \in \Phi$.

THEOREM 4.8. *Let Φ be a finite subset of \mathbf{F}_n and let β be a book on Φ . Then the following conditions are equivalent:*

- (1) β is strictly coherent;
- (2) β extends to a faithful state of \mathbf{F}_n .

Proof. The direction (2) \Rightarrow (1) is trivial. Thus, let β be strictly coherent and fix an enumeration g_1, g_2, \dots of $\mathbf{F}_n \setminus \{\Phi, 0, 1\}$. The construction above determines subsets $\Phi = \Phi_0 \subseteq \Phi_1 \subseteq \Phi_2 \subseteq \dots$ of \mathbf{F}_n and a sequence $\{s_i\}_{i \geq 1}$ of states of \mathbf{F}_n such that:

- (i) Each s_i is a Δ_i -faithful state of \mathbf{F}_n ;
- (ii) For all $n > m$, $s_m(e) = s_n(e)$ for all $e \in \Phi_m$.

By construction of the Φ_i 's, for every $f \in \mathbf{F}_n$ there exists an $m \geq 0$ such that $f \in \Phi_m$ and hence, by (ii), $s_m(f) = s_n(f)$ for all $n > m$. Therefore, $\{s_i(f)\}_{i \geq 0}$ is a Cauchy sequence. This gives that $\{s_i\}_{i \geq 0}$ is pointwise convergent. Define $s : \mathbf{F}_n \rightarrow [0, 1]$ as follows: for each $f \in \mathbf{F}_n$,

$$s(f) = \lim_{i \rightarrow \infty} s_i(f).$$

Let us prove that s is a state. Clearly $s(1) = 1$. If $a \oplus b = 0$ then, for all $i \geq 0$, $s_i(a \oplus b) = s_i(a) + s_i(b)$ and by the continuity of $+$, $s(a \oplus b) = \lim_{i \rightarrow \infty} s_i(a \oplus b) = \lim_{i \rightarrow \infty} s_i(a) + s_i(b) = \lim_{i \rightarrow \infty} s_i(a) + \lim_{i \rightarrow \infty} s_i(b) = s(a) + s(b)$. By construction, s extends β since so does each s_n . There remains to be proved that s is faithful. We will provide two proofs of this fact.

(Proof 1). By [11, Theorem 5.2] s is faithful iff for each finite subset Ψ of \mathbf{F}_n the restriction of s to Ψ is strictly coherent. Recalling the above construction, let i_0 be the minimum index such that $\Psi \subseteq \Phi_{i_0}$. Thus, the restriction of s to Ψ coincides with the restriction of s_{i_0} to Ψ and the restriction of s_{i_0} to Φ_{i_0} is strictly coherent. The claim immediately follows because strict coherence is preserved for books contained in a strictly coherent one.

(Proof 2). Let $1 > f > 0$. If $f \in \Phi$ there is nothing to prove. Conversely, assume that $f = g_i$ for some i . Therefore, for all $j \geq i$, $s_j(f) = \alpha > 0$. Thus, $s(f) = \lim_{i \rightarrow \infty} s_i(f) = \alpha > 0$ and the claim is settled. □

§5. Coherence, strict coherence and provability in Łukasiewicz logic. Propositional Łukasiewicz logic (\mathbb{L} in symbols) is the logical calculus having MV-algebras as its equivalent algebraic semantics. Formulas of Łukasiewicz logic will be denoted by lower case Greek letter and $\mathbb{L}(m)$ will stand for the set of formulas in a language with m propositional variables. A complete axiomatization of \mathbb{L} can be found in [3, Definition 4.3.1]. A formula φ is said to be a *theorem*, in symbols $\vdash \varphi$, if φ can be deduced from the axioms of \mathbb{L} and by its unique rule of modus ponens. A *theory* Θ is a deductively closed set of formulas. A theory Θ of $\mathbb{L}(m)$ is said to be *finitely axiomatizable* if for some (necessarily satisfiable) formula $\theta \in \mathbb{L}(m)$, Θ is the smallest theory of $\mathbb{L}(m)$ which contains θ .

By Proposition 2.1, valuations of the Łukasiewicz language $\mathbb{L}(m)$ are in one-one correspondence with homomorphisms of \mathbf{F}_m to $[0, 1]_{MV}$ as well as with points of the m -cube $[0, 1]^m$. Thus, a formula $\varphi \in \mathbb{L}(m)$ is a *tautology* if $h(\varphi) = 1$ for all homomorphisms

$h : \mathbf{F}_m \rightarrow [0, 1]_{MV}$ iff the onset of f_φ coincides with $[0, 1]^m$, where f_φ is the unique McNaughton function determined by φ [24].

For every $\mathcal{X} \subseteq [0, 1]^m$ and theory Θ we write

$$\text{Th}(\mathcal{X}) = \{\psi \in \mathbf{L}(m) \mid (\forall x \in \mathcal{X}) f_\psi(x) = 1\}$$

and

$$\text{Mod}(\Theta) = \{x \in [0, 1]^m \mid (\forall \psi \in \Theta) f_\psi(x) = 1\}.$$

Given two (not necessarily finitely axiomatizable) theories Θ_1 and Θ_2 , we write $\Theta_1 \models \Theta_2$, if $\text{Mod}(\Theta_1) \subseteq \text{Mod}(\Theta_2)$.

Following [27, Definition 3.9], two rational polyhedra \mathcal{P} and \mathcal{Q} of $[0, 1]^m$ are said to be \mathbb{Z} -homeomorphic (in symbols, $\mathcal{P} \cong_{\mathbb{Z}} \mathcal{Q}$) if there exists a homeomorphism $\eta : \mathcal{P} \rightarrow \mathcal{Q}$ such that both η and η^{-1} , as maps from $\mathbb{R}^m \rightarrow \mathbb{R}^m$ are \mathbb{Z} -maps, i.e., η and η^{-1} are piecewise linear with integer coefficients.

LEMMA 5.1 ([27, Theorem 3.20]). *For every $m = 1, 2, \dots$, the pair (Th, Mod) establishes a Galois connection between rational polyhedra of $[0, 1]^m$ and finitely axiomatizable theories of $\mathbf{L}(m)$. In particular:*

- (1) *For every finitely axiomatizable theory Θ of $\mathbf{L}(m)$, there exists a unique rational polyhedron \mathcal{P}_Θ of $[0, 1]^m$ such that $\text{Mod}(\text{Th}(\Theta)) \cong_{\mathbb{Z}} \mathcal{P}_\Theta$.*
- (2) *For each rational polyhedron \mathcal{P} of $[0, 1]^m$ there exists a unique finitely axiomatizable theory $\Theta_{\mathcal{P}}$ such that $\text{Mod}(\text{Th}(\Theta_{\mathcal{P}})) \cong_{\mathbb{Z}} \mathcal{P}$.*
- (3) *For \mathcal{P}_1 and \mathcal{P}_2 rational polyhedra, $\mathcal{P}_1 \subseteq \mathcal{P}_2$ iff $\text{Mod}(\Theta_{\mathcal{P}_1}) \subseteq \text{Mod}(\Theta_{\mathcal{P}_2})$ iff $\Theta_{\mathcal{P}_1} \models \Theta_{\mathcal{P}_2}$.*

In the rest of this section we will adopt the notation used in Lemma 5.1 above with the following exception: if $x \in ([0, 1] \cap \mathbb{Q})^m$, we denote by Θ_x the finitely axiomatizable theory $\Theta_{\{x\}}$.

Let Φ be a subset of \mathbf{F}_n of finite cardinality k and let β be a rational-valued book on Φ . As noted at the beginning of §3, $\{\beta\}$, \mathcal{C}_Φ and $\text{rb } \mathcal{C}_\Phi$ are rational polyhedra of $[0, 1]^k$. By Lemma 5.1, Θ_β , $\Theta_{\mathcal{C}_\Phi}$ and $\Theta_{(\text{rb } \mathcal{C}_\Phi)}$ are finitely axiomatizable.

The following lemma provides a first characterization of coherence and strict coherence in terms of deducibility.

LEMMA 5.2. *Let Φ be a finite subset of \mathbf{F}_n and let β be a rational-valued book on Φ . Thus the following conditions hold:*

- (1) *β is coherent iff $\Theta_\beta \models \Theta_{\mathcal{C}_\Phi}$.*
- (2) *β is strictly coherent iff $\Theta_\beta \models \Theta_{\mathcal{C}_\Phi}$ and $\Theta_\beta \not\models \Theta_{(\text{rb } \mathcal{C}_\Phi)}$.*

Proof. (1) follows from [26, Theorem 2.1], Lemma 5.1 and the definition of \models . Indeed, β is coherent iff $\beta \in \mathcal{C}_\Phi$ iff $\{\beta\} \subseteq \mathcal{C}_\Phi$.

As for (2), Theorem 3.4 shows that β is strictly coherent iff $\beta \in \text{ri}\mathcal{C}_\Phi = \mathcal{C}_\Phi \setminus \text{rb } \mathcal{C}_\Phi$ iff $\beta \in \mathcal{C}_\Phi$ and $\beta \notin \text{rb } \mathcal{C}_\Phi$ iff $\Theta_\beta \models \Theta_{\mathcal{C}_\Phi}$ (by (1) above) and $\{\beta\} \not\subseteq \text{rb } \mathcal{C}_\Phi$. By Lemma 5.1 (3) this condition is equivalent to $\Theta_\beta \not\models \Theta_{(\text{rb } \mathcal{C}_\Phi)}$. □

To characterize coherence and strict coherence in terms of provability in Łukasiewicz logic, we prepare.

PROPOSITION 5.3. *There exists an effective procedure Π to compute, for each rational polytope \mathcal{P} of $[0, 1]^k$, a formula $\Pi_{\mathcal{P}}$ which axiomatizes $\Theta_{\mathcal{P}}$.*

Proof. First, compute a regular complex Δ supporting \mathcal{P} (see [7, Chap. 6.2.2. and Theorem 6.5]). Notice that, $\text{ext } \mathcal{P} \subseteq V(\Delta)$ and let $\hat{h}_1, \dots, \hat{h}_q$ be the normalized Schauder hats at the vertices v_1, \dots, v_q of Δ . For $j = 1, \dots, q$ let Π_j be the Łukasiewicz formulas computed from \hat{h}_j (see [24]). Let further,

$$\Pi_{\mathcal{P}} = \bigoplus_{j=1}^q \Pi_j.$$

Since each \hat{h}_j is a member of $\mathbb{L}(k)$, $\Pi_{\mathcal{P}}$ belongs to $\mathbb{L}(k)$. There remains to be proved that $\Pi_{\mathcal{P}}$ axiomatizes $\Theta_{\mathcal{P}}$. To this purpose, let us prove that

$$x \in \mathcal{P} \text{ iff } h_x(\Pi_{\mathcal{P}}) = 1.$$

As a matter of fact, by Lemma 2.4(3) the onese of the McNaughton function $\Pi_{\mathcal{P}}$ is \mathcal{P} . Thus the claim is settled. □

COROLLARY 5.4. *There exists an effective procedure Π which computes, for each $\Phi = \{f_1, \dots, f_k\} \subseteq \mathbb{F}_n$ and for each $\beta \in [0, 1]^k$, formulas Π_{Φ} , $\Pi_{(\text{rb } \Phi)}$ and Π_{β} of $\mathbb{L}(k)$ which respectively axiomatize $\Theta_{\mathcal{C}_{\Phi}}$, $\Theta_{(\text{rb } \mathcal{C}_{\Phi})}$ and Θ_{β} .*

Proof. As the reader will recall from §3, \mathcal{C}_{Φ} is a polytope. Thus, \mathcal{C}_{Φ} , $\{\beta\}$ are rational polytopes of $[0, 1]^k$, whence Π_{Φ} and Π_{β} are computed as in Proposition 5.3.

As for $\Pi_{(\text{rb } \Phi)}$, $\text{rb } \mathcal{C}_{\Phi}$ is not convex. However, it can be realized as the finite union of the faces F_1, \dots, F_l of \mathcal{C}_{Φ} . Each face F_i is a rational polytope, whence Proposition 5.3 yields Łukasiewicz formulas Π_1, \dots, Π_l such that $x \in F_i$ iff $h_x(\Pi_i) = 1$. Thus, let

$$\Pi_{(\text{rb } \Phi)} = \bigvee_{i=1}^l \Pi_i.$$

Finally, $x \in \text{rb } \mathcal{C}_{\Phi}$ iff exists $i = 1, \dots, n$ such that $x \in F_i$ iff $h_x(F_i) = 1$ iff $h_x(\Pi_{(\text{rb } \Phi)}) = 1$. □

In the light of the above corollary, we may write Π_{Φ} , $\Pi_{(\text{rb } \Phi)}$ and Π_{β} without danger of confusion. In the following characterization, for every Łukasiewicz formula ψ , we write ψ^n for $\psi \odot \dots \odot \psi$ (n -times).

THEOREM 5.5. *Let Φ be a finite set of \mathbb{F}_n and let β be a book on Φ . Then the following conditions hold:*

- (1) β is coherent iff there exists a nonzero $n \in \mathbb{N}$ such that $\vdash (\Pi_{\beta})^n \rightarrow \Pi_{\Phi}$.
- (2) β is strictly coherent iff there exists a nonzero $n \in \mathbb{N}$ such that $\vdash (\Pi_{\beta})^n \rightarrow \Pi_{\Phi}$ and for all nonzero $n \in \mathbb{N}$, $\not\vdash (\Pi_{\beta})^n \rightarrow \Pi_{(\text{rb } \Phi)}$.

Proof. Both claims follow from Lemma 5.2, Corollary 5.4, the completeness theorem of Łukasiewicz calculus and Łukasiewicz deduction theorem stating that $\varphi \vdash \psi$ iff there exists a nonzero $n \in \mathbb{N}$ such that $\vdash \varphi^n \rightarrow \psi$ (see [3, Sec. 4]). We will prove (1) since the proof of (2) is essentially the same.

By Lemma 5.2, β is coherent iff $\Theta_{\beta} \models \Theta_{\mathcal{C}_{\Phi}}$ iff (from Lemma 5.4) $\Pi_{\beta} \models \Pi_{\Phi}$. The completeness theorem of Łukasiewicz calculus shows that $\Pi_{\beta} \models \Pi_{\Phi}$ iff $\Pi_{\beta} \vdash \Pi_{\Phi}$ iff $\vdash (\Pi_{\beta})^n \rightarrow \Pi_{\Phi}$ for some $n > 0$. □

§6. Conclusion. In this article we have presented geometrical, measure-theoretical and logical characterizations for the strict coherence of books on Łukasiewicz infinite-valued events. Our first result shows that, for any finite subset Φ of a finitely generated free MV-algebra \mathbf{A} , the set of all strictly coherent books on Φ coincides with the relative interior of the polytope of all coherent ones; the second characterization is a de Finetti-like theorem: a book on Φ is strictly coherent if and only if it extends to a faithful state of \mathbf{A} . Finally, our last theorem gives a characterization of coherence and strict coherence in terms of the provability relation of propositional Łukasiewicz logic.

We believe that this last result is interesting both from the logical and philosophical perspective as it may shed a light on an *intuitive reading* of propositional Łukasiewicz logic. Specifically, it is of particular interest to put forward a comparison between the role of Łukasiewicz logic prompted by Theorem 5.5 in theories of uncertain reasoning and the semantics proposed in [20]. There, the author, investigating the problem of *artificial precision* in theories of vagueness based on real numbers as degrees of truth, presents Łukasiewicz logic as a suitable formal system to handle vague predicates.³

In our future work we will mainly focus on extending the results of this article to more general algebraic structures. Particularly promising seems to be the class of *finitely presented* MV-algebras (see [21] and [27, Theorem 6.3]). Further, we will address the problem of determining an NP-algorithm to check strict coherence for Łukasiewicz events. The solution of this problem would immediately yield that for each finite set Φ of Łukasiewicz events, $\mathcal{K}_{\Phi}^{\mathbb{Q}}$ is NP-complete (see [11, Sec. 7]).

Acknowledgments. The author is grateful to the two referees for their careful reading and valuable comments. He is also much indebted to Lluís Godó, Hykel Hosni and Daniele Mundici for several conversations on the subject of this article. The author acknowledges partial support by the Spanish Ramón y Cajal research program RYC-2016-19799; the Spanish FEDER/MINECO project TIN2015- 71799-C2-1-P and the SYSMICS project (EU H2020-MSCA-RISE-2015, Project 689176).

BIBLIOGRAPHY

- [1] Carnap, R. (1950). *The Logical Foundations of Probability*. Chicago: University of Chicago Press.
- [2] Chang, C.C. (1958). Algebraic analysis of many-valued logics. *Transactions of the American Mathematical Society*, **88**, 467–490.
- [3] Cignoli, R., D’Ottaviano, I. M. L., & Mundici, D. (2000). *Algebraic Foundations of Many-valued Reasoning*. Trends in Logic, Vol. 8. Dordrecht: Kluwer.
- [4] Cignoli, R. & Marra, V. (2012). Stone duality for real-valued multisets. *Forum Mathematicum*, **24**(6), 1317–1331.
- [5] de Finetti, B. (1931). Sul significato soggettivo della probabilità. *Fundamenta Mathematicae*, **17**, 298–329. Translated into English as “On the subjective meaning of probability.” In Monari, P., and Cocchi, D., editors. *Probabilità e Induzione*. Bologna: Clueb, pp. 291–321, 1993.
- [6] de Finetti, B. (1974). *Theory of Probability*, Vol. 1. New York: Wiley.
- [7] Desiderata, F. P. & Shamos, M. I. (1985). *Computational Geometry – An Introduction*. New York: Springer-Verlag.

³ The author wishes to thank Eduardo Barrio for pointing out this to him.

- [8] Ewald, G. (1996). *Combinatorial Convexity and Algebraic Geometry*. New York: Springer-Verlag.
- [9] Flaminio, T., Godo, L., & Hosni, H. (2014). On the logical structure of de Finetti's notion of event. *Journal of Applied Logic*, **12**(3), 279–301.
- [10] Flaminio, T., Hosni, H., & Lapenta, S. (2018). Convex MV-algebras: Many-valued logics meet decision theory. *Studia Logica*, **106**(5), 913–945.
- [11] Flaminio, T., Hosni, H., & Montagna, F. (2018). Strict coherence on many-valued events. *The Journal of Symbolic Logic*, **83**(1), 55–69.
- [12] Flaminio, T. & Kroupa, T. (2015). States of MV-algebras. In Fermüller, C., Cintula, P., and Noguera, C., editors. *Handbook of Mathematical Fuzzy Logic - volume 3*. Studies in Logic, Mathematical Logic and Foundations, Vol. 58, Chapter XVII. London: College Publications.
- [13] Gaifman, H. (1964). Concerning measures on Boolean algebras. *Pacific Journal of Mathematics*, **14**(1), 61–73.
- [14] Hähnle, R. (1994). Many-valued logic and mixed integer programming. *Annals of Mathematics and Artificial Intelligence*, **12**(3–4), 231–263.
- [15] Kelley, J. L. (1959). Measures on Boolean Algebras. *Pacific Journal of Mathematics*, **9**(4), 1165–1177.
- [16] Kemeny, J. G. (1955). Fair bets and inductive probabilities. *The Journal of Symbolic Logic*, **20**(3), 263–273.
- [17] Kroupa, T. (2006). Every state on semisimple MV-algebra is integral. *Fuzzy Sets and Systems*, **157**(20), 2771–2787.
- [18] Kroupa, T. (2012). States in Łukasiewicz logic corresponds to probabilities of rational polyhedra. *International Journal of Approximate Reasoning*, **53**, 435–446.
- [19] Kühn, J. & Mundici, D. (2007). De Finetti theorem and Borel states in $[0, 1]$ -valued algebraic logic. *International Journal of Approximate Reasoning*, **46**(3), 605–616.
- [20] Marra, V. (2014). The problem of artificial precision in theories of vagueness: A note on the rôle of maximal consistency. *Erkenntnis*, **79**(5), 1015–1026.
- [21] Marra, V. & Spada, L. (2013). Duality, projectivity and unification in Łukasiewicz logic and MV-algebras. *Annals of Pure and Applied Logic*, **164**(3), 192–210.
- [22] McMullen, P. & Shephard, G. C. (1971). *Convex Polytopes and the Upper Bound Conjecture*. London Mathematical Society Lecture Note Series, Vol. 3. London: Cambridge University Press.
- [23] McNaughton, R. (1951). A theorem about infinite-valued sentential logic. *The Journal of Symbolic Logic*, **16**, 1–13.
- [24] Mundici, D. (1994). A constructive proof of McNaughton's theorem in infinite-valued logic. *The Journal of Symbolic Logic*, **58**(2), 596–602.
- [25] Mundici, D. (1995). Averaging the truth-value in Łukasiewicz logic. *Studia Logica*, **55**(1), 113–127.
- [26] Mundici, D. (2006). Bookmaking over infinite-valued events. *International Journal of Approximate Reasoning*, **43**(3), 223–240.
- [27] Mundici, D. (2011). *Advanced Łukasiewicz Calculus and MV-algebras*. Trends in Logic 35, Springer, 2011.
- [28] Mundici, D. (2011). Finite axiomatizability in Łukasiewicz logic. *Annals of Pure and Applied Logic*, **162**, 1035–1047.
- [29] Panti, G. (2009). Invariant measures on free MV-algebras. *Communications in Algebra*, **36**(8), 2849–2861.
- [30] Paris, J. (2001). A note on the Dutch Book method. In De Cooman, G., Fine, T., and Seidenfeld, T., editors. *Proceedings of the Second International Symposium*

on Imprecise Probabilities and their Applications. ISIPTA 2001. Ithaca, NY: Shaker Publishing Company, pp. 301–306.

[31] Paris, J. B. & Vencovska, A. (2015). *Pure Inductive Logic*. Cambridge, UK: Cambridge University Press.

[32] Shimony, A. (1955). Coherence and the axioms of confirmation. *The Journal of Symbolic Logic*, **20**(1), 1–28.

[33] Todorčević, S. (1997). *Topics in Topology*. Lecture Notes in Mathematics. Berlin: Springer.

[34] Weatherson, B. (2003). From classical to intuitionistic probability. *Notre Dame Journal of Formal Logic*, **44**(2), 111–123.

IIIA - CSIC

CAMPUS DE LA UNIVERSIDAD AUTÒNOMA DE BARCELONA S/N

08193 BELLATERRA, SPAIN

E-mail: tommaso@iiia.csic.es