

Vitamin D deficiency and its associated risk factors in children and adolescents in southern Iran

Forough Saki, Mohammad Hossein Dabbaghmanesh*, Gholamhossein Ranjbar Omrani and Marzieh Bakhshayeshkaram

Shiraz Endocrinology and Metabolism Research Center, Nemazee Hospital, Shiraz University of Medical Sciences, PO Box 71345-1744, Shiraz, Islamic Republic of Iran

Submitted 23 August 2014: Final revision received 20 March 2015: Accepted 13 May 2015: First published online 8 June 2015

Abstract

Objectives: To evaluate prevalence of vitamin D deficiency and its associated factors in southern Iranian children.

Design: Cross-sectional study. Anthropometric and pubertal characteristics were assessed by a trained physician. Physical activity and sun exposure were evaluated using standard questionnaires. Body composition measurements were performed using dual-energy X-ray absorptiometry. Serum Ca, P alkaline phosphatase and 25-hydroxyvitamin D (25(OH)D) were assessed in all children. Statistical analysis was done using the statistical software package IBM SPSS Statistics 18.0.

Subjects: Iranian children (n 477) aged 9–18 years.

Setting: Fars Province, Iran, 2011.

Results: Of the children, 81.3% were 25(OH)D deficient. There was no significant difference in 25(OH)D concentration between boys and girls ($P=0.3$). 25(OH)D concentration was associated with BMI ($r=-0.1$, $P=0.02$), pubertal status ($r=-0.08$, $P=0.04$) and sun exposure ($r=0.10$, $P=0.04$). Fat mass index was associated with 25(OH)D concentration ($r=-0.13$, $P=0.03$), but not lean mass index ($P=0.86$). In multiple regression analysis with adjustment for confounding factors, age and puberty were found to be independently associated with 25(OH)D concentration ($P=0.008$ and $P=0.006$); there was a significant correlation between exercise and 25(OH)D concentration after adjustment for either BMI ($P=0.01$) or fat mass index ($P=0.02$).

Conclusions: 25(OH)D deficiency is highly prevalent among children in the south of Iran. It is related to insufficient sun exposure, low physical activity, advancing age and pubertal stage. Measures should be taken to improve the health of southern Iranian children in this critical age group by preventing 25(OH)D deficiency.

Keywords
Vitamin D
Puberty
Exercise
Risk factors
Sun exposure

Vitamin D is a fat-soluble vitamin responsible for enhancing gastrointestinal absorption of Ca, Mg, phosphate and Zn⁽¹⁾. It has an important role in bone mineralization, as it increases the absorption of Ca in the small intestine, promotes osteoblastic activity and maintains serum concentration of Ca and P in the normal range⁽²⁾. Vitamin D may prevent insulin-dependent diabetes mellitus, hypertension and cancer⁽¹⁾. Vitamin D deficiency results in rickets, osteomalacia and low bone density⁽³⁾.

Vitamin D status varies widely between different countries in Europe, Asia and the Middle East⁽⁴⁾. It may be caused by differences in the amount of sunlight exposure, intake of dietary vitamin D and use of supplementary vitamin D⁽⁵⁾. Vitamin D deficiency is very prevalent in children in Australia, the Middle East, Africa, India and

South America^(2,6,7). In Iran, in spite of using supplementary vitamin D in breast-fed infants, vitamin D deficiency is common among children⁽⁸⁾. Prevalence of vitamin D deficiency in children under 12 years of age was reported as 38.3–46.0% in the northern and central part of Iran in 2008^(3,8,9), but no data are available about the prevalence of vitamin D deficiency in Iranian teenagers or in the south of Iran. Serum 25-hydroxyvitamin D (25(OH)D) concentration depends on many personal and environmental factors⁽¹⁰⁾. Understanding these factors helps to predict vitamin D deficiency in children. Most studies have studied only a few of these factors, such as sun exposure^(3,4,7,11,12). There is some confusion in the existing literature regarding the correlation between vitamin D deficiency and pubertal stage or BMI^(11–18) and there are

*Corresponding author: Email dabbaghm@sums.ac.ir

also limited data on the correlation of body composition (fat or lean mass) with vitamin D status⁽¹⁰⁾.

The present study was conducted to bridge the existing gap in knowledge on vitamin D status and Ca metabolism in children and teenagers living in southern Iran, the controversies about the association of vitamin D status with its various determinants and the lack of data about the associations of vitamin D status with body composition over a wide range of ages in children.

Materials and methods

To evaluate vitamin D status in Fars Province, we performed a cross-sectional study on healthy Iranian children aged 9–18 years in Kawar, an urban area located 50 km east of Shiraz, the capital city of Fars Province, in the autumn and winter of 2011. An age-stratified, randomly selected sample was drawn of 7.5% of pupils from all the elementary, secondary and other schools in the area, to provide a representative group of 477 participants for this investigation. Children diagnosed with bone disease, disorders of Ca metabolism, malabsorptive disorders and chronic granulomatous disease, as well as those using certain medications (such as anticonvulsants or medically prescribed vitamin D preparations), were excluded from the study.

The Ethics Committee of Shiraz University of Medical Sciences approved our study. All parents and children who participated in our study provided signed informed consent.

Anthropometric measurements and pubertal assessment

A trained general physician performed the physical examination of children, including weight, height and pubertal stage. Weight was measured with a standard scale (Seca, Germany) while the children were wearing light clothing and without shoes, and rounded to the nearest 0.1 kg. Height was measured with a wall-mounted stadiometer on barefoot children and rounded to the nearest 0.5 cm. BMI was calculated by the standard method (weight/height², kg/m²). Pubertal stage was determined according to Tanner's five-stage classification⁽¹⁹⁾. BMI Z-score was calculated using the growth charts of the Centers for Disease Control and Prevention (www.cdc.gov/growthcharts).

Assessment of physical activity and sun exposure

Physical activity on at least 3 d/week is recommended for children and adolescents by the American College of Sports Medicine⁽²⁰⁾. Children and their parents were asked about the number of days per week they undertook physical activity (walking, recreational activity or sports) and placed in groups with acceptable or unacceptable physical activity (≥ 3 or < 3 d/week). Children and their parents were also asked about their average exposure to

sunshine per day and classified into three groups (those exposed for < 15 min/d, 15–30 min/d and > 30 min/d).

Body composition measurements

Total fat mass (grams), total lean mass (grams) and total body fat percentage were determined by dual-energy X-ray absorptiometry (Hologic Discovery QDR instrument, USA). Fat mass index was calculated as fat mass/height² (kg/m²) and lean mass index as lean mass/height² (kg/m²). Assessments were performed with the children wearing standardized clothing and without shoes. The CV in our laboratory was 0.7% for fat mass and 1.9% for lean mass and fat percentage.

Biochemical variables

A 5 ml venous blood sample was drawn from all participants in the hormonal research laboratory of Shiraz University of Medical Sciences. All samples were centrifuged, separated, and serum and plasma stored at -20°C until analysis. Serum 25(OH)D was assayed by HPLC (Young Lin 9100 system, South Korea; inter- and intra-assay CV 3.3% and 5.1%, respectively) according to current Endocrine Society clinical practice guidelines for the evaluation, treatment and prevention of vitamin D deficiency (vitamin D deficiency was defined as serum 25(OH)D < 20 ng/ml (< 50 nmol/l); serum 25(OH)D values between 21 and 29 ng/ml were defined as vitamin D insufficiency⁽⁵⁾). Serum Ca, P and alkaline phosphatase were measured by colorimetric assays using an auto-analyser (Biosystems SA, Spain). Normal range for serum corrected Ca was 8.5–10.5 mg/dl, and for P was 3.7–5.4 mg/dl for children < 16 years old and 2.5–4.5 mg/dl for those aged ≥ 16 years.

Statistical analysis

Data are presented as means and standard deviations. Normality of data distribution was evaluated with the Kolmogorov–Smirnov test. Student's *t* test was used for the comparison of normally distributed data and the Mann–Whitney test for comparison of non-normally distributed data. Correlations between normally distributed parameters were determined using Pearson's test and Spearman's ranking test for non-normally distributed data. Comparison of qualitative data was carried out using the χ^2 test. Variables with significant correlations with serum concentration of 25(OH)D in univariate analysis were identified. Multivariate binary logistic regression analysis was then performed using these variables to assess their independent predictive effect on serum 25(OH)D concentration; collinearity was assessed by variance inflation factor, with a variance inflation factor value < 5 considered as non-collinearity. The linear regression analyses used data that were normally distributed (assessed by Q–Q plots), or data normalized before use, with homogeneity of variance and reported independence of predictive effects. $P < 0.05$ was considered significant. Statistical

analysis was carried out using the statistical software package IBM SPSS Statistics 18.0.

Results

Four hundred and seventy-seven children aged 9–18 years were included in the present study; 49% of them were female (236 girls and 241 boys). General characteristics and a summary of serum biochemical variables are shown in Table 1. Boys were heavier and taller, and had more sun exposure than girls ($P=0.009$, $P<0.001$ and $P<0.001$, respectively).

Vitamin D status and body composition

Mean serum 25(OH)D concentration of the children in autumn and winter was 15.2 (SD 5.6) ng/ml. Four per cent of the children had normal vitamin D status (serum 25(OH)D >30 ng/ml), 15% had 25(OH)D insufficiency (serum 25(OH)D = 20–30 ng/ml), 82% of the children were vitamin D deficient (serum 25(OH)D <20 ng/ml) and 13% had severe vitamin D deficiency (serum 25(OH)D <10 ng/ml; see Table 2). Mean total body fat percentage was 22.6 (SD 8.1), mean fat mass index was 4.2 (SD 2.9) kg/m² and mean lean mass index was 14.4 (SD 10.6) kg/m² in the study population (see Table 3).

Factors associated with vitamin D status

Vitamin D status was associated with BMI ($r=-0.10$, $P=0.02$), pubertal status ($r=-0.08$, $P=0.04$) and sun exposure ($r=0.10$, $P=0.04$). Fat mass index was associated with 25(OH)D concentration ($r=-0.13$, $P=0.03$), but lean mass index showed no association with 25(OH)D concentration ($P=0.86$).

There was no significant difference in 25(OH)D concentration between boys and girls ($P=0.3$). Vitamin D status was not associated with age ($r=-0.07$, $P=0.15$) or physical activity ($r=0.10$, $P=0.08$). On multiple regression analysis, there were independent predictive effects of age, sun exposure, physical activity, pubertal state and fat mass index on 25(OH)D concentration (Table 4). All variance inflation factor values were ≤ 5 (range: 1.05–3.19). Age remained associated with 25(OH)D concentration after adjustment for sun exposure, physical activity, pubertal state and fat mass index ($P=0.008$). The association between puberty and 25(OH)D concentration persisted ($P=0.006$) after adjustment for age, physical activity, sun exposure and fat mass index, but the association between BMI and fat mass index and 25(OH)D concentration was abolished by adjustment for age, sun exposure, physical activity and puberty ($P=0.06$ and $P=0.14$, respectively). The association between exercise and 25(OH)D concentration remained significant after adjustment for BMI ($P=0.01$) and fat mass index ($P=0.02$).

Table 1 General characteristics of the studied population: Iranian children aged 9–18 years, Fars Province, autumn and winter of 2011

Variable	Total		Boys		Girls		P value*
	Mean	SD	Mean	SD	Mean	SD	
Weight (kg)	43.0	13.4	45.0	14.5	42.0	12.0	0.009
Height (cm)	154.0	14.3	157.4	15.9	150.8	11.7	<0.001
BMI (kg/m ²)	17.7	3.2	17.5	3.1	17.9	3.3	0.29
Age (years)	13.8	2.7	13.8	2.6	13.8	2.8	0.93
BMI Z-score	-0.7	1.2	-0.7	1.3	-0.7	1.2	0.69
Ca (mg/dl)	9.9	0.5	10.0	0.6	9.8	0.4	0.009
P (mg/dl)	4.1	0.7	4.1	0.9	4.0	0.6	0.31
ALP (IU/l)	370	185	442	181	325	173	<0.001
25(OH)D (ng/ml)	15.2	5.6	15.7	5.7	14.7	5.3	0.07

ALP, alkaline phosphatase; 25(OH)D, 25-hydroxyvitamin D.

*P value refers to the comparison of each variable between sexes.

Table 2 Vitamin D status (based on serum 25(OH)D) in the studied population*: Iranian children aged 9–18 years, Fars Province, autumn and winter of 2011

Vitamin D status	Total		Boys		Girls	
	n	%	n	%	n	%
Normal (>30 ng/ml)	19	4	12	5	7	3
Vitamin D insufficiency (20–30 ng/ml)	70	15	40	17	30	13
Vitamin D deficiency (10–20 ng/ml)	325	68	164	68	161	68
Severe vitamin D deficiency (<10 ng/ml)	63	13	25	10	38	16
Total	477	100	241	100	236	100

25(OH)D, 25-hydroxyvitamin D.

*Vitamin D status was not dependent on sex ($P=0.145$).

Discussion

Vitamin D status

The present study showed a remarkably high prevalence of vitamin D deficiency in children aged 9–18 years in southern Iran (83% in boys and 86% in girls). 25(OH)D concentration was independently associated with age, exercise and sun exposure, and with pubertal status after adjustment for age, physical activity, sun exposure and fat mass index.

To date, three cross-sectional studies have been conducted to evaluate the prevalence of vitamin D deficiency in the northern (Tehran) and central (Isfahan) parts of Iran^(3,8,9). One showed an 86% prevalence of vitamin D deficiency in Tehran in children aged 9–12 years⁽⁸⁾; another study in children aged 7–12 years in Tehran showed that 53.6% of girls and 11.3% of boys had serum 25(OH)D <20 ng/ml⁽³⁾; and in children aged 6–7 years in Isfahan, the prevalence of serum 25(OH)D <20 ng/ml was 5% in girls and 10% in boys⁽⁹⁾.

The present study appears to be the first to have assessed vitamin D status in children aged 9–18 years in Fars

Table 3 Body composition of the studied population: Iranian children aged 9–18 years, Fars Province, autumn and winter of 2011

Variable	Total		Boys		Girls		P value*
	Mean	SD	Mean	SD	Mean	SD	
Total body fat percentage	22.0	8.9	17.4	6.7	28.4	5.9	<0.001
Fat mass index (kg/m ²)	4.2	2.9	3.4	3.2	5.3	2.0	<0.001
Lean mass index (kg/m ²)	14.4	10.6	15.5	13.8	12.9	1.6	0.04
Total lean mass (g)	32 219	9858	35 419	11 292	28 938	6719	<0.001
Total fat mass (g)	10 085	5545	8113	4758	12 098	5577	<0.001

*P value refers to the comparison of each variable between sexes.

Table 4 Factors associated with vitamin D status in the studied population: Iranian children aged 9–18 years, Fars Province, autumn and winter of 2011

Factor	P value of univariate analysis	P and adjusted r^2 value of multiple regression analysis*
BMI	0.02	$P=0.06$, $r^2=0.35$
Tanner stage	0.04	$P=0.006$, $r^2=0.35$
Sun exposure	0.04	$P=0.03$, $r^2=0.35$
Fat mass index	0.03	$P=0.14$, $r^2=0.32$
Lean mass index	0.861	–
Sex	0.3	–
Age	0.16	$P=0.008$, $r^2=0.35$
Physical activity	0.082	$P=0.001$, $r^2=0.35$

*All factors with $P<0.2$ in univariate analysis were included in multiple regression analysis (mode: Enter).

Province, in the south of Iran, and shows a high prevalence of vitamin D deficiency in southern Iran, at 84%. Differences in age of participants, in season of sampling for serum 25(OH)D and in methods used to measure serum 25(OH)D concentration (RIA *v.* HPLC) may explain the difference in findings between the present and previous studies in Iran. However, most of the reported data has indicated a high prevalence of vitamin D deficiency in Iran, which needs to be recognized and corrected, probably through suitable health ministry programmes.

Factors associated with serum 25-hydroxyvitamin D concentrations

Children's age

Our study showed that children's age was inversely related to serum 25(OH)D concentration after adjustment for physical activity, pubertal status, sun exposure and fat mass index, an association not previously recognized during childhood and adolescence. Previous reports studied children within a narrow age range, such as adolescents or toddlers, limiting the ability of such studies to reveal any negative effect of age^(10,21–26). Some reports in Korea and Europe have also shown that younger adolescents had higher serum 25(OH)D concentrations than older ones^(5,21–26), while Willis *et al.* showed that age was negatively associated with 25(OH)D concentration in pre-pubertal females, although this was abolished by

controlling for fat-free soft tissue mass⁽¹⁰⁾. The only study on a wider range of younger ages was that of Weng *et al.* in the USA⁽²⁷⁾, who found an inverse relationship between 25(OH)D concentration and age in American children and adolescents after adjustment for confounding factors such as puberty and vitamin D intake. However, they did not evaluate sun exposure or outdoor activity although they did suggest these should be evaluated in future studies⁽²⁷⁾, a suggestion supported by the fact that younger children tend to have more physical activity, spend more time outdoors (school and playground) and have a healthier diet (enriched by dairy foods under parental control) than older children^(22,23). Indeed, we showed that the inverse relationship between age and vitamin D status persists after adjusting for sun exposure and physical activity, while further work is needed to explain the reasons for decreases in serum 25(OH)D in older children, since this appears to be independent of the effects of physical activity, pubertal status, sun exposure or obesity.

Sun exposure

Shiraz is located at a latitude of 29.6°N, longitude of 52.5°E and 1506 m elevation above sea level, with relatively good sun exposure, and our study revealed that sun exposure was positively associated with serum 25(OH)D concentration ($P=0.04$). Exposure of skin to sunlight is a major source of vitamin D for man⁽⁵⁾; although it is reduced by a variety of factors including increased skin pigmentation, topical usage of sunscreen and changes in latitude, not all these could be adjusted for in the present study^(2,5). Differences in housing may be another factor influencing serum 25(OH)D concentration. Some children may have a yard or garden attached to the house where they may privately remove some of their outer clothes and get more sun⁽²⁴⁾.

BMI and body composition

Our study showed that BMI and fat mass index had an inverse association with serum 25(OH)D concentration, but not lean mass index. Many studies have shown a negative correlation between BMI and vitamin D status^(24–29); however, there is little information on the association between fat mass index and 25(OH)D concentration, which a previous study showed to be more

consistent in its complications than BMI⁽³⁰⁾. Fat mass index has also been shown to be associated with vitamin D status in 10-year-old urban South African children⁽³¹⁾, but in a narrow age range limited to pre-pubertal children. As originally suggested, vitamin D may well be sequestered in fat stores, reducing its bioavailability⁽³²⁾, as supported by the findings of large-scale Mendelian randomization studies⁽³³⁾. A further factor causing reduced 25(OH)D concentrations with increased fat mass could be the fact that increased leptin released from excess body fat can inhibit renal activation of vitamin D⁽³⁴⁾. Some studies also suggest reverse confounding due to reduced outdoor activity in obesity and to lower dietary vitamin D intakes in obese children who are dieting to lose weight, which also reduces 25(OH)D values^(35,36); and indeed our study does show that the association of BMI or fat mass index with 25(OH)D concentration disappears after adjustment for factors such as age, puberty, physical activity and sun exposure. Consistent with our results, a US study using multivariable models showed that neither lean mass nor fat mass index was associated with vitamin D status⁽²⁷⁾, a finding that may be due to the use of appropriate adjustments of the analyses for the confounding effects of age and puberty since prior studies reporting associations between fat mass and hypovitaminosis D may have been confounded by the effects of age, puberty and sunlight, that were not allowed for.

Puberty

Our study revealed a negative correlation between Tanner stage of puberty and 25(OH)D concentration in both boys and girls, which persisted after adjusting for age, sun exposure, physical activity and fat mass index, while some previous studies showed that serum 25(OH)D concentration decreased with increasing Tanner stage in boys (but not in girls), with the greatest proportion of vitamin D insufficiency being found during Tanner stages 4–5⁽¹⁷⁾. As in our results, some other studies have shown negative associations between pubertal status and serum 25(OH)D concentration in both sexes^(15,18,37,38), although increasing age, lesser outdoor sport and greater demands on vitamin D stores for skeletal mineralization could contribute to this finding^(17,39). It is necessary to consider pubertal stage as it is associated with a remarkable change in the lifestyle of children in terms of daily sun exposure and outdoor physical activity; clothing and percentage of body surface area exposed may contribute significantly to change in 25(OH)D concentration in this stage⁽⁴⁰⁾.

Exercise

The present study showed that exercise and sun exposure have a positive association with vitamin D status after adjusting the analysis for BMI or fat mass index, as previously suggested by a study in Saudi adolescents that showed physical activity was directly associated with 25(OH)D concentration, independent of sunshine exposure⁽⁴¹⁾. Physical activity was shown to be associated with

improved local bone mass, decreased Ca excretion and increased Ca absorption in a further study, while exercise could reduce body weight and BMI, thereby reducing vitamin D sequestration into fatty tissues allowing higher serum 25(OH)D values⁽⁴²⁾.

Study limitations

Our study was cross-sectional in nature and the various vitamin D-related genetic factor axes (known to contribute independently to growth and probably also to pubertal status) were not examined, so that these are possible relevant factors. Future studies in this area should therefore ensure that changes in vitamin D status during childhood and adolescence can be adjusted for the effects of all the other relevant factors, such as age, clothing, vitamin D-gene variant axes and leptin secretion.

Conclusion

The prevalence of vitamin D deficiency was 82% among children in the south of Iran and 25(OH)D concentration was related to sun exposure, physical activity, age and pubertal stage of the children. Health improvement strategies should be developed to prevent vitamin D deficiency during this critical period, although further studies are needed to determine the pathophysiology of the decrease in serum 25(OH)D concentration seen during and after puberty, since it appears to be independent of the effects of physical activity, pubertal status, fat mass index or exposure to sunshine.

Acknowledgements

Acknowledgements: The authors would like to thank Dr Nasrin Shokrpour for editorial assistance and Mrs Sareh Roosta for statistical analysis at Center for Development of Clinical Research of Nemazee Hospital. *Financial support:* This work was supported by Shiraz University of Medical Sciences (grant number 89–5127). The funder had no role in the design, analysis or writing of this article. Shiraz University of Medical Sciences funded our research just as a thesis for graduate students. *Conflict of interest:* None. *Authorship:* F.S. formulated the research questions, designed the study, analysed the data and wrote the article. M.H.D. formulated the research questions, designed the study and analysed data. G.R.O. carried out the study. M.B. formulated the research question. *Ethics of human subject participation:* The study was approved by the Ethics Committee of Shiraz University of Medical Sciences. All parents and children who participated provided signed informed consent.

References

1. Holick MF (2003) Vitamin D: a millennium perspective. *J Cell Biochem* **88**, 296–307.

2. Holick MF (2007) Vitamin D deficiency. *N Engl J Med* **357**, 266–281.
3. Rabbani A, Alavian SM, Motlagh ME et al. (2009) Vitamin D insufficiency among children and adolescents living in Tehran, Iran. *J Trop Pediatr* **55**, 189–191.
4. Lips P (2007) Vitamin D status and nutrition in Europe and Asia. *J Steroid Biochem Mol Biol* **103**, 620–625.
5. Holick MF, Binkley NC, Bischoff-Ferrari HA et al. (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* **96**, 1911–1930.
6. Marwaha RK, Tandon N, Reddy DR et al. (2005) Vitamin D and bone mineral density status of healthy schoolchildren in northern India. *Am J Clin Nutr* **82**, 477–482.
7. Thacher TD, Fischer PR, Strand MA et al. (2006) Nutritional rickets around the world: causes and future directions. *Ann Trop Paediatr* **26**, 1–16.
8. Neyestani TR, Hajifaraji M, Omidvar N et al. (2012) High prevalence of vitamin D deficiency in school-age children in Tehran, 2008: a red alert. *Public Health Nutr* **15**, 324–330.
9. Ardestani PM, Salek M, Keshteli AH et al. (2010) Vitamin D status of 6- to 7-year-old children living in Isfahan, Iran. *Endokrynol Pol* **61**, 377–382.
10. Willis CM, Laing EM, Hall DB et al. (2007) A prospective analysis of plasma 25-hydroxyvitamin D concentrations in white and black prepubertal females in the southeastern United States. *Am J Clin Nutr* **85**, 124–130.
11. Zhu Z, Zhan J, Shao J et al. (2012) High prevalence of vitamin D deficiency among children aged 1 month to 16 years in Hangzhou, China. *BMC Public Health* **12**, 126.
12. Gordon CM, DePeter KC, Feldman HA et al. (2004) Prevalence of vitamin D deficiency among healthy adolescents. *Arch Pediatr Adolesc Med* **158**, 531–537.
13. Buyukinan M, Ozen S, Kokkun S et al. (2012) The relation of vitamin D deficiency with puberty and insulin resistance in obese children and adolescents. *J Pediatr Endocrinol Metab* **25**, 83–87.
14. Razzaghy-Azar M & Shakiba M (2010) Assessment of vitamin D status in healthy children and adolescents living in Tehran and its relation to iPTH, gender, weight and height. *Ann Hum Biol* **37**, 692–701.
15. Aypak C, Türedi O & Yüce A (2014) The association of vitamin D status with cardiometabolic risk factors, obesity and puberty in children. *Eur J Pediatr* **173**, 367–373.
16. Breen ME, Laing EM, Hall DB et al. (2011) 25-Hydroxyvitamin D, insulin-like growth factor-I, and bone mineral accrual during growth. *J Clin Endocrinol Metab* **96**, E89–E98.
17. Ginty F, Cavadini C, Michaud PA et al. (2004) Effects of usual nutrient intake and vitamin D status on markers of bone turnover in Swiss adolescents. *Eur J Clin Nutr* **58**, 1257–1265.
18. Fuleihan G, Nabulsi M, Choucair M et al. (2001) Hypovitaminosis D in healthy schoolchildren. *Pediatrics* **107**, E53.
19. Tanner JM, & Whitehouse RH (1976) Clinical longitudinal standards for height, weight, height velocity, weight velocity, and stages of puberty. *Arch Dis Child* **51**, 170–179.
20. Pate RR, Pratt M, Blair SN et al. (1995) Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* **273**, 402–407.
21. Lee YA, Kim HY, Hong H et al. (2014) Risk factors for low vitamin D status in Korean adolescents: the Korea National Health and Nutrition Examination Survey (KNHANES) 2008–2009. *Public Health Nutr* **17**, 764–771.
22. Choi HS, Oh HJ, Choi H et al. (2011) Vitamin D insufficiency in Korea – a greater threat to younger generation: the Korea National Health and Nutrition Examination Survey (KNHANES) 2008. *J Clin Endocrinol Metab* **96**, 643–651.
23. Ginde AA, Liu MC & Camargo CA Jr (2009) Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004. *Arch Intern Med* **169**, 626–632.
24. Kumar J, Muntner P, Kaskel FJ et al. (2009) Prevalence and associations of 25-hydroxyvitamin D deficiency in US children: NHANES 2001–2004. *Pediatrics* **124**, 362–370.
25. Alemzadeh R, Kichler J, Babar G et al. (2008) Hypovitaminosis D in obese children and adolescents: relationship with adiposity, insulin sensitivity, ethnicity, and season. *Metabolism* **57**, 183–191.
26. Shin YH, Shin HJ & Lee YJ (2013) Vitamin D status and childhood health. *Korean J Pediatr* **56**, 417–423.
27. Weng FL, Shults J, Leonard MB et al. (2007) Risk factors for low serum 25-hydroxyvitamin D concentrations in otherwise healthy children and adolescents. *Am J Clin Nutr* **86**, 150–158.
28. Tolppanen AM, Fraser A, Fraser WD et al. (2012) Risk factors for variation in 25-hydroxyvitamin D₃ and D₂ concentrations and vitamin D deficiency in children. *J Clin Endocrinol Metab* **97**, 1202–1210.
29. Jeong SJ (2013) Factor affecting the vitamin D status in South Korean children. *J Food Nutr Sci* **1**, 7–12.
30. Horan M, Gibney E, Molloy E et al. (2015) Methodologies to assess paediatric adiposity. *Ir J Med Sci* **184**, 53–68.
31. Poopedi MA, Norris SA & Pettifor JM (2011) Factors influencing the vitamin D status of 10-year-old urban South African children. *Public Health Nutr* **14**, 334–339.
32. Wortsman J, Matsuoka LY, Chen TC et al. (2000) Decreased bioavailability of vitamin D in obesity. *Am J Clin Nutr* **72**, 690–693.
33. Vimalaewaran KS, Berry DJ, Lu C et al. (2013) Causal relationship between obesity and vitamin D status: bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS Med* **10**, e1001383.
34. Tsuji K, Maeda T, Kawane T et al. (2010) Leptin stimulates fibroblast growth factor 23 expression in bone and suppresses renal 1 α ,25-dihydroxyvitamin D₃ synthesis in leptin-deficient mice. *J Bone Miner Res* **25**, 1711–1723.
35. Harel Z, Flanagan P, Forcier M et al. (2011) Low vitamin D status among obese adolescents: prevalence and response to treatment. *J Adolesc Health* **48**, 448–452.
36. Dong Y, Pollock N, Stallmann-Jorgensen IS et al. (2010) Low 25-hydroxyvitamin D levels in adolescents: race, season, adiposity, physical activity, and fitness. *Pediatrics* **125**, 1104–1111.
37. Guillemand J, Cabrol S, Allemandou A et al. (1995) Vitamin D-dependent seasonal variation of PTH in growing male adolescents. *Bone* **17**, 513–516.
38. Cashman KD (2002) Calcium intake, calcium bioavailability and bone health. *Br J Nutr* **87**, Suppl. 2, S169–S177.
39. Cashman KD (2007) Vitamin D in childhood and adolescence. *Postgrad Med J* **83**, 230–235.
40. Gannagé-Yared MH, Chemali R, Yaacoub N et al. (2000) Hypovitaminosis D in a sunny country: relation to lifestyle and bone markers. *J Bone Miner Res* **15**, 1856–1862.
41. Al-Musharaf S, Al-Othman A, Al-Daghri NM et al. (2012) Vitamin D deficiency and calcium intake in reference to increased body mass index in children and adolescents. *Eur J Pediatr* **171**, 1081–1086.
42. Michael YL, Smit E, Seguin R et al. (2011) Serum 25-hydroxyvitamin D and physical performance in postmenopausal women. *J Womens Health (Larchmt)* **20**, 1603–1608.