

# Embryos, DOHaD and David Barker

T. P. Fleming<sup>1\*</sup>, M. A. Velazquez<sup>1,†</sup> and J. J. Eckert<sup>2</sup>

<sup>1</sup>Centre for Biological Sciences, University of Southampton, Southampton, UK

<sup>2</sup>Faculty of Medicine, University of Southampton, Southampton, UK

The early embryo and periconceptual period is a window during which environmental factors may cause permanent change in the pattern and characteristics of development leading to risk of adult onset disease. This has now been demonstrated across small and large animal models and also in the human. Most evidence of periconceptual ‘programming’ has emerged from maternal nutritional models but also other *in vivo* and *in vitro* conditions including assisted reproductive treatments, show consistent outcomes. This short review first reports on the range of environmental *in vivo* and *in vitro* periconceptual models and resulting long-term outcomes. Second, it uses the rodent maternal low protein diet model restricted to the preimplantation period and considers the stepwise maternal-embryonic dialogue that comprises the induction of programming. This dialogue leads to cellular and epigenetic responses by the embryo, mainly identified in the extra-embryonic cell lineages, and underpins an apparently permanent change in the growth trajectory during pregnancy and associates with increased cardiometabolic and behavioural disease in adulthood. We recognize the important advice of David Barker some years ago to investigate the sensitivity of the early embryo to developmental programming, an insight for which we are grateful.

Received 27 February 2015; Revised 27 March 2015; Accepted 5 April 2015; First published online 8 May 2015

**Key words:** assisted reproductive treatment (ART), blastocyst, maternal low protein diet, primitive endoderm, trophoctoderm

## Introduction

Research in recent years has identified a close association between early embryonic development and the processes that lead to adverse programming of postnatal phenotype and the risks of adult disease – the framework of the Developmental Origins of Health and Disease (DOHaD) hypothesis and the mission of this journal. Evidence from both clinical and animal models points to a vulnerability or at least sensitivity of the early embryo to environmental conditions that can trace a pathway culminating in the setting of long-term body characteristics and disease risk.<sup>1–3</sup> This short review provides a summary of this evidence, considers what future directions we need to take to explore the relationship further, and recognizes the insight of David Barker in identifying the relevance of the periconceptual period for future health and encouraging investigations into this area.

## Embryonic and periconceptual environment and developmental programming

Several types of environmental cue may be sensed by early embryos and/or by the gametes before fertilization and lead to permanent changes in the developmental programme culminating in an altered postnatal phenotype associate with increased disease risk. These come from both *in vivo* conditions, including poor maternal or paternal nutrition and health

status, and *in vitro* conditions, present mainly from assisted reproductive treatments (ART) usually to overcome infertility in the human but also to promote reproductive efficiency in domestic animals. These environmental cues are within the normal range of experience. For example, poor nutrition does not necessarily mean famine or starvation and includes both under-nutrition with reduced protein or energy levels and over-rich nutrient levels as in diets high in fat and energy constituents. Moreover, evidence of embryo and periconceptual programming has been forthcoming from across mammalian species, including rodents, large domestic animals and the human. There have been several recent reviews of embryo and periconceptual developmental programming.<sup>3–8</sup>

## Evidence from maternal nutrition

We have used as our main model mouse maternal low protein diet limited only to the preimplantation period after mating and with normal diet for the remainder of pregnancy and postnatally (Emb-LPD; 9% casein). This diet, *v.* isocaloric normal protein diet controls (NPD; 18% casein), caused foetal growth to be increased and, in female offspring, excess weight and increased adiposity through to late adult life. In addition, adult cardiovascular dysfunction occurred in both sexes, notably hypertension, attenuated arterial dilatation capacity and increased levels of angiotensin converting enzyme, coupled with abnormal hyperactive behaviour pattern in female offspring.<sup>9–11</sup> Emb-LPD also caused adult cardiometabolic disease in rats.<sup>12</sup> Moreover, a similar adult cardiometabolic and behavioural phenotype occurred in response to maternal Emb-LPD when administered exclusively during the period of oocyte maturation.<sup>13</sup> Sheep models comprising maternal

\*Address for correspondence: T. P. Fleming, Centre for Biological Sciences, University of Southampton, Southampton SO16 6YD, UK.  
(Email [tpf@soton.ac.uk](mailto:tpf@soton.ac.uk))

† Present address: School of Agriculture, Food and Rural Development, Newcastle University, Newcastle Upon Tyne NE1 7RU, UK.

periconceptual under-nutrition and normal nutrition thereafter also result in cardiometabolic and behavioural dysfunction in adult offspring.<sup>14–16</sup> Similarly, restricting the supply of B-vitamins and methionine during the periconceptual period in sheep results in adverse cardiometabolic health in postnatal offspring.<sup>17</sup> In the human, although equivalent maternal under-nutritional data sets do not exist, it has been shown that people conceived and experienced early gestation during periods of famine in the Netherlands during WW2 or in China during the later cultural revolution had increased risks of cardiometabolic and behavioural disease as adults.<sup>1,18,19</sup>

A substantial literature similarly shows that maternal over-nutrition and obesity increase health risks of both the pregnant mother herself but also of her offspring later in life.<sup>20–23</sup> The majority of evidence stems from maternal obesity experienced throughout pregnancy (and lactation) (reviewed in<sup>24</sup>). However, reduced fertility linked to obesity and involving difficulty to ovulate and establish a pregnancy suggests that, in addition to the nutritional impact on foetal growth during ongoing pregnancies, over-nutrition also profoundly affects the developmental capacity of the oocyte and early embryo before implantation, similar to the murine low protein diet models discussed above. This concept of a periconceptual sensitivity to environmental challenge holds up across a number of different species. For example, diet induced obesity affects ovulation, fertilization, developmental capacity and differentiation in farm animals and rodents (rats, mice, rabbits) alike.<sup>25–27</sup>

In many studies it is difficult to distinguish whether the dietary effects were induced during final gamete development and maturation, preimplantation embryo development and/or even after implantation had occurred during foetal stages. This is because development has continued within the same mother thus not accounting for potential ongoing consequences of the nutritional regime after change to a control diet. This may compromise our mechanistic understanding. Although embryo transfer can have its own pitfalls regarding foetal and postnatal health risks, it is the only tool that allows us to define periconceptual exposure to different environments. Thus, reciprocal embryo transfer designs between control and challenged mothers clearly show that timing of maternal diet challenge is critical in determining the nature of consequences during foetal and postnatal life following maternal high fat diet (HFD).<sup>28</sup> For example, while pre-gestational (10–12 weeks before mating) or gestational (from mating to term) exposure to a maternal HFD leads to impaired foetal and placental growth, only a HFD exposure during gestation resulted in obesity and impaired glucose tolerance in offspring.<sup>28</sup> Recent work focusing on gamete origin and environment is showing that the final stages of oocyte and also sperm maturation are prone to programming and induction of long-term consequences for future health. For example, obtaining oocytes or zygotes from obese (or diabetic) mothers either before or just after fertilization and transferring the resulting embryos to non-obese foster mothers fed a control diet has shown that oocyte metabolism, spindle integrity and gene expression are all targets of nutritional

programming. Furthermore, REDOX state, ER stress, mitochondrial function and even mitochondrial number can be compromised up to at least the blastocyst stage.<sup>29–33</sup> Epigenetic mis-regulation as a consequence of parental obesity (often linked with a diabetic or pre-diabetic state) has also been demonstrated recently in gametes. A number of such epigenetic faults can persist not only in blastocysts but even in offspring organs through several generations.<sup>34–36</sup> Moreover, this appears to be transmittable in a sex-specific manner.

While it is also well demonstrated that maternal obesity profoundly impacts on offspring health and development in the human (reviewed in<sup>23</sup>), the effect of maternal obesity on oocyte and early embryo physiology is less well characterized. Recent data from human preimplantation embryos suggest that maternal BMI similarly impacts on blastocyst metabolic profiles, developmental competence and differentiation,<sup>37</sup> as seen in animal models. Since human embryos are only accessible during fertility treatment procedures where the oocyte is removed from the maternal environment before fertilization, such impact of maternal obesity is most likely to have been triggered during final oocyte development and maturation.

Good evidence for a role of paternal obesity is also starting to emerge, either through metabolically or epigenetically altered sperm themselves or indirectly via the seminal fluids and their interaction with the maternal reproductive tract.<sup>38–40</sup> Developing preventative strategies based, for example, on controlled weight loss programmes, are beginning to emerge yet with varying success with regards to fully protecting offspring health without compromise.<sup>6,20,27,35,41</sup>

### *Evidence from maternal sickness*

Severe maternal infection and subsequent sickness during the periconceptual and early pregnancy periods are known to increase the miscarriage rate.<sup>42</sup> However, whether, like nutritional models, such maternal conditions at a milder level may cause adverse programming of the embryo that affects postnatal health has received very little attention. To test such possibilities, we used a mouse model in which bacterial endotoxin at varying but relatively low levels were administered intra-peritoneally to mothers on a single occasion, the morning after mating.<sup>43</sup> Endotoxin treatment caused a transient and typical sickness response in dams with elevated serum pro-inflammatory cytokine release over 1–2 days. However, this treatment, distinct from nutritional models discussed above, had minimal influence on postnatal cardiometabolic or behavioural health but did affect the offspring innate immune system. Thus, adult offspring challenged with endotoxin themselves showed an attenuated cytokine response that was inversely related to the dose received by their mothers when they were zygotes. This model suggests innate immunity may be programmed from embryo environment *in vivo* with pathogen-rich maternal conditions acting to suppress offspring immunity, a mechanism to protect against auto-immune damage.<sup>43</sup> Different *in vivo* conditions may

therefore affect early embryos in different ways and distinct long-term consequences.

### **Evidence from ART**

Several million children have now been born by ART but epidemiological evaluation of the effect of gamete and embryo *in vitro* manipulations on their health status through to adulthood remains unclear. This is because it is difficult to discriminate between effects mediated through parental infertility and the genetic status of children, the obstetric and perinatal treatments received, and the actual IVF and ART treatments undertaken. Coupled with this, gradual advances in technologies and the general reduction in multiple embryo transfer complicate broad comparisons with the health of naturally conceived children. Nevertheless, in several recent reviews of ART children health, the increased risk of adverse perinatal outcome (notably growth restriction and prematurity), congenital abnormalities and imprinting disorders remain a concern of the ART treatment.<sup>2,44–46</sup> Studies assessing the health of ART children have documented a susceptibility to cardiometabolic disorders including high blood pressure, increased glucose levels and adiposity, and altered growth rates in early years, with growth velocity associating with increased blood pressure.<sup>47–50</sup> Significantly, birthweight and continuing body weight during early infancy can associate with the type of commercial culture medium used in ART, a legacy from 9 months previously and clear demonstration of the influence of the period around conception.<sup>51</sup> Also, birthweight can be heavier in cryopreserved embryos,<sup>52</sup> likely mediated either by an effect of the freezing process on the embryo or by the more natural uterine steroidal environment in a frozen embryo transfer cycle. There is also limited evidence of increased neurodevelopmental delay, cerebral palsy and prevalence to clinical depression in ART children but the influence of confounding factors may be contributory.<sup>53</sup>

These concerns from epidemiological studies on ART children indicate some similarities to the adverse cardiometabolic and behavioural health outcomes identified from periconceptional and embryo exposures to poor maternal nutrition. Animal studies have been able to overcome the limitations in interpreting the causes of health disorders in ART offspring with confounders such as parental infertility, genetic variation and perinatal complications controlled for. In well-controlled mouse studies, preimplantation embryo culture and subsequent transfer is sufficient to induce cardiometabolic disorders in adult offspring including relative hypertension,<sup>54</sup> glucose intolerance and insulin resistance,<sup>55,56</sup> abnormal hepatic and fat metabolomes<sup>57</sup> and behavioural deficits including memory loss.<sup>58,59</sup> Interestingly, combinations of different environments during development, such as embryo culture and later maternal malnutrition, act synergistically to modify postnatal phenotype.<sup>60</sup> A mouse study of particular importance demonstrated that postnatal growth rates could be increased or decreased by subtle *in vitro* manipulation of energy

levels during the pronuclear zygote stage after fertilization.<sup>61</sup> Similar effects of embryo culture have been demonstrated in larger animals.<sup>62</sup>


### **From embryo environment to adult phenotype, what are the links?**

#### **Wise advice**

As discussed above, from both *in vivo* nutritional and *in vitro* ART-related studies, across animal and human data sets, there is consistent evidence that periconceptional environment is critical in the long-term setting of postnatal phenotype. Why might events around the time of conception be so pivotal in DOHaD? My (TPF) first discussion on this question was with David Barker back in the late 1990s. At the time, our research lab focused mainly on *intrinsic* developmental mechanisms of the early embryo leading to blastocyst morphogenesis and segregation of early embryonic and extra-embryonic cell lineages. David suggested to me that *extrinsic* factors were likely to be modulating these early inherent steps in embryogenesis with far greater impact on long-term potential. There are good reasons why these insightful comments might be true. The pre-implantation period is characterized by diversification of two extraembryonic lineages, trophoblast (TE) and primitive endoderm (PE), on the outside and inside, respectively, of the expanding blastocyst with the epiblast progenitor of the entire foetus located between them (Fig. 1). TE and PE subsequently construct the chorio-allantoic placenta and yolk sac placenta, respectively, with responsibility in provision of nutrients during post-implantation pregnancy. If early embryo environment can generate diverse adult phenotypes as reviewed above, perhaps the characteristics of nascent placental lineages may be subject to external cues to optimize their efficiency dependent upon nutrient availability – a concept commonly known today as developmental plasticity? Moreover, the close signalling mediated between maternal and embryonic cells to coordinate implantation<sup>63</sup> and the epigenetic restructuring of the chromatin occurring during preimplantation development<sup>64</sup> provides a natural window when such plasticity could be induced in the embryo and propagated, through altered gene expression, into later pregnancy. David Barker's comments led to new research investigating the relevance of periconceptional environment in developmental programming as a consequence.

#### **Maternal signalling, embryo sensing and lineage responses**

If, as considered above, the early embryo constitutes a suitable window for *in vivo* maternal environmental information to influence and optimize the pattern of future development, such plasticity must comprise a series of mechanisms both to permit embryo sensing of the external milieu and to activate appropriate responses to the developmental programme. This dialogue between mother and early embryo has been a focus of our laboratory in recent years using the mouse Emb-LPD model.<sup>3</sup>



**Fig. 1.** Diverse environmental conditions during the preimplantation period can impact on the early embryo and affect cell lineages at the blastocyst stage with detrimental effects in postnatal health of the resultant offspring. (a) Mid-plane confocal microscopy section displaying Cdx2 staining in nuclei of trophectoderm cells (pale blue) from a mouse blastocyst. Cells of the epiblast and primitive endoderm (dotted white contour) were stained with DAPI (dark blue nuclei).

We find, during the period of preimplantation development, that Emb-LPD is able to alter the metabolite composition of maternal serum, which in turn modifies the composition of the uterine lumen. Coupled with this, embryos at the blastocyst stage are equipped with signalling pathways that can ‘read’ these metabolic cues and then change their developmental trajectory as a result, most evident in the extra-embryonic lineages. These changes in developmental plan can best be described as compensatory, optimizing the nutrient provision from mother to embryo to protect growth. Most interesting is that this dialogue at least in mice has enduring consequences. It appears that some key decisions are made in developmental plasticity at this time and that they are sufficiently hard-wired to persist through the entire gestation period. We believe this dialogue to be the core component of embryo programming in this model leading to the adult onset disease phenotype described above. This is because the increased foetal growth response identified in Emb-LPD offspring is a biomarker of

future disease – birthweight of Emb-LPD or sustained LPD throughout gestation is positively correlated with later adult weight, blood pressure and behavioural activity.<sup>9</sup> Moreover, the Emb-LPD adult metabolism is energy-storage in character with increased insulin receptor and IGF-1R expression in white adipose tissue,<sup>11</sup> further stimulating weight increase and disease risk. We briefly summarize critical features in this maternal-embryonic dialogue below, a fuller account is reviewed elsewhere.<sup>3</sup>

The protein composition of maternal diet, as would be anticipated, changes the metabolite composition of maternal serum, and, following Emb-LPD, amino acids (AAs) and insulin concentrations are reduced while glucose concentration rises; this is evident during the period of preimplantation development and particularly during blastocyst morphogenesis, and identified in rat<sup>12</sup> as well as mouse<sup>65</sup> models. Embryo metabolism and development are sensitive to all of these metabolites<sup>66–69</sup> and therefore offers the potential for dietary composition to be sensed by embryos. Indeed, microanalysis of mouse Emb-LPD uterine fluid at the time of blastocyst formation shows a similar reduction in AA concentration as found in the serum, as well as attendant changes in the AA profile within the blastocysts.<sup>65</sup> Most relevant to the embryo sensing mechanism that may activate programming is that the concentration of the branched chain AAs (BCAAs), leucine, isoleucine and valine, are significantly depleted in Emb-LPD rat and mouse serum and mouse uterine fluid.<sup>12,65</sup> That these metabolite changes are indeed sensed by mouse Emb-LPD blastocysts has been demonstrated by quantitative immunoblotting of blastocyst mTORC1 signalling.<sup>65</sup> This pathway is the major cellular mechanism of nutrient sensing to regulate growth rate, mediated through extracellular BCAA and insulin levels.<sup>70</sup> Thus, Emb-LPD blastocysts exhibit significantly depleted mTORC1 signalling compared with control blastocysts from NPD fed mothers.<sup>65</sup> This, we believe, comprises the Emb-LPD sensing mechanism required for induction of developmental plasticity in periconceptual nutritional programming.

The response mechanisms by mouse extra-embryonic lineages following Emb-LPD sensing are diverse and include increased proliferation and endocytosis in the TE as well as increased motility and invasiveness of trophoblast cells during the implantation process, evidenced in *in vitro* outgrowths.<sup>65,71</sup> The stimulation in endocytosis activity and proliferation is also apparent in the Emb-LPD PE lineage, evaluated in embryoid bodies following embryonic stem cell derivation.<sup>71,72</sup> Such changes in cellular endocytosis are comprehensive, including increased uptake of extracellular ligands, increased numbers of lysosomes and increased expression and activity of the relevant endocytic receptor, megalin, brought about through Rho-A modulation of the actin cytoskeleton.<sup>71</sup> These responses are compensatory, permitting increased uptake of nutrients in conditions of poor maternal diet and establishing more efficient chorio-allantoic and yolk sac placentas, evident throughout pregnancy to protect foetal growth.<sup>9,73</sup> The responses are

regulated through the integration of epigenetic and cellular mechanisms. For example, the stimulated proliferation within the PE lineage is accompanied by reduced gene expression of the *Gata6* transcription factor that acts to induce PE differentiation.<sup>72</sup> Thus, the *Gata6* promoter in the outer PE layer of Emb-LPD embryoid bodies exhibits histone H3 and H4 hypoacetylation and reduced RNA polymerase binding, characteristics that would suppress expression.<sup>72</sup>

That the induction and subsequent legacy of altered developmental plasticity derives from the maternal-embryonic dialogue can be demonstrated by embryo transfer of Emb-LPD blastocysts into control NPD recipients resulting in the stimulated foetal growth phenotype that leads to cardiometabolic disease in later life.<sup>9</sup> In a more refined *in vitro* model, we have now cultured embryos through cleavage in medium comprising low BCAA and insulin concentrations to mimic the Emb-LPD maternal environment and found, after transfer and gestation, offspring to exhibit the increased growth and relative hypertensive state typical of Emb-LPD progeny.<sup>74</sup>

## Conclusions

We have identified across different mammalian species including the human that periconceptual or preimplantation environment can alter the developmental programme and have long-lasting consequences for foetal growth and development and postnatal health and disease risk. There is also good evidence that the nature of the environmental condition can induce diverse changes affecting a range of physiological systems, for example the different legacies found from maternal dietary and sickness models. This indicates there is specificity in programming and that it is not merely a stress response with a single default consequence. While many examples of embryo programming are likely to be the result of a direct perturbation under extreme conditions, others can be viewed as true biological processes, conserved during evolution, to drive developmental plasticity and optimize offspring phenotype based upon cues acting in a 'predictive' manner. The temporal steps in programming of embryos through to adult disease are usually poorly understood and in only limited models do we have evidence of the sequence of events involved, such as the Emb-LPD model. This gap in our knowledge needs to be filled. Embryo programming should be considered an integrated phenomenon involving multiple biological components – for example, Emb-LPD programming involves physiological, epigenetic, cell biological and metabolic mechanisms. Epigenetic changes may facilitate the driving of cellular and physiological responses to environmental conditions by embryos to provide a coherent plan for developmental plasticity. Clearly, identifying the major inductive factors that may initiate adverse embryo programming (such as BCAAs and insulin in Emb-LPD) will permit a preventative strategy to protect future health, most important in human ART. Lastly, our thanks to David Barker for his wisdom in advising research into embryo developmental programming, his influence was pivotal in the progress made in this field.

## Acknowledgements

This work was supported through awards from the Biotechnology and Biological Sciences Research Council [BB/I001840/1; BB/F007450/1], The Medical Research Council [G9800781], the NICHD National Cooperative Program [U01 HD044635] and the EU-FP7 EpiHealth and EpiHealthNet programmes to TPF.

## Conflicts of Interest

None.

## Ethical Standards

All cited research involving animals was conducted under UK Home Office project and personal licences and local ethical approval.

## References

- Roseboom TJ, Painter RC, van Abeelen AF, Veenendaal MV, de Rooij SR. Hungry in the womb: what are the consequences? Lessons from the Dutch famine. *Maturitas*. 2011; 70, 141–145.
- Hart R, Norman RJ. The longer-term health outcomes for children born as a result of IVF treatment: Part I—General health outcomes. *Hum Reprod Update*. 2013; 19, 232–243.
- Fleming TP, Watkins AJ, Sun C, *et al*. Do little embryos make big decisions? How maternal dietary protein restriction can permanently change an embryo's potential, affecting adult health. *Reprod Fertil Dev*. 2015; 27, 684–692.
- Fleming TP, Velazquez MA, Eckert JJ, Lucas ES, Watkins AJ. Nutrition of females during the peri-conceptual period and effects on foetal programming and health of offspring. *Anim Reprod Sci*. 2012; 130, 193–197.
- El Hajj N, Haaf T. Epigenetic disturbances in *in vitro* cultured gametes and embryos: implications for human assisted reproduction. *Fertil Steril*. 2013; 99, 632–641.
- Zhang S, Rattanatray L, Morrison JL, *et al*. Maternal obesity and the early origins of childhood obesity: weighing up the benefits and costs of maternal weight loss in the periconceptual period for the offspring. *Exp Diabetes Res*. 2011; 2011, 585749.
- Jungheim ES, Moley KH. Current knowledge of obesity's effects in the pre- and periconceptual periods and avenues for future research. *Am J Obstet Gynecol*. 2010; 203, 525–530.
- McMillen IC, MacLaughlin SM, Muhlhausler BS, *et al*. Developmental origins of adult health and disease: the role of periconceptual and foetal nutrition. *Basic Clin Pharmacol Toxicol*. 2008; 102, 82–89.
- Watkins AJ, Ursell E, Panton R, *et al*. Adaptive responses by mouse early embryos to maternal diet protect fetal growth but predispose to adult onset disease. *Biol Reprod*. 2008; 78, 299–306.
- Watkins AJ, Lucas ES, Torrens C, *et al*. Maternal low-protein diet during mouse pre-implantation development induces vascular dysfunction and altered renin-angiotensin-system homeostasis in the offspring. *Br J Nutr*. 2010; 103, 1762–1770.
- Watkins AJ, Lucas ES, Wilkins A, Cagampang FR, Fleming TP. Maternal periconceptual and gestational low protein diet affects mouse offspring growth, cardiovascular and adipose phenotype at 1 year of age. *PLoS One*. 2011; 6, e28745.

12. Kwong WY, Wild AE, Roberts P, Willis AC, Fleming TP. Maternal undernutrition during the preimplantation period of rat development causes blastocyst abnormalities and programming of postnatal hypertension. *Development*. 2000; 127, 4195–4202.
13. Watkins AJ, Wilkins A, Cunningham C, et al. Low protein diet fed exclusively during mouse oocyte maturation leads to behavioural and cardiovascular abnormalities in offspring. *J Physiol*. 2008; 586, 2231–2244.
14. Torrens C, Snelling TH, Chau R, et al. Effects of pre- and periconceptional undernutrition on arterial function in adult female sheep are vascular bed dependent. *Exp Physiol*. 2009; 94, 1024–1033.
15. Todd SE, Oliver MH, Jaquiere AL, Bloomfield FH, Harding JE. Periconceptional undernutrition of ewes impairs glucose tolerance in their adult offspring. *Pediatr Res*. 2009; 65(4), 409–413.
16. Hernandez CE, Matthews LR, Oliver MH, Bloomfield FH, Harding JE. Effects of sex, litter size and periconceptional ewe nutrition on offspring behavioural and physiological response to isolation. *Physiol Behav*. 2010; 101, 588–594.
17. Sinclair KD, Allegrucci C, Singh R, et al. DNA methylation, insulin resistance, and blood pressure in offspring determined by maternal periconceptional B vitamin and methionine status. *Proc Natl Acad Sci U S A*. 2007; 104, 19351–19356.
18. Li Y, Jaddoe VW, Qi L, et al. Exposure to the Chinese famine in early life and the risk of hypertension in adulthood. *J Hyperten*. 2011; 29(6), 1085–1092.
19. Xu MQ, Sun WS, Liu BX, et al. Prenatal malnutrition and adult schizophrenia: further evidence from the 1959–1961 Chinese famine. *Schizophr Bull*. 2009; 35, 568–576.
20. Lane M, Zander-Fox DL, Robker RL, McPherson NO. Periconception parental obesity, reproductive health, and transgenerational impacts. *Trends Endocrinol Metab*. 2015; 26, 84–90.
21. Tenenbaum-Gavish K, Hod M. Impact of maternal obesity on fetal health. *Fetal Diag Ther*. 2013; 34, 1–7.
22. O'Reilly JR, Reynolds RM. The risk of maternal obesity to the long-term health of the offspring. *Clin Endocrinol*. 2013; 78, 9–16.
23. Ruager-Martin R, Hyde MJ, Modi N. Maternal obesity and infant outcomes. *Early Hum Dev*. 2010; 86, 715–722.
24. Alfaradhi MZ, Ozanne SE. Developmental programming in response to maternal overnutrition. *Front Genet*. 2011; 2, 27.
25. Picone O, Laigre P, Fortun-Lamothe L, et al. Hyperlipidic hypercholesterolemic diet in prepubertal rabbits affects gene expression in the embryo, restricts fetal growth and increases offspring susceptibility to obesity. *Theriogenology*. 2011; 75, 287–299.
26. Dunning KR, Russell DL, Robker RL. Lipids and oocyte developmental competence: the role of fatty acids and beta-oxidation. *Reproduction*. 2014; 148, R15–27.
27. Nicholas LM, Morrison JL, Rattanaray L, et al. Differential effects of exposure to maternal obesity or maternal weight loss during the periconceptional period in the sheep on insulin signalling molecules in skeletal muscle of the offspring at 4 months of age. *PLoS One*. 2013; 8, e84594.
28. Sasson IE, Vitins AP, Mainigi MA, Moley KH, Simmons RA. Pre-gestational vs gestational exposure to maternal obesity differentially programs the offspring in mice. *Diabetologia*. 2015; 58, 615–624.
29. Luzzo KM, Wang Q, Purcell SH, et al. High fat diet induced developmental defects in the mouse: oocyte meiotic aneuploidy and fetal growth retardation/brain defects. *PLoS One*. 2012; 7, e49217.
30. Turner N, Robker RL. Developmental programming of obesity and insulin resistance: does mitochondrial dysfunction in oocytes play a role? *Mol Hum Reprod*. 2015; 21, 23–30.
31. Igosheva N, Abramov AY, Poston L, et al. Maternal diet-induced obesity alters mitochondrial activity and redox status in mouse oocytes and zygotes. *PLoS One*. 2010; 5, e10074.
32. Grindler NM, Moley KH. Maternal obesity, infertility and mitochondrial dysfunction: potential mechanisms emerging from mouse model systems. *Mol Hum Reprod*. 2013; 19, 486–494.
33. Wu LL, Russell DL, Wong SL, et al. Mitochondrial dysfunction in oocytes of obese mothers: transmission to offspring and reversal by pharmacological endoplasmic reticulum stress inhibitors. *Development*. 2015; 142, 681–691.
34. Ge ZJ, Luo SM, Lin F, et al. DNA methylation in oocytes and liver of female mice and their offspring: effects of high-fat-diet-induced obesity. *Environ Health Perspect*. 2014; 122, 159–164.
35. Wei Y, Yang CR, Wei YP, et al. Enriched environment-induced maternal weight loss reprograms metabolic gene expression in mouse offspring. *J Biol Chem*. 2015; 290, 4604–4619.
36. Wei Y, Schatten H, Sun QY. Environmental epigenetic inheritance through gametes and implications for human reproduction. *Hum Reprod Update*. 2015; 21, 194–208.
37. Leary C, Leese HJ, Sturmey RG. Human embryos from overweight and obese women display phenotypic and metabolic abnormalities. *Hum Reprod*. 2015; 30, 122–132.
38. Bromfield JJ, Schjenken JE, Chin PY, et al. Maternal tract factors contribute to paternal seminal fluid impact on metabolic phenotype in offspring. *Proc Natl Acad Sci U S A*. 2014; 111, 2200–2205.
39. McPherson NO, Fullston T, Aitken RJ, Lane M. Paternal obesity, interventions, and mechanistic pathways to impaired health in offspring. *Ann N Nutr Metab*. 2014; 64, 231–238.
40. Watkins AJ, Sinclair KD. Paternal low protein diet affects adult offspring cardiovascular and metabolic function in mice. *Am J Physiol Heart Circ Physiol*. 2014; 306, H1444–1452.
41. Nicholas LM, Rattanaray L, Morrison JL, et al. Maternal obesity or weight loss around conception impacts hepatic fatty acid metabolism in the offspring. *Obesity (Silver Spring)*. 2014; 22, 1685–1693.
42. Cram LF, Zapata MI, Toy EC, Baker B, 3rd. Genitourinary infections and their association with preterm labor. *Am Fam Physician*. 2002; 65, 241–248.
43. Williams CL, Teeling JL, Perry VH, Fleming TP. Mouse maternal systemic inflammation at the zygote stage causes blunted cytokine responsiveness in lipopolysaccharide-challenged adult offspring. *BMC Biol*. 2011; 9, 49.
44. Brison DR, Roberts SA, Kimber SJ. How should we assess the safety of IVF technologies? *Reprod Biomed Online*. 2013; 27, 710–721.
45. Fauser BC, Devroey P, Diedrich K, et al. Health outcomes of children born after IVF/ICSI: a review of current expert opinion and literature. *Reprod Biomed Online*. 2014; 28, 162–182.
46. Lazaraviciute G, Kauser M, Bhattacharya S, Haggarty P. A systematic review and meta-analysis of DNA methylation levels and imprinting disorders in children conceived by IVF/ICSI

- compared with children conceived spontaneously. *Hum Reprod Update*. 2014; 20, 840–852.
47. Ceelen M, van Weissenbruch MM, Vermeiden JP, van Leeuwen FE, Delemarre-van de Waal HA. Cardiometabolic differences in children born after in vitro fertilization: follow-up study. *J Clin Endocrinol Metab*. 2008; 93, 1682–1688.
  48. Ceelen M, van Weissenbruch MM, Roos JC, et al. Body composition in children and adolescents born after in vitro fertilization or spontaneous conception. *J Clin Endocrinol Metab*. 2007; 92, 3417–3423.
  49. Ceelen M, van Weissenbruch MM, Prein J, et al. Growth during infancy and early childhood in relation to blood pressure and body fat measures at age 8–18 years of IVF children and spontaneously conceived controls born to subfertile parents. *Hum Reprod*. 2009; 24, 2788–2795.
  50. Sakka SD, Loutradis D, Kanaka-Gantenbein C, et al. Absence of insulin resistance and low-grade inflammation despite early metabolic syndrome manifestations in children born after in vitro fertilization. *Fertil Steril*. 2010; 94, 1693–1699.
  51. Zandstra H, Van Montfoort AP, Dumoulin JC. Does the type of culture medium used influence birthweight of children born after IVF? *Hum Reprod*. 2015; 30, 530–542.
  52. Vergouw CG, Kosteljik EH, Doejaaren E, et al. The influence of the type of embryo culture medium on neonatal birthweight after single embryo transfer in IVF. *Hum Reprod*. 2012; 27, 2619–2626.
  53. Hart R, Norman RJ. The longer-term health outcomes for children born as a result of IVF treatment. Part II–Mental health and development outcomes. *Hum Reprod Update*. 2013; 19, 244–250.
  54. Watkins AJ, Platt D, Papenbrock T, et al. Mouse embryo culture induces changes in postnatal phenotype including raised systolic blood pressure. *Proc Natl Acad Sci U S A*. 2007; 104, 5449–5454.
  55. Scott KA, Yamazaki Y, Yamamoto M, et al. Glucose parameters are altered in mouse offspring produced by assisted reproductive technologies and somatic cell nuclear transfer. *Biol Reprod*. 2010; 83, 220–227.
  56. Donjacour A, Liu X, Lin W, Simbulan R, Rinaudo PF. In vitro fertilization affects growth and glucose metabolism in a sex-specific manner in an outbred mouse model. *Biol Reprod*. 2014; 90, 80.
  57. Feuer SK, Donjacour A, Simbulan RK, et al. Sexually dimorphic effect of in vitro fertilization (IVF) on adult mouse fat and liver metabolomes. *Endocrinology*. 2014; 155, 4554–4567.
  58. Ecker DJ, Stein P, Xu Z, et al. Long-term effects of culture of preimplantation mouse embryos on behavior. *Proc Natl Acad Sci U S A*. 2004; 101, 1595–1600.
  59. Fernandez-Gonzalez R, Moreira P, Bilbao A, et al. Long-term effect of in vitro culture of mouse embryos with serum on mRNA expression of imprinting genes, development, and behavior. *Proc Natl Acad Sci U S A*. 2004; 101, 5880–5885.
  60. Strata F, Giritharan G, Sebastiano FD, et al. Behavior and brain gene expression changes in mice exposed to preimplantation and prenatal stress. *Reprod Sci*. 2015; 22, 23–30.
  61. Banrezes B, Sainte-Beuve T, Canon E, et al. Adult body weight is programmed by a redox-regulated and energy-dependent process during the pronuclear stage in mouse. *PLoS One*. 2011; 6, e29388.
  62. Sinclair KD, Young LE, Wilmot I, McEvoy TG. In-utero overgrowth in ruminants following embryo culture: lessons from mice and a warning to men. *Hum Reprod*. 2000; 15(Suppl. 5), 68–86.
  63. Fritz R, Jain C, Armant DR. Cell signaling in trophoblast-uterine communication. *Int J Dev Biol*. 2014; 58, 261–271.
  64. Cantone I, Fisher AG. Epigenetic programming and reprogramming during development. *Nat Struct Mol Biol*. 2013; 20, 282–289.
  65. Eckert JJ, Porter R, Watkins AJ, et al. Metabolic induction and early responses of mouse blastocyst developmental programming following maternal low protein diet affecting life-long health. *PLoS One*. 2012; 7, e52791.
  66. Lane M, Gardner DK. Differential regulation of mouse embryo development and viability by amino acids. *J Reprod Fertil*. 1997; 109, 153–164.
  67. Kaye PL, Gardner HG. Preimplantation access to maternal insulin and albumin increases fetal growth rate in mice. *Hum Reprod*. 1999; 14, 3052–3059.
  68. Martin PM, Sutherland AE. Exogenous amino acids regulate trophectoderm differentiation in the mouse blastocyst through an mTOR-dependent pathway. *Dev Biol*. 2001; 240, 182–193.
  69. Martin PM, Sutherland AE, Van Winkle LJ. Amino acid transport regulates blastocyst implantation. *Biol Reprod*. 2003; 69, 1101–1108.
  70. Wang X, Proud CG. Nutrient control of TORC1, a cell-cycle regulator. *Trends Cell Biol*. 2009; 19, 260–267.
  71. Sun C, Velazquez MA, Marfy-Smith S, et al. Mouse early extra-embryonic lineages activate compensatory endocytosis in response to poor maternal nutrition. *Development*. 2014; 141, 1140–1150.
  72. Sun C, Denisenko O, Sheth B, et al. Epigenetic regulation of histone modifications and Gata6 gene expression induced by maternal diet in mouse embryoid bodies in a model of developmental programming. *BMC Dev Biol*. 2015; 15, 3.
  73. Coan PM, Vaughan OR, McCarthy J, et al. Dietary composition programmes placental phenotype in mice. *J Physiol*. 2011; 589(Pt 14), 3659–3670.
  74. Velazquez MA, Sheth B, Marfy-Smith S, Eckert J, Fleming TP. Insulin and branched-chain amino acid depletion during mouse in vitro preimplantation embryo development alters postnatal growth and cardiovascular physiology. World Congress of Animal Reproduction 2014, Edinburgh, UK. *Reproduction Abstracts*. 2014; 1, P096.