

# CONTINUITY OF HILBERT–KUNZ MULTIPLICITY AND F-SIGNATURE

THOMAS POLSTRA AND ILYA SMIRNOV 

**Abstract.** We establish the continuity of Hilbert–Kunz multiplicity and F-signature as functions from a Cohen–Macaulay local ring  $(R, \mathfrak{m}, k)$  of prime characteristic to the real numbers at reduced parameter elements with respect to the  $\mathfrak{m}$ -adic topology.

## §1. Introduction

In this paper we consider the problem of determining when two rings have similar singularities. As a motivating example let  $R = k[[x_1, \dots, x_n]]$  be the power series ring in  $n$  variables over a field  $k$  and  $f \in R$ . If we add a term of sufficiently high order to  $f$ , will the new singularity be similar to that of  $f$ ?

The origin of the above question can be traced at least as far back as 1956. Samuel [Sam56] proved if the Jacobian ideal  $J(f)$  of  $f$  is  $\mathfrak{m}$ -primary and if  $f - g \in (x_1, \dots, x_n)J(f)^2$ , then there is an automorphism  $\varphi: R \rightarrow R$  such that  $\varphi(f) = g$ . In particular, Samuel’s result asserts that if  $f$  has an isolated singularity and  $g$  is sufficiently close to  $f$ , then the rings  $R/(f)$  and  $R/(g)$  are isomorphic. Samuel’s result was furthered by Hironaka [Hir65]: if  $R/I$  is a reduced and equidimensional isolated singularity, then there is an integer  $e > 0$  such that for any ideal  $J$  that also defines a reduced and equidimensional singularity of same dimension and if  $I \equiv J \pmod{(x_1, \dots, x_n)^e}$  then  $R/I \cong R/J$ .

More recently, Cutkosky and Srinivasan have extended Samuel’s result to include all complete intersection prime ideals and Hironaka’s theorem to all reduced equidimensional ideals. See [CS93] for precise statements.

However, when  $f \in k[[x_1, \dots, x_n]]$  does not have an isolated singularity, the results of Samuel, Hironaka, and Cutkosky and Srinivasan do not allow

---

Received July 15, 2017. Revised November 14, 2018. Accepted November 19, 2018.

2010 Mathematics subject classification. 13D40, 13A35, 13H15, 14B05.

Polstra was supported in part by NSF Postdoctoral Research Fellowship DMS #1703856.

© 2018 *Foundation Nagoya Mathematical Journal*

us to relate the singularities of  $R/(f)$  with the singularities of  $R/(g)$ . To compare the singularity of  $R/(f)$  with the singularity of  $R/(g)$  we first require a formal notion of “closeness” of singularities. For example, one could require the Hilbert–Samuel functions of  $R/(f)$  and  $R/(g)$  to coincide. Srinivas and Trivedi [ST96] have shown if  $I = (f_1, \dots, f_h)$  is a complete intersection<sup>1</sup>, or more generally if  $I$  is a parameter ideal in a generalized Cohen–Macaulay ring, then the associated graded rings of  $I$  and an ideal  $J = (f_1 + \epsilon_1, \dots, f_h + \epsilon_h)$  will be isomorphic, provided  $\epsilon_1, \dots, \epsilon_h$  are in a sufficiently large power of the maximal ideal.

We now turn our attention to numerical invariants in positive characteristic. In particular, we introduce and explore the behavior of F-pure threshold, Hilbert–Kunz multiplicity, and F-signature with respect to  $\mathfrak{m}$ -adic topologies. In what follows, let  $(R, \mathfrak{m})$  be a local ring of prime characteristic  $p$ . Let  $F: R \rightarrow R$  denote the Frobenius endomorphism. Given ideal  $I \subseteq R$  and  $e \in \mathbb{N}$  let  $I^{[p^e]} = (i^{p^e} \mid i \in I)$  be the expansion of  $I$  along  $F^e$ . We shall always assume  $R$  is F-finite, meaning that  $F$  is a finite map.

The study of F-pure thresholds was initiated by Takagi and Watanabe [TW04]. If  $(R, \mathfrak{m}, k)$  is regular then the F-pure threshold of an element  $f \in R$  can be defined as the limit  $\text{fpt}(f) = \lim_{e \rightarrow \infty} (\nu_e(f)/p^e)$  where  $\nu_e(f) = \max\{t \in \mathbb{N} \mid f^t \notin \mathfrak{m}^{[p^e]}\}$ . Two elementary observations concerning F-pure thresholds are that if  $f, g \in R$  then  $\text{fpt}(f + g) \leq \text{fpt}(f) + \text{fpt}(g)$  and that if  $f \in \mathfrak{m}^{[p^{e_0}]}$  then  $\text{fpt}(f) \leq 1/p^{e_0}$ . From these observations it follows that the F-pure threshold function is continuous as a function from  $R \rightarrow \mathbb{R}$ : specifically, if  $f \in R$  and  $\delta > 0$  then there exists  $N \in \mathbb{N}$  such that for all  $g \in R$  such that  $f - g \in \mathfrak{m}^N$ ,  $|\text{fpt}(f) - \text{fpt}(g)| < \delta$ . Continuity of the F-pure threshold function motivates studying continuity properties of Hilbert–Kunz multiplicity and F-signature and sheds light onto the problem of relating the singularities of  $R/(f)$  with  $R/(g)$  when  $f$  and  $g$  are suitably close.

In 1983, Monsky, building on the earlier work of Kunz [Kun69, Kun76], defined a new invariant, the Hilbert–Kunz multiplicity of  $R$ , as the limit

$$e_{\text{HK}}(R) = \lim_{e \rightarrow \infty} \frac{\lambda(R/\mathfrak{m}^{[p^e]})}{p^{ed}}.$$

Monsky [Mon83] showed that this limit always exists. Values of  $e_{\text{HK}}(R)$  dictate the severity of the singularities of  $R$ . Most notable is that under mild

<sup>1</sup>This result was recently reproved by Adamus and Patel [AP17] using a more combinatorial method.

hypotheses  $e_{\text{HK}}(R) \geq 1$  with equality if and only if  $R$  is regular [WY00]. It has also been shown in [BE04, AE08] that small enough values of  $e_{\text{HK}}(R)$  imply  $R$  is Gorenstein and strongly F-regular. To be able to relate the singularities of  $R/(f)$  and  $R/(g)$  when  $f$  and  $g$  are suitably close it is natural to investigate the following question, which was originally asked to the second author by Luis Núñez-Betancourt:

QUESTION 1.1. Let  $(R, \mathfrak{m}, k)$  be a regular local ring of prime characteristic. Is the Hilbert–Kunz multiplicity function defined by  $f \mapsto e_{\text{HK}}(R/(f))$  a continuous function in the  $\mathfrak{m}$ -adic topology? Namely, if  $f \in R$  is an element and we are given  $\delta > 0$ , can we find an integer  $N$  such that for all  $g \in R$  such that  $f - g \in \mathfrak{m}^N$ ,

$$|e_{\text{HK}}(R/(f)) - e_{\text{HK}}(R/(g))| < \delta?$$

We now turn our attention to F-signature. Given finitely generated  $R$ -module  $M$  we let  $\text{frk}(M)$  denote the largest rank of a free  $R$ -module  $F$  for which there is a surjection  $M \rightarrow F$ . The F-signature of  $(R, \mathfrak{m}, k)$  was defined by Huneke and Leuschke [HL02], conceptualizing the earlier work [SVdB97] of Smith and Van den Bergh, as the limit

$$s(R) = \lim_{e \rightarrow \infty} \frac{\text{frk}(R^{1/p^e})}{\text{rank}(R^{1/p^e})}.$$

The number  $\text{frk}(R^{1/p^e})$  will be denoted by  $a_e(R)$  and referred to as the  $e$ th Frobenius splitting number of  $R$ . Tucker [Tuc12] showed existence of the above limit for all local rings<sup>2</sup>. F-signature is a measurement of singularities: in particular,  $s(R) \leq 1$  with equality if and only if  $R$  is regular by [HL02] and  $s(R) > 0$  if and only if  $R$  is strongly F-regular by [AL03]. As with Hilbert–Kunz multiplicity, it is natural to relate the singularities of  $R/(f)$  with  $R/(g)$  by investigating continuity properties of F-signature with respect to the  $\mathfrak{m}$ -adic topology.

QUESTION 1.2. Let  $(R, \mathfrak{m}, k)$  be a regular local ring of prime characteristic. Is the F-signature function defined by  $f \mapsto s(R/(f))$  a continuous function with respect to the  $\mathfrak{m}$ -adic topology? Namely, if  $f \in R$  is an element and we are given  $\delta > 0$ , can we find an integer  $N$  such that for all  $g \in R$

<sup>2</sup>The existence of the above limit has been recently established without the local hypothesis in [DPY16].

such that  $f - g \in \mathfrak{m}^N$

$$|s(R/(f)) - s(R/(g))| < \delta?$$

We are able to answer Questions 1.1 and 1.2, provided that  $f$  has no multiple factors, that is,  $R/(f)$  is reduced. Our main contribution is the following theorem:

**THEOREM 1.3.** *Let  $(R, \mathfrak{m}, k)$  be a complete Cohen–Macaulay  $F$ -finite ring of prime characteristic  $p > 0$ . If  $(f_1, \dots, f_c)$  is a parameter ideal such that  $R/(f_1, \dots, f_c)$  is reduced then for any  $\delta > 0$  there exists an integer  $N > 0$  such that for any  $\epsilon_1, \dots, \epsilon_c \in \mathfrak{m}^N$ :*

- (1) (Corollary 3.7)  $|e_{\text{HK}}(R/(f_1, \dots, f_c)) - e_{\text{HK}}(R/(f_1 + \epsilon_1, \dots, f_c + \epsilon_c))| < \delta$ ; and
- (2) (Theorem 3.12)  $|s(R/(f_1, \dots, f_c)) - s(R/(f_1 + \epsilon_1, \dots, f_c + \epsilon_c))| < \delta$ .

We show that Hilbert–Kunz multiplicity and F-signature functions are the uniform limit of functions which are locally constant with respect to the  $\mathfrak{m}$ -adic topology. Naively, one might also expect the Hilbert–Kunz multiplicity and F-signature functions are locally constant with respect to the  $\mathfrak{m}$ -adic topology:  $e_{\text{HK}}(R/(f)) = e_{\text{HK}}(R/(g))$  and  $s(R/(f)) = s(R/(g))$  can be forced by making  $f - g$  is sufficiently small. We point that this cannot hold for Hilbert–Kunz multiplicity in Example 4.1 and for F-signature in Example 4.2.

Similar to a number of recent developments in the study of invariants in prime characteristic rings, the proof of Theorem 1.3 comes by showing uniform convergence of continuous functions. We get our uniform convergence result by using the methods of Polstra and Tucker [PT18], where it was shown that suitable Noether normalizations provide a kind of “canonical” convergence estimate. The essential reason is that when  $R$  is a module-finite generically separable extension of a regular local ring  $A$ , then  $R[A^{1/p}] \cong R^{\oplus p^d}$ . The “canonicity” in our construction comes from discriminants, and we show that the discriminant of  $R/(f)$  does not “change” much when  $f$  changes slightly (Lemma 3.3). This allows us to show that one can control the convergence rate of Hilbert–Kunz function uniformly, independent of a small perturbation (Theorem 3.5).

## §2. Preliminary results

**DEFINITION 2.1.** Let  $A$  be a ring and  $R$  a finite  $A$ -algebra which is free as an  $A$ -module. Let  $e_1, \dots, e_n$  be a basis of  $R$  as an  $A$ -module.

The discriminant of  $R$  is

$$D_A(R) = \det \begin{pmatrix} \operatorname{Tr}(e_1^2) & \operatorname{Tr}(e_1 e_2) & \cdots & \operatorname{Tr}(e_1 e_n) \\ \operatorname{Tr}(e_2 e_1) & \operatorname{Tr}(e_2^2) & \cdots & \operatorname{Tr}(e_2 e_n) \\ \vdots & \vdots & \cdots & \vdots \\ \operatorname{Tr}(e_n e_1) & \operatorname{Tr}(e_n e_2) & \cdots & \operatorname{Tr}(e_n^2) \end{pmatrix},$$

where  $\operatorname{Tr}(r)$  denotes the trace of the multiplication map  $\times r$  on  $R$ . Up to multiplication by a unit of  $A$ , the discriminant is independent of the choice of basis.

If  $A$  is a regular local ring and  $R$  is a finite Cohen–Macaulay extension of  $A$ , then  $R$  is necessarily a free  $A$ -module by the Auslander–Buchsbaum formula, and thus the discriminant of  $R$  over  $A$  is well defined. We begin by recording some well-known and useful facts concerning discriminants.

**PROPOSITION 2.2.** *Let  $A$  be a ring and  $R$  a finitely generated  $A$ -algebra which is free as an  $A$ -module.*

- (1) *If  $A$  is a normal domain, then  $D_A(R) \in A$ .*
- (2) *If  $I$  is an ideal of  $A$  then the image of  $D_A(R)$  in  $R/IR$  is  $D_{A/I}(R/IR)$ .*
- (3) *If  $A$  is a domain then the ring  $R$  is generically separable over  $A$  if and only if  $D_A(R) \neq 0$ .*

*Proof.* We refer the reader to [Hoca, p. 200] for a proof of (1). For (2), if  $e_1, \dots, e_n \in R$  form a basis of  $R$  as an  $A$ -module then the images of  $e_1, \dots, e_n$  form a basis of  $R/IR$  as an  $A/IA$ -module. Moreover, if  $x \in R$  and  $\bar{x}$  denotes the image of  $x$  in  $R/IR$ , then  $\operatorname{Tr}(\bar{x}) = \overline{\operatorname{Tr}(x)}$ . Hence the image of  $D_A(R)$  in  $R/IR$  is indeed  $D_{A/I}(R/IR)$ .

For (3), as a first step, let  $L$  be the fraction field of  $A$ . We observe that  $D_A(R) = D_L(R \otimes_A L)$ , since the decomposition of an element  $a = \sum a_i e_i$  over  $A$  can be considered as a decomposition of  $a \otimes 1 = \sum a_i e_i \otimes 1$  of an element  $a \otimes 1$  over  $L$  and  $e_i \otimes 1$  still form a basis of  $R \otimes_A L$  over  $L$ . Thus we can replace  $R$  by  $R \otimes_A L$  and assume  $A = L$  is a field. It is well known that  $D_L(R) \neq 0$  if  $R$  is a finite separable field extension of  $L$ . By linearity this can be extended to the product and show one direction.

For the other direction, we first show that  $R$  is reduced. Since the nilradical is an ideal, it is a vector space over  $L$ , so we may extend a basis of it to a basis of  $R$  over  $L$ . If  $x$  is a nilpotent element, then  $\operatorname{Tr}(xe) = 0$  for any element  $e \in R$ , because the trace of a nilpotent matrix is nilpotent. This shows that  $\operatorname{Tr}$  gives a degenerate bilinear form and  $D_L(R) = 0$ . Since  $R$  is

reduced and finite over  $L$ , it is a direct sum of fields. Again, by linearity we can see that each of these fields must be separable over  $L$ .  $\square$

Let  $(R, \mathfrak{m}, k)$  be a complete local ring of positive characteristic  $p > 0$  with the residue field  $k$ . Cohen’s structure theorem shows that there is an injection  $\phi: k \rightarrow R$  such that the induced map to  $R/\mathfrak{m}$  is an isomorphism. We call  $\phi(k)$  a coefficient field. Moreover, Cohen’s structure theorem asserts that for any system of parameters  $x_1, \dots, x_d$  the  $k$ -subalgebra of  $R$  generated by the system of parameters,  $k[x_1, \dots, x_d]$ , is isomorphic to a polynomial ring. Since  $R$  is complete, it therefore contains the completion  $A = k[[x_1, \dots, x_d]]$  and, since  $x_1, \dots, x_d$  form a system of parameters,  $R$  is module-finite over  $A$ . Of course, there is a lot of freedom in choosing a system of parameters; the Cohen–Gabber structure theorem [GO08, Théorème 7.1]<sup>3</sup> asserts that  $x_1, \dots, x_d$  can be chosen so that  $R$  is generically separable over  $k[[x_1, \dots, x_d]]$ .

**THEOREM 2.3.** *Let  $(R, \mathfrak{m}, k)$  be a reduced complete local ring of equidimension  $d$  and of prime characteristic  $p$ . There exist a system of parameters  $x_1, \dots, x_d$  of  $R$  and coefficient field  $k \subseteq R$  such that  $R$  is finite and generically separable over the power series ring  $k[[x_1, \dots, x_d]]$ .*

It is useful to keep in mind that generic separability is equivalent to generic étaleness.

### §3. Main theorem and its applications

We say that a sequence  $f_1, \dots, f_c \in \mathfrak{m}$  is a parameter sequence if  $\dim R/(f_1, \dots, f_{i+1}) < \dim R/(f_1, \dots, f_i)$  for all  $i$ . We use  $\underline{f}$  to denote a sequence of elements  $f_1, \dots, f_c$ . If  $I$  is an ideal, we say  $\underline{f} \in (I)^{\oplus c}$  if each element is in  $I$ . We add sequences componentwise, so  $\underline{f} + \underline{g}$  is the sequence  $f_1 + g_1, \dots, f_c + g_c$ . We will use  $\lambda(M)$  to denote the length of a module, and  $\mu(M)$  to denote the minimal number of generators.

We borrow the following property observed in [ST96, Lemma 1].

**LEMMA 3.1.** *Let  $(R, \mathfrak{m}, k)$  be a local ring and  $\underline{f} \in (\mathfrak{m})^{\oplus c}$  be a parameter sequence. Then for any ideal  $I$  such that  $\lambda(R/(I, \underline{f}))$  is finite, there exists  $N > 0$  such that for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$*

$$(I, \underline{f}) = (I, \underline{f} + \underline{\epsilon}).$$

*In particular,  $\lambda(R/(I, \underline{f})) = \lambda(R/(I, \underline{f} + \underline{\epsilon}))$  for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ .*

<sup>3</sup>We refer the reader to [KS18] for an elementary proof of Theorem 2.3.

*Proof.* Take  $N$  such that  $\mathfrak{m}^N \subseteq (I, \underline{f})$ . Then  $(I, \underline{f}) = (I, \underline{f} + \underline{\epsilon})$  for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ . □

As observed in [ST96, Lemma 2], a small enough perturbation of a system of parameters remains a system of parameters.

**COROLLARY 3.2.** *Let  $(R, \mathfrak{m}, k)$  be a local ring. Suppose  $\underline{f} \in (\mathfrak{m})^{\oplus c}$  is a part of a system of parameters. Then there exists  $N \in \mathbb{N}$  such that for each  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ ,  $\underline{f} + \underline{\epsilon}$  is a part of a system of parameters.*

*Proof.* Let  $d$  be the dimension of the ring. We may complete  $\underline{f}$  to a full system of parameters  $\underline{f}, y_1, \dots, y_{d-c}$ . Setting  $I = (y_1, \dots, y_{d-c})$  in the previous lemma we obtain  $N$  such that for any  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$

$$(f_1, \dots, f_c, y_1, \dots, y_{d-c}) = (f_1 + \epsilon_1, \dots, f_c + \epsilon_c, y_1, \dots, y_{d-c}).$$

In particular  $f_1 + \epsilon_1, \dots, f_c + \epsilon_c, y_1, \dots, y_{d-c}$  are  $d$  elements that generate an  $\mathfrak{m}$ -primary ideal, so they form a system of parameters. □

**LEMMA 3.3.** *Let  $(R, \mathfrak{m}, k)$  be a complete Cohen–Macaulay local ring with coefficient field  $k \subseteq R$  and let  $f_1, \dots, f_c, x_1, \dots, x_{d-c}$  be a system of parameters for  $R$ . Let  $A = k[[T_1, \dots, T_{d-c}]]$  be the power series ring in  $d - c$  variables over the field  $k$ . There exists  $C \geq 0$  such that for all  $n \geq 1$  and all  $\underline{\epsilon} \in (\mathfrak{m}^{Cn})^{\oplus c}$ ,*

$$f_1 + \epsilon_1, \dots, f_c + \epsilon_c, x_1, \dots, x_{d-c}$$

*is a system of parameters and*

$$D_A(R/(\underline{f})) \equiv D_A(R/(\underline{f} + \underline{\epsilon})) \pmod{\mathfrak{m}_A^n},$$

*where  $R/(\underline{f})$  and  $R/(\underline{f} + \underline{\epsilon})$  are  $A$ -algebras by mapping  $T_i \mapsto x_i$  and identifying the coefficient field of  $A$  with the induced coefficient fields of  $R/(\underline{f})$  and  $R/(\underline{f} + \underline{\epsilon})$ .*

*Proof.* We can choose  $C$  as in Lemma 3.1, so for all  $\underline{\epsilon} \in (\mathfrak{m}^C)^{\oplus c}$  we have  $(\underline{f} + \underline{\epsilon}, x_1, \dots, x_{d-c}) = (\underline{f}, x_1, \dots, x_{d-c})$  and thus  $\underline{f} + \underline{\epsilon}, x_1, \dots, x_{d-c}$  is a system of parameters. Hence  $R/(\underline{f})$  and  $R/(\underline{f} + \underline{\epsilon})$  are free  $A$ -modules of the same rank  $\lambda_A(R/(\underline{f}, x_1, \dots, x_{d-c}))$ . Moreover, if  $r_1, \dots, r_n \in R$  are chosen so that their images in  $R/(\underline{f})$  serve as a basis for  $R/(\underline{f})$  as an  $A$ -module, then for each  $\underline{\epsilon} \in (\mathfrak{m}^C)^{\oplus c}$ , because the images of  $r_1, \dots, r_n$  form a basis of  $R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}_A) = R/(\underline{f}, x_1, \dots, x_{d-c})$ , the images of  $r_1, \dots, r_n$  in  $R/(\underline{f} + \underline{\epsilon})$  are a basis of  $R/(\underline{f} + \underline{\epsilon})$  too.

Observe that for each  $n \geq 1$ ,  $\mathfrak{m}^{Cn} \subseteq (x_1, \dots, x_{d-c})^n + (f)$ . Thus for each  $\epsilon \in (\mathfrak{m}^{Cn})^{\oplus c}$ ,  $R/(\underline{f}, \mathfrak{m}_A^n) = R/(\underline{f} + \epsilon, \mathfrak{m}_A^n)$  as  $A/\mathfrak{m}_A^n$ -algebras. Then by part (2) of Lemma 2.2

$$\begin{aligned} D_A(R/(\underline{f})) \pmod{\mathfrak{m}_A^n R/(\underline{f})} &= D_{A/\mathfrak{m}_A^n}(R/(\underline{f}, \mathfrak{m}_A^n)) \\ &= D_{A/\mathfrak{m}_A^n}(R/(\underline{f} + \epsilon, \mathfrak{m}_A^n)) \\ &= D_A(R/(\underline{f} + \epsilon)) \pmod{\mathfrak{m}_A^n R/(\underline{f} + \epsilon)}, \end{aligned}$$

where the middle equality holds since we have a common basis  $r_1, \dots, r_n \in R$  for  $R/(\underline{f} + \epsilon)$  and  $R/(\underline{f})$  as  $A$ -modules. □

Suppose  $(R, \mathfrak{m}, k)$  satisfies the hypotheses of Lemma 3.3. The following corollary shows that if  $f \in \mathfrak{m}$  is a parameter element and  $R/(f)$  is generically separable over a regular ring  $A$ , then  $R/(g)$  is generically separable over the same regular local ring  $A$ , provided  $g$  is a small enough perturbation of  $f$ .

**COROLLARY 3.4.** *Let  $(R, \mathfrak{m}, k)$  be a complete Cohen–Macaulay local ring with coefficient field  $k \subseteq R$ . Let  $f_1, \dots, f_c, x_1, \dots, x_{d-c}$  be a system of parameters for  $R$ . Let  $A = k[[T_1, \dots, T_{d-c}]]$  be the power series ring in  $d-c$  variables over the field  $k$  and consider  $R/(\underline{f})$  is an  $A$ -algebra by mapping  $T_i \mapsto x_i$  and identifying the coefficient field of  $A$  with the induced coefficient field of  $R/(\underline{f})$ . Suppose that  $A \subseteq R/(\underline{f})$  is generically separable. Then there exists  $N \geq 1$  such that for any  $\epsilon \in (\mathfrak{m}^N)^{\oplus c}$ ,  $A \subseteq R/(\underline{f} + \epsilon)$  is generically separable as an  $A$ -algebra given by mapping  $T_i \mapsto x_i$  and identifying the coefficient field of  $A$  with the induced coefficient field of  $R/(\underline{f} + \epsilon)$ .*

*Proof.* The assumption  $R/(\underline{f})$  is generically separable over  $A$  is equivalent to  $D_A(R/(\underline{f})) \neq 0$  by Part (1) of Lemma 2.2. Let  $C \geq 0$  as given by Lemma 3.3. By Krull’s intersection theorem we can choose  $n \geq 1$  so that  $D_A(R/(\underline{f})) \not\equiv 0 \pmod{\mathfrak{m}_A^n}$ . Then for any  $\epsilon \in (\mathfrak{m}^{Cn})^{\oplus c}$ ,  $D_A(R/(\underline{f} + \epsilon)) \equiv D_A(R/(\underline{f})) \pmod{\mathfrak{m}_A^n} \neq 0$ , so  $D_A(R/(\underline{f} + \epsilon)) \neq 0$ . □

### 3.1 Continuity of Hilbert–Kunz multiplicity

It is time to set up the uniform convergence machinery. We closely follow [PT18, Lemma 2.3, Theorem 3.2]. Let  $(R, \mathfrak{m}, k)$  be an F-finite local ring and  $I$  be an  $\mathfrak{m}$ -primary ideal. Then  $F_*^e R \otimes_R R/I \cong F_*^e R/I^{[p^e]}$  and we may compute

$$\lambda_R(F_* R \otimes_R R/I) = [k : k^p] \lambda_R(R/I^{[p]}).$$



**THEOREM 3.5.** *Let  $(R, \mathfrak{m}, k)$  be an  $F$ -finite Cohen–Macaulay local ring of prime characteristic  $p$  and of dimension  $d + c$ . If  $\underline{f} = f_1, \dots, f_c$  is a parameter sequence such that  $\hat{R}/(\underline{f})\hat{R}$  is reduced, then there exists a constant  $C \geq 0$  and an integer  $N \geq 1$  such that for any  $e \geq 1$  and  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$*

$$|\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^{e+1}]}) - p^d \lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]}))| \leq Cp^{e(d-1)}.$$

*Proof.* Without loss of generality, we may assume  $R$  is complete. Because  $R$  is Cohen–Macaulay,  $R/(\underline{f})$  is equidimensional and reduced. By Theorem 2.3 there exists a coefficient field  $k \subseteq R/(\underline{f})$  and parameters  $x_1, \dots, x_d \in R$  such that  $x_1, \dots, x_d$  is a system of parameters for  $R/(\underline{f})$  and  $R/(\underline{f})$  is module-finite and generically separable over the regular local ring  $A := k[[x_1, \dots, x_d]]$ .

We observe that the coefficient field can be lifted from  $R/(\underline{f})$  to  $R$ . Essentially, the proof of [Mat86, Theorem 28.3] shows that a coefficient field is determined by the lifts of a  $p$ -basis of  $k$  (see also [Hocb, Theorem, p. 12]). Thus we may just lift the  $p$ -basis to  $R$ .

By Lemma 6.5 of [HH90] (the proof still holds if we replace  $R^\infty$  by  $R^{1/p}$ ), we have an exact sequence

$$0 \rightarrow R/(\underline{f})[A^{1/p}] \rightarrow (R/(\underline{f}))^{1/p} \rightarrow M \rightarrow 0$$

where  $0 \neq c = D_A(R/(\underline{f}))$  annihilates  $M$ .

By Lemma 3.3 and Corollary 3.4, we can find  $N$  such that  $A \subseteq R/(\underline{f} + \underline{\epsilon})$  is still generically separable. Therefore we have an exact sequence

$$0 \rightarrow R/(\underline{f} + \underline{\epsilon})[A^{1/p}] \rightarrow (R/(\underline{f} + \underline{\epsilon}))^{1/p} \rightarrow M_\epsilon \rightarrow 0$$

where  $c_\epsilon = D_A(R/(\underline{f} + \underline{\epsilon}))$  annihilates  $M_\epsilon$  and  $c_\epsilon - c \in \mathfrak{m}_A^N$ .

By [Kum69]  $A^{1/p}$  is a free  $A$ -module of rank  $[k : k^p]p^d$ , so tensoring the above sequence with  $\mathfrak{m}^{[p^e]}$  we get that

$$\begin{aligned} & [k : k^p] \lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^{e+1}]}) \\ & - [k : k^p] p^d \lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]})) \leq \lambda(M_\epsilon / \mathfrak{m}^{[p^e]} M_\epsilon). \end{aligned}$$

Since  $M_\epsilon$  is a  $R/(\underline{f} + \underline{\epsilon}, c_\epsilon)$ -module and a quotient of  $(R/(\underline{f} + \underline{\epsilon}))^{1/p}$ ,

$$\lambda(M_\epsilon / \mathfrak{m}^{[p^e]} M_\epsilon) \leq \lambda(R/(\underline{f} + \underline{\epsilon}, c_\epsilon, \mathfrak{m}^{[p^e]})) \mu((R/(\underline{f} + \underline{\epsilon}))^{1/p}).$$

Furthermore, it easily follows from the inclusion  $(x_1, \dots, x_d) \subseteq \mathfrak{m}$  that

$$\mu((R/(\underline{f} + \underline{\epsilon}))^{1/p}) \leq \lambda((R/(x_1^p, \dots, x_d^p, \underline{f} + \underline{\epsilon}))^{1/p}),$$

thus

$$\begin{aligned} \mu((R/(\underline{f} + \underline{\epsilon}))^{1/p}) &\leq [k : k^p]\lambda(R/(x_1^p, \dots, x_d^p, \underline{f} + \underline{\epsilon})) \\ &\leq [k : k^p]p^d\lambda(R/(x_1, \dots, x_d, \underline{f})), \end{aligned}$$

where the last inequality follows by filtration.

Now, we can complete  $0 \neq c$  to a system of parameters  $c, t_1, \dots, t_{d-1}$  in  $A$  and, after increasing  $N$  if necessary, by Lemma 3.3 we can assume that  $c_{\underline{\epsilon}}, t_1, \dots, t_{d-1}$  is still a system of parameters and that  $(\underline{f} + \underline{\epsilon}, c_{\underline{\epsilon}}, t_1, \dots, t_{d-1}) = (\underline{f}, c, t_1, \dots, t_{d-1})$ . Therefore

$$\begin{aligned} \lambda(R/(\underline{f} + \underline{\epsilon}, c_{\underline{\epsilon}}, \mathfrak{m}^{[p^e]})) &\leq \lambda(R/(\underline{f} + \underline{\epsilon}, c_{\underline{\epsilon}}, t_1^{p^e}, \dots, t_{d-1}^{p^e})) \\ &\leq p^{e(d-1)}\lambda(R/(\underline{f}, c, t_1, \dots, t_{d-1})). \end{aligned}$$

Let

$$C = p^d\lambda(R/(\underline{f}, c, t_1, \dots, t_{d-1}))\lambda(R/(x_1, \dots, x_d, \underline{f}))$$

and observe that we found one of the two required inequalities:

$$\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^{e+1}]}) - p^d\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]})) \leq Cp^{e(d-1)}.$$

For the other inequality, we can repeat the proof for the exact sequence

$$0 \rightarrow (R/(\underline{f}))^{1/p} \xrightarrow{c} R/(\underline{f})[A^{1/p}] \rightarrow L \rightarrow 0$$

where the map is multiplication by  $c = D_A(R/(\underline{f}))$ . Again the cokernel is annihilated by  $c$ , because  $R/(\underline{f})[A^{1/p}] \subseteq (R/(\underline{f}))^{1/p}$ . Moreover, we get for each  $\underline{\epsilon}$  a short exact sequence

$$0 \rightarrow (R/(\underline{f} + \underline{\epsilon}))^{1/p} \xrightarrow{c_{\underline{\epsilon}}} R/(\underline{f} + \underline{\epsilon})[A^{1/p}] \rightarrow L_{\underline{\epsilon}} \rightarrow 0.$$

Most of the proof carries through as above, except that now we estimate

$$\begin{aligned} \lambda(L_{\underline{\epsilon}}/\mathfrak{m}^{[p^e]}L_{\underline{\epsilon}}) &\leq \mu(R/(\underline{f})[A^{1/p}])\lambda(R/(\underline{f} + \underline{\epsilon}, c_{\underline{\epsilon}}, \mathfrak{m}^{[p^e]})) \\ &= [k : k^p]p^d\lambda(R/(\underline{f} + \underline{\epsilon}, c_{\underline{\epsilon}}, \mathfrak{m}^{[p^e]})) \end{aligned}$$

and from there obtain

$$\begin{aligned} p^d\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]})) - \lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^{e+1}]}) & \\ \leq (p^d[k : k^p]\lambda(R/(\underline{f}, c, t_1, \dots, t_{d-1})))p^{e(d-1)}. & \quad \square \end{aligned}$$

Following the treatment in [Tuc12], we obtain the following corollary:

COROLLARY 3.6. *Let  $(R, \mathfrak{m}, k)$  be an  $F$ -finite Cohen–Macaulay local ring of prime characteristic  $p$  and of dimension  $d + c$ . If  $\underline{f}$  is a parameter sequence of length  $c$  and such that  $\hat{R}/(\underline{f})\hat{R}$  is reduced, then there exist a constant  $C \geq 0$  and an integer  $N \geq 1$  such that for any  $e \geq 1$  and  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$*

$$|e_{\text{HK}}(R/(\underline{f} + \underline{\epsilon})) - p^{-ed}\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]}))| \leq Cp^{-e}.$$

*Proof.* Using induction on  $q'$  as in the proof of [Tuc12, Proposition 3.4] we may obtain from Theorem 3.5 that

$$\begin{aligned} & |\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^{e+e'}]})) - p^{e'd}\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]}))| \\ & \leq Cp^{(e+e'-1)(d-1)}(1 + p + \dots + p^{e'-1}). \end{aligned}$$

Thus, dividing by  $p^{(e+e')d}$  we get that

$$\begin{aligned} & |p^{-(e+e')d}\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^{e+e'}]})) - p^{-ed}\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]}))| \\ & \leq C \frac{(1 + p + \dots + p^{e'-1})}{p^{d-1}p^{e+e'}}. \end{aligned}$$

Note that

$$C \frac{(1 + p + \dots + p^{e'-1})}{p^{d-1}p^{e+e'}} = \frac{C(p^{e'} - 1)}{(p - 1)p^{d-1}p^{e'}}p^{-e} \leq Cp^{-e},$$

so the claim follows after letting  $e' \rightarrow \infty$ . □

Continuity of Hilbert–Kunz multiplicity follows from Corollary 3.6.

COROLLARY 3.7. *Let  $(R, \mathfrak{m}, k)$  be a Cohen–Macaulay  $F$ -finite local ring of prime characteristic  $p$  and dimension  $d + c$ . If  $\underline{f}$  is a parameter sequence of length  $c$  and such that  $\hat{R}/(\underline{f})\hat{R}$  is reduced, then for any  $\delta > 0$  there exists an integer  $N > 0$  such that for any  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$*

$$|e_{\text{HK}}(R/(\underline{f})) - e_{\text{HK}}(R/(\underline{f} + \underline{\epsilon}))| < \delta.$$

*Proof.* By Lemma 3.1, for any given  $e$  there exists  $N$  such that

$$\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]})) = \lambda(R/(\underline{f}, \mathfrak{m}^{[p^e]}))$$

for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ . By Corollary 3.6, we may further assume that for some constant  $C > 0$

$$\left| e_{\text{HK}}(R/(\underline{f} + \underline{\epsilon})) - \frac{1}{p^{ed}}\lambda(R/(\underline{f} + \underline{\epsilon}, \mathfrak{m}^{[p^e]})) \right| < Cp^{-e}.$$

Thus

$$|e_{\text{HK}}(R/(\underline{f} + \underline{\epsilon})) - e_{\text{HK}}(R/(\underline{f}))| < 2Cp^{-e}$$

and the claim follows. □

### 3.2 Continuity of F-signature

To establish continuity of F-signature we will need an effective way of comparing the  $e$ th Frobenius splitting numbers of rings which are the homomorphic image of a common ring. We begin by recalling an effective method which allows us to measure Frobenius splitting numbers in rings which are approximately Gorenstein [Hoc77].

LEMMA 3.8. *Let  $(R, \mathfrak{m}, k)$  be an  $F$ -finite approximately Gorenstein ring, that is there exists a descending chain of irreducible  $\mathfrak{m}$ -primary ideals  $\{I_t\}$  cofinal with  $\{\mathfrak{m}^t\}$ . Let  $u_t$  generate the socle modulo  $I_t$ .*

- (1) *Then for each  $e$  there exists  $t_e$  such that  $a_e(R) = [k : k^{p^e}] \lambda(R/(I_t^{[p^e]} :_R u_t^{p^e}))$  for all  $t \geq t_e$ .*
- (2) *If  $R$  is Cohen–Macaulay and  $J \subseteq R$  is an ideal of  $R$  isomorphic to the canonical module of  $R$  and  $x_1 \in J$  is not a zero divisor, then  $I_t$  can be taken to be  $(x_1^{t-1}J, x_2^t, \dots, x_d^t)$  where  $x_2, \dots, x_d$  is a system of parameters for  $R/J$ . Moreover,  $u$  generates the socle mod  $(J, x_2, \dots, x_d)$  then  $u_t := (x_1 \cdots x_d)^{t-1}$  generates the socle mod  $I_t$ .*
- (3) *Under the assumptions and notation of (2), there exists  $t_e$  such that*

$$a_e(R) = [k : k^{p^e}] \lambda(R/(J, x_2^t, \dots, x_d^t)^{[p^e]} :_R (x_2 \cdots x_d)^{tp^e})$$

*for all  $t \geq t_e$ .*

We refer the reader to [PT18, Lemma 6.2] and the proof of [PT18, Corollary 6.6] for details.

LEMMA 3.9. *Let  $(R, \mathfrak{m}, k)$  be a complete Cohen–Macaulay domain of dimension at least  $c + 1$ , not necessarily of prime characteristic. Let  $\underline{f}$  be a regular sequence of length  $c$  and such that  $R/(\underline{f})$  is generically Gorenstein. Then there exists a canonical ideal  $J \subseteq R$  such that the ideal  $J + (\underline{f})$  of  $R/(\underline{f})$  is the canonical module of  $R/(\underline{f})$ . Moreover, there exists  $N \in \mathbb{N}$  such that for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ , the ideal  $(\underline{f} + \underline{\epsilon}, J)/(\underline{f})$  of  $R/(\underline{f} + \underline{\epsilon})$  is the canonical module of  $R/(\underline{f} + \underline{\epsilon})$ .*

*Proof.* Recall that  $R$  has a canonical ideal  $\omega$  if and only if it is generically Gorenstein [BH93, Proposition 3.3.18]. We also recall that  $R$  is generically Gorenstein if and only if  $R_{P_i}$  is Gorenstein for any minimal prime  $P_i$  of  $R$ .

Let  $S = R \setminus \cup P_i$  where  $P_i$  are the minimal primes of  $(\underline{f})$ . Then  $S^{-1}\omega \subseteq S^{-1}R$  is a canonical module of  $S^{-1}R$  and, since  $S^{-1}R$  is Gorenstein,  $S^{-1}\omega = (u)$  is a principal ideal, and we may assume that  $u \in \omega$ . This means that there exists  $s \in S$  such that for some ideal  $J$  in  $R$ ,

$$s\omega = uJ \subseteq uR.$$

Since  $R$  is a domain,  $\omega \cong J$ . Furthermore, we claim that  $J \not\subseteq \cup P_i$ . Namely, if  $J \subseteq P_i$  for some  $i$ , then  $s\omega = uJ \subseteq uP_i \subseteq \omega P_i$ . But then localizing at  $P_i$  we obtain that  $\omega R_{P_i} \subseteq P_i \omega R_{P_i}$ , which contradicts Nakayama’s lemma.

We verify that  $J$  satisfies the assertions of the theorem. Since  $J$  is not contained in any minimal prime of  $(\underline{f})$ ,  $\text{ht}(J, \underline{f}) \geq c + 1$ . Moreover,  $J$  has height 1 and  $R$  is Cohen–Macaulay, so  $\text{ht}(\underline{f}, J)R/J \geq c$  and  $\underline{f}$  must be a parameter sequence in  $R/J$ . Since  $R/J$  is Gorenstein [BH93, Proposition 3.3.18], the images of  $\underline{f}$  form a regular sequence. Hence

$$\frac{J + (\underline{f})}{(\underline{f})} = \frac{J}{(\underline{f}) \cap J} = \frac{J}{(\underline{f})J} = JR/(\underline{f})$$

is a canonical module of  $R/(\underline{f})$ . By Corollary 3.2 we can choose a neighborhood  $\mathfrak{m}^N$  such that the images of  $\underline{f} + \underline{\epsilon}$  are still a part of a system of parameters in the Gorenstein ring  $R/J$  for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$  and the statement follows. □

**COROLLARY 3.10.** *Let  $(R, \mathfrak{m}, k)$  be a complete Cohen–Macaulay  $F$ -finite domain of prime characteristic  $p$ . Let  $\underline{f}$  be a regular sequence of length  $c$  such that  $R/(\underline{f})$  is generically Gorenstein. Then for any integer  $e$  there exists an integer  $N$  such that  $a_e(R/(\underline{f})) = a_e(R/(\underline{f} + \underline{\epsilon}))$  for any  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ .*

*In particular, if  $R$  is also normal, then the functions  $a_e: R \rightarrow \mathbb{R}$ ,  $f \mapsto a_e(R/f)$ , are locally constant at parameter elements with respect to the  $\mathfrak{m}$ -adic topology.*

*Proof.* If  $\dim(R) = c$ , then  $R/(\underline{f})$  is Artinian and, thus, has a splitting if and only if it is a field. If we take  $\underline{\epsilon} \in (\mathfrak{m}^2)^{\oplus c}$ , then  $(\underline{f}) = \mathfrak{m}$  if and only if  $(\underline{f} + \underline{\epsilon}) = \mathfrak{m}$ . So the Frobenius splitting numbers of  $R/(\underline{f})$  and  $R/(\underline{f} + \underline{\epsilon})$  will be the same, either 0 or  $[k : k^{p^e}]$ , the latter of which occurs if and only if  $(\underline{f}) = \mathfrak{m}$ .

Suppose  $\dim(R) = d + c > c$  and choose  $J$  and  $N$  as in Lemma 3.9. If  $J = R$  replace  $J$  by a principal ideal  $(g)$  such that  $\underline{f}, g$  is a regular sequence. Let  $x_2, \dots, x_d$  be a system of parameters for  $R/(\underline{f}, J)$ . If necessary, increase

$N$  so that  $\mathfrak{m}^N \subseteq \mathfrak{m}^{[p^e]}$  and  $\mathfrak{m}^N \subseteq (\underline{f}, J, x_2, \dots, x_d)$ . In particular, we have for each  $\epsilon \in \mathfrak{m}^N$  that  $x_2, \dots, x_d$  is a system of parameters for  $R/(\underline{f} + \epsilon, J)$ .

For each  $\epsilon \in (\mathfrak{m}^N)^{\oplus c}$ ,  $(\underline{f} + \epsilon, J)/(\underline{f})$  is the canonical module of  $R/(\underline{f} + \epsilon)$  by Lemma 3.9; hence

$$J + (\underline{f} + \epsilon, x_2, \dots, x_d) = J + (\underline{f}, x_2, \dots, x_d)$$

is an irreducible  $\mathfrak{m}$ -primary ideal [BH93, Proposition 3.3.18]. Let  $u \in R$  generate the socle mod  $(\underline{f}, J, x_2, \dots, x_d) = (\underline{f} + \epsilon, J, x_2, \dots, x_d)$ . For each  $t \in \mathbb{N}$  let  $I_t = (J, x_2^t, \dots, x_d^t)$  and  $u_t = (x_2 \cdots x_d)^{t-1}u$ . By Lemma 3.8 there is for each  $\epsilon \in (\mathfrak{m}^N)^{\oplus c}$  a  $t_\epsilon$  such that

$$a_e(R/(\underline{f} + \epsilon)) = [k : k^{p^e}] \lambda \left( \frac{R}{(\underline{f} + \epsilon, (I_{t_\epsilon}^{[p^e]} :_R u_{t_\epsilon}^{p^e}))} \right)$$

and

$$a_e(R/(\underline{f})) = [k : k^{p^e}] \lambda \left( \frac{R}{(\underline{f}, (I_{t_\epsilon}^{[p^e]} :_R u_{t_\epsilon}^{p^e}))} \right).$$

Moreover, we have  $\mathfrak{m}^{[p^e]} \subseteq (\underline{f} + \epsilon, (I_{t_\epsilon}^{[p^e]} :_R u_{t_\epsilon}^{p^e}))$ . Since  $N$  was chosen so that  $\mathfrak{m}^N \subseteq \mathfrak{m}^{[p^e]}$ , we have

$$(\underline{f}, (I_{t_\epsilon}^{[p^e]} :_R u_{t_\epsilon}^{p^e})) = (\underline{f} + \epsilon, (I_{t_\epsilon}^{[p^e]} :_R u_{t_\epsilon}^{p^e})).$$

Therefore for each  $\epsilon \in (\mathfrak{m}^N)^{\oplus c}$ ,  $a_e(R/(\underline{f})) = a_e(R/(\underline{f} + \epsilon))$ . □

We need less assumptions for an inequality.

LEMMA 3.11. *Let  $(R, \mathfrak{m}, k)$  be a complete  $F$ -finite ring of positive characteristic  $p > 0$ . Then for any parameter sequence  $\underline{f}$  of length  $c$  and a fixed positive integer  $e$  there exists integer  $N$  such that for all  $\epsilon \in (\mathfrak{m}^N)^{\oplus c}$*

$$a_e(R/(\underline{f} + \epsilon)) \leq a_e(R/(\underline{f})).$$

*Proof.* Since  $R$  is complete, we can represent it as a quotient of a regular local ring  $(S, \mathfrak{m})$ ,  $R = S/I$ . Then by [EY11, Proposition 3.1]

$$a_e(R/(\underline{f})) = [k : k^{p^e}] \lambda_S \left( \frac{(I, \underline{f})^{[p^e]} :_S (I, \underline{f}) + \mathfrak{m}^{[p^e]}}{\mathfrak{m}^{[p^e]}} \right).$$

Let  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$  for an arbitrary  $N$ . Then

$$(I, \underline{f} + \underline{\epsilon})^{[p^e]} :_S (I, \underline{f} + \underline{\epsilon}) \subseteq ((I, \underline{f})^{[p^e]} + \mathfrak{m}^N) :_S (I, \underline{f} + \underline{\epsilon}).$$

Since  $\epsilon_i$  are in  $\mathfrak{m}^N$ ,

$$((I, \underline{f})^{[p^e]} + \mathfrak{m}^N) :_S (I, \underline{f} + \underline{\epsilon}) = ((I, \underline{f})^{[p^e]} + \mathfrak{m}^N) :_S (I, \underline{f}).$$

By Krull’s intersection theorem

$$\bigcap_{n=1}^{\infty} ((I, \underline{f})^{[p^e]} + \mathfrak{m}^n) :_S (I, \underline{f}) = ((I, \underline{f})^{[p^e]} :_S (I, \underline{f})),$$

so by Chevalley’s lemma we could have chosen  $N$  so that

$$((I, \underline{f})^{[p^e]} + \mathfrak{m}^N) :_S (I, \underline{f}) \subseteq ((I, \underline{f})^{[p^e]} :_S (I, \underline{f}) + \mathfrak{m}^{[p^e]}).$$

In which case  $a_e(R/(\underline{f} + \underline{\epsilon})) \leq a_e(R/(\underline{f}))$ . □

**THEOREM 3.12.** *Let  $(R, \mathfrak{m}, k)$  be a Cohen–Macaulay  $F$ -finite ring of prime characteristic  $p$  and dimension  $d + c$ . If  $\underline{f}$  is a parameter sequence of length  $c$  such that  $\hat{R}/(\underline{f})\hat{R}$  is reduced then for any  $\delta > 0$  there exists an integer  $N > 0$  such that for any  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$*

$$|s(R/(\underline{f})) - s(R/(\underline{f} + \underline{\epsilon}))| < \delta.$$

*Proof.* We may assume  $R$  is complete. By Theorem 2.3 we may choose parameters  $x_1, \dots, x_d \in R$  such that  $x_1, \dots, x_d$  is a system of parameters for  $R/(\underline{f})$  and  $R/(\underline{f})$  is module-finite and generically separable over the regular local ring  $A := k[[x_1, \dots, x_d]]$ . In which case we have a short exact sequence

$$0 \rightarrow R/(\underline{f})[A^{1/p}] \rightarrow (R/(\underline{f}))^{1/p} \rightarrow M \rightarrow 0$$

and  $0 \neq c = D_A(R/(\underline{f}))$  annihilates  $M$ .

Let  $M$  and  $M_{\underline{\epsilon}}$  be as in the proof of Theorem 3.5. In which case, there are isomorphisms  $R/(\underline{f} + \underline{\epsilon})[A^{1/p}] \cong \bigoplus^{p^d [k:k^p]} (R/(\underline{f} + \underline{\epsilon}))$  and short exact sequences

$$0 \rightarrow R/(\underline{f} + \underline{\epsilon})[A^{1/p}] \rightarrow (R/(\underline{f} + \underline{\epsilon}))^{1/p} \rightarrow M_{\underline{\epsilon}} \rightarrow 0.$$

Apply the exact functor  $(-)^{1/p^e}$  to the above to get the exact sequence

$$0 \rightarrow \bigoplus^{p^d [k:k^p]} ((R/(\underline{f} + \underline{\epsilon}))^{1/p^e}) \rightarrow (R/(\underline{f} + \underline{\epsilon}))^{1/p^{e+1}} \rightarrow M_{\underline{\epsilon}}^{1/p^e} \rightarrow 0.$$

By [PT18, Lemma 2.1]

$$\text{frk}((R/(\underline{f} + \underline{\epsilon}))^{1/p^{e+1}}) \leq p^d[k : k^p] \text{frk}((R/(\underline{f} + \underline{\epsilon}))^{1/p^e}) + \mu((M_{\underline{\epsilon}})^{1/p^e}),$$

or, equivalently,

$$a_{e+1}(R/(\underline{f} + \underline{\epsilon})) \leq p^d[k : k^p] a_e(R/(\underline{f} + \underline{\epsilon})) + \mu((M_{\underline{\epsilon}})^{1/p^e}).$$

Techniques used in the proof of Theorem 3.5 give the existence of a constant  $C$  such that for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$

$$\mu((M_{\underline{\epsilon}})^{1/p^e}) = \lambda(M_{\underline{\epsilon}}/\mathfrak{m}^{[p^e]}M_{\underline{\epsilon}}) \leq Cp^{e(d-1)}.$$

Let  $L$  and  $L_{\underline{\epsilon}}$  be as in the proof of Theorem 3.5. Similarly, we can also bound

$$p^d[k : k^p] \text{frk}((R/(\underline{f} + \underline{\epsilon}))^{1/p^e}) - \text{frk}((R/(\underline{f} + \underline{\epsilon}))^{1/p^{e+1}}) \leq \lambda(L_{\underline{\epsilon}}/\mathfrak{m}^{[p^e]}L_{\underline{\epsilon}})$$

and can once again obtain constant  $C$  independent of  $\underline{\epsilon}$  so that

$$|a_{e+1}(R/(\underline{f} + \underline{\epsilon})) - p^d[k : k^p] a_e(R/(\underline{f} + \underline{\epsilon}))| < Cp^{e(d-1)}.$$

Since  $\text{rank } R^{1/p^e} = p^{ed}[k : k^{p^e}]$  [Kun76, Proposition 2.3], as explained in Corollary 3.7, this gives us a uniform convergence statement: there exist  $D, N > 0$  such that for all  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$

$$\left| s(R/(\underline{f} + \underline{\epsilon})) - \frac{1}{\text{rank } R^{1/p^e}} a_e(R/(\underline{f} + \underline{\epsilon})) \right| < \frac{D}{p^e}.$$

Now, if  $s(R/(\underline{f})) = 0$ , for any given  $\delta > 0$  we can find  $e$  such that  $a_e(R/(\underline{f})) < \delta/2$  and  $D/p^e < \delta/2$ . By Lemma 3.11, we can find  $N$  such that for any  $\underline{\epsilon} \in (\mathfrak{m}^N)^{\oplus c}$ ,

$$a_e(R/(\underline{f} + \underline{\epsilon})) \leq a_e(R/(\underline{f})) < \delta/2.$$

Thus, in this case the claim follows from the bound

$$s(R/(\underline{f} + \underline{\epsilon})) \leq \frac{1}{\text{rank } R^{1/p^e}} a_e(R/(\underline{f} + \underline{\epsilon})) + \frac{D}{p^e} < \delta.$$

If  $s(R/(\underline{f})) > 0$ , then  $R/(\underline{f})$  is strongly  $F$ -regular and, in particular, is a domain [AL03, main result]. Thus  $(\underline{f})$  is a prime ideal, so  $R/(\underline{f})$  is then generically Gorenstein. The statement now follows by employing Corollary 3.10 and using the methods of the proof of Corollary 3.7.  $\square$



**§4. Examples and questions**

In this section we want to present examples relevant to the results in Section 3. The first example shows that we cannot expect Hilbert–Kunz multiplicity to be locally constant at reduced parameter elements.

EXAMPLE 4.1. Consider an element  $f = xy$  in  $R = k[[x, y, t]]$ . An easy computation shows that  $e_{\text{HK}}(R/(xy)) = 2$ , while we will show that  $e_{\text{HK}}(R/(xy + t^n)) = 2 - 2/n$  for  $n \geq 2$ . Thus:

- (1) there is no neighborhood of  $xy$  such that  $e_{\text{HK}}$  is constant;
- (2) of course,  $e(R/(xy + t^n)) = 2 = e(R/(xy))$  for all  $n \geq 2$ ;
- (3) moreover, we have  $\text{Gr}_{(x,y,t)}(R/(xy + t^n)) = k[x, y, t]/(xy) = \text{Gr}_{(x,y,t)}(R/(xy))$  for all  $n \geq 3$ .

In order to compute the Hilbert–Kunz multiplicity we will compute the length of

$$k[[x, y, t]]/(xy - t^n, x^{p^e}, y^{p^e})$$

by counting monomials. It is easy to see that a basis is given by monomials of the form  $x^a t^n, y^a t^b$  where  $a < p^e - \lfloor b/n \rfloor$  and  $b < p^e$ . Thus when  $e \rightarrow \infty$  the number of such monomials is

$$2 \sum_{b=0}^{p^e-1} (p^e - \lfloor b/n \rfloor) \approx 2p^{2e} - 2 \frac{p^e(p^e - 1)}{n} \approx \left(2 - \frac{2}{n}\right) p^{2e}.$$

EXAMPLE 4.2. Essentially the same example works for F-signature. Clearly,  $k[[x, y, t]]/(x^2 + y^2)$  is not normal, so the F-signature is 0. On the other hand,  $k[[x, y, t]]/(x^2 + y^2 + t^n)$  is the  $A_{n-1}$  singularity, its F-signature is known to be  $1/n$  [HL02, Example 18].

By comparison, it is not that hard to see that Hilbert–Samuel multiplicity is locally constant at parameter elements in regular local ring. In fact, Srinivas and Trivedi [ST96, Corollary 5] showed that the entire Hilbert function is locally constant.

**4.1 Cohen–Macaulayness**

The Cohen–Macaulay assumption of our main results may not be completely necessary. For a possible extension, one could recall that Srinivas and Trivedi [ST96, Corollary 5] show continuity of the Hilbert–Samuel multiplicity if  $R$  is generalized Cohen–Macaulay, that is,  $\lambda(H_m^i(R)) < \infty$  for all  $i < \dim R$ .

We also want to give two examples that show that one cannot relax Cohen–Macaulayness too much. The first example comes from [ST96, Example 1].

EXAMPLE 4.3. Let  $R$  be the localization at  $\mathfrak{m} = (x, y, z)$  of  $k[x, y, z]/(xy, xz)$ . Note that  $R$  is not equidimensional. Because  $\dim R/(y) = 1$ , the two multiplicities coincide and we easily see that

$$e_{\text{HK}}(R/(y)) = e(R/(y)) = e(k[x, z]_{\mathfrak{m}}/(xz)) = 2.$$

On the other hand,  $R/(y + x^N) = k[x, z]_{\mathfrak{m}}/(x^{N+1}, xz)$  has the only minimal prime  $(x)$  for all  $N$ . Hence, by the associativity formula,

$$e_{\text{HK}}(R/(y + x^N)) = e(R/(y + x^N)) = e(R/(x))\lambda(R_{(x)}) = 1.$$

PROPOSITION 4.4. Let  $R$  be the subring  $k[[x^3, x^2y, y^3, y^2z, z^3, z^2x]] \subseteq k[[x, y]]$ . Observe that  $x^3, y^3, z^3$  form a system of parameters and consider a family of ideals  $J_k = (x^3, y^3 + z^{3k})$ . Then:

- (1)  $e_{\text{HK}}(R/(x^3, y^3)) = e(R/(x^3, y^3)) = 11$ ;
- (2)  $e_{\text{HK}}(R/J_k) = e(R/J_k) \neq 11$  for all  $k \not\equiv 1 \pmod{3}$ .

*Proof.* First, we observe that  $\dim R/(x^3, y^3) = \dim R/J_k = 1$ , so multiplicity equals to Hilbert–Kunz multiplicity.

We compute  $e(R/(x^3, y^3))$  by using the associativity formula. Observe that

$$P = (x^3, x^2y, y^3, y^2z, z^2x) = \sqrt{(x^3, y^3)}$$

is prime. Thus the associativity formula gives that  $e_{\text{HK}}(R/(x^3, y^3)) = e_{\text{HK}}(R/P)\lambda(R_P/(x^3, y^3))$ .

The quotient  $R/P \cong k[z^3]$  is regular, so  $e_{\text{HK}}(R/P) = 1$ . Now,

$$\widehat{R}_P/(x^3, y^3) \cong k(z^3)[[x^3, y^3, x^2y, y^2/z^2, x/z]]/(x^3, y^3)$$

and it is easy to compute the basis of this ring over  $k(z^3)$ :

$$1, x/z, y^2/z^2, x^2/z^2, x^2y, xy^2, x^2y^2/z, x^3y/z, y^4/z, xy^4/z^2, x^4y/z^2.$$

Therefore,  $e(R/(x^3, y^3)) = 11$ .

The computation for  $J_k$  is similar but a bit more complicated. First of all, if  $k \not\equiv 1 \pmod{3}$  (so  $2k + 1$  is not divisible by 3), then

$$Q = (x^3, x^2y, z^2x, y^3 - z^{3k}) = \sqrt{J_k}$$

is prime as can be seen by taking the presentation

$$k[[a, b]]/(a^3 - b^{2k+1}) \rightarrow R/Q \cong k[[y^2z, z^3]]/(y^3 - z^{3k}).$$

In this case,  $e(R/Q) = 3$ , so  $e(R/J_k)$  is divisible by 3 and we already know that it cannot be equal to 11. □

Thus even Hilbert–Samuel multiplicity does not behave well in this example. Observe that  $xy^2 = x^3y^3/x^2y$  and  $xyz = y^3z^2x/y^2z$ , so the normalization of  $R$  is the Veronese subring

$$V = k[[x^3, x^2y, xy^2, y^3, y^2z, yz^2, z^3, z^2x, zx^2]].$$

Because  $xy^2 \cdot y^{3n} \notin R$  for all  $n$ , the quotient  $S/R$  does not have finite length. On the other hand,  $x^3y^3V \subseteq R$ , so  $\dim S/R$  must equal to 1.

Therefore since  $\text{depth } V = 2$ , from the exact sequence of local cohomology we have

$$0 \rightarrow H_m^0(S/R) \rightarrow H_m^1(R) \rightarrow 0 \rightarrow H_m^1(S/R) \rightarrow H_m^2(R) \rightarrow 0 = H_m^2(S).$$

Therefore,  $\lambda(H_m^1(R)) < \infty$ , but  $\lambda(H_m^2(R)) = \infty$ . This example shows that the assumptions of Srinivas and Trivedi are close to being necessary.

### 4.2 Questions

The techniques used to prove the main theorems of the paper, which are summarized in Theorem 1.3, rely on the assumption that the parameter sequence  $f_1, \dots, f_h$  forms a reduced ideal.

QUESTION 4.5. Let  $(R, \mathfrak{m}, k)$  be a complete Cohen–Macaulay F-finite ring of prime characteristic  $p$ . If  $(f_1, \dots, f_c)$  is a parameter ideal, but  $R/(f_1, \dots, f_c)$  is not reduced, then given  $\delta > 0$  does there exist an integer  $N > 0$  such that for any  $\epsilon_1, \dots, \epsilon_h \in \mathfrak{m}^N$ :

- (1)  $|\text{e}_{\text{HK}}(R/(f_1, \dots, f_c)) - \text{e}_{\text{HK}}(R/(f_1 + \epsilon_1, \dots, f_c + \epsilon_c))| < \delta$ ; and
- (2)  $\text{s}(R/(f_1 + \epsilon_1, \dots, f_c + \epsilon_c)) < \delta$ ? Note that  $\text{s}(R/(f_1, \dots, f_c)) = 0$  since  $R/(f_1, \dots, f_c)$  is not reduced.

Originally, the second author hoped to approach Question 1.1, by passing to the associated graded ring. Namely, Srinivas and Trivedi [ST96, Theorem 3] show  $\text{Gr}_{\mathfrak{m}}(R/(f))$  is a locally constant function of  $f$  in  $\mathfrak{m}$ -adic topology, which immediately implies that Hilbert–Samuel multiplicity is locally constant.

However the Hilbert–Kunz multiplicities of a local ring and its associated graded ring need not be equal ([WY00, p. 302] and also Example 4.1). Moreover, Hilbert–Kunz multiplicity is not an invariant of the associated graded ring, that is, as we see in Example 4.1, it may happen that  $e_{\text{HK}}(R) \neq e_{\text{HK}}(S)$  even though  $\text{Gr}_{\mathfrak{m}_R}(R) \cong \text{Gr}_{\mathfrak{m}_S}(S)$ . Investigation of the following question could yield an explanation.

QUESTION 4.6. Does the convergence rate of the Hilbert–Kunz multiplicity come from the associated graded ring? Namely, is there a constant  $C$  that depends only on the associated graded ring of  $S$  (not  $S!$ ) such that

$$\left| e_{\text{HK}}(S) - \frac{1}{p^{ed}} \lambda(S/(\mathfrak{m})^{[p^e]}) \right| < \frac{C}{p^e}$$

for all  $e$ ?

This question can be stated even more explicitly: consider the family of ideals in  $\text{Gr}_{\mathfrak{m}}(R)$

$$J_e = \bigoplus_{n \geq 0} \frac{\mathfrak{m}^{[p^e]} \cap \mathfrak{m}^n + \mathfrak{m}^{n+1}}{\mathfrak{m}^{n+1}}.$$

By the definition,  $\lambda(R/\mathfrak{m}^{[p^e]}) = \lambda(\text{Gr}_{\mathfrak{m}}(R)/J_e)$ . Because  $J_e^{[p]} \subseteq J_{e+1}$ , we can show the existence of  $e_{\text{HK}}(R)$  directly in  $\text{Gr}_{\mathfrak{m}}(R)$  [PT18, Theorem 4.3]. However, in order to get the convergence rate, the current techniques [PT18, Corollary 4.5] require us to find a  $p^{-1}$ -linear map  $\psi$  on  $\text{Gr}_{\mathfrak{m}}(R)$  such that  $\psi(J_{e+1}) \subseteq J_e$  for all  $e$ .

Because for a fixed  $e$  the individual Hilbert–Kunz function  $R/(f, \mathfrak{m}^{[p^e]})$  is clearly continuous (Lemma 3.1), we can employ the uniform convergence machinery to show that a positive answer to Question 4.6 gives a positive answer to Question 1.1. However, we currently do not see how to approach Question 4.6, so we had to search for other methods.

We also want to offer two questions motivated by intuitive understanding of singularities: we expect that singularity will not get worse after a sufficiently small perturbation.

QUESTION 4.7. Let  $(R, \mathfrak{m}, k)$  be a local ring of prime characteristic  $p$ . Is there  $N$  such that for all  $\epsilon \in \mathfrak{m}^N$   $e_{\text{HK}}(R/(f)) \geq e_{\text{HK}}(R/(f + \epsilon))$ ?

QUESTION 4.8. Let  $(R, \mathfrak{m}, k)$  be a local ring of prime characteristic  $p$ . Is there  $N$  such that for all  $\epsilon \in \mathfrak{m}^N$   $s(R/(f)) \leq s(R/(f + \epsilon))$ ?

We remark that Srinivas and Trivedi [ST96, Corollary 2] showed that the analogue of Question 4.7 holds for Hilbert–Samuel multiplicity. They prove that if  $(R, \mathfrak{m}, k)$  is local then one can always find a small neighborhood  $\mathfrak{m}^N$  such that  $e(R/(f)) \geq e(R/(f + \epsilon))$  for all  $\epsilon \in \mathfrak{m}^N$ .

### 4.3 Comments on F-pure thresholds

A conjecture related to the results of this paper is the Ascending Chain Condition conjecture for F-pure thresholds, very recently resolved by Sato [Sat17, Sat18], which states that if  $(R, \mathfrak{m}, k)$  is a regular local ring and  $f_1, f_2, f_3, \dots$  are elements of  $R$  such that

$$\text{fpt}(f_1) \leq \text{fpt}(f_2) \leq \text{fpt}(f_3) \leq \dots$$

then  $\text{fpt}(f_n) = \text{fpt}(f_{n+1})$  for all  $n \gg 0$ . The conjecture is motivated by Shokurov’s ACC conjecture for log canonical thresholds [Sho92], which was also resolved in the smooth case [dFEM10, HMX14]. As the F-pure threshold function is continuous with respect to the  $\mathfrak{m}$ -adic topology, it follows that if the ACC conjecture holds then for each  $f \in R$  there exists  $N \in \mathbb{N}$  such that for each  $\epsilon \in \mathfrak{m}^N$ ,  $\text{fpt}(f + \epsilon) \geq \text{fpt}(f)$ . Hernández, Núñez-Betancourt, and Witt [HNnW17] recently investigated this particular implication of the ACC conjecture. Their techniques establish that if the Jacobian ideal of an element  $f \in R$  is  $\mathfrak{m}$ -primary, then  $\text{fpt}(f) = \text{fpt}(g)$  for all  $g$  sufficiently close to  $f$ . Their result is also recovered by Samuel’s theorem from [Sam56] mentioned in the introduction.

A natural analogue to the ACC conjecture for Hilbert–Kunz multiplicity would be a descending chain condition. That is given a sequence of elements  $f_1, f_2, f_3, \dots$  in a regular local ring  $(R, \mathfrak{m}, k)$  such that  $e_{\text{HK}}(R/(f_1)) \geq e_{\text{HK}}(R/(f_2)) \geq e_{\text{HK}}(R/(f_3)) \geq \dots$  is it necessarily the case that  $e_{\text{HK}}(R/(f_n)) = e_{\text{HK}}(R/(f_{n+1}))$  for all  $n \gg 0$ ? Work of Monsky [Mon98] shows the natural analogue of the ACC conjecture for Hilbert–Kunz multiplicity does not hold.

EXAMPLE 4.9. In [Mon98] Monsky computed the Hilbert–Kunz multiplicity of a family of hypersurfaces of the form  $R_\alpha = k[[x, y, z]]/(z^4 + xyz^2 + (x^3 + y^3)z + \alpha x^2 y^2)$  where  $\alpha \in k$ , an algebraically closed field of characteristic 2. His computations show that there is a sequence  $\alpha_n$  such that  $e_{\text{HK}}(R_{\alpha_n}) = 3 + 4^{-n}$ , which gives an infinite decreasing sequence.

We suspect there is a family of hyperplanes, similar to that of Example 4.9, whose F-signatures form an infinite increasing sequence.

**Acknowledgments.** We want to thank Ian Aberbach, Bhargav Bhatt, Dale Cutkosky, and Mel Hochster for the discussion of this work. In particular, Bhargav Bhatt has helped shape Lemma 3.3 and Proposition 4.4 was suggested by Mel Hochster. This project started while the first author was visiting University of Michigan, we thank Mel Hochster and University of Michigan for making that trip possible.

We are grateful to Linquan Ma for pointing an inaccuracy in an earlier draft of this paper.

#### REFERENCES

- [AE08] I. M. Aberbach and F. Enescu, *Lower bounds for Hilbert–Kunz multiplicities in local rings of fixed dimension*, Michigan Math. J. **57** (2008), 1–16. Special volume in honor of Melvin Hochster.
- [AL03] I. M. Aberbach and G. J. Leuschke, *The F-signature and strong F-regularity*, Math. Res. Lett. **10**(1) (2003), 51–56.
- [AP17] J. Adamus and A. Patel, *On finite determinacy of complete intersection singularities*, preprint, 2017, arXiv:1705.08985.
- [BE04] M. Blickle and F. Enescu, *On rings with small Hilbert–Kunz multiplicity*, Proc. Amer. Math. Soc. **132**(9) (2004), 2505–2509 (electronic).
- [BH93] W. Bruns and J. Herzog, *Cohen–Macaulay Rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, Cambridge, 1993.
- [CS93] S. D. Cutkosky and H. Srinivasan, *An intrinsic criterion for isomorphism of singularities*, Amer. J. Math. **115**(4) (1993), 789–821.
- [DPY16] A. De Stefani, T. Polstra and Y. Yao, *Globalizing F-invariants*, preprint, 2016, arXiv:1608.08580.
- [EY11] F. Enescu and Y. Yao, *The lower semicontinuity of the Frobenius splitting numbers*, Math. Proc. Cambridge Philos. Soc. **150**(1) (2011), 35–46.
- [dFEM10] T. de Fernex, L. Ein and M. Mustața, *Shokurov’s ACC conjecture for log canonical thresholds on smooth varieties*, Duke Math. J. **152**(1) (2010), 93–114.
- [GO08] O. Gabber and F. Orgogozo, *Sur la p-dimension des corps*, Invent. Math. **174**(1) (2008), 47–80.
- [HMX14] C. D. Hacon, J. McKernan and C. Xu, *ACC for log canonical thresholds*, Ann. of Math. (2) **180**(2) (2014), 523–571.
- [HNnW17] D. J. Hernández, L. Núñez-Betancourt and E. E. Witt, *Local  $\mathfrak{m}$ -adic constancy of F-pure thresholds and test ideals*, Math. Proc. Cambridge Philos. Soc. **164**(2) (2018), 285–295.
- [Hir65] H. Hironaka, “On the equivalence of singularities. I”, in *Arithmetical Algebraic Geometry (Proc. Conf. Purdue Univ., 1963)*, Harper & Row, New York, 1965, 153–200.
- [Hoc77] M. Hochster, *Cyclic purity versus purity in excellent Noetherian rings*, Trans. Amer. Math. Soc. **231**(2) (1977), 463–488.

- [Hocb] M. Hochster, *The structure theory of complete local rings*. Supplementary notes to Math 615 Winter 2017, available at <http://www.math.lsa.umich.edu/~hochster/615W17/615.html>.
- [Hoca] M. Hochster, *Foundations of tight closure*. Lecture notes available at <http://www.math.lsa.umich.edu/~hochster/mse.html>.
- [HH90] M. Hochster and C. Huneke, *Tight closure, invariant theory, and the Briançon–Skoda theorem*, J. Amer. Math. Soc. **3**(1) (1990), 31–116.
- [HL02] C. Huneke and G. J. Leuschke, *Two theorems about maximal Cohen–Macaulay modules*, Math. Ann. **324**(2) (2002), 391–404.
- [Kun69] E. Kunz, *Characterizations of regular local rings of characteristic  $p$* , Amer. J. Math. **91** (1969), 772–784.
- [Kun76] E. Kunz, *On Noetherian rings of characteristic  $p$* , Amer. J. Math. **98**(4) (1976), 999–1013.
- [KS18] K. Kurano and K. Shimomoto, *An elementary proof of Cohen–Gabber theorem in the equal characteristic  $p > 0$  case*, Tohoku Math. J. (2) **70**(3) (2018), 377–389.
- [Mat86] H. Matsumura, *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics **8**, Cambridge University Press, Cambridge, 1986. Translated from the Japanese by M. Reid.
- [Mon83] P. Monsky, *The Hilbert–Kunz function*, Math. Ann. **263**(1) (1983), 43–49.
- [Mon98] P. Monsky, *Hilbert–Kunz functions in a family: point- $S_4$  quartics*, J. Algebra **208**(1) (1998), 343–358.
- [PT18] T. Polstra and K. Tucker,  *$F$ -signature and Hilbert–Kunz multiplicity: a combined approach and comparison*, Algebra Number Theory **12**(1) (2018), 61–97.
- [Sam56] P. Samuel, *Algébricité de certains points singuliers algébroides*, J. Math. Pures Appl. (9) **35** (1956), 1–6.
- [Sat17] K. Sato, *Ascending chain condition for  $F$ -pure thresholds on a fixed strongly  $F$ -regular germ*, preprint, 2017, arXiv:1710.05331.
- [Sat18] K. Sato, *Ascending chain condition for  $F$ -pure thresholds with fixed embedding dimension*, preprint, 2018, arXiv:1805.07066.
- [Sho92] V. V. Shokurov, *Three-dimensional log perestroikas*, Izv. Ross. Akad. Nauk Ser. Mat. **56**(1) (1992), 105–203.
- [SVdB97] K. E. Smith and M. Van den Bergh, *Simplicity of rings of differential operators in prime characteristic*, Proc. Lond. Math. Soc. (3) **75**(1) (1997), 32–62.
- [ST96] V. Srinivas and V. Trivedi, *The invariance of Hilbert functions of quotients under small perturbations*, J. Algebra **186**(1) (1996), 1–19.
- [TW04] S. Takagi and K.-i. Watanabe, *On  $F$ -pure thresholds*, J. Algebra **282**(1) (2004), 278–297.
- [Tuc12] K. Tucker,  *$F$ -signature exists*, Invent. Math. **190**(3) (2012), 743–765.
- [WY00] K.-i. Watanabe and K.-i. Yoshida, *Hilbert–Kunz multiplicity and an inequality between multiplicity and colength*, J. Algebra **230**(1) (2000), 295–317.

Thomas Polstra  
*Department of Mathematics*  
*University of Utah*  
*Salt Lake City, USA*  
polstra@math.utah.edu

Ilya Smirnov  
*Department of Mathematics*  
*Stockholm University*  
*S-106 91, Stockholm*  
*Sweden*  
smirnov@math.su.se