

Associations between monthly rainfall and mortality in cattle due to East Coast fever, anaplasmosis and babesiosis

Research Article

Cite this article: Chepkwony R, Castagna C, Heitkönig I, van Bommel S, van Langevelde F (2020). Associations between monthly rainfall and mortality in cattle due to East Coast fever, anaplasmosis and babesiosis. *Parasitology* **147**, 1743–1751. <https://doi.org/10.1017/S0031182020001638>

Received: 20 April 2020

Revised: 26 July 2020

Accepted: 1 August 2020


First published online: 10 September 2020

Key words:

Animal diseases; indigenous cattle; tick-borne diseases; ticks; time series analysis

Author for correspondence:

Richard Chepkwony,
E-mail: rchepkwony74@gmail.com

Richard Chepkwony^{1,2,3}, Carolina Castagna², Ignas Heitkönig², Severine van Bommel^{3,4} and Frank van Langevelde^{2,5} 

¹Kenya Wildlife Service, P.O. Box 40241-00100, Nairobi, Kenya; ²Wageningen University, Wildlife Ecology and Conservation Group, P.O. Box 47, 6700 AA, Wageningen, The Netherlands; ³Wageningen University, Strategic Communications Group, P.O. Box 8130, 6700 EW, Wageningen, The Netherlands; ⁴University of Queensland, School of Agriculture and Food Sciences, Gatton, QLD 4343, Australia and ⁵School of Life Sciences, Westville Campus, University of KwaZulu-Natal, Durban 4000, South Africa

Abstract

Weather conditions can impact infectious disease transmission, causing mortalities in humans, wild and domestic animals. Although rainfall in dry tropical regions is highly variable over the year, rainfall is thought to play an important role in the transmission of tick-borne diseases. Whether variation in rainfall affects disease-induced mortalities, is, however, poorly understood. Here, we use long-term data on monthly rainfall and Boran cattle mortality (1998–2017) to investigate associations between within-year variation in rainfall and cattle mortalities due to East Coast fever (ECF), anaplasmosis and babesiosis in Laikipia, Kenya, using ARIMAX modelling. Results show a negative correlation between monthly rainfall and cattle mortality for ECF and anaplasmosis, with a lag effect of 2 and 6 months, respectively. There was no association between babesiosis-induced mortalities and monthly rainfall. The results of this study suggest that control of the tick-borne diseases ECF and anaplasmosis to reduce mortalities should be intensified during rainy periods after the respective estimated time lags following dry periods.

Introduction

Weather conditions are thought to play an important role in the transmission of many infectious diseases that cause significant mortalities in humans, wild and domestic animals in both temperate and tropical countries. For instance, the outbreaks of diseases such as trypanosomiasis are correlated with rainfall that provides favourable conditions for survival and reproduction of trypanosomes (Rogers and Randolph, 2006). Also, avian influenza is positively influenced by rainfall (Brown *et al.*, 2017). Temperature, relative humidity and precipitation have been shown to play a vital role in the incidences of dengue fever (Gubler *et al.*, 2001), Rift Valley fever (Maloo, 1993; Anyamba *et al.*, 2009; Lwande *et al.*, 2015) and malaria (Wangdi *et al.*, 2010).

Tick-borne diseases (TBDs), such as tick-borne encephalitis (Subak, 2003), boutonneuse fever (de Sousa *et al.*, 2006) and lyme disease (Rogers and Randolph, 2006), are known to be also positively correlated to rainfall. In dry tropical areas, rainfall patterns are often highly variable. Ticks reproduce massively at the onset of any rainfall episode and slowly develop into nymphs in 3 weeks to 1 month. The nymphal and adult stages have been shown to be the stages where pathogens are transmitted from the tick to the mammalian host, resulting in diseases and probably mortalities (Jongejan & Uilenberg, 2004; Hechemy *et al.*, 2006; Rogers and Randolph, 2006; Dantas-Torres, 2015). Variations in tick densities over time depend on three factors: fecundity, length of time between the stages (conditioned partly by the waiting time of successive hosts and environmental conditions) and mortality rate of the ticks. These three processes lead to tick densities that follow a certain seasonality and which result in a more or less predictable annual pattern. Moreover, the growth of the vegetation during these periods attracts reservoir mammalian hosts facilitating transmission of pathogens (Norval *et al.*, 1988; Cumming, 2002; Awa *et al.*, 2015). On the other hand, as more forage is available during periods of rainfall for mammalian hosts that may be negatively affected by TBDs, their body condition (and immunity levels) also improve, which reduces their susceptibility.

TBDs cause severe economic losses and are associated with 60–80% of the livestock mortalities affecting livelihoods (Chepkwony *et al.*, 2018). To date, few studies on the effects of TBDs on livestock mortality have been conducted in dry tropical areas. Identification of the environmental determinants of TBDs and their associated mortalities is of great importance to understand the incidences of TBDs, especially in dry tropical areas with highly variable rainfall patterns. Rainfall may affect ticks directly and indirectly by its effect on relative humidity and vegetation. Whether variation in rainfall affects disease-induced mortalities is, however, poorly understood. In the absence of tick monitoring in many regions in the tropics, livestock mortalities, which are indications of effective pathogen transmission, may provide understanding about the associations between within-year variation in rainfall and the incidences of TBDs (Gachohi *et al.*, 2012; Fouque and Reeder, 2019).




Fig. 1. Map of Kenya and the location study site Olpejeta Conservancy (OPC), in Laikipia, Kenya. OPC is located to the south of Laikipia county and borders Mutara ranch to the West and Naro-Moru township to the South East.

Although there is a need to determine if there are any associations between rainfall and TBDs-induced mortalities, long-term data are generally lacking (Heuer *et al.*, 2004; Keeling and Rohani, 2008; Kołodziej-Sobocińska, 2019). Here, we use long-term data on Boran cattle mortality to determine associations between monthly rainfall and three TBDs, namely East Coast fever (ECF), anaplasmosis and babesiosis, to further improve our understanding about the extent weather conditions play in the dynamics of pathogen transmission and for use in disease control. We expect that high monthly rainfall would lead to an increase in mortalities induced by these TBDs. ECF causes most mortalities (65%; Kanyari and Kagira, 2000) in cattle of all TBDs, as the vector, the tick *Rhipicephalus appendiculatus*, is one of the most abundant tick species in East Africa, infesting both wild and domestic animals (Zieger *et al.*, 1998; Wesonga *et al.*, 2006; Swai *et al.*, 2009; Keesing *et al.*, 2013; Mwamuye *et al.*, 2016). Babesiosis accounts for 5.1% and anaplasmosis for 4.5% of the mortalities in cattle due to TBDs (Kanyari and Kagira, 2000), and these two TBDs share the same main vector, the tick *R. decoloratus*.

Methods

Study area

The data on cattle mortality were collected in Olpejeta Conservancy (OPC), a facility for commercial livestock production and wildlife

management in Laikipia County, Kenya. Laikipia county is located between 0°04'60.00"N and 36°39'59.99"E (Fig. 1). The county covers ~10 000 km² and the rainfall is typically bimodal and unpredictable throughout the year. Laikipia county has an estimated human population of ~400 000 living in either urban or rural areas and relies on wildlife conservation, livestock production and crop farming for their livelihoods (Government of Kenya, 2016). The interfaces between human, wildlife and domestic animals coupled with the local environmental conditions in the area may provide all-year-round conditions for tick survival, the spread of TBDs with resulting mortalities with major ramifications on livestock production and wildlife conservation (Odadi *et al.*, 2011; Keesing *et al.*, 2013; Allan *et al.*, 2017; VanderWaal *et al.*, 2017). OPC is an intensively managed protected area in Laikipia County and covers 364 km² (Fischhoff *et al.*, 2007). In OPC, ticks are managed in domestic animals through weekly acaricide applied via high-pressure spraying nozzles (Chepkwony *et al.*, 2018).

Data collection

We obtained monthly data on domestic cattle mortalities from OPC for a period of 19 years beginning from October 1998 to October 2017. The data are used at OPC to monitor performances of commercial cattle ranching, one of the key land use activities in the property. Cattle diseases in OPC were monitored through

surveillance for signs and symptoms to enable early detection, prevention, control and management. Diagnosis of a disease is done at several levels: (i) clinical signs or symptoms of the disease enabling to make a tentative diagnosis are based on signs that are pathognomonic for a particular disease. For example, for ECF, symptoms may include reduced foraging, dullness, and swollen lymph nodes of an affected animal. (ii) Aided diagnosis (laboratory, histology, culture and sensitivity, polymerase chain reaction (PCR), serology). Here, samples collected from an animal were subjected to a series of tests for confirmatory diagnosis, with more than 95% accuracy. (iii) Post-mortem analysis to ascertain the cause of death. Additionally, the employees monitored tick loads and body condition to infer other causes of death such as injuries, poisoning, snake bites and predation.

To control ticks in cattle, acaricides are applied weekly using high-pressure nozzle spray-races. Besides ECF, anaplasmosis and babesiosis, other causes of cattle mortalities, namely anthrax, brucellosis, dysentery and stillbirths, among others, were recorded. When the disease was not confirmed *via* laboratory diagnostics, it was generally classified in the data as non-specified. Other forms of mortalities recorded included depredation from wild carnivores, poisoning by plants or chemicals, bloat, debility, accidents occasioned through falling-off cliffs while grazing in the wild, snares and trampling by other members while aggregated in places such as in chemical spraying-race or supplemental feeding areas.

We also have monthly data on herd size of cattle. We linearly interpolated missing datasets for the herd sizes for the years 2015 and 2016 using the data-sets of herd sizes of the previous and next year. Cattle mortality per TBD was calculated as the percentage of animals diagnosed with ECF, anaplasmosis or babesiosis per month from the corresponding herd size. We obtained monthly rainfall (mm) data from OPC's Kamok meteorological station spanning over five decades since 1965. We used the monthly rainfall over the period that coincided with the mortality data.

Testing for associations between monthly rainfall and cattle mortalities

As our data contained the time series of mortalities and rainfall, we used Autoregressive Integrated Moving Average (ARIMA) models (Huang and Shih, 2003; Gaur and Gaur, 2006). In these models, the percentage of cattle mortalities per month per TBD (ECF, anaplasmosis and babesiosis) was used as the response variable and monthly rainfall as the predictor variable. Time series analysis allows us to understand whether the underlying structure of monthly rainfall can explain the observed mortality observations and it tests whether there is a seasonal pattern in mortality that can be related to monthly rainfall with or without a lag effect.

We excluded all outliers due to known causes from the analysis since they significantly affected the model performance (Wangdi *et al.*, 2010). Our sample size was 228 observations of monthly cattle mortalities per TBD. ARIMA uses the parameters p , d and q , where p represents the autoregressive (AR) order, d stands for the differencing order and q for the moving average (MA) order. AR orders specify which previous values from the time series are used to predict current values. Our ARIMAX model using the predictor variable rainfall can be specified as:

$$Z_t + \sum_{p=1}^P \phi_p Z_{t-p} = \mu + \sum_{i=1}^I \sum_{n=1}^N \beta_i n X_{i,t-n} + \sum_{q=1}^Q \theta_q \varepsilon_{t-q}$$

where Z_t is the mortality due to the TBD at month t , Z_{t-p} is the mortality in the previous month with a lag of p (i.e. AR), μ is the intercept, $X_{i,t-n}$ is the independent variable i (here rainfall) with a lag of n months, and ε_{t-q} is the error at lag q (i.e. MA). ϕ_p , β_i and

θ_q are the model parameters for the AR term, the independent variable and the MA term, respectively (Monamele *et al.*, 2017). MA orders specify the deviations from the series mean for previous values and are used to predict current values. With monthly rainfall as an exogenous variable (X) to make predictions in the ARIMA models, the models are then referred to as ARIMAX. Following the procedures described by Huang and Shih (2003), we developed the ARIMAX models for the TBDs using the Forecasting Expert Modeler in SPSS 25 IBM statistics (Huang and Shih, 2003; Gaur and Gaur, 2006). The Expert Modeller was used to fit the best model for the time series data using the maximum likelihood. The Expert Modeller identified and estimated the best-fitting Autoregressive Integrated Moving Average (ARIMA) or exponential smoothing model for the dependent variable with one or more AR terms and/or one or more MA terms. The ARIMAX model assumed that all other conditions, such as control and preventive measures including vaccination of infected animals, do not vary systematically over time. We did not distinguish a training and test set in the data as monthly rainfall in a semi-arid region such as Laikipia is highly variable. For training the model, we used a long period of data, but short-term predictions of the mortalities for say 1 year are not reliable due to the highly variable monthly rainfall.

As a general rule in time series analysis, stationarity or non-stationarity of the series is indicated by the existence of a sigmoidal pattern with sharp points at the peak of the sequence plots indicating seasonality. If an increasing trend is observed, then the pattern is deemed as exhibiting a non-stationary trend (Geweke & Porter-Hudak 1983). When the data are non-stationary, differencing will be necessary to 'stationary' it through either log-transformation, deflating or raising-to-some-power to convert the data to a form where its local random variations over time vary around zero (Huang and Shih, 2003). The stationarity of the data was checked by the autocorrelation function (ACF) and the partial autocorrelation function (PACF). We then used the seasonally adjusted factor (SAF) to determine the peaks of seasonal variations of the three TBDs. We used the Ljung-Box (modified Box-Pierce) test to evaluate if the model was correctly specified (Huang and Shih, 2003; Clement, 2014). Besides, we used the mean absolute percentage error (MAPE) to characterize the model, which is a test of the smoothness of the time series. The lower the MAPE, the more reliable is the model as MAPE is a measure of error (Huang and Shih, 2003; Clement, 2014).

Results

Livestock mortalities

Depredation by wild carnivores was responsible for 23% of the cattle mortalities over the entire period, while the three TBDs, namely ECF, anaplasmosis and babesiosis, accounted for 20.4% of the mortalities, which made them the second most common cause of mortalities for the Boran cattle at OPC (ECF: 13.2%, anaplasmosis: 4.0%, babesiosis: 3.2%). The population of Boran cattle varied from 4591 to 8368 cattle over the years we studied. The time-series of the mortalities due to ECF, anaplasmosis and babesiosis are shown in Fig. 2. The unusual observations in May and June 2005 were attributed to the OPC management's decision to change the chemical acaricides used for tick control from 'Triatix[®]' to 'Delete EC[®]', causing a spike in cattle mortality. In August 2005, there were some reported cases of resistance by blue ticks (vector for anaplasmosis and babesiosis) to the applied acaricide leading to increases in anaplasmosis and babesiosis-induced mortalities. Because this affected the models as a confounding factor not related to monthly rainfall, we treated them as outliers and they were removed from the analysis.

Table 1. Summary of the ARIMAX models with rainfall as a predictor variable for the three tick-borne diseases East Coast fever (ECF), babesiosis and anaplasmosis

			Estimate	s.e.	<i>t</i>	<i>P</i>
ECF	Constant		0.070	0.010	6.880	<0.001
	AR	Lag 1	0.171	0.070	2.437	0.016
		Lag 2	0.179	0.071	2.532	0.012
Rainfall	Delay		2			
	Numerator	Lag 0	0.000	4.317×10^{-5}	-2.934	0.004
		Denominator	Lag 1	1.536	0.024	63.823
		Lag 2	-0.935	0.031	-30.137	<0.001
Babesiosis	AR	Lag 1	0.830	0.051	16.179	<0.001
	MA	Lag 1	0.368	0.080	4.595	<0.001
		Lag 6	-0.329	0.070	-4.684	<0.001
Anaplasmosis	Constant		0.021	0.004	4.728	<0.001
	AR	Lag 1	0.378	0.070	5.412	<0.001
		Lag 2	0.195	0.071	2.764	0.006
Rainfall	Delay		6			
	Numerator	Lag 0	-7.768×10^{-5}	3.526×10^{-5}	-2.203	0.029

Table 2. Summary of ARIMAX model statistics with rainfall as a predictor variable for the three tick-borne diseases East Coast fever (ECF), babesiosis and anaplasmosis (see Table 1 for the model performance)

Model	Number of predictors	Model Fit statistics			Ljung-Box Q		
		Stationary R^2	MAPE	Normalized BIC	Statistics	df	<i>P</i>
ECF	1	0.183	88.634	-5.091	18.859	16	0.276
Babesiosis	0	0.520	91.777	-6.924	32.454	15	0.006
Anaplasmosis	1	0.275	45.561	-7.304	13.675	16	0.623

MAPE, Mean absolute percentage error; BIC, Bayesian information criterion.

Associations between rainfall intensity and cattle mortalities

The ARIMAX models show that ECF and Anaplasmosis followed rainfall with a lag of 2 months and 6 months for ECF and anaplasmosis, respectively (Tables 1 and 2; Figs 2 and 3). We found negative correlations between rainfall and the predicted mortalities due to ECF (rainfall with a lag of 2 months; Spearman rank correlation = -0.135, $N = 224$, $P = 0.04$) and anaplasmosis (rainfall with a lag of 6 months; Spearman rank correlation = -0.461, $N = 222$, $P < 0.001$). The lag of 2 months refers to the amount of rainfall 2 months earlier (the same for the lag of 6 months). Given the negative correlation between rainfall and ECF or anaplasmosis, our results suggest that the occurrence of these diseases is low during a dry period when the earlier amount of rainfall (2 or 6 months earlier) was relatively high, and the occurrence was high during a rainy period when there was earlier a dry period. We can thus conclude that there were more mortalities in a rainy period after a dry period (after 2 and 6 months for ECF and anaplasmosis, respectively). Babesiosis could not be correlated to rainfall. In the Discussion, we will mainly deal with the association with monthly rainfall. For both ECF and anaplasmosis, the number of AR orders in the model is 2, whereas this number is 1 for babesiosis (Table 1). For example, an AR order of 2 specifies that the mortalities of the series 2 months in the past be used to predict the current mortality. For babesiosis, the number of MA orders in the model is 1, which means that the deviations from the mean mortality of the series from each of the last months are considered when predicting current mortality.

The Ljung-Box test indicated that the models for ECF and anaplasmosis were correctly specified ($P > 0.05$) (Table 2). For each TBD, cattle mortalities seem to be stationary with a constant variance oscillating around 0 (Fig. 4).

Discussion

The aim of this study was to understand the association between monthly rainfall and mortalities in Boran cattle due to the three TBDs ECF, anaplasmosis and babesiosis. We found a clear association between monthly rainfall and the occurrence of mortalities due to ECF and anaplasmosis with a disease-specific lag after rainfall. However, this association was negative, whereas we expected that high monthly rainfall would lead to an increase in mortalities due to TBDs. The negative association between monthly rainfall and cattle mortalities associated with ECF and anaplasmosis had a 2-months and 6-months lag after a rainfall event for ECF and anaplasmosis, respectively. For babesiosis, we did not find an association with monthly rainfall.

Most studies indicate that *R. appendiculatus* numbers (vector tick for ECF) are higher during the rainy season (Okello-Onen *et al.*, 1999; Wesonga *et al.*, 2006; Laisser *et al.*, 2015). The nymphs of *R. appendiculatus* were found to be abundant from mid rainy seasons to shortly after the rainy seasons (McCulloch *et al.*, 1968). Fyumagwa *et al.* (2007) and Zieger *et al.* (1998) found, however, that the nymphs were more abundant during the dry season and others found no clear seasonality pattern for




Fig. 2. Monthly Boran cattle mortality as a percentage of the total herd sizes for (A) East Coast fever (ECF), (B) babesiosis, and (C) anaplasmosis in Olpejeta Conservancy, Laikipia, Kenya. The outliers were: May 2005 for ECF, May and August 2005 for both babesiosis and anaplasmosis.




Fig. 3. Observed cattle mortalities due to East Coast fever (ECF) and Anaplasmosis at Olpejeta Conservancy, Kenya from October 1998 to October 2017. T best fit of the ARIMAX models (blue) with the upper confidence limits (UCL).

R. appendiculatus (Zieger *et al.*, 1998; Rogers and Randolph, 2006; Fyumagwa *et al.*, 2007; Walker *et al.*, 2014). An explanation for the negative correlation we found could be that the nymphs were indeed more frequent during the dry season and that these numbers were overall higher than the adult numbers, as Zieger *et al.* (1998) have found. For *R. decoloratus* (vector tick for Anaplasmosis), a higher number was found during the wet period (Wesonga *et al.*, 2006; Walker, 2011). However, Okello-Onen *et al.* (1999) did not find a seasonal effect probably due to the high densities of this tick species, because the control interventions by farmers are still low as compared to other areas such as in Kenya. So, for both vector tick species, there is no clear relation between rainfall and their densities. In this study, we recognize that there are many environmental factors involved in tick dynamics and consequently in TBDs that may confound our findings, such as temperature, relative humidity, vegetation, host density, wildlife densities (contact between wildlife and domestic animals are expected to increase ECF cases, Allan *et al.*, 2017), the presence of tick predators and control practices (acaricide spraying). Unfortunately, we do not have these data during the same time period for which we have data on rainfall.

Different TBDs have different aetiology that influences their transmission, the incubation and subsequent infectiousness in humans and animals. ECF caused by protozoan *Theileria parva* has been shown to take a period between 10 days and 3 weeks from the transmission, development into full-blown infections (Gachohi *et al.*, 2012). Because of its short incubation period and

effect on an animal, ECF is considered as a more lethal and high impact disease than other TBDs such as anaplasmosis and babesiosis. So, if ECF starts shortly after the onset of rainfall it makes the disease prevalent to be high during the dry season. Anaplasmosis also has its own aetiology with its development that may take up to 4–5 months between the period of transmission in an animal, incubation and subsequent effect and manifestation in an animal (Gachohi *et al.*, 2012). Anaplasmosis, therefore, takes a long latency or incubation than either ECF or babesiosis. Moreover, the time between infection with a disease and the time the infected animal dies, i.e. the time of death, may vary between the TBDs. Several studies have found the time of death for ECF-infected cattle to be around 26 days, with ranges from 11 to 44 days (Brocklesby, 1962; Radley *et al.*, 1974; Robson *et al.*, 1977). For anaplasmosis, the time of death is currently unknown. As a proxy, the incubation period varied from 7 to up to 100 days (Theiler, 1911; Pipano *et al.*, 1992; Kocan *et al.*, 2003). These differences in aetiology between the TBDs may probably explain the differences in the lag period of 6 months that this study found for anaplasmosis compared to the lag period of 2 months for ECF. The absence of a correlation between babesiosis and monthly rainfall may be due to the small number of babesiosis-induced mortality cases in the Boran cattle populations (Fig. 2B). Babesiosis is mainly detected in Boran cattle from August 2004 until February 2007 as it is not associated with the changes in monthly rainfall.

The effect of the different TBDs on livestock may also be influenced by an animal's immunity levels or responses, and are often




Fig. 4. Correlograms of the residual autocorrelation function (ACF) and partial autocorrelation function (PACF) of the suggested ARIMAX models for cattle mortalities due to East Coast fever (ECF) (top panels), babesiosis (panels in the middle) and anaplasmosis (lower panels) at the Olpejeta Conservancy, Kenya from October 1998 to October 2017. The dark grey lines indicate 95% confidence bands.

influenced by an animal's body condition (Fyumagwa *et al.*, 2007). When an animal loses condition through parasite infestation or lack of forage and water such as during drought, the animal may easily succumb to the disease, increasing the risk of mortality (Swai *et al.*, 2006). During the high rainfall season when forage is abundant, the animal's body condition may improve considerably boosting the animal's immune response and decreasing its mortality risk. In addition to the aetiology of the TBDs, the negative relationships between the Boran cattle mortalities and monthly rainfall may be also attributed to the lower immunity levels of the animal during the dry months (Potkanski, 1994; Kanyari and Kagira, 2000; Subak, 2003; VanderWaal *et al.*, 2017). Yet another explanation could be that during the dry season contact between herds at OPC increases near water resources, which causes more potential for pathogen spread than during the wet season (VanderWaal *et al.*, 2017).

Furthermore, previous studies in China showed that monthly rainfall was positively correlated to monthly notification of Japanese encephalitis, spread by either ticks or mosquitoes, and their lag times varied from 1 to 2 months (Bi *et al.*, 2007). Although this study did not account for the resultant mortalities, it was nevertheless consistent with our observations that tick-borne disease transmission is influenced by environmental variables (rainfall) and the pathogen infection has lagged effect due to vector biology and perhaps immunity levels in hosts.

We used mortality data of Boran cattle to determine if there is an association between the long-term mortalities due to the three TBDs and monthly rainfall. However, one shortcoming of this study is that we did not have data to compare how many animals

were infected and recovered from each infection due to these TBDs. We also made the general assumption that the control interventions for TBDs were relatively uniform and were based on consistent use of acaricides over the period, which was based on the information obtained from the ranch management (Chepkwony *et al.*, 2018). Although Boran cattle are widely bred in most ranches and pastoral areas of local communities, extrapolation of our results to more extensive and less-controlled herds that are common in the Eastern Africa region may be limited due to different conditions such as control practices. Moreover, the impact of the diseases may differ between age classes. For example, ECF primarily affects calves (Latif *et al.*, 1995). These differences are beyond the scope of our paper.

Our study shows a negative correlation between rainfall intensity and TBDs-induced mortalities in Boran cattle. Our findings suggest that Boran cattle mortalities are probably amplified in the rainy period after dry periods. The analysis presented in this study may have some implications for the prevention and control of TBDs. We suggest that warnings of enhanced risk should take into account knowledge of the lag times of the diseases and the measured rainfall conditions in relation to the risk of the spread of pathogens. Tick control strategies to reduce TBD-induced mortalities may be intensified, such as increasing the frequencies of the application of chemicals (acaricides), during rainy periods after the respective estimated time lags following dry periods, and perhaps by improving the body condition of cattle. This study contributes to a better understanding of the occurrence of mortalities in Boran cattle due to three TBDs, namely ECF, anaplasmosis and babesiosis, and to our ability to predict

ECF- and anaplasmosis-induced mortalities based on monthly rainfall, a prerequisite for disease control in livestock.

Acknowledgements. We wish to thank the Kenya Wildlife Service for authorising the research to be conducted in Laikipia and its staff for providing security during the fieldwork. We also thank Samuel Mutisya, Alfred Kibungei and Joseph Mathenge for providing part of the data and counter-checking the data for errors and omissions. We sincerely thank the entire management of Olpejeta Conservancy Ltd, Laikipia, Kenya, for allowing fieldwork in their conservancy.

Financial support. This study was funded by the Wildlife Ecology and Conservation Group of Wageningen University, The Netherlands, and the INREF Programme 'Environmental Virtual Observatories for Connective Action (EVOCA)'.

Conflict of interests. The authors declare that they have no conflict of interests.

Ethical standards. Not applicable.

References

- Allan BF, Tallis H, Chaplin-Kramer R, Hockett S, Kowal VA, Musengezi J, Okanga S, Ostfeld RS, Schieltz J and Warui CM (2017) Can integrating wildlife and livestock enhance ecosystem services in central Kenya? *Frontiers in Ecology and the Environment* **15**, 328–335.
- Anyamba A, Chretien J-P, Small J, Tucker CJ, Formenty PB, Richardson JH, Britch SC, Schnabel DC, Erickson RL and Linthicum KJ (2009) Prediction of a Rift Valley fever outbreak. *Proceedings of the National Academy of Sciences* **106**, 955–959.
- Awa D, Adakal H, Luogbou N, Wachong K, Leinyuy I and Achukwi M (2015) Cattle ticks in Cameroon: is *Rhipicephalus* (*Boophilus*) *microplus* absent in Cameroon and the Central African region? *Ticks and Tick-Borne Diseases* **6**, 117–122.
- Bi P, Zhang Y and Parton KA (2007) Weather variables and Japanese encephalitis in the metropolitan area of Jinan City, China. *Journal of Infection* **55**, 551–556.
- Brocklesby D (1962) The febrile reaction in fatal East Coast fever – A review of 150 cases. *Bulletin of Epizootic Diseases of Africa* **10**, 49–54.
- Brown I, Mulatti P, Smietanka K, Staubach C, Willeberg P, Adlhoeh C, Candiani D, Fabris C, Zancanaro G and Morgado J (2017) Avian influenza overview October 2016–August 2017. *EFSA Journal* **15**, e05018.
- Chepkwony R, Van Bommel S and Van Langevelde F (2018) Citizen science for development: potential role of mobile phones in information sharing on ticks and tick-borne diseases in Laikipia, Kenya. *NJAS – Wageningen Journal of Life Sciences* **86–87**, 123–135.
- Clement EP (2014) Using normalized Bayesian information criterion (BIC) to improve box-Jenkins model building. *American Journal of Mathematics and Statistics* **4**, 214–221.
- Cumming GS (2002) Comparing climate and vegetation as limiting factors for species ranges of African ticks. *Ecology* **83**, 255–268.
- Dantas-Torres F (2015) Climate change, biodiversity, ticks and tick-borne diseases: the butterfly effect. *International Journal for Parasitology: Parasites and Wildlife* **4**, 452–461.
- de Sousa R, Luz T, Parreira P, Santos-Silva M and Bacellar F (2006) Boutonneuse fever and climate variability. *Annals of the New York Academy of Sciences* **1078**, 162–169.
- Fischhoff IR, Sundaesan SR, Cordingley J and Rubenstein DI (2007) Habitat use and movements of plains zebra (*Equus burchelli*) in response to predation danger from lions. *Behavioral Ecology* **18**, 725–729.
- Fouque F and Reeder JC (2019) Impact of past and on-going changes on climate and weather on vector-borne diseases transmission: a look at the evidence. *Infectious Diseases of Poverty* **8**, 1–9.
- Fyumagwa RD, Runyoro V, Horak IG and Hoare R (2007) Ecology and control of ticks as disease vectors in wildlife of the Ngorongoro Crater, Tanzania. *South African Journal of Wildlife Research* **37**, 79–90.
- Gachohi J, Skilton R, Hansen F, Ngumi P and Kitala P (2012) Epidemiology of East Coast fever (*Theileria parva* infection) in Kenya: past, present and the future. *Parasites & Vectors* **5**, 194.
- Gaur AS and Gaur SS (2006) *Statistical Methods for Practice and Research: A Guide to Data Analysis Using SPSS*. Newbury Park, California: Sage Publications Inc.
- Geweke J and Porter-Hudak S (1983) The estimation and application of long memory time series models. *Journal of Time Series Analysis* **4**, 221–238.
- Government of Kenya (2016) Laikipia County Integrated Development Plan (CIDP). Available at <https://africaopendata.org/dataset/2013-2017-laikipia-county-integrated-development-plan-cidp>.
- Gubler DJ, Reiter P, Ebi KL, Yap W, Nasci R and Patz JA (2001) Climate variability and change in the United States: potential impacts on vector- and rodent-borne diseases. *Environmental Health Perspectives* **109**, 223–233.
- Hechemy KE, Oteo JA, Raoult D, Silverman DJ and Blanco JR (2006) A century of rickettsiology: emerging, reemerging rickettsioses, clinical, epidemiologic, and molecular diagnostic aspects and emerging veterinary rickettsioses: an overview. *Annals of the New York Academy of Sciences* **1078**, 1–14.
- Heuer C, Boehle W, Rutagwenda T, Jongejan F, Unger F, Kivaria FM and Okello-Onen J (2004) Endemic stability for *Theileria parva* infections in Ankole calves of the Ankole ranching scheme, Uganda. *Onderstepoort Journal of Veterinary Research* **71**, 189–195.
- Huang S-J and Shih K-R (2003) Short-term load forecasting via ARMA Model identification including non-Gaussian process considerations. *IEEE Transactions on Power Systems* **18**, 673–679.
- Jongejan F and Uilenberg G (2004) The global importance of ticks. *Parasitology* **129**, S3.
- Kanyari P and Kagira J (2000) The role of parasitic diseases as causes of mortality in cattle in a high potential area of central Kenya: a quantitative analysis. *The Onderstepoort Journal of Veterinary Research* **67**, 157–161.
- Keeling MJ and Rohani P (2008) Modeling infectious diseases in humans and animals. *Clinical Infectious Diseases* **47**, 864–865.
- Keesing F, Allan BF, Young TP and Ostfeld RS (2013) Effects of wildlife and cattle on tick abundance in central Kenya. *Ecological Applications* **23**, 1410–1418.
- Kocan KM, De la Fuente J, Guglielme AA and Meléndez RD (2003) Antigens and alternatives for control of *Anaplasma marginale* infection in cattle. *Clinical Microbiology Reviews* **16**, 698–712.
- Kołodziej-Sobocińska M (2019) Factors affecting the spread of parasites in populations of wild European terrestrial mammals. *Mammal Research* **64**, 301–318.
- Laisser ELK, Chenyambuga SW, Msalya G, Kipanyula MJ, Mdegela RH, Karimuribo ED, Mwilawa AJ and Kusiluka LJK (2015) Knowledge and perception on ticks, tick-borne diseases and indigenous cattle tolerance to East Coast fever in agro-pastoral communities of Lake Zone in Tanzania. *Livestock Research for Rural Development* **27**, Article 64. Retrieved September 15, 2020, from <http://www.lrrd.org/lrrd27/4/lais27064.htm>
- Latif AA, Rowlands GJ, Punyua DK, Hassan SM and Capstick PB (1995) An epidemiological study of tick-borne diseases and their effects on productivity of zebu cattle under traditional management on Rusinga Island, western Kenya. *Preventive Veterinary Medicine* **22**, 169–181.
- Lwande OW, Paul GO, Chiyo PI, Ng'ang'a E, Otieno V, Obanda V and Evander M (2015) Spatio-temporal variation in prevalence of Rift Valley fever: a post-epidemic serum survey in cattle and wildlife in Kenya. *Infection Ecology & Epidemiology* **5**, 30106.
- Maloo SH (1993) *Epidemiological Studies of Vector-Borne Diseases with Consequent Development of Preventive Medicine Programmes for Small-Holder Dairy Farmers in Coastal Kenya* (PhD thesis). The University of Glasgow.
- McCulloch B, Kalaye WJ, Tungaraza R and Mbasha EMS (1968) A study of the life history of the tick *Rhipicephalus appendiculatus* the main vector of East Coast fever-with reference to its behaviour under field conditions and with regard to its control in Sukumaland, Tanzania. *Bulletin of Epizootic Diseases of Africa* **16**, 477–500.
- Monamele GC, Vernet MA, Nsaibirni R, Bigna JJR, Kenmoe S, Njankou MR and Njouom R (2017) Associations between meteorological parameters and influenza activity in a subtropical country: Case of five sentinel sites in Yaounde-Cameroon. *PloS one* **12**, e0186914.
- Mwamuye MM, Kariuki E, Omondi D, Kabii J, Odongo D, Masiga D and Villinger J (2016) Occurrence of novel and emergent tick-borne pathogens in a Kenyan biodiversity hotspot. *International Journal of Infectious Diseases* **45**, 350.
- Norval RAI, Sutherst RW, Kurki J, Gibson JD and Kerr JD (1988) The effect of the brown ear-tick *Rhipicephalus appendiculatus* on the growth of Sanga and European breed cattle. *Veterinary Parasitology* **30**, 149–164.
- Odadi WO, Karachi MK, Abdulrazak SA and Young TP (2011) African wild ungulates compete with or facilitate cattle depending on season. *Science* **333**, 1753–1755.

- Okello-Onen J, Tukahirwa EM, Perry BD, Rowlands GJ, Nagda SM, Musisi G, Bode E, Heinonen R, Mwayi W and Opuda-Asibo J** (1999) Population dynamics of ticks on indigenous cattle in a pastoral dry to semi-arid rangeland zone of Uganda. *Experimental & Applied Acarology* **23**, 79–88.
- Pipano E, Krigel Y, Markovics A and Shkap V** (1992) The effect of oxytetracycline treatment on immunity induced by *Anaplasma centrale*. *Veterinary Microbiology* **31**, 81–87.
- Potkanski T** (1994) *Property Concepts, Herding Patterns and Management of Natural Resources among the Ngorongoro and Salei Maasai of Tanzania* (No. 6; Pastoral Land Tenure Series). London: International Institute for Environment and Development.
- Radley D, Brown C, Burridge M, Cunningham M, Peirce M and Purnell R** (1974) East coast fever: quantitative studies of *Theileria parva* in cattle. *Experimental Parasitology* **36**, 278–287.
- Robson J, Pedersen V, Odeke G, Kamya E and Brown C** (1977) East coast fever immunisation trials in Uganda: field exposure of zebu cattle immunized with three isolates of *Theileria parva*. *Tropical Animal Health and Production* **9**, 219–231.
- Rogers DJ and Randolph SE** (2006) Climate change and vector-borne diseases. *Advances in Parasitology* **62**, 345–381.
- Subak S** (2003) Effects of climate on variability in Lyme disease incidence in the Northeastern United States. *American Journal of Epidemiology* **157**, 531–538.
- Swai ES, Mtui PF, Mbise AN, Kaaya E, Sanka P and Loomu PM** (2006) Prevalence of gastro intestinal parasite infections in Maasai cattle in Ngorongoro district, Tanzania. *Livestock Research for Rural Development* **18**, 18–27.
- Swai ES, Karimuribo ED, Kambarage DM and Moshy WE** (2009) A longitudinal study on morbidity and mortality in youngstock smallholder dairy cattle with special reference to tick borne infections in Tanga region, Tanzania. *Veterinary Parasitology* **160**, 34–42.
- Theiler A** (1911) *Further Investigations Into Anaplasmosis of South African Cattle*. Pretoria: Government Printer and Stationery Office.
- VanderWaal K, Gilbertson M, Okanga S, Allan BF and Craft ME** (2017) Seasonality and pathogen transmission in pastoral cattle contact networks. *Royal Society Open Science* **4**, 170808.
- Walker AR** (2011) Eradication and control of livestock ticks: biological, economic and social perspectives. *Parasitology* **138**, 945–959.
- Walker JG, Klein EY and Levin SA** (2014) Disease at the wildlife-livestock interface: Acaricide use on domestic cattle does not prevent transmission of a tick-borne pathogen with multiple hosts. *Veterinary Parasitology* **199**, 206–214.
- Wangdi K, Singhasivanon P, Silawan T, Lawpoolsri S, White NJ and Kaewkungwal J** (2010) Development of temporal modelling for forecasting and prediction of malaria infections using time-series and ARIMAX analyses: a case study in endemic districts of Bhutan. *Malaria Journal* **9**, 251.
- Wesonga FD, Orinda GO, Ngae GN and Grootenhuis J** (2006) Comparative tick counts on game, cattle and sheep on a working game ranch in Kenya. *Tropical Animal Health and Production* **38**, 35–42.
- Zieger U, Cauldwell A and Horak IG** (1998) Dynamics of free-living ixodid ticks on a game ranch in the Central Province, Zambia. *The Onderstepoort Journal of Veterinary Research* **65**, 49–59.