

More Local Volume dwarf galaxy candidates

I.D.Karachentsev¹ and V.E.Karachentseva² and K.V.Vladimirova³ and K.A.Kozyrev⁴

¹Special Astrophysical Observatory, Nizhnij Arkhiz, Zelenchukskiy region, Karachai-Cherkessian Republic, 369167, Russia, and ²Main Astronomical Observatory, National Academy of Sciences of Ukraine, Kiev, 03143, Ukraine, and ³St. Petersburg University, 199034, St. Petersburg, Russia, and ⁴Kazan (Volga Region) Federal University, 420000, Kazan, Russia

Abstract

We present the results of searching for new dwarf galaxies in the Local Volume. We found 40 satellite candidates in the double-virial-radius regions of 20 Milky Way-like and LMC-like galaxies in the southern sky using DESI Legacy Imaging Surveys, 10 of which were known but not clearly associated with the Local Volume previously. Among the 40 satellite candidates, 8 are supposed members of the NGC 6744 group and 13 are located in the vicinity of the Sombrero galaxy. Based on seven companions to the giant spiral galaxy NGC 6744 with measured radial velocities, we estimate that the total mass of the group is $M_T = (1.88 \pm 0.71) \times 10^{12} M_\odot$ and the total mass-to- K -luminosity ratio $M_T/L_K = (16.1 \pm 6.0) M_\odot/L_\odot$. We reproduce a distribution of 68 early- and late-type galaxies in the Local Volume situated around the Sombrero, noting their strong morphological segregation and also the presence of a foreground diffuse association of dwarf galaxies at 8 degrees to SE from the Sombrero.

Keywords: dwarf galaxies: halo galaxies : local volume

1. Introduction

The creation of the most complete and representative sample of galaxies in the nearest volume of the universe provides an important observational basis for testing various cosmological models on small scales. This sample, limited by a distance of ~ 10 Mpc, was systematically created over the course of recent decades. This has been facilitated by the emergence of increasingly in-depth surveys of large sky areas in optical and radio ranges. Data on distances, radial velocities, luminosity, and other characteristics of nearby galaxies are collected in the Updated Nearby Galaxy Catalogue (UNGC), (Karachentsev et al. 2013), which is regularly updated with new objects. Currently, the Local Volume (LV) galaxy database, available at <http://www.sao.ru/lv/lvgdb>, contains more than 1500 galaxies. Among them, about 90% dwarf galaxies have a luminosity lower than the luminosity of the Large Magellanic Cloud (LMC).

The most recent and significant source of population growth for the LV is DESI Legacy Imaging Surveys (Dey et al. 2019), which covers about 1/3 of the entire sky. Using data from this multi-colour survey, we have found 98 new candidates to the LV members in regions of the known nearby groups (Karachentsev and Kaisina 2022; Karachentsev et al. 2023a; Karachentseva et al. 2023), as well as in the general field in direction to the Local Void (Karachentsev et al. 2023b). Most of these new dwarf galaxies were discovered in the northern hemisphere of the sky. The appearance of the tenth version of the DESI surveys (DR10) provided us the opportunity to continue searching for nearby dwarf galaxies in

the southern sky. The results of our efforts are the subject of this article.

2. Searching for dwarf satellites around bright southern galaxies

We have not included in our program a survey of the vicinity of the very nearby galaxies: NGC 55, NGC 253, NGC 300, Cen A, and M 83, since these regions have been studied in detail by other authors using DESI survey and images obtained with large telescopes (Okamoto et al. 2024; Crnojević et al. 2019; Crosby et al. 2024a; Müller et al. 2015, 2024; Martínez-Delgado et al. 2021; Martínez-Delgado et al. 2024; Mutlu-Pakdil et al. 2022; Jones et al. 2024a; McNanna et al. 2024; Sand et al. 2024).

A list of galaxies around which we conducted our searches is presented in Table 1. Its columns contain: (1) — name of the host galaxy; (2) — equatorial coordinates of the galaxy in degrees; (3) — galaxy integral luminosity in the K -band, expressed in the Sun luminosity units; (4) — distance to the galaxy in Mpc; (5) — the method by which the distance was determined: TRGB — by the tip of the red giant branch, SN — by the luminosity of supernovae, TF — by the Tully-Fisher relationship (Tully et al. 2013) between the 21-cm line width and luminosity of a galaxy; (6) — the number of new candidates for satellites around the host galaxy that we have found; (7) — the size of the square in degrees within which the search for dwarf galaxies was conducted. The center of the square coincided with the host galaxy, and the square sides were oriented along RA and Dec.

For each host galaxy, we estimated the virial radius of its halo using the relation

$$(R_v/215 \text{ kpc}) = (M_T/10^{12} \times M_\odot)^{1/3}, \quad (1)$$

proposed by Tully (2015). Here, the total mass of the galaxy halo, M_T , was determined from its integrated K -band luminosity as

$$M_T/L_K = 32 \times M_\odot/L_\odot. \quad (2)$$

As Karachentsev and Kashibadze (2021) have shown, only $\sim 2/3$ of the group members are located within the virial radius, and the rest of the group population is distributed between R_v and the radius of the zero-velocity sphere $R_0 \simeq 3R_v$, which separates the group volume from the general expanding field of galaxies.

Searches for new companions around large galaxies have mostly been undertaken inside the virial radius of the galaxy halo (Carlsten et al. 2022). We conducted a wider search, covering a $2R_v$ square centered on the galaxy under consideration. The median luminosity of the galaxies in our sample is $\log(L_K/L_\odot) = 9.47$, which corresponds to the LMC luminosity.

When searching for new satellites around a host galaxy, we focused on low surface brightness objects with a major angular diameter $a > 20''$. Preference was given to dwarf galaxies with a hint of granular structure and a weak brightness gradient toward the center. We also took into account the environment of the satellite candidate in order to exclude, if possible, cases of association of the dwarf with a more distant group of galaxies. For each candidate for the Local Volume objects, the maximum apparent angular diameter, apparent axial ratio, and morphological type were determined.

As a result, 40 candidate companions were detected in a total survey area of 320 square degrees around 20 host galaxies. There is a noticeable trend that the number of companions decreases rapidly with decreasing luminosity of the host galaxy. Data from deep searches for satellites conducted with large telescopes (Davis et al. 2024; Carlin et al. 2024) show that there is, on average, just under one companion per a host galaxy with a luminosity similar to the LMC luminosity.

The images of 30 candidates for new members of the Local Volume that we discovered, taken from the DESI Legacy Imaging Surveys, are shown in Figs. 1–3. The first of them contains dwarf galaxies in the vicinity of the southern host galaxies of moderate (sub-Milky Way) luminosity: NGC 628, NGC 1313, NGC 3621, NGC 5068, NGC 7090 and NGC 7424. Fig 2 and Fig 3 present, respectively, dwarf objects around the high-luminosity galaxies, NGC 6744 and NGC 4594 (“Sombrero”). In addition to them, we have noted 10 more dwarf galaxies that were known earlier with ESO, PGC, KK and KKs names, but were not considered as members of the LV. Each image size in Figs. 1–3 is $2' \times 2'$. North is up, east is left.

The summary list of 40 supposed dwarf satellites is given in Table 2, which contains the following data: (1) — galaxy name; (2) — equatorial coordinates; (3, 4) — angular diameter in arc minutes and apparent axial ratio of the galaxy; (5) — apparent magnitude of the galaxy in the B -band, determined from g - and r -magnitudes in the DESI survey as $B = g + 0.313 \times (g - r) + 0.227$, or estimated visually for diffuse objects with a clump-like structure; (6) — morphological type of the dwarf galaxy: Irr — irregular, Im — Magellanic, BCD — blue compact dwarf, Sph — spheroidal, Tr — transitional between Irr and Sph. Below we note some features of the detected dwarf galaxies.

ESO 472-015. A relatively bright dwarf galaxy with a heliocentric velocity of $V_h = 655 \pm 46 \text{ km s}^{-1}$, at a projected separation of $R_p = 90 \text{ kpc}$ from the host galaxy NGC 24, which has $V_h = 553 \pm 2 \text{ km s}^{-1}$ according to HyperLEDA (Makarov et al. 2014).

PGC 01242. This is a BCD galaxy with a radial velocity of $V_h = 670 \pm 2 \text{ km s}^{-1}$. The angular size is related to a faint extended

halo, which is also clearly visible in the UV-band according to GALEX data (Martin et al. 2005).

Dw 0128-4116. An irregular dwarf with asymmetric halo.

Dw 0312-6643. This dwarf spheroidal galaxy of the low surface brightness and semi-resolved into stars seems to be a satellite of peculiar spiral galaxy NGC 1313.

Dw 1105-3041, KK 101, Dw 1128-3225. Probable satellites of spiral galaxy NGC 3621.

Dw 1231-1216. A dwarf galaxy with granular structure and faint asymmetric halo; a probable satellite of NGC 4594 (“Sombrero”).

Dw 1245-1332. A dwarf spheroidal galaxy with a radial velocity of $V_{LG} = 783 \text{ km s}^{-1}$ measured by Crosby et al. (2024b).

Dw 1253-0613. A dwarf spheroidal galaxy with granular structure; it belongs probably to a foreground diffuse association of dwarfs near DDO 148.

Dw 1301-1627. A dSph galaxy with granular structure, a probable member of association of dwarfs around DDO 161.

KK 186. A transition type (Tr) dwarf with a granular structure. Together with the four following objects in Table 2, it forms a retinue of the spiral galaxy NGC 5068.

Dw 1321-1935. A nucleated dSph, projected onto a very distant cluster of galaxies.

Dw 1851-6400. A Tr-type dwarf with a granular structure, a probable satellite of the giant spiral galaxy NGC 6744.

Dw 2140-5544. A spheroidal dwarf with a granular structure; located in a dense field of interstellar cirrus.

Dw 2256-4107. A satellite of the spiral galaxy NGC 7424, half-destroyed by the host.

3. Suite of dwarfs around the galaxy NGC 6744

The Sc spiral galaxy NGC 6744 is the most extended object in the LV. Its optical Holmberg’s linear diameter reaches 60 kpc, and its angular momentum is twice that of the Milky Way (Karachentsev and Zozulia 2023). Among NGC 6744’s neighbours, seven galaxies in the known catalogues (NGC, IC, ESO) are associated with NGC 6744, judging by their radial velocities. Searches for fainter dwarf galaxies (Karachentseva and Karachentsev 2000) and (Karachentsev et al. 2020b) led to the discovery of three and two low surface brightness probable companions, respectively. Carlsten et al. (2022) and Hunt et al. (2024) added four more objects to the list of members of the NGC 6744 group. Carlsten et al. (2022) estimated their distances from surface brightness fluctuations (sbf). Our searches for dwarf galaxies in the wider vicinity of NGC 6744 revealed eight more new candidates for this group: Dw 1844-6232, Dw 1851-6400, Dw 1851-6511, Dw 1851-6356, Dw 1851-6355, Dw 1903-6405, KKs 73, and KKs 74. The last two of these were known previously but were not included in the LV database.

A summary list of 25 supposed members of the NGC 6744 group is presented in Table 3. Its rows contain: (1) — galaxy name; (2,3) — supergalactic coordinates in degrees; (4) — morphological type; (5) — distance to the galaxy in Mpc; (6) — method by which the distance was estimated; for two galaxies (IC 4824 and ESO 141-042), the distances were determined from their radial velocities in the Numerical Action Method (NAM) model (Shaya et al. 2017; Kourkchi et al. 2020), which takes into account the local field of peculiar velocities; (7) — projected separation of the galaxy from NGC 6744 under the assumption that all members of the group are located at a radial distance of $D = 9.51 \text{ Mpc}$,

measured by Anand et al. (2021) using the TRGB; (8) — radial velocity of the galaxy relative to the center of the Local Group; (9) — the total mass of the group estimated via individual members based on their projected separation R_p and the difference in radial velocities ΔV relative to NGC 6744 from the relation

$$M_T/M_\odot = (16/\pi) \times G^{-1} \times \Delta V^2 \times R_p, \quad (3)$$

where G is the gravitational constant. This relation assumes a chaotic orientation of the satellite orbits with an average orbital eccentricity $e = 0.7$ (Karachentsev and Kashibadze 2021).

The perimeter of our search area is shown in Fig. 4. The distribution of supposed members of the NGC 6744 group is presented in supergalactic coordinates to reduce large-scale distortions due to the convergence of meridians in the {RA, Dec} system. Previously known members of the group are shown as circles, and objects we have discovered are marked with asterisks. Spheroidal dwarf galaxies without signs of star formation are highlighted with red symbols. The radial velocities of the galaxies (in km s^{-1}) are indicated by numbers. The large circle centered on NGC 6744 corresponds to the virial radius of the group, 296 kpc or 1.78° .

The distribution of galaxies shows a tendency towards the presence of substructures. The asymmetric distribution of spheroidal dwarf galaxies is noteworthy: all nine Sph objects are located on the right half of the figure. We have not found an explanation for this feature. In general, such asymmetry is not uncommon among galaxy groups. There is quite an extensive literature discussing both observational manifestations of the lopsidedness of satellite galaxy systems (Brainerd and Samuels 2020; Heesters et al. 2024) and their comparison with the results of numerical simulations of the structure of groups (Pawlowski et al. 2017; Wang et al. 2021). It is also noteworthy that the distribution of potential satellites around NGC 6744 appears to be flattened in the diagonal direction, reminiscent of the issue of satellite planes seen around the Milky Way, the Andromeda nebulae and other nearby major galaxies (Ibata et al. 2013; Pawlowski and Kroupa 2020; Müller et al. 2021; Karachentsev and Kroupa 2024). The few available velocities even appear consistent with a kinematic trend: higher velocities than the host on the top-left side, lower on the bottom-right side. Future velocity measurements of the remaining supposed members of the group could easily confirm or disprove this assumption.

As follows from the data in the last column of Table 3, the average estimate of the total mass of the NGC 6744 group based on seven satellites is $\langle M_T \rangle = (1.88 \pm 0.71) \times 10^{12} M_\odot$. With the integrated luminosity of the group $\Sigma L_K = 1.17 \times 10^{11} L_\odot$, the ratio of the total mass to the total luminosity of the group is $(16.1 \pm 6.0) M_\odot/L_\odot$, which is two times smaller than the average ratio $(31 \pm 6) M_\odot/L_\odot$ obtained for the ensemble of 25 groups in the LV (Karachentsev and Kashibadze 2021). This difference is leveled out if we consider only the LV groups with spiral host galaxies, for which the total mass-to-K-luminosity ratio is $(17.4 \pm 2.8) M_\odot/L_\odot$ according to Karachentsev and Kashibadze (2021).

4. The surroundings of NGC 4594

The galaxy NGC 4594 = M 104 (“Sombrero”) has the highest luminosity, $\log(L_K/L_\odot) = 11.32$, among the LV population. The distance to it, $D = (9.55 \pm 0.46)$ Mpc, has been determined by McQuinn et al. (2016) via TRGB method. The search for satellites around Sombrero was undertaken by various teams (Karachentsev et al. 2020b; Javanmardi et al. 2016; Karachentsev et al. 2000; Carlsten et al. 2022; Crosby et al. 2024a).

Carlsten et al. (2022) used images obtained with the 3.6 meter Canadian-French-Hawaii Telescope to discover 11 new satellites of Sombrero in a region with a radius of 0.6° or ~ 100 kpc. The belonging of these dwarfs to the Sombrero group was confirmed by measuring their distances using the surface brightness fluctuation (sbF) method. The most significant addition to the number of dwarf galaxies in the Sombrero neighborhood was given by the study of Crosby et al. (2024a). Using images obtained with the Subaru Supreme-Cam, they discovered 40 new potential satellites of Sombrero within a circle of $\sim 2^\circ$ radius. Surface photometry was performed for these galaxies and integral properties of the galaxies were determined. Of these 40 dwarfs, 27 were classified as satellites of Sombrero with high confidence.

Looking at a wider region, Karachentsev et al. (2020a), identified 15 supposed satellites of the Sombrero with measured radial velocities. Their average projected separation relative to the Sombrero is 431 kpc, and their radial velocity dispersion is 204 km s^{-1} . Applying relations (1) and (3) to this ensemble of satellites yields an estimate of the total mass of the group of $M_T = (15.5 \pm 4.9) \times 10^{12} M_\odot$ and a virial radius of $R_v = 538$ kpc or 3.23° .

Using the DR10 data of DESI Legacy Imaging Surveys, we undertook searches for new satellites of Sombrero in the region : RA= [183, 197] $^\circ$, Dec= [-5, -19] $^\circ$, and found 13 dwarf galaxies presented in Table 2. We also re-examined images of 27 dwarf galaxies from the Crosby et al. (2024a) list, identified as Sombrero satellites with high confidence. In our estimation, 25 of them look like real members of the Sombrero group. But we considered two galaxies as belonging to the distant background: UGCA 287 due to its estimated distance of $D_{TF} = 20.5$ Mpc (Karachentsev and Nasonova 2013), and the galaxy dw 1241-1008 due to its texture.

It should be noted that the Sombrero group is located near the equator of the Local Supercluster, projecting onto the Virgo Southern Extension filament (Tully 1982; Kourkchi and Tully 2017). Many galaxies in this filament have radial velocities close to that of Sombrero ($V_{LG} = 912$) km s^{-1} , but are located at distances of 15–20 Mpc, typical for members of the Virgo cluster. The presence of the rich background makes it difficult to single out Sombrero’s satellites.

We selected from the LV database (<http://www/lv/lv gdb>) the galaxies with distance estimates $D < 12$ Mpc, which are located in the square of 14° by 14° around Sombrero. This region extends over $2R_v$, encompassing the bulk of the galaxy halo. A list of 68 of these galaxies is presented in Table 4. Its columns contain: (1) — galaxy name as it indicated in UNGC; (2,3) — equatorial coordinates, J2000.0, in degrees; (4) — galaxy distance in Mpc; (5) — method used to determine the distance; (6) — radial velocity in the Local Group rest frame (in km s^{-1}); (7) — morphological type; (8) — references to the distance estimates. All radial velocities are taken from the HyperLEDA (Makarov et al. 2014) with the addition of a new radial velocity for UGCA 287 from Crosby et al. (2024b).

The distribution of the LV galaxies in the considered sky region is presented in Figure 5. The galaxies of early types (Sph, dE) and late types (Irr, Im, Tr, BCD, S) are shown by red and blue symbols, respectively. Supposed foreground galaxies with $D < 7.3$ Mpc are indicated with open circles. The dwarf galaxies we discovered are shown as blue and red asterisks according to their morphological type. The large circle of 3.23° radius corresponds to R_v . The radial velocities of galaxies located outside the dense central zone are indicated by numbers (in km s^{-1}).

The distribution of early-type dwarf galaxies shows the expected stronger concentration towards the host galaxy. The distribution of suspected satellites appears somewhat asymmetric. On the right side of this map, outside the virial circle, only one probable satellite of Sombrero has been detected. It is KKSG 27 with $V_{LG} = 1128 \text{ km s}^{-1}$.

In the lower left corner of Fig. 5, a group of foreground galaxies stands out with a typical radial velocity of $V_{LG} \sim 600 \text{ km s}^{-1}$ and a distance of $D \sim 6 \text{ Mpc}$. This association of dwarf galaxies around DDO 161 likely includes also the two spheroidal dwarfs, Dw 1301-1627 and Dw 1309-1721, we discovered. Three more new Sph dwarfs are located in the upper left part of the figure near DDO 148 ($V_{LG} = 1170 \text{ km s}^{-1}$). It remains unclear whether these galaxies are members of the Sombrero group or belong to outskirts of the Virgo cluster. Obviously, for a better understanding the structure and kinematics of the Sombrero group, it is necessary to measure more distances and radial velocities of galaxies in the considered sky region.

5. Conclusions

We searched for new dwarf companions around 20 southern galaxies of the Local Volume having distances within (4 - 11) Mpc and $\log(L_K/L_\odot)$ luminosities in the range of (8.1–11.3). For half of the host galaxies, no new companions were detected in an area inside of two virial radii. We found eight new companion candidates for the spiral galaxy NGC 6744, which is the most extended galaxy in the LV. The population of NGC 6744 group currently includes 25 candidate members. Based on seven supposed satellites of NGC 6744 with measured radial velocities, the total mass of the group is estimated to be $(1.88 \pm 0.71) \times 10^{12} M_\odot$. The accuracy of determining the total mass of this group may be noticeably increased when measuring radial velocities of its other members.

Our search for dwarf galaxies in a 14° by 14° square centered on M 104 (Sombrero) has resulted in 13 new dwarfs. Most of these are likely associated with the Sombrero itself and with a foreground diffuse group of dwarf galaxies around DDO 161. We also rediscovered 25 of the 27 objects recently found by Crosby et al. (2024b) as members of the Sombrero group with high confidence. A deep 21-cm survey of this region with sufficiently high angular resolution, like the Wallaby survey (Koribalski et al. 2020), could elucidate the structure and kinematics of the galaxy ensemble in this intricate sky region adjacent to the Virgo Cluster.

Acknowledgements

We are grateful to an anonymous referee for constructive comments, in particular for drawing our attention to the peculiarity of the configuration of satellites around the NGC 6744. This work has made use of DESI Legacy Imaging Surveys data, the HyperLEDA, and the revised version of the Local Volume galaxies database.

Funding Statement. This work was supported by the Russian Science Foundation grant 24-12-00277.

Data Availability Statement. The data on which this work is based is publicly available and is detailed in the corresponding tables of the manuscript.

References

- Anand, G. S., Rizzi, L., Tully, R. B., et al. 2021, The Extragalactic Distance Database: The Color-Magnitude Diagrams/Tip of the Red Giant Branch Distance Catalog. *Astrophysical Journal*, 162(2), 80.
- Brainerd, T. G. & Samuels, A. 2020, Lopsided Satellite Distributions around Isolated Host Galaxies. *Astrophysical Journal L*, 898(1), L15.
- Carlin, J. L., Sand, D. J., Mutlu-Pakdil, B., et al. 2024, A census of dwarf galaxy satellites around LMC-mass galaxy NGC 2403. *arXiv e-prints*, arXiv:2409.17437.
- Carlsten, S. G., Greco, J. P., Beaton, R. L., & Greene, J. E. 2020, Wide-field Survey of Dwarf Satellite Systems around 10 Hosts in the Local Volume. *Astrophysical Journal*, 891(2), 144.
- Carlsten, S. G., Greene, J. E., Beaton, R. L., Danieli, S., & Greco, J. P., et al. 2022, The Exploration of Local Volume Satellites (ELVES) Survey: A Nearly Volume-limited Sample of Nearby Dwarf Satellite Systems. *Astrophysical Journal*, 933(1), 47.
- Crnjević, D., Sand, D. J., Bennet, P., et al. 2019, The Faint End of the Centaurus A Satellite Luminosity Function. *Astrophysical Journal*, 872(1), 80.
- Crosby, E., Jerjen, H., Müller, O., Pawlowski, M. S., Mateo, M., & Lelli, F. 2024a, A New dwarf galaxy candidates in the sphere of influence of the Sombrero galaxy. *Monthly Notices of the Royal Astronomical Society*, 527a(3), 9118–9131.
- Crosby, E., Mateo, M., Escala, I., Jerjen, H., Müller, O., & Pawlowski, M. S. 2024b, IFUM Integrated Field Spectroscopy of Ten M104 Satellite Galaxy Candidates. *arXiv e-prints*, arXiv:2412.02060.
- Davis, A. B., Garling, C. T., Nierenberg, A. M., et al. 2024, The LBT Satellites of Nearby Galaxies Survey (LBT-SONG): The Diffuse Satellite Population of Local Volume Hosts. *arXiv e-prints*, arXiv:2409.03999.
- Dey, A., Schlegel, D. J., Lang, D., Blum, R., et al. 2019, Overview of the DESI Legacy Imaging Surveys. *Astronomical Journal*, 157(5), 168.
- Heesters, N., Jerjen, H., Müller, O., Pawlowski, M. S., & Jamie Kanehisa, K. 2024, Lopsided distribution of MATLAS and ELVES dwarf satellite systems around isolated host galaxies. *Astronomy & Astrophysics*, 690, A110.
- Hunt, L. K., Annibali, F., Cuillandre, J. C., Ferguson, A. M. N., et al. 2024, Euclid: Early Release Observations – Deep anatomy of nearby galaxies. *arXiv e-prints*, arXiv:2405.13499.
- Ibata, R. A., Lewis, G. F., Conn, A. R., et al. 2013, A vast, thin plane of co-rotating dwarf galaxies orbiting the Andromeda galaxy. *Nature*, 493(7430), 62–65.
- Javanmardi, B., Martinez-Delgado, D., Kroupa, P., et al. 2016, DGSAT: Dwarf Galaxy Survey with Amateur Telescopes. I. Discovery of low surface brightness systems around nearby spiral galaxies. *Astronomy & Astrophysics*, 588, A89.
- Jones, M. G., Sand, D. J., Karunakaran, A., et al. 2024a, Gas and Star Formation in Satellites of Milky Way Analogs. *Astrophysical Journal*, 966a(1), 93.
- Jones, M. G., Sand, D. J., Mutlu-Pakdil, B., et al. 2024b, Corvus A: A Low-mass, Isolated Galaxy at 3.5 Mpc. *Astrophysical Journal Lett.*, 971b(2), L37.
- Karachentsev, I. & Kashibadze, O. 2021, Tracing the local volume galaxy halo-to-stellar mass ratio with satellite kinematics. *Astronomische Nachrichten*, 342(999), 999–1023.
- Karachentsev, I. D. & Kaisina, E. I. 2022, New Local Volume Dwarf Galaxy Candidates from the DESI Legacy Imaging Surveys. *Astrophysical Bulletin*, 77(4), 372–387.
- Karachentsev, I. D., Kaisina, E. I., & Karachentseva, V. E. 2023a, The pride of lions around Messier 105. *Monthly Notices of the Royal Astronomical Society*, 521a(1), 840–849.
- Karachentsev, I. D., Karachentseva, V. E., Kaisin, S. S., & Kaisina, E. I. 2023b, A Search for New Dwarf Galaxies Outside the Nearby Groups. *Astrophysics*, 66b(4), 441–451.
- Karachentsev, I. D., Karachentseva, V. E., Suchkov, A. A., & Grebel, E. K. 2000, Dwarf galaxy candidates found on the SERC EJ sky survey. *Astronomy & Astrophysics Suppl.*, 145, 415–423.
- Karachentsev, I. D. & Kroupa, P. 2024, A co-rotating gas and satellite structure

- around the interacting galaxy pair NGC 4490/85. *Monthly Notices of the Royal Astronomical Society*, 528(2), 2805–2811.
- Karachentsev, I. D., Makarov, D. I., & Kaisina, E. I. 2013, Updated Nearby Galaxy Catalog. *Astronomical Journal*, 145(4), 101.
- Karachentsev, I. D., Makarova, L. N., Brent Tully, R., Anand, G. S., Rizzi, L., & Shaya, E. J. 2020,a Distance and mass of the M 104 (Sombrero) group. *Astronomy & Astrophysics*, 643a, A124.
- Karachentsev, I. D., Makarova, L. N., Tully, R. B., Rizzi, L., & Shaya, E. J. 2018, TRGB Distances to Galaxies in Front of the Virgo Cluster. *Astrophysical Journal*, 858(1), 62.
- Karachentsev, I. D. & Nasonova, O. G. 2013, Intense look at Virgo Southern Extension. *Monthly Notices of the Royal Astronomical Society*, 429(3), 2677–2686.
- Karachentsev, I. D., Riepe, P., & Zilch, T. 2020,b A Dozen new Dwarf Galaxy Candidates in the Local Volume. *Astrophysics*, 63b(1), 5–14.
- Karachentsev, I. D. & Zozulia, V. D. 2023, Orientation of the spins of galaxies in the local volume. *Monthly Notices of the Royal Astronomical Society*, 522(3), 4740–4747.
- Karachentseva, V. E. & Karachentsev, I. D. 2000, A supplementary list of southern nearby dwarf galaxy candidates. *Astronomy & Astrophysics Suppl.*, 146, 359–363.
- Karachentseva, V. E., Karachentsev, I. D., Kaisina, E. I., & Kaisin, S. S. 2023, The M 101 galaxy group as a node in a nearby cosmic filament. *Astronomy & Astrophysics*, 678, A16.
- Kashibadze, O. G., Karachentsev, I. D., & Karachentseva, V. E. 2018, Surveying the Local Supercluster Plane. *Astrophysical Bulletin*, 73(2), 124–141.
- Koribalski, B. S., Staveley-Smith, L., Westmeier, T., et al. 2020, WALLABY – an SKA Pathfinder H I survey. *Astrophysics and Space Science*, 365(7), 118.
- Kourkchi, E., Courtois, H. M., Graziani, R., Hoffman, Y., Pomarède, D., Shaya, E. J., & Tully, R. B. 2020, Cosmicflows-3: Two Distance-Velocity Calculators. *Astronomical Journal*, 159(2), 67.
- Kourkchi, E. & Tully, R. B. 2017, Galaxy Groups Within 3500 km s⁻¹. *Astrophysical Journal*, 843(1), 16.
- Makarov, D., Prugniel, P., Terekhova, N., Courtois, H., & Vauglin, I. 2014, HyperLEDA. III. The catalogue of extragalactic distances. *Astronomy & Astrophysics*, 570, A13.
- Martin, D. C., Fanson, J., Schiminovich, D., et al. 2005, The Galaxy Evolution Explorer: A Space Ultraviolet Survey Mission. *Astrophysical Journal Lett.*, 619(1), L1–L6.
- Martínez-Delgado, D., Makarov, D., Javanmardi, B., et al. 2021, Tracing satellite planes in the Sculptor group. I. Discovery of three faint dwarf galaxies around NGC 253. *Astronomy & Astrophysics*, 652, A48.
- Martínez-Delgado, D., Stein, M., Pawlowski, M. S., Makarov, D., Makarova, L., Donatiello, G., & Lang, D. 2024, Tracing satellite planes in the Sculptor group: II. Discovery of five faint dwarf galaxies in the DESI Legacy Survey. *arXiv e-prints*, arXiv:2405.03769.
- McNanna, M., Bechtol, K., Mau, S., et al. 2024, A Search for Faint Resolved Galaxies Beyond the Milky Way in DES Year 6: A New Faint, Diffuse Dwarf Satellite of NGC 55. *Astrophysical Journal*, 961(1), 126.
- McQuinn, K. B. W., Skillman, E. D., Dolphin, A. E., Berg, D., & Kennicutt, R. 2016, The Distance to M104. *Astronomical Journal*, 152(5), 144.
- Müller, O., Jerjen, H., & Binggeli, B. 2015, New dwarf galaxy candidates in the Centaurus group. *Astronomy & Astrophysics*, 583, A79.
- Müller, O., Pawlowski, M. S., Lelli, F., et al. 2021, The coherent motion of Cen A dwarf satellite galaxies remains a challenge for Λ CDM cosmology. *Astronomy & Astrophysics*, 645, L5.
- Müller, O., Pawlowski, M. S., Revaz, Y., Venhola, A., Rejkuba, M., Hilker, M., & Lutz, K. 2024, A too-many-dwarf-galaxy-satellites problem in the M 83 group. *Astronomy & Astrophysics*, 684, L6.
- Mutlu-Pakdil, B., Sand, D. J., Crnojević, D., et al. 2022, Hubble Space Telescope Observations of NGC 253 Dwarf Satellites: Three Ultra-faint Dwarf Galaxies. *Astrophysical Journal*, 926(1), 77.
- Okamoto, S., Ferguson, A. M. N., Arimoto, N., et al. 2024, Uncovering the Ghostly Remains of an Extremely Diffuse Satellite in the Remote Halo of NGC 253. *Astrophysical Journal Lett.*, 967(2), L24.
- Pawlowski, M. S., Ibata, R. A., & Bullock, J. S. 2017, The Lopsidedness of Satellite Galaxy Systems in Λ CDM Simulations. *Astrophysical Journal*, 850(2), 132.
- Pawlowski, M. S. & Kroupa, P. 2020, The Milky Way's disc of classical satellite galaxies in light of Gaia DR2. *Monthly Notices of the Royal Astronomical Society*, 491(2), 3042–3059.
- Sand, D. J., Mutlu-Pakdil, B., Jones, M. G., et al. 2024, Three Quenched, Faint Dwarf Galaxies in the Direction of NGC 300: New Probes of Reionization and Internal Feedback. *arXiv e-prints*, arXiv:2409.16345.
- Shaya, E. J., Tully, R. B., Hoffman, Y., & Pomarède, D. 2017, Action Dynamics of the Local Supercluster. *Astrophysical Journal*, 850(2), 207.
- Tully, R. B. 1982, The Local Supercluster. *Astrophysical Journal*, 257, 389–422.
- Tully, R. B. 2015, Galaxy Groups. *Astronomical Journal*, 149(2), 54.
- Tully, R. B., Courtois, H. M., Dolphin, A. E., et al. 2013, Cosmicflows-2: The Data. *Astronomical Journal*, 146(4), 86.
- Wang, P., Libeskind, N. I., Pawlowski, M. S., et al. 2021, The Lopsided Distribution of Satellites of Isolated Central Galaxies. *Astrophysical Journal*, 914(2), 78.




Figure 1. Reproduction of images of 11 nearby dwarf galaxy candidates from the DESI Legacy Imaging Surveys, found near the southern LV galaxies of sub-Milky Way luminosity. Each image size is $2' \times 2'$. North is at the top, east is on the left.




Figure 2. Reproduction of images of six nearby dwarf galaxy candidates from the DESI Legacy Imaging Surveys, found near the giant galaxy NGC 6744. Each image size is $2' \times 2'$. North is at the top, east is on the left.




Figure 3. Reproduction of images of 13 nearby dwarfs from the DESI Legacy Imaging Surveys, found in the vicinity of the Sombrero galaxy. Each image size is $2' \times 2'$. North is at the top, east is on the left.




Figure 4. The galaxy group region around NGC 6744 in supergalactic coordinates. Previously known members of the group are shown as circles, new candidate members are shown as asterisks. Spheroidal dwarf galaxies are indicated by red symbols. The numbers indicate the radial velocity of the galaxies in km s^{-1} . The circle centered on NGC 6744 corresponds to the virial radius of the group of 296 kpc.




Figure 5. Distribution of the LV dwarfs around the Sombrero galaxy. The galaxies of early types (Sph, dE) and late types (Irr, Im, Tr, BCD, Sm, Sdm) are shown by red and blue symbols, respectively. Supposed foreground galaxies with $D < 7.3$ Mpc are indicated with open circles. The dwarf galaxies we discovered are shown as blue and red asterisks according to their morphological type.

Table 2. New candidate dwarf galaxies in the Local Volume .

Name	RA (2000.0) Dec	a'	b/a	B	Type
(1)	(2)	(3)	(4)	(5)	(6)
ESO 472-015	00 06 48.4 –24 56 41	1.53	0.56	16.24	Im
PGC 73354	00 17 01.4 –23 42 30	0.77	0.57	16.89	BCD
PGC 01242	00 19 11.5 –22 40 06	3.06	0.51	15.38	BCD
Dw 0128-4116	01 28 43.0 –41 16 34	0.68	0.52	17.68	Irr
Dw 0312-6643	03 12 21.1 –66 43 16	0.64	0.63	21.0	Sph
Dw 1105-3041	11 05 47.5 –30 41 19	0.77	0.84	20.0	Sph
KK 101	11 16 44.6 –32 39 04	1.06	0.72	17.56	Sph
Dw 1128-3225	11 28 58.1 –32 25 23	0.47	0.55	20.5	Irr
Dw 1231-1216	12 31 06.0 –12 16 12	0.72	0.73	18.7	Tr
Dw 1244-0842	12 44 01.9 –08 42 47	0.56	0.58	19.6	Sph
Dw 1244-0951	12 44 05.0 –09 51 00	0.24	0.75	21.8	Sph
Dw 1245-1332	12 45 54.5 –13 32 31	0.87	0.70	18.8	Sph
Dw 1246-0533	12 46 07.2 –05 33 25	0.75	0.89	18.8	Sph
Dw 1249-0928	12 49 15.1 –09 28 08	0.51	0.60	21.5	Tr
Dw 1252-0506	12 52 20.9 –05 06 18	0.80	0.79	19.0	Sph
Dw 1253-0613	12 53 16.1 –06 13 55	0.91	0.95	18.6	Sph
Dw 1258-0948	12 58 24.0 –09 48 54	0.53	0.68	21.4	Sph
Dw 1259-1429	12 59 01.4 –14 29 02	0.66	0.68	19.1	Sph
Dw 1259-1407	12 59 54.2 –14 07 23	0.61	0.90	20.7	Irr
Dw 1301-1627	13 01 28.1 –16 27 40	0.76	0.61	18.9	Sph
Dw 1309-1721	13 09 18.0 –17 21 54	0.52	0.78	19.9	Sph
KK 186	13 09 45.6 –23 32 35	1.25	0.65	17.40	Tr
Dw 1314-1817	13 14 35.0 –18 17 20	0.83	0.81	18.00	Irr
Dw 1321-2202	13 21 04.1 –22 01 59	0.84	0.74	18.06	Sph
Dw 1321-1935	13 21 59.0 –19 35 31	0.89	0.94	18.62	Sph
Dw 1324-2017	13 24 54.2 –20 17 53	0.41	0.70	20.5	Irr
Dw 1844-6232	18 44 54.5 –62 32 49	0.46	0.86	20.04	Sph
Dw 1851-6400	18 51 10.3 –64 00 29	0.53	0.90	18.28	Tr
Dw 1851-6511	18 51 39.1 –65 11 38	0.56	0.93	21.5	Sph
Dw 1851-6356	18 51 42.2 –63 56 06	0.28	0.85	21.8	Sph
Dw 1851-6355	18 51 43.0 –63 55 12	0.38	0.80	21.0	Sph
Dw 1903-6405	19 03 39.4 –64 05 56	0.32	0.81	21.7	Sph
KKs 73	19 21 46.2 –60 41 00	1.24	0.64	16.90	Irr
KKs 74	19 27 43.2 –61 00 38	1.11	0.41	17.60	Tr
Dw 2140-5544	21 40 59.3 –55 44 49	0.88	0.61	18.90	Sph
ESO 346-007	22 53 23.7 –38 47 59	1.60	0.68	15.06	BCD
Dw 2256-4107	22 56 13.9 –41 07 43	3.00	0.38	17.0	Tr
Dw 2256-4250	22 56 38.2 –42 50 38	1.42	0.68	17.2	Irr
ESO 290-028	22 57 09.2 –42 48 18	5.63	0.10	14.52	Sdm
PGC 580285	23 10 38.1 –40 59 11	0.83	0.76	17.51	Irr

Table 1. Southern host galaxies in the Local Volume.

Name	RA, Dec	$\log(L_K/L_\odot)$	D	meth	N	deg
(1)	(2)	(3)	(4)	(5)	(6)	(7)
NGC 0024	02.48–24.96	9.48	7.31	TRGB	1	3.1
NGC 0045	03.52–23.18	9.33	6.64	TRGB	2	3.0
NGC 0625	23.77–41.44	8.96	4.02	TRGB	1	3.7
ESO 300-014	47.41–41.03	9.30	9.80	TF	0	2.0
NGC 1313	49.56–66.50	9.57	4.31	TRGB	1	5.6
NGC 1311	50.03–52.18	8.43	5.55	TRGB	0	1.8
NGC 1592	67.43–27.41	8.17	9.10	TF	0	0.9
NGC 1637	70.37–02.86	10.07	9.29	SN	0	3.8
NGC 1800	76.61–31.95	9.04	8.00	TF	0	2.0
NGC 2188	92.54–34.11	9.37	8.39	TRGB	0	2.5
NGC 3037	147.85–27.01	8.90	10.28	NAM	0	1.4
NGC 3621	169.57–32.81	10.34	6.64	TRGB	3	6.5
NGC 4594	190.00–11.62	11.32	9.55	TRGB	13	7.0
NGC 5068	199.73–21.04	9.73	5.15	TRGB	5	5.3
NGC 6744	287.44–63.86	10.91	9.51	TRGB	8	7.1
IC 5052	313.03–69.20	9.27	5.50	TRGB	0	3.5
NGC 7090	324.12–54.56	10.03	9.51	TRGB	1	3.6
NGC 7424	344.33–41.07	10.08	9.86	TF	5	3.6
IC 5332	353.62–36.10	9.74	9.01	TRGB	0	3.1
NGC 7713	354.06–37.94	9.46	8.05	TRGB	0	2.8

Table 3. Suggested members of the NGC 6744 galaxy group .

Name	SGL	SGB	Type	D	meth	R_p	V_{LG}	M_T
	deg	deg		Mpc		kpc	km s ⁻¹	10 ¹² M_{\odot}
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
Dw 1844-6232	205.261	11.719	Sph	9.51	mem	509	-	-
NGC 6684	205.807	9.110	Sa	10.23	TRGB	435	720	0.10
Dw 1851-6400	206.022	10.280	Tr	9.51	mem	345	-	-
Dw 1851-6356	206.080	10.353	Sph	9.51	mem	335	-	-
Dw 1851-6355	206.082	10.369	Sph	9.51	mem	335	-	-
Dw 1851-6511	206.093	9.095	Sph	9.51	mem	396	-	-
ESO 104-022	206.523	9.480	Irr	8.95	TRGB	302	654	0.96
dw 1859-64	206.943	10.246	Irr	9.51	mem	193	-	-
dw 1901-63	207.192	11.021	Sph	9.51	mem	184	-	-
Dw 1903-6405	207.409	10.175	Sph	9.51	mem	120	-	-
N 6744dwTBGa	207.700	10.990	Sph	7.61	sbf	121	-	-
KKs 70	207.784	10.292	Irr	7.31	sbf	55	-	-
dw 1907-63	207.842	10.549	Sph	7.99	sbf	51	-	-
KKs 71	207.993	10.515	Irr	9.51	mem	28	-	-
EDwC1	208.042	10.558	Sph	9.51	mem	31	-	-
NGC 6744	208.101	10.382	Sc	9.51	TRGB	0	706	0
ESO 104-044	208.254	10.010	Irr	9.73	TRGB	66	614	0.65
KKs 72	208.394	10.364	Irr	9.51	mem	63	-	-
N 6744dwTBGb	208.453	10.553	Tr	9.51	mem	65	-	-
IC 4824	208.630	12.119	Im	10.30	NAM	301	811	3.91
AM1909-613	208.722	12.287	Sm	9.51	mem	333	821	5.20
ESO 141-042	208.955	11.819	Sm	9.70	NAM	332	769	1.55
KKs 73	209.811	13.419	Irr	9.51	mem	578	-	-
KKs 74	210.507	13.002	Tr	9.51	mem	590	-	-
IC 4870	210.838	8.086	BCD	8.51	TRGB	592	740	0.81

Table 4. LV galaxies within RA=[183.0 – 197.0], Dec=[–5.0, –19.0]

Name	RA, °	Dec, °	<i>D</i> , Mpc	meth	<i>V</i> _{LG} km s ^{–1}	Type	Ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Corvus A	183.690	–16.397	3.48	TRGB	297	Tr	[1]
KKSG 27	185.524	–9.800	9.08	bTF	1128	Im	[2]
dw 1234-1238	188.704	–12.640	9.55	mem	–	Sph	–
dw 1234-1142	188.704	–11.707	9.55	mem	–	Tr	–
LV J1235-1104	188.914	–11.067	10.00	TF	1003	BCD	[3]
dw 1235-1216	188.929	–12.273	9.55	mem	–	Irr	–
dw 1235-1155	188.967	–11.931	9.55	mem	–	Sph	–
dw 1237-1125	189.298	–11.433	7.51	sbf	–	dE	[4]
dw 1237-1006	189.292	–10.116	9.55	mem	–	Sph	–
KKSG 29	189.309	–10.498	9.82	TRGB	562	Irr	[5]
KKSG 30	189.400	–8.867	9.73	TRGB	917	Irr	[6]
KKSG 31	189.640	–10.490	9.55	mem	–	Sph	–
dw 1238-1043	189.675	–10.725	9.55	mem	–	Sph	–
dw 1238-1102	189.743	–11.04	9.55	mem	–	Irr	–
dw 1239-1227	189.788	–12.454	9.55	mem	–	Sph	–
dw 1239-1159	189.788	–11.987	11.33	sbf	–	Sph	[4]
NGC 4594DW1	189.814	–11.719	9.42	sbf	1171	dE	[7]
dw 1239-1154	189.842	–11.907	9.55	mem	–	Sph	–
dw 1239-1240	189.875	–12.675	9.55	mem	–	Sph	–
NGC 4594-DGSAT-3	189.887	–11.227	7.87	sbf	–	Sph	[4]
dw 1239-1026	189.962	–10.438	9.55	mem	–	Irr	–
Sombrero DwA	189.965	–11.341	9.71	sbf	–	Sph	[4]
KKSG 32	189.979	–11.747	9.00	sbf	–	Sph	[4]
NGC 4594	189.996	–11.623	9.55	TRGB	894	S0a	[8]
SUCD 1	190.013	–11.668	9.55	mem	1109	dE	–
KKSG 33	190.037	–12.365	9.55	mem	–	Sph	–
dw 1240-1118	190.039	–11.314	8.79	sbf	–	dE	[4]
NGC 4597	190.054	–5.799	10.10	TF	912	Sm	[9]
dw 1240-1140	190.073	–11.679	9.55	mem	1097	dE	–
dw 1241-1131	190.262	–11.529	9.55	mem	–	Sph	–
Sombrero DwB	190.300	–11.892	11.19	sbf	–	Sph	[4]
KKSG 34	190.329	–11.928	9.02	sbf	–	Sph	[4]
dw 1241-1055	190.408	–10.926	9.55	mem	–	Sph	–
dw 1241-1234	190.450	–12.570	9.55	mem	–	Sph	–
dw 1242-1116	190.683	–11.274	9.55	mem	–	Sph	–
dw 1242-1309	190.688	–13.166	9.55	mem	–	Sph	–
PGC 042730	190.704	–12.391	9.55	mem	828	dE	–
dw 1242-1107	190.733	–11.128	9.55	mem	–	Sph	–
dw 1242-1010	190.738	–10.169	9.55	mem	–	Irr	–

Table 4. LV galaxies within RA=[183.0 – 197.0], Dec=[–5.0, –19.0]

Name	RA, °	Dec, °	<i>D</i> , Mpc	meth	<i>V</i> _{LG} km s ^{–1}	Type	Ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
Dw 1244-0842	191.008	–8.713	9.55	mem	–	Sph	–
dw 1244-1043	191.062	–10.722	9.55	mem	–	Sph	–
dw 1244-1246	191.071	–12.773	9.55	mem	–	Sph	–
dw 1244-1135	191.125	–11.585	9.55	mem	–	Irr	–
dw 1244-1238	191.225	–12.636	9.55	mem	–	Sph	–
dw 1245-1041	191.279	–10.699	9.55	mem	–	Tr	–
Dw 1245-1332	191.477	–13.542	9.55	mem	783	Sph	–
dw 1246-1240	191.558	–12.679	9.55	mem	–	Sph	–
dw 1246-1108	191.679	–11.138	9.55	mem	–	Sph	–
dw 1246-1142	191.742	–11.707	9.55	mem	–	Sph	–
Dw 1247-0824	191.854	–8.408	9.55	mem	1036	BCD	–
KKSG 37	192.004	–12.655	9.55	mem	–	Sph	–
Dw 1248-0915	192.160	–9.256	9.55	mem	–	Sph	–
DDO 148	192.180	–5.254	9.00	TF	1170	Sm	[9]
NGC 4700	192.282	–11.411	7.30	TF	1222	Sdm	[9]
PGC 974136	192.498	–10.757	9.55	mem	–	Im	–
Dw 1252-0904	193.013	–9.075	9.55	mem	–	Irr	–
KKSG 38	193.382	–5.928	10.40	TF	953	Irr	[9]
KKSG 39	193.425	–6.084	6.80	bTF	1303	Irr	[2]
DDO 153	193.490	–12.1086	9.84	TRGB	731	Im	[6]
PGC 963198	193.691	–11.661	6.15	TF	672	Im	[2]
Dw 1258-0948	194.600	–9.815	9.55	mem	–	Sph	–
MCG-02-33-75	194.618	–10.577	8.70	TF	1087	Sm	[9]
Dw 1259-1735	194.918	–17.596	6.03	mem	–	Irr	–
KK 176	194.985	–19.413	7.28	TRGB	618	Irr	[5]
UGCA 319	195.560	–17.238	5.75	TRGB	555	Irr	[5]
DDO 161	195.820	–17.423	6.03	TRGB	545	Sm	[5]
PGC 886203	196.266	–17.258	5.61	NAM	547	Irr	[2]
MCG-03-34-2	196.986	–16.689	7.90	TF	765	BCD	[9]

Notes: [1] — (Jones et al. 2024b); [2] — present paper;
 [3] — (Kashibadze et al. 2018); [4] — (Carlsten et al. 2022);
 [5] — (Karachentsev et al. 2018); [6] — (Karachentsev et al. 2020a);
 [7] — (Carlsten et al. 2020); [8] — (McQuinn et al. 2016);
 [9] — (Karachentsev and Nasonova 2013);