

particularly when compared to the overarching influence of the general factor.

This work was supported by the ÚNKP-22-2 New National Excellence Program of the Ministry for Culture and Innovation from the source of the National Research, Development and Innovation Fund (grant number ÚNKP-22-2-I-ELTE-854).

Disclosure of Interest: None Declared

O0078

Associations between psychiatric diagnoses in parents and psychiatric, behavioral, psychosocial outcomes in their offspring: a Swedish population-based register study

M. Zhou^{1*}, C. T. Lageborn¹, A. Sjölander¹, H. Larsson¹, B. D'Onofrio², M. Landén¹, P. Lichtenstein¹ and E. Pettersson¹

¹Karolinska Institutet, Stockholm, Sweden and ²Indiana University, Bloomington, United States

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.202

Introduction: Children with parents with psychiatric diagnoses have an increased probability for not only the same condition as their parent, but also for other conditions and behavioral and psychosocial problems. Whereas many studies have focused on parental severe mental illness due to their significant impairment, less attention has been paid to more common disorders despite their higher prevalence. In addition, because most past research only included one exposure or one outcome at a time, it remains difficult to examine and compare broad patterns of intergenerational transmission.

Objectives: To examine associations between six parental psychiatric diagnoses in parents, and a broad range of psychiatric diagnoses, psychotropic medications, criminality, suicide, violent victimization, accidents, and school and labor performance in their offspring.

Methods: Based on Swedish national registers, we linked all individuals born in Sweden between 1970 and 2000 to their biological parents ($N = 3\,286\,293$). We used a matched cohort design, analyzed with stratified Cox regression and conditional logistic regressions to examine associations between six psychiatric diagnoses in the parents, and 32 outcomes in their offspring. All exposed and unexposed children were followed from their date of birth to the date of emigration from Sweden, the death, or 31 December 2013 when the offspring were 14-44 years old.

Results: In terms of absolute risk, most children who had parents with psychiatric diagnoses were not diagnosed in specialist care themselves, as the proportion of having any of the 16 types of psychiatric conditions ranged from 22.17% (exposed to parental depression) to 25.05% (exposed to parental drug-related disorders) at the end of follow-up. Nevertheless, in terms of relative risk, all six parental psychiatric diagnoses increased the probability of all 32 outcomes in their offspring, with the Hazard Ratio ranging from 1.04 to 8.91 for time-to-event outcomes, and the Odds Ratio ranging from 1.29 to 3.36 for binary outcomes. Some specificities were observed for parental psychotic and substance misuse diagnoses, which strongly predicted offspring psychotic-like and externalizing-related outcomes, respectively.

Conclusions: The intergenerational transmission of parental psychiatric conditions appeared largely transdiagnostic, even for non-psychiatric outcomes in offspring. Given the broad spectrum of associations with the outcomes, service providers (e.g., psychiatrists, teachers, and social workers) should consider clients' broader psychiatric family history when predicting prognosis and planning interventions/treatment.

Disclosure of Interest: None Declared

O0079

The Prevalence of Attention Deficit Hyperactivity Disorder in Children and Adolescents: An Umbrella Review of Global Evidence

G. Ayano¹, L. Tsegay², Y. Gizachew³, S. Demelash⁴ and R. Alati^{1*}

¹Curtin University, Perth, Australia; ²Aksum University, Aksum; ³Bethel Medical College, Addis Ababa, Ethiopia and ⁴Maastricht University, Maastricht, Netherlands

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.203

Introduction: From recent epidemiological studies to emerging epidemiological evidence, it becomes evident that numerous primary studies have investigated the prevalence of ADHD in children and adolescents. Additionally, several systematic reviews and meta-analyses have explored this subject. The objective of this umbrella review is to offer a robust synthesis of evidence derived from these systematic reviews and meta-analyses

Objectives: To conduct a comprehensive umbrella review that synthesizes emerging epidemiological evidence regarding the prevalence of ADHD in children and adolescents, drawing insights from numerous primary studies as well as systematic reviews and meta-analyses.

Methods: We conducted a systematic search across multiple databases, including PubMed, Web of Science, PsychINFO, and Scopus, to identify relevant studies. The study was preregistered with PROSPERO (registration number: CRD42023389704). To assess the quality of these studies, we utilized the Measurement Tool to Assess Systematic Reviews (AMSTAR). We employed an inverse variance-weighted random-effects meta-analysis to combine prevalence estimates from the included studies.

Results: The final analysis incorporated thirteen meta-analytic systematic reviews, encompassing 588 primary studies and a total of 3,277,590 participants. A random-effects meta-analysis of these studies revealed that the global prevalence of ADHD in children and adolescents stood at 8.0% (95% CI: 6.0%–10%). Notably, the prevalence estimate was twice as high in boys (10%) compared to girls (5%). Among the three subtypes of ADHD, the inattentive type (ADHD-I) emerged as the most prevalent, followed by the hyperactive type (ADHD-HI) and the combined type (ADHD-C).

Conclusions: The comprehensive umbrella review findings emphasize the high prevalence of ADHD in children and adolescents, with a notable gender disparity, wherein boys are twice as likely to be affected compared to girls. These results underscore the urgency of prioritizing prevention, early identification, and treatment strategies for ADHD in children and adolescents.

Disclosure of Interest: None Declared