

cambridge.org/raf

## Research Paper

**Cite this article:** dos Reis JC *et al* (2020).Assessing the economic viability of integrated crop–livestock systems in Mato Grosso, Brazil. *Renewable Agriculture and Food Systems* **35**, 631–642. <https://doi.org/10.1017/S1742170519000280>

Received: 11 November 2018

Revised: 27 April 2019

Accepted: 28 June 2019

First published online: 11 September 2019

**Key words:**


Amazon; cattle; Cerrado; land use; soy; sustainable agriculture

**JEL:**

Q00; Q10; Q14

**Author for correspondence:**

Júlio César dos Reis,

E-mail: [julio.reis@embrapa.br](mailto:julio.reis@embrapa.br)Júlio César dos Reis<sup>1</sup> , Mariana Y. T. Kamoi<sup>2</sup>, Daniel Latorraca<sup>3</sup>, Rafael F. F. Chen<sup>3</sup>, Miqueias Michetti<sup>3</sup>, Flávio Jesus Wruck<sup>1</sup>, Rachael D. Garrett<sup>4</sup>, Judson Ferreira Valentim<sup>5</sup>, Renato de Aragão Ribeiro Rodrigues<sup>6</sup> and Saulo Rodrigues-Filho<sup>7</sup><sup>1</sup>Embrapa Agrosilvopastoral, Sinop, MT, Brazil; <sup>2</sup>Network Technology Transfer - ILPF, Brasília, DF, Brazil; <sup>3</sup>Mato Grosso Institute of Agriculture Economics (IMEA), Cuiabá, MT, Brazil; <sup>4</sup>Departments of Environmental System Science and Humanities, Social, and Political Science, ETH Zürich, 8092 Zürich, Switzerland; <sup>5</sup>Embrapa Acre, Rio Branco, AC, Brazil; <sup>6</sup>Embrapa Soils, Rio de Janeiro, RJ, Brazil and <sup>7</sup>Sustainable Development Center, University of Brasília, DF, Brazil**Abstract**

Population growth and rising incomes have led to increasing global demand for meat products. Meeting this demand without converting remaining natural ecosystems or further degrading ecosystems is one of the largest global sustainability challenges. A critical step to overcoming this challenge is to increase the productivity of livestock grazing systems, which occupy the largest land area of any type of agriculture globally. Integrated crop–livestock systems (iCL), which re-couple crop and livestock production at the farm scale, have been considered a promising strategy to tackle this challenge by restoring degraded pasturelands and providing supplemental nutrition to livestock. However, few studies have analyzed the economic viability of such systems, especially in Brazil, an important player in global food systems. This paper presents an economic analysis of iCL in Mato Grosso, Brazil, the largest grain and beef producer in the country, which spans the ecologically diverse Amazon, Cerrado and Pantanal biomes. We compare the economic performance of an integrated soybean/corn and beef cattle system to a continuous crop (soybean/corn) system and a continuous livestock (beef cattle) production system from 2005 to 2012. We use empirical case study data to characterize a ‘typical’ farm for each production system within the study region. We find that the integrated crop–livestock system has a higher annual net present value (NPV) per hectare (ha) than continuous cropping or livestock under a range of discount rates. However, under a scenario of substantially higher crop prices, the continuous cropping outperforms iCL. While iCL is not feasible in all regions of the Amazon and Cerrado, our results indicate that in places where the biophysical and market conditions are suitable for production, it could be a highly profitable way to intensify cattle production and potentially spare land for other uses, including conservation. Nevertheless, additional credit and technical support may be needed to overcome high upfront costs and informational barriers to increase iCL areas as a sustainable development strategy for agriculture in the Amazon and Cerrado regions.

**Introduction**

Agriculture is the main economic activity in many low-to-moderate income countries (World Bank, 2017; FAOSTAT, 2018) and employs a large number of workers worldwide (UNEP, 2011; ECLAC, 2017; FAOSTAT, 2018). In Brazil, crop and livestock production contributes substantially to economic growth—roughly 23% of the gross domestic product (GDP) as of 2016 (USD 336.9 billion) (MAPA, 2017a; 2017b). However, it has also been associated with high levels of greenhouse gas emissions (GHGs) and environmental degradation (Graziano da Silva, 2010; Vilela *et al.*, 2011; MAPA, 2017a; 2017b), as well as increasing income inequality in rural areas (Abramovay, 2000; Graziano da Silva and Campanhola, 2004; Balsan, 2006). Beef cattle production, in particular, has been associated with very low incomes and high levels of land degradation, abandonment and deforestation (Margulis, 2004; Fearnside, 2005; Garrett *et al.*, 2017a). In this context, there has been a growing impetus to develop alternative agricultural models that achieve higher productivity and incomes, while reducing environmental impacts, most notably deforestation and GHGs (Nair, 1991; Porfirio-Da-Silva, 2007; Graziano da Silva, 2010; Lemaire *et al.*, 2014; Reis *et al.*, 2016). Improving the sustainability of agriculture in Brazil is a key component of the country’s plan to achieve their emissions reduction targets.

Considering this challenge, two agricultural models that have been encouraged by the Brazilian government, mainly in the Amazon and Cerrado region, are integrated crop–livestock systems (iCL) and integrated crop–livestock–forestry systems

© Cambridge University Press 2019

**CAMBRIDGE**  
UNIVERSITY PRESS

(iCLF)<sup>a</sup> (Brasil, 2012). These types of production systems aim to improve the sustainability of agriculture production through the integration of various types of agricultural production (i.e. crops, livestock and forestry) in the same area, via intercropping, or rotations, to obtain synergies among agroecosystem components (Nair, 1991; Macedo, 2009; Balbino *et al.*, 2011; Lemaire *et al.*, 2014).

Integrated systems represent a strategy to intensify resource uses—labor, land and capital, to increase productivity, while also diversifying production and sparing land for conservation or other uses (Franzluebbers, 2007; Herrero *et al.*, 2010; Lemaire *et al.*, 2014; Reis *et al.*, 2016). Production diversification has the additional benefit of reducing market risk, since farmers have opportunities to manage their product portfolio to take advantage of agricultural market price fluctuations (Herrero *et al.*, 2010; Lazzarotto *et al.*, 2010).

A key feature of integrated systems, mainly iCL, is that they can be used to recover degraded pastures (Kluthcouski *et al.*, 2003; Macedo, 2009; Vilela *et al.*, 2011; Salton *et al.*, 2014) by using residual fertility from the crop rotation to restore soil quality and finance further system improvements (Vilela *et al.*, 2011; Costa *et al.*, 2012). Prior studies in Brazil, especially in the Cerrado, have also shown that iCL systems can increase production efficiency since they contribute to: (i) improvements in soil quality; (ii) water conservation; (iii) an increase of animal performance; and (iv) a reduction in GHGs per unit of food produced (Kluthcouski *et al.*, 2003; Macedo, 2009; Vilela *et al.*, 2011; Salton *et al.*, 2014). What is less understood is how economically viable these productions systems are in the Legal Amazon region of Brazil, particularly in light of their potentially high initial investment costs (Gil *et al.*, 2018) and (Appendix 1). This lack of generalized information about the economic performance of iCL in the country's largest cattle and crop production region may help explain its low adoption rates, despite fairly high levels of government support (Martha Júnior *et al.*, 2011; Vilela *et al.*, 2011; De Oliveira *et al.*, 2013; Salton *et al.*, 2014; Reis *et al.*, 2016).

The aim of this paper is to conduct a comprehensive economic viability analysis of iCL vs a 'typical' (as defined below) continuous crop or livestock farm in the Brazilian Legal Amazon state of Mato Grosso, which is the country's largest producer of soybean and cattle. The evaluation process focuses on assessing the return on investment (ROI) of these systems to inform both producers' decision-making processes as well as bank financial evaluations for funding iCL projects. The integrated system evaluated in this study pertains to soybeans double cropped with corn, followed by pasture and beef cattle grazing, which is the most common integrated system in the Brazilian Amazon and Cerrado (Nair, 1991; Macedo, 2009; Balbino *et al.*, 2011; Lemaire *et al.*, 2014). Our analysis relies on experimental data for a period of 7 years: 2005–2012. In addition to conducting a specific assessment of the case of Mato Grosso, the methods used here can inform future efforts to evaluate the economic viability and returns of iCL at broader scales.

## Material and methods

### Case selection

Our analysis focuses on two representative crop and livestock regions in the state of Mato Grosso, Brazil, one of the largest

<sup>a</sup>In this paper we will concentrate our analysis in integrated crop and livestock systems because this is the integrated system most adopted in Brazil, mainly in Brazilian Cerrado and the Amazon region.

agricultural frontiers in the world (IBGE, 2017; IMEA, 2019; MAPA, 2017a, 2017b). Pastures occupy a majority of the area, followed by soybeans, which are often followed by corn during the course of a single year. Our livestock data were acquired from the municipality of Alta Floresta, in the North region of the state (Fig. 1b), which had the fifth largest cattle herd of the state (706,500 animals) in 2016. Our cropping data were acquired from the municipality of Santa Carmem, in the Mid-North region of the state (Fig. 1a), where about 40% of the soy production occurred in 2016.

The great concentration of agricultural production in the focal livestock and crop regions makes these cases globally important. Yet, they may not be generalizable to all regions within the state, which contains a great deal of climate, soil and institutional variability. The state spans three ecological biomes: the Amazon, Cerrado and Pantanal. Since colonization of the region did not begin in earnest until 1960, it is still a highly dynamic environment characterized by agricultural systems across a range of farm sizes and technology levels.

### Defining a 'typical' crop and livestock farm in Mato Grosso

We defined the 'typical' crop and livestock systems for the North and Mid-North regions of Mato Grosso for the year 2005<sup>b</sup> using farm observations, meetings with local agricultural experts, including farmers, retailers, technicians, consultants, trading managers and data from the Mato Grosso Institute of Agricultural Economics (IMEA). IMEA carries out a comprehensive yearly economic survey focusing on the main agricultural commodities in Mato Grosso: soybean, corn, cotton and beef cattle. These surveys are performed in all Mato Grosso regions using focus group meetings that include farmers and representatives from agricultural organizations and businesses. The purpose of these meetings is to gather up-to-date information about costs, revenue, productivity, investments, farm size, management practices, labor and infrastructure for each commodity across farms in the state.

Based on these data we determined that the typical farm size in Mato Grosso is 700 ha of cultivated land area. The typical crop farm is defined by an intensive and specialized production system with two crop seasons per year: soybean (*Glycine max*) (October–February) and corn (*Zea mays*) (February–June/July). The initial investment required for the operation of this continuous soybean/corn system was USD 765.63<sup>c</sup> ha<sup>-1</sup>, excluding the land acquisition cost.<sup>d</sup> This farm possesses a high level of technology in all production stages with high investment in infrastructure and inputs. As a consequence, it has high soybean productivity levels (av. 3.12 MT ha<sup>-1</sup>), as well as high production costs (av. USD 530.45 ha<sup>-1</sup>) (Table 1). Most soybean production in the region is exported through multinational traders. As of 2005, corn area in the state was still limited, but most production is marketed through domestic channels.

In contrast, the typical livestock farm is characterized by traditional cattle ranches with a low level of technology, low productivity

<sup>b</sup>This year was selected to allow comparison of economic results given that the integrated system experiment started at 2005.

<sup>c</sup>2005 prices. Conversion using exchange data from official Brazilian Govern database provided by Research Institute of Economic Applied (IPEA): <http://www.ipeadata.gov.br/Default.aspx>.

<sup>d</sup>The perspective of the analysis was to evaluate the productive activity performed in the area.




Fig. 1. (a) Crop concentration in Mato Grosso, 2016. (b) Livestock concentration in Mato Grosso, 2016.

and large areas. Farmers do not invest in sophisticated infrastructure, only basic equipment, such as a corral, troughs and fences. Also, farmers do not invest in pasture management. As a consequence, in the dry season, they have difficulties providing adequate nutrition to their herd. The most common cattle breed is Zebu cattle (*Bos taurus indicus*) and pasture is *Urochloa brizantha* cv. Marandu. In contrast to soybeans, the cattle are mainly sold for internal

markets and this activity is less responsive of international prices and exchange rates. The initial investment required for the operation of a continuous traditional livestock system was USD 173.73 ha<sup>-1</sup>, excluding the land acquisition cost.

Integrated crop and livestock systems are still somewhat rare in the study region, so it was not possible to use observations and expert knowledge to characterize these systems. Instead, we draw

**Table 1.** Productivity, operating and inputs cost for a typical integrated crop-livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012

	2005/2006	2006/2007	2007/2008	2008/2009	2009/2010	2010/2011	2011/2012
<b>Soybean</b>							
Soybean productivity (MT/ha)							
iCL typical farm	3.58	3.72	3.34	3.63	3.70	3.77	3.56
Crop typical farm	3.14	3.14	3.14	3.03	3.07	3.33	3.01
Operational cost (inputs, work force and machinery) (USD/ha)							
iCL typical farm	274.94	149.77	194.26	408.10	233.59	271.58	282.71
Crop typical farm	375.19	432.65	520.91	631.64	440.84	532.49	779.46
Input cost (USD/ha)							
iCL typical farm	165.19	98.83	128.53	280.70	157.49	172.76	180.62
Crop typical farm	319.29	368.18	443.29	537.52	340.94	435.80	704.73
<b>Corn</b>							
Corn productivity (MT/ha)							
iCL typical farm	2.19	5.04	4.08	2.82	–	1.95	4.80
Crop typical farm	4.63	4.63	4.63	5.07	4.00	3.99	6.22
Operational cost (inputs, work force and machinery) (USD/ha)							
iCL typical farm	26.19	57.56	61.82	69.06	64.61	61.52	109.52
Crop typical farm	225.10	259.57	312.52	378.96	309.30	408.05	459.87
Input cost (USD/ha)							
iCL typical farm	16.30	37.63	39.28	48.79	47.21	50.96	79.88
Crop typical farm	183.43	211.52	254.67	308.80	246.29	338.25	400.28
<b>Cattle</b>							
Cattle productivity (kg/ha)							
iCL typical farm	162.00	372.00	360.00	216.00	402.00	399.00	411.00
Livestock typical farm	324.00	–	–	–	–	336.00	–
Operational cost (inputs, work force and machinery) (USD/ha)							
iCL typical farm	897.01	1181.71	1376.48	1003.18	2353.35	2495.65	3679.37
Livestock typical farm	92.39	108.68	127.78	142.87	138.07	164.78	182.24

<sup>a</sup>Meetings to collect data on the livestock typical farm were accomplished every five years.

<sup>b</sup>iCL typical farm operating cost includes animal acquisition, which is the most important input cost of livestock. In the livestock typical farm, this value is not computed because farmers produce their herds.

our data from the first iCL experiment established by the Brazilian Agricultural Research Corporation (Embrapa) on a farm called Dona Isabina in the municipality of Santa Carmen in 2005. The farm has 2000 ha cultivated with soybean, corn and rice (*Oryza sativa*) in rotation and crop sequences. However, the iCL experiment occurred on just 100 ha of the site. The soils in the test site are yellow Oxisols and the topography is flat, with very little slope. The average altitude is 386 m, average annual rainfall of 2064 mm with a dry season from June to September and average temperatures of 27.6°C. To establish pasture rotations and crop sequences, the area of 100 ha was divided into five parcels of 20 ha, bounded by fences. The area in which the experiment was implemented had already been cultivated with soybeans in the summer and pearl millet (*Pennisetum glaucum*) as a cover crop after the soybean harvest. Scaling this area up to 700 ha (to match the size of typical crop and livestock farms in the region) we calculated an initial investment of USD 863.38 ha<sup>-1</sup>, excluding land acquisition costs.

Each parcel was cultivated with pastures (*Urochloa brizantha* cv. Marandu and *Urochloa brizantha* cv. BRS Piata). The land use of the iCL system followed an annual rotation of crops: soybean or rice in the summer (October–February) and corn or beans (*Phaseolus vulgaris*) immediately following (February–June). The second crop was intercropped with grass pastures. After the second crop harvest, the cattle were allowed to graze on the pastures that remained, which provided them with additional nutrition during the dry season (June to September) when there is low forage availability.

In the first five years of the experiment the herd was a mixture of male and female Zebu cattle acquired in the region. These animals were sold for slaughter when they reached weight of 480 kg. In the last two years, only males were raised, but still slaughtered when they reached 480 kg. The only supplementation used all year long was mineral salt with an average consumption of 90 g day<sup>-1</sup> during the rainy season and 120 g day<sup>-1</sup> during the

dry season. In the dry season, the cattle also received sorghum silage (*Sorghum bicolor*), soybean residues, corn and rice produced in the farm processing unit. In all modules, mangers for supplementation and watering were available.

### Economic indicators

We used an economic viability analysis approach to compare the economic results of the three agricultural systems (Buarque, 1984; Gitman and Zutter, 2014). This method is established in the economic literature as an instrument to evaluate the economic potential of any investment decision (Buarque, 1984; Lapponi, 2013; Gitman and Zutter, 2014). We used data from IMEA to generate typical crop and livestock farm and survey data to generate the iCL farm. Taking into account the lack of available economic performance data for agriculture systems, the use of IMEA and experimental data are the only feasible approaches for establishing the time-series data required to carry out the economic viability analysis presented. The results can be useful for farmers, helping them compare different investment options, as well as for funding agents since they can evaluate different complex agriculture systems using comparable indicators. Since prior studies have identified that a lack of technical information on the economic performance of iCL for both farmers and financiers is a key constraint for farmer adoption (Martha Júnior *et al.*, 2011; Vilela *et al.*, 2011; De Oliveira *et al.*, 2013; Salton *et al.*, 2014; Reis *et al.*, 2016; Cortner *et al.*, 2019), our approach may help enable wider scale adoption of this technology. The financial accounting approach used here, which is based on observed outcomes, is also a useful complement to process models, which predict outcomes based on inputs (e.g. Gil *et al.*, 2018 for the same region).

We used the following five indicators to assess economic viability and potential economic returns of the iCL system, continuous soy/corn system, and continuous beef cattle system over 7 years (2005–2012): (i) internal rate of return (IRR), (ii) net present value (NPV), (iii) return on investment (ROI), (iv) profitability index (PI) and (v) payback (Gitman and Zutter, 2014).<sup>e</sup>

*Cash flow:* To calculate each of these five indicators we first needed to estimate the real cash flow (CF) based on 2005 prices. Following (Lapponi, 2013):

$$CF_t = FCO_t + \Delta I + \Delta CG_t \quad (1)$$

In which:  $FCO_t$  = Operating cash flow;  $\Delta I$  = Net investment in assets;  $\Delta CG_t$  = Net investment in working capital.

Apart from the relationship between costs and revenues, cash flow results take into account interest deductions, taxes and labor charges to demonstrate the cash generation potential of each system.<sup>f</sup> As a measure of yearly profitability, we used the net operating profit after income tax (NOPAT).<sup>g</sup> It represents the net profit that the system generates to remunerate both the funding entity and the producer (Assaf Neto, 2011; Lapponi, 2013; Gitman and Zutter, 2014). As an inflation indicator, we used the broad consumer price index (IPCA) provided by the Brazilian Institute of Geography and Statistics (IBGE), which is the official inflation index in Brazil.

<sup>e</sup>Annual results from indicators NPV (annual net present value- NPVA) and ROI (annual return of investment—ROIA) were calculated and displayed to become easier the comparison between the three systems.

<sup>f</sup>The share of working capital was disregarded and the assets' flow was incorporated into the operating result observed in the last year of assessment.

<sup>g</sup>For construction of the NOPAT, see the supplementary material.

*Investment value:* Except for the land value, which was not incorporated into the cash flow, all other infrastructure elements required for production activities were considered as if they had been purchased in the initial year of all production systems, 2005. A market survey was conducted with consultants and equipment retailers to collect prices data in the Mid-North region in 2005, taking into account the infrastructure needed to set up each farm system.

*Discount rate:* The discount rate defines the present value of future returns (Buarque, 1984; Gitman and Zutter, 2014). Choosing a discount rate is one of the most controversial points in economic investment analysis because the choice of incorrect values can lead to suboptimal results and decisions (Buarque, 1984; Lapponi, 2013). The project economic evaluation literature defines the discount rate as the opportunity cost of investment, which means that it should reflect the expected return value for alternative available investments with similar risk to the activity being analyzed (Buarque, 1984; Lapponi, 2013; Gitman and Zutter, 2014). This approach, although it incorporates correctly the perspective of the discount rate to be used, is limited by the lack of investment alternatives that can serve as a reference (Buarque, 1984).

As a result, the official savings rate is more commonly used in many agricultural investment evaluations, since it represents a low-risk and low return alternative investment option (Buarque, 1984; Gitman and Zutter, 2014). In other cases, the economy basic interest rate or long-term interest rates has been used, also indicating low-risk investment alternatives, but with higher returns. An important issue regarding the use of these rates as a reference is no consideration of the investor's profile for defining the interest rate to be used.

Given these drawbacks, our study uses the weighted average cost of capital (WACC), to adjust the variables that make up the investment opportunity cost based on the agent's profile, as well as the level of risk associated with the business being evaluated<sup>h</sup>. The WACC is more appropriate for this evaluation since it considers an agent's decision about which percentage of investment will be funding as well as incorporates market risks of alternative investments (Buarque, 1984; Lapponi, 2013; Gitman and Zutter, 2014). The WACC rate was built taking into account the financial market conditions in Mid-North region in 2005.

### Incorporating changing land use and market dynamics

Since our study analyzed the economic viability of the three systems over a 7-year period it was necessary to incorporate changes in land use and markets that were occurring over that period. These dynamics include the growing importance of corn as a second crop<sup>i</sup> (resulting in an increase in the on-farm area allocated to integrated crop–livestock systems), changes in the marketing arrangements used by farmers, and a dynamic macroeconomic environment in which real prices for soybean, corn and beef were changing frequently due to growth in demand and exchange rate variations.

Data from IMEA show that corn as a second harvest crop grew by 14.87% per year in the period 2008–2012. To simulate dynamics of land use in the integrated crop–livestock farm, the growth of corn second harvest area in the typical continuous crop farm was used to define the growth of the integrated

<sup>h</sup>For construction of the WACC, see the supplementary material.

<sup>i</sup>According to IMEA, for the 2007/2008 crop year the corn area in Mato Grosso was 1,670,800 hectares and 796,500 hectares for Mid North region. In the 2009/2010 this area increasing to 1,948,020 hectares in the state and 964,000 hectares for Mid North region. The crop year with a more expressive planted area was 2012/2013, in which were planted 3,702,053 hectares in the state with 1,830,318 hectares in the Mid North region.




Fig. 2. Average commodity prices in Mato Grosso from 2005 to 2017.

system area<sup>j</sup> (Kluthcouski *et al.*, 2003; Macedo, 2009; Balbino *et al.*, 2011; Vilela *et al.*, 2011).

Interviews with farmers and specialized consultants who worked in the North and Mid-North regions in 2005 identified that the most common soybean marketing practice used during that time was to sell their harvest over three periods: (i) 25% of production was sold in advance, from August to October, (ii) 50% of production was sold from November to April, during the harvesting and immediate post-harvest period and (iii) 25% of production was sold from May to July, the period of preparation for another harvest. To adjust the revenue dynamics to the trading practices of that period, the crop sales process was adjusted according to the moment of the soybean harvest<sup>k</sup>. Soy sale prices for each period are calculated as the average of the prices observed during the months of soy trading. Similarly, corn sale prices are calculated as the average of the prices observed from September to November, the main months for corn trading.

Of particular importance, soybean prices were very low in 2005 and 2006, while production costs remained high (CEPEA, 2007). However, after 2007 the soybean price steadily increased, a trajectory influenced by China's consolidation as the main Brazilian soybean importer (Fig. 2). In 2012, the soybean price in the Mid-North of Mato Grosso—USD 28.94 per sack (60 kg), was three times higher than the value observed in 2005—USD 9.55 per sack; (IMEA, 2016). Nonetheless, in 2009, the financial crisis complicated production and trading. The devaluation of the Brazilian currency during this period (9% in one year), led to increased crop production costs (10% in 2009), largely as a result of fertilizer imports, while soybean prices remained low (IMEA, 2019).

In contrast, corn prices increased during 2010–2012 as a consequence of financial crisis of 2009, since corn production is oriented toward domestic consumption and is not as influenced by global commodity markets. The same domestic market orientation and price dynamics can be seen in the prices for beef, which achieved a historic high price in 2011, USD 54.40 per “arroba” (30 kg of live weight). However, a considerable portion of Mato Grosso's beef production is exported, 22.1% on average in the last 5 years (MAPA, 2017b), destined mainly for EU, Russia, China and Middle East (MAPA, 2017b; IMEA, 2019).

<sup>j</sup>The most common practice is to plant corn intercropping with pasture to recover soil quality and provide food for cattle during the driest period of the year in the region, from June to September.

<sup>k</sup>Only the soybeans trading process was taken into consideration, once the corn, at that moment, did not present the economic relevance observed currently.

## Results

### Productivity

The average cattle productivity in the iCL farm (331.71 kg ha<sup>-1</sup>) was 5 times higher than the typical livestock farm (63.3 kg ha<sup>-1</sup>) (IMEA, 2019) due to the availability of higher quality pasture during the dry period of the year. The productivity of soybean in the iCL farm was also on average 16% higher than crop typical farm during the whole study period (Table 1). On the other hand, the input cost of iCL system was 62% lower than the continuous crop farm. Taking into account the high contribution of fertilizers to input costs, this association between higher productivity and lower input cost is likely related to the positive influence of the integrated systems on soil nutrient availability (Carvalho *et al.*, 2010; Garrett *et al.*, 2017b). Further systematic measurement of soil nutrient availability is needed to confirm this hypothesis. A different result was observed with corn. Since corn had little economic importance at the time that the iCL experiment was started at the Dona Isabina farm and the main objective was to provide agronomic benefits for pasture, low productivity corn seeds were used. Moreover, in 2009 and 2010 there was an intense dry period at the crop germination stage which affected productivity.

### Cash flows<sup>l</sup>

The iCL system had the largest investment costs (negative cash flow in years one and two), but also had the largest positive cash flows throughout the remainder of the study period, achieving a positive result of USD 654.04 ha<sup>-1</sup> in 2012 compared to USD 460.85 ha<sup>-1</sup> for continuous cropping and USD 27.59 ha<sup>-1</sup> for continuous livestock (Fig. 3). Macroeconomic fluctuations explain most of the changes in cash flows over the study period. In particular, soybean and beef prices increased during the study period (Fig. 2).

During June to September most continuous livestock farms have to sell off part of their herd, since they do not have conditions to feed them (IMEA, 2016; Valentim, 2016; Gil *et al.*, 2018; Reis *et al.*, 2019), which is thought to cause declines in the local cattle price. The higher pasture productivity obtained in the iCL system on the Dona Isabina farm, translated to higher cattle productivity (331.71 kg ha<sup>-1</sup> annual average) (Table 1), and enabled this farm to keep their animals during the annual dry

<sup>l</sup>For a detailed cash flow description, see supplement material.




Fig. 3. Discounted cash flow of a typical integrated crop-livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012.




Fig. 4. NOPAT of a typical integrated crop-livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012.

season. Indeed, the pasture management strategy implemented at Dona Isabina provided an annual increase of 14% in cattle productivity over the seven years. Moreover, in 2012, the annual cattle productivity was 2.5 times higher than its cattle productivity in 2005 (Table 1). The seasonal dilemma of traditional cattle ranches also enabled the Dona Isabina farm to acquire animals at a low price during the dry season and sell them in periods when prices were high. The seasonal advantage and the high cattle productivity largely explain the better economic results of iCL vs continuous cattle (Fig. 3).

The iCL farm also resulted in higher cash flows than the continuous crop farm (Fig. 3), due to the combination of higher productivity and, on average, 62% lower production costs and 51% lower operating costs (Table 1). The large reduction in production costs can be attributed to lower fertilizer needs due to improved soil fertility from both manure and nitrogen fixing legumes in the pasture.

The economic fragility of traditional livestock is evidenced by the smaller cash flow throughout the study period (on average USD 23,131.62 vs USD 109,164.24 for continuous cropping and USD 204,318.97 for the integrated system).

The iCL farm also outperforms the continuous cropping and continuous livestock systems in terms of the NOPAT (Fig. 4). This indicator, which can be interpreted as the annual system capacity to provide economic return after taxes and financial expenses (e.g. interest on debt), indicated that the iCL farm provided a greater money supply than the continuous crop and livestock systems throughout the study period, aside from the initial year.

Another economic indicator widely used in the project analysis approach is the recovery period of the investment (the number of years of positive cash flows it takes to repay the initial investment and negative cash flows), known in the literature as the payback period. The iCL farm recovered the investment after 4 years (Fig. 5) while the continuous crop did not recover their investment until year 6. The livestock system recovered the investment after 5 years. In the end of seventh year, the continuous crop and livestock farms had an accumulated cash flow of USD 228,207.46 and USD 40,313.76, respectively. However, the iCL farm had accumulated USD 825,868.81.

### Economic viability indicators

The cash flow of all systems provides useful information to elaborate the set of economic viability indicators displayed in Table 2. Across all indicators (NPV, internal rate of return, payback, profitability index and ROI) the iCL system performs substantially better than the continuous crop and livestock farms. The exception is the higher upfront investment cost per hectare. The livestock farm has the worst performance across all indicators.

### Scenario Analysis

#### Different Interest Rates

All the results presented above are quite sensitive to the discount rate. Here, we used the Center-West Constitutional Fund rate,




Fig. 5. Payback of a typical integrated crop-livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012.

Table 2. Economic viability indicators for a typical integrated crop–livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012

Indicators	Crop typical farm	Livestock typical farm	iCL typical farm
WACC	9.66%	9.18%	9.53%
Investment (USD) ha <sup>-1</sup>	USD 765.63	USD 173.73	USD 863.38
NPV (USD) ha <sup>-1</sup>	USD 66.73	USD 5.22	USD 674.17
NPVA (USD) ha <sup>-1</sup>	USD 13.56	USD 1.04	USD 136.25
IRR	11.32%	10.01%	22.16%
ROI	10.98%	9.64%	18.94%
ROIA	1.2%	0.42%	8.58%
Profitability index	1.09	1.03	1.78

8.75%, to construct the WACC, as well as the entire set of economic viability indicators because in 2005 there was no specific government loan program to encourage iCL in Brazil. However, in 2010, Brazilian Government implemented the low carbon agriculture plan (ABC plan) with incentives and low interest rates for more sustainable agricultural systems, including iCL. The ABC plan offered interest rates of 5.5% in 2010 (Brasil, 2012). However, the performance of iCL relative to the other systems does not change if we use the ABC plan rate or the basic interest rate of Brazilian economy (SELIC) in 2005 (19.24%), a rate used to evaluated investments in stock market (Table 3).

### Different Prices

Between 2005 and 2006, soybean prices were very low in the global market (Fig. 2). Both soybean and corn prices peaked in 2010 and then again in 2016. To capture the effects of these higher prices we used the average soybean and corn prices observed in the Mid-North region between 2013 and 2017. For consistency, the cattle prices were also adjusted to the average prices in the Mid-North region between 2013 and 2017. Moreover, the corn planted area was increased, to match the growth in the average farm-level planted area in the Mid-North in the last 10 years (2007–2017: 46.44%). All other conditions were kept unchanged.

As a result of these scenario adjustments the continuous crop system overtook iCL as a better investment (Table 4).

### Discussion

#### High profitability and greater profit stability of iCL under a range of scenarios offsets its high upfront costs

Despite its low uptake compared to continuous cropping system or traditional extensive ranching, our results indicate that iCL is a substantially better land use investment than continuous crop or livestock systems from a financial perspective under existing crop price scenarios. It both increases the productivity of pasture areas and reduces reliance on external inputs in cropping areas, contributing to higher overall profitability. One reason for the low uptake of iCL is that farmers accurately perceive the system to have high upfront costs and they are uncertain as to how long it will take for the system to pay back this investment (Martha Júnior *et al.*, 2011; Costa *et al.*, 2012; Cortner *et al.*, 2019). However, our results show that the payback period is only 4 years for the iCL system, less than that of continuous cropping—6 years, or continuous livestock—5 years.

If payback time is considered as an investment risk indicator (Assaf Neto, 2011; Gitman and Zutter, 2014), then iCL actually demonstrates lower economic risk than continuous crop or livestock systems (Muniz *et al.*, 2007; Lazzarotto *et al.*, 2010). The iCL system also shows lower variations in profit and NPV under different price and interest rate scenarios. Given the high fluctuations in prices that have occurred in grain commodity prices over the 2000s and their inverse relationship to domestic beef prices, iCL allows farmers the opportunity to buffer their losses when one system suffers due to major price changes. However, the positive returns on continuous cropping are likely a market barrier to the adoption of iCL in regions that are highly suitable for soybean and corn production.

#### Economic performance of continuous cropping is highly dependent on exchange rates and world prices

Due to its dependence on external markets for both sales and fertilizers, the performance of continuous cropping was strongly influenced by the prevailing exchange rate and international commodity prices, the same main drivers of deforestation in the Amazon (Rodrigues-Filho *et al.*, 2015). When the currency was devalued, Brazilian crops became more competitive in global



**Table 3.** Simulation with different interest rates—economic viability indicators for a typical integrated crop-livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012

Indicators	Crop typical farm		Livestock typical farm		iCL typical farm	
	SELIC (19.24%)	ABC plan (5.5%)	SELIC (19.24%)	ABC plan (5.5%)	SELIC (19.24%)	ABC plan (5.5%)
WACC	13.86%	8.36%	19.24%	5.5%	13.73%	8.24%
Investment (USD) ha <sup>-1</sup>	765.63	765.63	173.73	173.73	863.38	863.38
NPV (USD) ha <sup>-1</sup>	(89.98)	124.33	(45.18)	31.52	393.73	776.62
NPVA (USD) ha <sup>-1</sup>	(20.89)	24.17	(12.27)	5.55	91.07	150.38
IRR	11.31%	11.31%	10.01%	10.01%	22.15%	22.15%
ROI	11.84%	10.71%	14.22%	8.04%	19.99%	18.61%
ROIA	(1.77%)	2.17%	(4.21%)	2.41%	5.50%	9.58%
Profitability index	0.88	1.16	0.74	1.18	1.45	1.89

**Table 4.** Simulation with different crop prices—economic-financial viability indicators for a typical integrated crop-livestock, continuous crop and continuous livestock farm in Mato Grosso from 2005 to 2012

Indicators	Crop typical farm	iCL typical farm
WACC	9.66%	9.53%
Investment (USD) ha <sup>-1</sup>	765.63	863.38
NPV (USD) ha <sup>-1</sup>	761.38	52.70
NPVA (USD) ha <sup>-1</sup>	154.66	10.66
IRR	30.54%	10.86%
ROI	21.01%	10.46%
ROIA	10.35%	0.84%
Profitability index	1.99	1.06

markets, but, also faced higher production costs (Table 1). In 2008 and 2011, when the exchange rate increased substantially, production costs were particularly high.

Moreover, the high profitability of cropping under current price scenarios may explain why the most common strategy of iCL in this region has been the ‘third harvest’, in which a farmer produces soybean in the first harvest and plants corn intercropped with pasture. Recent research by Embrapa found that 83% of integrated systems in Brazil are iCL and the same pattern can be observed in Mato Grosso (Embrapa; Rede iLPF, 2017). Furthermore, the ‘third harvest’ strategy represents around 50% of iCL in Mato Grosso (Embrapa; Rede iLPF, 2017). As the results show, iCL can reduce external input dependence and improve the economic viability of farming in the region.

#### Extensive livestock ranching traps farmers in a cycle of low income due to dry season losses

The cash flow restrictions faced by traditional extensive livestock producers make it difficult for ranchers to take advantage of the livestock market. These farmers have few alternatives than selling part of their herd in the dry season, which limits their cash flow and, as a consequence, their capacity to generate revenue. The lack of economic competitiveness of extensive livestock relative to cropping or iCL explains why over the last decade in Mato Grosso many pasture areas have been overtaken by cropland (Macedo *et al.*, 2012; Lapola *et al.*, 2014).

Given the existing low returns of continuous livestock systems, and future potential changes in climate that will further reduce pasture productivity in Mato Grosso (Gil *et al.*, 2018), it will be even more imperative to help farmers adopt improved pasture management practices, such as iCL to maintain their livelihoods, or else abandon production entirely. iCL would also help reduce the GHGs from livestock (Gil *et al.*, 2018) and provide new funding opportunities, which have been connected with use and adoption of sustainable practices such as ABC plan.

#### Low interest loans are key to the viability of establishing all three systems

Using the SELIC interest rate scenario of 19.24%, only iCL was still economically viable. Using the ABC interest rate of 5.5% doubled the NPV of iCL. Continuous cropping showed a huge deficit in the SELIC interest rate scenario, indicating the relationship between technological levels and financial obligations. These results underscore the importance of public policies to provide attractive funding plans with low interest rates to agriculture. However, in recent years, because of economic and political crises, the interest rates provided by the ABC program increased to 8.5% in 2016/2017 and 7.5% in 2017/2018 (MAPA, 2017b).

#### Conclusion

The challenge of protecting the environment, while generating income and reducing social inequality, requires the identification of agricultural strategies that enable the sustainable intensification of production. Given the growing international concern about the environmental impacts of agricultural activities in the Brazilian Amazon and Cerrado, as well as the importance of Brazilian agriculture in world food systems, the promotion of sustainable agricultural practices in Brazil is of global relevance.

This work, in addition to presenting an alternative to the current model of agriculture, sought to advance understanding of the economic performance of iCL as a sustainable intensification strategy compared to traditional continuous crop and livestock systems. Our results showed that iCL had higher levels of productivity, profitability and ROI and lower payback periods and economic risk than the continuous crop and livestock systems under existing prices and exchange rates over a 7 year period between 2005 and 2012. However, under higher crop prices,

continuous cropping provides better economic results than the integrated system.

The case study approach used here is necessary and useful in the absence of a large sample of iCL farms from which to draw data, but does not guarantee that the results are representative of all potential iCL farms in the region. In order to assess how generalizable our results are to northern Mato Grosso and the rest of the Legal Amazon, a wider sample of farms across the region needs to be considered. As iCL continues to be adopted, these types of surveys will become increasingly possible.

Finally, the financial performance of iCL, though potentially important for decisions to adopt or not adopt these systems, are not the only outcomes that are relevant to farmers and policy makers. Systematic measurement of environmental indicators, such as soil fertility, GHGs and water consumption on iCL farms in the study region are needed. Further research should also explore the tradeoffs between economic and environmental outcomes in integrated systems (e.g. Gil *et al.*, 2018). Since farmers are often motivated by non-monetary objectives and integrated systems entail major changes in management complexity, debt financing and farm aesthetics, better understanding of their cultural appropriateness is needed (Garrett *et al.*, 2017b; Cortner *et al.*, 2019). Given the multifaceted and dynamic reality associated with agriculture, it is vital to assess the social and environmental benefits across a wider range of farms and regions, as well as climate and macroeconomic scenarios. The evaluation of any agricultural system's potential to promote sustainable development must be based on models and assessments that capture the interrelations between different system components—economic, social and environmental—at broader spatial scales beyond the farm (Garrett and Rausch, 2016).

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S1742170519000280>.

**Acknowledgement.** The authors acknowledge Embrapa Agrosilvopastoral, Mato Grosso Institute of Agricultural Economics and the National Rural Learning Service-MT for the support and funding of this research. Moreover, we thank the Work Group of iCLF systems of Embrapa Agrosilvopastoral for fundamental collaboration in development of this economic analysis. We also thank all the colleagues and reviewers who helped with important advices for the text. Special thanks to Aisten Baldan for fundamental help with the reference search and Paula Emilia Pimentel for help with figures. Errors and limitations remaining are the responsibility of the authors.

## References

- Abramovay R** (2000) Agricultura, Diferenciação Social e Desempenho Econômico. *Seminário: Desafios Da Pobreza Rural No Brasil*. Rio de Janeiro-RJ, Brasil.
- Assaf Neto A** (2011) *Estrutura e Análise de Balanço: Um Enfoque Econômico-Financeiro*. 5o Edição. ed. São Paulo-SP, Brasil: Atlas.
- Balbino LC, Barcellos AO and Stones LF** (2011) *Marco referencial: integração Lavoura- Pecuária-Floresta*. 1o. ed. Brasília-DF, Brasil: Embrapa.
- Balsan R** (2006) Impactos decorrentes da modernização da agricultura Brasileira. *Campo - Territ. Rev. Geogr. Agrária* **1**, 123–151.
- Brasil** (2012) Plano Setorial de Mitigação e Adaptação às Mudanças Climáticas para Consolidação da Economia de Baixa Emissão de Carbono na Agricultura – PLANO ABC, Ministério da Agricultura, Pecuária e Abastecimento-MAPA. Brasília-DF, Brasil.
- Buarque C** (1984) *Avaliação Econômica de Projetos*. 1o. ed. Rio de Janeiro-RJ, Brasil: Editora Campus.
- Carvalho JLN, Raucí GS, Cerri CEP, Bernoux M, Feigl BJ, Wruck FJ and Cerri CC** (2010) Impact of pasture, agriculture and crop-livestock systems on soil C stocks in Brazil. *Soil & Tillage Research* **110**, 175–186. <https://doi.org/10.1016/j.still.2010.07.011>.
- CEPEA** (2007) Centro de Estudos Avançados em Economia Aplicada-ESALQ/USP, Relatório de Custos de Agrícolas-Soja. Piracicaba-SP, Brasil.
- Cortner O, Garrett RD, Valentim JF, Ferreira J, Niles MT, Reis J and Gil J** (2019) Perceptions of integrated crop-livestock systems for sustainable intensification in the Brazilian Amazon. *Land Use Policy* **82**, 841–853. <https://doi.org/10.1016/j.landusepol.2019.01.006>.
- Costa FP, de Almeida RG, de A Pereira M, Kichel AN and Macedo MCM** (2012) Avaliação econômica de sistemas de integração lavoura-pecuária-floresta voltados para a recuperação de áreas degradadas em Mato Grosso do Sul. VII Congresso Latinoamericano de Sistemas Agroflorestais Para a Produção Pecuária Sustentável, pp. 523–527.
- De Oliveira P, Freitas RJ, Kluthcouski J, Ribeiro AA, Adriano L, Cordeiro M, Teixeira LP, Augusto R, Castro D, Vilela L and Balbino LC** (2013) Evolução de Sistemas de Integração Lavoura-Pecuária-Floresta (ILPF): estudo de caso da Fazenda Santa Brígida, Ipameri, GO, pp. 1–51.
- de Oliveira CAO, Bremm C, Anghinoni I, De Moraes A, Kunrath TR and De Faccio Carvalho PC** (2014) Comparison of an integrated crop-livestock system with soybean only: economic and production responses in southern Brazil. *Renewable Agriculture and Food System* **29**, 230–238. <https://doi.org/10.1017/S1742170513000410>.
- ECLAC** (2017) Economic Commission for Latin America and the Caribbean—Database. Available at <https://estadisticas.cepal.org/cepalstat/Portada.html> (Accessed 5 November 2017).
- Embrapa; Rede iLPF** (2017) ILPF em Números. Available at <https://www.embrapa.br/web/rede-ilpf/ilpf-em-numeros> (Accessed 10 August 2018).
- FAOSTAT** (2018) Food and Agriculture Organization of the United Nations. Available at <http://www.fao.org/faostat/en/#data> (Accessed 12 December 2018).
- Fearnside PM** (2005) Deforestation in Brazilian Amazonia: history, rates, and consequences. *Conservation Biology* **19**, 680–688. <https://doi.org/10.1111/j.1523-1739.2005.00697.x>.
- Franzluubbers AJ** (2007) Integrated crop-livestock systems in the southeastern USA. *Agronomy Journal* **99**, 361–372. <https://doi.org/10.2134/agronj2006.0076>.
- Garrett RD and Rausch LL** (2016) Green for gold: social and ecological trade-offs influencing the sustainability of the Brazilian soy industry. *Journal of Peasant Studies* **43**, 461–493. <https://doi.org/10.1080/03066150.2015.1010077>.
- Garrett R, Niles M, Gil J, Dy P, Reis J and Valentim J** (2017a) Policies for reintegrating crop and livestock systems: a comparative analysis. *Sustainability* **9**, 473. <https://doi.org/10.3390/su9030473>.
- Garrett RD, Niles MT, Gil JDB, Gaudin A, Chaplin-Kramer R, Assmann A, Assmann TS, Brewer K, de Faccio Carvalho PC, Cortner O, Dynes R, Garbach K, Kebreab E, Mueller N, Peterson C, Reis JC, Snow V and Valentim J** (2017b) Social and ecological analysis of commercial integrated crop livestock systems: current knowledge and remaining uncertainty. *Agricultural Systems* **155**, 136–146.
- Gil JDB, Garrett RD, Rotz A, Daioglou V, Valentim J, Pires GF, Costa MH, Lopes L and Reis JC** (2018) Tradeoffs in the quest for climate smart agricultural intensification in Mato Grosso, Brazil. *Environmental Research Letters* **13**, 064025.
- Gitman LJ and Zutter CJ** (2014) *Principles of Managerial Finance—Global Edition*, 14o. ed. New York, USA: Pearson Education.
- Graziano da Silva J** (2010) Os desafios das agriculturas brasileiras. *Seminário: A Agricultura Brasileira: Desempenho, Desafios e Perspectivas*, Instituto de Pesquisa Economia Aplicada – IPEA, Brasília-DF, Brasil.
- Graziano da Silva J and Campanhola C** (2004) *O Novo Rural Brasileiro: Novas Atividades Rurais*. 1o. ed. Brasília-DF, Brasil: Embrapa Informação Tecnológica.
- Herrero M, Thornton PK, Notenbaert AM, Wood S, Msangi S, Freeman HA, Bossio D, Dixon J, Peters M, Steeg JVD, Lynam J and Rao PP** (2010) Smart investments in sustainable crop-livestock systems: revisiting mixed crop-livestock systems. *Science* **327**, 822–825. <https://doi.org/10.1126/science.1183725>.
- IBGE** (2017) Instituto Brasileiro de Geografia e Estatística. Available at <https://sidra.ibge.gov.br> (Accessed 10 August 2018).
- IMEA** (2016) Relatório de Rentabilidade da Pecuária. Cuiabá- MT, Brasil.

- IMEA (2019) Instituto Matogrossense de Economia Agropecuária. Available at <http://www.imea.com.br/imea-site/relatorios-mercado> (Accessed 14 April 2019).
- Kluthcouski J, Stone LF and Aidar H (2003) *Integração Lavoura-Pecuária*, 1o. ed. Santo Antônio de Goiás-GO, Brasil: Embrapa Arroz e Feijão.
- Lapola DM, Martinelli LA, Peres CA, Ometto JPHB, Ferreira ME, Nobre CA, Aguiar APD, Bustamante MMC, Cardoso MF, Costa MH, Joly CA, Leite CC, Moutinho P, Sampaio G, Strassburg BBN and Vieira ICG (2014) Pervasive transition of the Brazilian land-use system. *Nature Climate Change* 4, 27–35. <https://doi.org/10.1038/nclimate2056>.
- Lapponi J (2013) *Projetos de Investimento na Empresa*, 1o. ed. Rio de Janeiro-RJ, Brasil: Elsevier Brasil.
- Lazzarotto JJ, dos Santos ML, de Lima JE and de Moraes A (2010) Financial viability and risks of integrated crop-livestock systems in the state of Paraná. *Organizações Rurais & Agroindustriais* 12, 113–130.
- Lemaire G, Franzluebbers A, de F Carvalho PC and Dedieu B (2014) Integrated crop-livestock systems: strategies to achieve synergy between agricultural production and environmental quality. *Agriculture, Ecosystems & Environment* 190, 4–8. <https://doi.org/10.1016/j.agee.2013.08.009>.
- Macedo MCM (2009) Integração lavoura e pecuária: O estado da arte e inovações tecnológicas. *Revista Brasileira de Zootecnia* 38, 133–146. <https://doi.org/10.1590/S1516-35982009001300015>.
- Macedo MN, DeFries RS, Morton DC, Stickler CM, Galford GL and Shimabukuro YE (2012) Decoupling of deforestation and soy production in the southern Amazon during the late 2000s. *Proceedings of the National Academy of Sciences* 109, 1341–1346. <https://doi.org/10.1073/pnas.1111374109>.
- MAPA (2017a) Ministério da Agricultura Pecuária e Abastecimento—Valor Bruto da Produção Agropecuária (VBP). Available at <http://www.agricultura.gov.br/assuntos/politica-agricola/valor-bruto-da-producao-agropecuaria-vbp> (Accessed 17 September 2018).
- MAPA (2017b) Ministério da Agricultura, Pecuária e Abastecimento—Agropecuária Brasileira em Números. Available at <http://www.agricultura.gov.br/assuntos/politica-agricola/agropecuaria-brasileira-em-numeros> (Accessed 20 August 2018).
- Margulis S (2004) Causes of deforestation of the Brazilian Amazon, World Bank Working Paper. Washington-DC, USA. <https://doi.org/10.1596/0-8213-5691-7>.
- Martha Júnior GB, Alves E and Contini E (2011) Dimensão econômica de sistemas de integração lavoura-pecuária. *Pesquisa Agropecuária Brasileira* 46, 1117–1126. <https://doi.org/10.1590/S0100-204X2011001000002>.
- Muniz L, Figueiredo R, Magnabosco C, Wander A and Júnior G (2007) Análise de risco da integração lavoura e pecuária com a utilização de System Dynamics. *XLV Congresso Da Sociedade Brasileira de Economia, Administração e Sociologia Rural*, Londrina, PR-Brasil.
- Nair PKR (1991) State-of-the-art of agroforestry systems. *Forest Ecology and Management* 45, 5–29. [https://doi.org/10.1016/0378-1127\(91\)90203-8](https://doi.org/10.1016/0378-1127(91)90203-8).
- Porfirio-Da-Silva V (2007) A Integração “Lavoura-pecuária-floresta” como Proposta de Mudança do Uso da Terra. *Seminário: Sistemas de Produção Agropecuária—Ciências Agrárias, Animais e Florestais—Universidade Tecnológica Federal Do Paraná*, Dois Irmãos, PR-Brasil.
- Reis J, Rodrigues R, Conceição M and Martins C (2016) Integração Lavoura-Pecuária-Floresta no Brasil: uma estratégia de agricultura sustentável baseada nos conceitos da Green Economy Initiative. *Sustentabilidade em Debate* 7, 58–73. <https://doi.org/10.18472/SustDeb.v7n1.2016.18061>.
- Reis JC, Kamoi MYT, Pedreira BC, Michetti M, Gimenez MA, Mombach MA and Silva NMF (2019) Aspectos econômicos da recuperação de pastagens na Amazônia. In Dias-Filho MB and Andrade CMS (eds), *Recuperação de Pastagens Degradadas Na Amazônia*. Brasília-DF: Embrapa-DF, p. 443.
- Rodrigues-Filho S, Verburg R, Bursztyn M, Lindoso D, Debortoli N and Vilhena AMG (2015) Election-driven weakening of deforestation control in the Brazilian Amazon. *Land Use Policy* 43, 111–118. <https://doi.org/10.1016/j.landusepol.2014.11.002>.
- Salton JC, Mercante FM, Tomazi M, Zanatta JA, Concencço G, Silva WM and Retore M (2014) Integrated crop-livestock system in tropical Brazil: toward a sustainable production system. *Agriculture, Ecosystems & Environment* 190, 70–79. <https://doi.org/10.1016/j.agee.2013.09.023>.
- UNEP (2011) *Towards a Green Economy: Pathways to Sustainable Development and Poverty Eradication—A Synthesis for Policy Makers*, United Nations Environment Programme. St-Martin-Bellevue, France.
- Valentim JF (2016) Desafios e Estratégias para a Recuperação de Pastagens Degradadas e Intensificação da Pecuária a Pasto na Amazônia Legal. In: Pereira DH and Pedreira BC (eds), *Simpósio de Pecuária Integrada*. Sinop-MT, Brasil: Fundação Uniselva, p. 45.
- Vilela L, Martha GB, Macedo MCM, Marchão RL, Guimarães R, Pulrolnik K and Maciel GA (2011) Sistemas de integração lavoura-pecuária na região do Cerrado. *Pesquisa Agropecuária Brasileira* 46, 1127–1138. <https://doi.org/10.1590/S0100-204X2011001000003>.
- World Bank (2017) World Bank. Available at <https://data.worldbank.org/indicator/sl.agr.empl.zs> (Accessed 20 October 2018).

## Appendix 1 Summary Literature Review: Economic analysis of integrated crop and livestock systems in Brazil

Authors	Focus of the analysis	Productive Systems	Period	Indicators	Main results
Muniz <i>et al.</i> (2007)	Economic viability and minimizing market risks	iCL in Goias, Brazil	3 years, using simulations	NPV and IRR	The iCL was economically viable in all scenarios considered
Lazzarotto <i>et al.</i> (2010)	Economic viability and minimizing market risks	iCL, continuous crop system (soybeans and corn in the summer and wheat in the winter) and continuous livestock (beef cattle) system in Paraná, Brazil	13 years, using simulations	NPV and IRR	In both situations (real and simulated) the iCL presented better economic results: NPV 103% higher than the crop system and 19.6% higher than the livestock system. Furthermore, the iCL presented lower probabilities to display negative NPV considering investment and prices fluctuations
de Oliveira <i>et al.</i> (2014)	Economic viability	iCL and continuous crop system (soybean) in Rio Grande do Sul, Brazil	12 years	Productivity and Gross Margin	The iCL presented better results, especially in years when the rainfall volume in crop development time was insufficient
Costa <i>et al.</i> (2012)	Economic viability, cash flows dynamics and higher investment requirements	iCL; iCLF with eucalyptus trees on simple lines (227 trees/ha) and iCLF with eucalyptus on simple lines (357 trees/ha), in Mato Grosso do Sul, Brazil	12 years, with real data for the first two years	NPV	The lower necessity of investing on the iCL to both iCLF in addition to a return on capital invested in a shorter period, indicate that system iCL tends to be a more suitable alternative to producers who deals with financial constraints and/or risk averse.
Martha Júnior <i>et al.</i> (2011)	Economic viability	iCL; continuous livestock system (beef cattle) and a continuous crop system (soybean) in Goias, Brazil	1 year	Net Revenue, Productivity and Entrepreneur Return Rate	The iCL was more economically attractive than the livestock system, but did not show better results than the soybean crop system. The ERR for the livestock system was negative (−1.55%), for the iCL the return rate was 26.7 and 55.9% for the soybean crop system
De Oliveira <i>et al.</i> (2013)	Economic viability	iCLF system in Goiás, Brazil	7 years, with real data for the first three years	NPV and IRR	Due to favorable crop prices scenario, the economic results were very positive: NPV annual of USD 269.53 ha to 2009 prices. For the IRR the value was 54.24%, well above the attractiveness minimum rate considered, which was 8.75%.

iCL, integrated crop and livestock system; iCLF, integrated crop, livestock and forest system; NPV, net present value; IRR, internal return rate; ERR, entrepreneur return rate.